

Additional file 5. The case of markers in Linkage disequilibrium

Principles

The objective is to derive $E[x_{cl}x_{cm}X_{il}X_{im}]$ in the situation of LD. As in Supplementary Material 1, we use $X_{im} = g_{imf} + g_{imd}$ where g_{imf} and g_{imd} are the “values” of the alleles transmitted to individual i by its father and its dam, with g_{imf} and $g_{imd} = (0 \text{ or } 1) - p_m$. Equivalent terms are defined for x_{cl} , x_{cm} and X_{il} .

$$E[x_{cl}x_{cm}X_{il}X_{im}] = \sum_{s \in \{f,d\}} \sum_{t \in \{f,d\}} \sum_{u \in \{f,d\}} \sum_{v \in \{f,d\}} E[g_{cls}g_{cmt}g_{ilu}g_{imv}]$$

The random variable M_{cls} is the allele individual c received from s at locus l . M_{cmt} , M_{ilu} and M_{imv} are defined similarly. More generally, $M_{\alpha lf}$ and $M_{\alpha ld}$ are the locus l allele individual α received from its father and $M_{\alpha ld}$, from its dam.

Four types of $g = (g_{cls}, g_{cmt}, g_{ilu}, g_{imv})$ vectors are encountered

$$\mathcal{T}_{s=t}^{u=v} : s = t \text{ and } u = v$$

$$\mathcal{T}_{s=t}^{u \neq v} : s = t \text{ and } u \neq v$$

$$\mathcal{T}_{s \neq t}^{u=v} : s \neq t \text{ and } u = v$$

$$\mathcal{T}_{s \neq t}^{u \neq v} : s \neq t \text{ and } u \neq v$$

As each of the indices s, t, u, v may be for d , a total of 16 different g vectors are possible.

In type 1, both alleles (belonging to locus m and l) of each couple of locus (one for c and one for i) are on the same chromosome (may be from the two fathers, the two dams, c 's father and i 's dam or i 's father and c 's dam).

In type 2, both alleles (belonging to locus m and l) of the candidate c are on the same chromosome, while alleles of the reference i are not on the same chromosome.

Type 3 is the reverse from type 2.

In type 4, alleles of locus m and l of both individuals c and i are on different chromosomes.

For each of these situations, we consider the IBD status between alleles at locus m on chromosomes ct and iv , and at locus l on chromosomes cs and iu .

We may have

$$\mathcal{S}_{ml} = \{ M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu} \} \text{ with a probability } \varphi_{ml}^{stuv}$$

$$\mathcal{S}_{m\bar{l}} = \{ M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \not\equiv M_{ilu} \} \text{ with a probability } \varphi_{m\bar{l}}^{stuv}$$

$$\mathcal{S}_{\bar{m}l} = \{ M_{cmt} \not\equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu} \} \text{ with a probability } \varphi_{\bar{m}l}^{stuv}$$

$$\mathcal{S}_{\bar{m}\bar{l}} = \{ M_{cmt} \not\equiv M_{imv} \text{ and } M_{cls} \not\equiv M_{ilu} \} \text{ with a probability } \varphi_{\bar{m}\bar{l}}^{stuv}$$

The computation of the probability φ_k^{stuv} depends on the type of g -vector.

$$E[g_{cls}g_{cmt}g_{ilu}g_{imv}] = \sum_{k \in \{ml, ml, ml, ml\}} \varphi_k^{stuv} E[g_{cls}g_{cmt}g_{ilu}g_{imv} | \mathcal{S}_k]$$

Expectations $E[g_{cls}g_{cmt}g_{ilu}g_{imv} | \mathcal{S}_k]$

Let $p_{AA} = (1 - p_m)(1 - p_l) + \Delta_{lm}$; $p_{AB} = (1 - p_m)p_l - \Delta_{lm}$; $p_{BA} = p_m(1 - p_l) - \Delta_{lm}$ and $p_{BB} = p_m p_l + \Delta_{lm}$. Under \mathcal{S}_{ml} , only 4 genotypes vectors are possible :

M_{cls}	M_{cmt}	M_{ilu}	M_{imv}	prob[$g_{cls}g_{cmt}g_{ilu}g_{imv} \mathcal{S}_{ml}$]				$g_{cls}g_{cmt}g_{ilu}g_{imv}$
				$s = t$ and $u = v$	$s = t$ and $u \neq v$	$s \neq t$ and $u = v$	$s \neq t$ and $u \neq v$	
A_l	A_m	A_l	A_m	p_{AA}	p_{AA}	p_{AA}	$(1 - p_m)(1 - p_l)$	$p_m^2 p_l^2$
A_l	B_m	A_l	B_m	p_{BA}	p_{BA}	p_{BA}	$p_m(1 - p_l)$	$p_l^2(1 - p_m)^2$
B_l	A_m	B_l	A_m	p_{AB}	p_{AB}	p_{AB}	$(1 - p_m)p_l$	$p_m^2(1 - p_l)^2$
B_l	B_m	B_l	B_m	p_{BB}	p_{BB}	p_{BB}	$p_m p_l$	$(1 - p_l)^2(1 - p_m)^2$

The expectation follows:

	$E[g_{cls}g_{cmt}g_{ilu}g_{imv} \mathcal{S}_{ml}]$
$s = t$ and $u = v$	$(1 - p_m)(1 - p_l)p_m p_l + \Delta_{lm}(1 - 2p_l)(1 - 2p_m)$
$s = t$ and $u \neq v$	$(1 - p_m)(1 - p_l)p_m p_l + \Delta_{lm}(1 - 2p_l)(1 - 2p_m)$
$s \neq t$ and $u = v$	$(1 - p_m)(1 - p_l)p_m p_l + \Delta_{lm}(1 - 2p_l)(1 - 2p_m)$
$s \neq t$ and $u \neq v$	$(1 - p_m)(1 - p_l)p_m p_l$

For other IBD status (\mathcal{S}_{ml} , \mathcal{S}_{ml} , \mathcal{S}_{ml}), it must be noted that all expectations of the product $g_{cls}g_{cmt}g_{ilu}g_{imv}$ are null in g -types different from $\mathcal{T}_{s=t}^{u=v}$. For instance, given $\mathcal{R} = \{\mathcal{S}_{mt}; s = t \text{ and } u \neq v\}$, $E[g_{cls}g_{cmt}g_{ilu}g_{imv} | \mathcal{R}] = E[g_{cls}g_{cmt}g_{ilu} \times E[g_{imv} | \mathcal{R}, g_{cls}, g_{cmt}, g_{ilu}] | \mathcal{R}]$.

But $E[g_{imv} | \mathcal{R}, g_{cls}, g_{cmt}, g_{ilu}] = E[g_{imv}] = 0$ (the variable g_{imv} is independent on the other g under \mathcal{R}).

M_{cls}	Genotypes			Genotypes probability for IBD status:			Value of $g_{cls}g_{cmt}g_{ilu}g_{imv}$
	M_{cmt}	M_{ilu}	M_{imv}	\mathcal{S}_{mt}	\mathcal{S}_{ml}	\mathcal{S}_{ml}	
A_l	A_m	A_l	A_m	$p_{AA}^2/[1 - p_m]$	$p_{AA}^2/[1 - p_l]$	p_{AA}^2	$p_m^2 p_l^2$
A_l	A_m	A_l	B_m	0	$p_{AA} p_{BA}/[1 - p_l]$	$p_{AA} p_{BA}$	$-p_l^2 p_m(1 - p_m)$
A_l	A_m	B_l	A_m	$p_{AA} p_{AB}/[1 - p_m]$	0	$p_{AA} p_{AB}$	$-p_m^2 p_l(1 - p_l)$
A_l	A_m	B_l	B_m	0	0	$p_{AA} p_{BB}$	$p_m(1 - p_m)^2 p_l(1 - p_l)$
A_l	B_m	A_l	A_m	0	$p_{BA} p_{AA}/[1 - p_l]$	$p_{BA} p_{AA}$	$-p_l^2 p_m(1 - p_m)$
A_l	B_m	A_l	B_m	p_{BA}^2/p_m	$p_{BA}^2/[1 - p_l]$	p_{BA}^2	$p_l^2(1 - p_m)^2$
A_l	B_m	B_l	A_m	0	0	$p_{BA} p_{AB}$	$p_m(1 - p_m) p_l(1 - p_l)$
A_l	B_m	B_l	B_m	$p_{BA} p_{BB}/p_m$	0	$p_{BA} p_{BB}$	$-(1 - p_m)^2 p_l(1 - p_l)$
B_l	A_m	A_l	A_m	$p_{AB} p_{AA}/[1 - p_m]$	0	$p_{AB} p_{AA}$	$-p_m^2 p_l(1 - p_l)$
B_l	A_m	A_l	B_m	0	0	$p_{AB} p_{BA}$	$p_m(1 - p_m) p_l(1 - p_l)$
B_l	A_m	B_l	A_m	$p_{AB}^2/[1 - p_m]$	p_{AB}^2/p_l	p_{AB}^2	$p_m^2(1 - p_l)^2$
B_l	A_m	B_l	B_m	0	$p_{AB} p_{BB}/p_l$	$p_{AB} p_{BB}$	$-(1 - p_l)^2 p_m(1 - p_m)$
B_l	B_m	A_l	A_m	0	0	$p_{BB} p_{AA}$	$p_m(1 - p_m) p_l(1 - p_l)$
B_l	B_m	A_l	B_m	$p_{BB} p_{BA}/p_m$	0	$p_{BB} p_{BA}$	$-(1 - p_m)^2 p_l(1 - p_l)$
B_l	B_m	B_l	A_m	0	$p_{BB} p_{AB}/p_l$	$p_{BB} p_{AB}$	$-(1 - p_l)^2 p_m(1 - p_m)$
B_l	B_m	B_l	B_m	p_{BB}^2/p_m	p_{BB}^2/p_l	p_{BB}^2	$(1 - p_l)^2(1 - p_m)^2$

	$E[g_{cls}g_{cmt}g_{ilu}g_{imv} s = t \text{ and } u = v]$
\mathcal{S}_{mt}	$\Delta_{lm}^2 \times [p_m^3 + (1 - p_m)^3] / [p_m(1 - p_m)]$
\mathcal{S}_{ml}	$\Delta_{lm}^2 \times [p_l^3 + (1 - p_l)^3] / [p_l(1 - p_l)]$
\mathcal{S}_{mt}	$\Delta_{lm}^2 \times (1 - 2p_m)(1 - 2p_l)$

Assembling all elements we get

$$E[x_{cl}x_{cm}x_{il}x_{im}] = [(1 - p_m)(1 - p_l)p_m p_l + \Delta_{lm}(1 - 2p_l)(1 - 2p_m)] \left[(\sum_{s \in \{f,d\}} \sum_{t \in \{f,d\}} \sum_{u \in \{f,d\}} \sum_{v \in \{f,d\}} \varphi_{ml}^{stuv}) - (\varphi_{ml}^{fada} + \varphi_{ml}^{fdaf} + \varphi_{ml}^{dfda} + \varphi_{ml}^{dffd}) \right] + \Delta_{lm}^2 \sum_{s \in \{f,d\}} \sum_{u \in \{f,d\}} [\varphi_{ml}^{ssuu} \times [p_m^3 + (1 - p_m)^3] / [p_m(1 - p_m)] + \varphi_{ml}^{ssuu} \times [p_l^3 + (1 - p_l)^3] / [p_l(1 - p_l)] + \varphi_{ml}^{ssuu} \times (1 - 2p_m)(1 - 2p_l)]$$

Probabilities of IBD status \mathcal{S}_k

We are interested in genes at loci l and m individuals i and c received from specific parents denoted by s and u or t and v .

The probability $p(M_{cmt} \equiv M_{imv}) = \varphi_m^{tv}$ (resp. $p(M_{cls} \equiv M_{ilu}) = \varphi_l^{su}$) equals the coancestry coefficient between t and v (resp. s and u). It is estimated the usual way.

We only need to develop $\varphi_{ml}^{stuv} = p(\mathcal{S}_{ml}) = p(M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu})$. Indeed, the other terms are given by:

$$\varphi_{mt}^{stuv} = p(M_{cmt} \equiv M_{imv}) - p(M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu}) = \varphi_m^{tv} - \varphi_{ml}^{stuv}$$

$$\varphi_{ml}^{stuv} = p(M_{cls} \equiv M_{ilu}) - p(M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu}) = \varphi_l^{su} - \varphi_{ml}^{stuv}$$

$$\varphi_{ml}^{stuv} = 1 - (\varphi_{ml}^{stuv} + \varphi_{mt}^{stuv} + \varphi_{ml}^{stuv}) = 1 - (\varphi_m^{tv} + \varphi_l^{su}) + \varphi_{ml}^{stuv}$$

We first examine the situation of a single locus (say l). Given the pedigree, different genealogical chains linking c and i are often possible. They will be indexed $pl = 1 \dots Nl_p$ (and $pm = 1 \dots Nm_p$ when considering the m locus). A chain is characterized by a shared ancestor a_{pl} and consists of two subchains (pl_c and pl_i) linking a_{pl} to c and to i . In graph theory, these subchains are made of edges with nodes (the individuals) linked by arcs (characterizing transmission events). Subchain pl_c (resp. pl_i) comprises one and only one parent of c (resp. i). The ancestor may be c or i themselves, in which case the chain is extended to one parent on this ancestor. The lengths (numbers of generations) of the pl_c and pl_i subchains will be noted $n_{a_{pl}c}$ and $n_{a_{pl}i}$. We will note $\alpha = \{\alpha_1, \alpha_2 \dots \alpha_{n_{a_{pl}c}-1}\}$ and $\beta = \{\beta_1, \beta_2 \dots \beta_{n_{a_{pl}i}-1}\}$ the lists of intermediate individuals belonging to $pl_c = \{c, \alpha, a_{pl}\}$ and $pl_i = \{i, \beta, a_{pl}\}$. We are interested in the ancestor allele $M_{a_{pl}lw}$ shared by individuals c and to i . Genders (f or d) of individual ω will be noted $\mathcal{G}(\omega)$.

The IBD status $M_{cls} \equiv M_{ilu}$ occurs if the allele at locus l carried by chromosomes transmitted by s to c and transmitted by u to i both come from the same ancestor allele carried by a_{pl} : $M_{cls} \equiv M_{ilu}$ if

$$\begin{cases} s = \mathcal{G}(\alpha_1) \text{ and } u = \mathcal{G}(\beta_1) \\ M_{cls} \equiv M_{\alpha_1 \mathcal{G}(\alpha_2)} \equiv M_{\alpha_2 \mathcal{G}(\alpha_3)} \dots \equiv M_{a_{pl}w} \text{ and } M_{ilu} \equiv M_{\beta_1 \mathcal{G}(\beta_2)} \equiv M_{\beta_2 \mathcal{G}(\beta_3)} \dots \equiv M_{a_{pl}w} \text{ with } wl = f \text{ or } d \end{cases}$$

In other words the $M_{a_{pl}wl}$ allele was transmitted :

- along the pl_c subchain to c via s (event $J_{pl_c, wl}^l = \{M_{cls} \equiv M_{\alpha_1 lG(\alpha_2)} \equiv M_{\alpha_2 lG(\alpha_3)} \cdots \equiv M_{a_{pl}wl}\}$) and
- along the pl_i subchain to i via u (event $J_{pl_i, wl}^l = \{M_{ilu} \equiv M_{\beta_1 lG(\beta_2)} \equiv M_{\beta_2 lG(\beta_3)} \cdots \equiv M_{a_{pl}wl}\}$)

$$prob(J_{pl_c, wl}^l) = prob(M_{cls} \equiv M_{\alpha_1 lG(\alpha_2)}) prob(M_{\alpha_1 lG(\alpha_2)} \equiv M_{\alpha_2 lG(\alpha_3)} | M_{cls} \equiv M_{\alpha_1 lG(\alpha_2)}) \cdots$$

Transmissions events between generations being independent, for a single locus (l in the current example), $prob(J_{pl_c, wl}^l) = \frac{1}{2}^{n_{a_{pl}c}}$ and $prob(J_{pl_i, wl}^l) = \frac{1}{2}^{n_{a_{pl}i}}$.

It must be noted that pl_c and pl_i may share edges. In these situations $prob(J_{pl_c, wl}^l \cap J_{pl_i, wl}^l) = prob(J_{pl_c, wl}^l) prob(J_{pl_i, wl}^l | J_{pl_c, wl}^l) \neq prob(J_{pl_c, wl}^l) prob(J_{pl_i, wl}^l)$: If n_s edges belong to both pl_c and pl_i , $prob(J_{pl_i, wl}^l | J_{pl_c, wl}^l) = \frac{1}{2}^{n_{a_{pl}i} - n_s}$. Thus $p(M_{cls} \equiv M_{ilu} | pl, wl) = \frac{1}{2}^{n_{a_{pl}c} + n_{a_{pl}i} - n_s}$

As there is two possible ancestral alleles ($M_{a_{pl}wl}$ with $wl = f$ or d) shared by chromosomes transmitted by s to c and transmitted by u to i , we finally get the classical $p(M_{cls} \equiv M_{ilu} | pl) = \frac{1}{2}^{n_{a_{pl}c} + n_{a_{pl}i} - 1 - n_s}$. IBD status $M_{cls} \equiv M_{ilu}$ due to the transmission by the different possible chains being exclusive, the unconditional probability is $p(M_{cls} \equiv M_{ilu}) = \sum_{pl} \frac{1}{2}^{n_{a_{pl}c} + n_{a_{pl}i} - 1 - n_s}$

When considering both loci l and m simultaneously, all possible pairs (pl, pm) of genealogical chains and parental origins of ancestor alleles linking c and i must be examined.

$$\varphi_{ml}^{stuv} = p(M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu}) = \sum_{pl, wl} \sum_{pm, wm} p(M_{cmt} \equiv M_{imv} \text{ and } M_{cls} \equiv M_{ilu} | pl, wl, pm, wm)$$

$$\varphi_{ml}^{stuv} = \sum_{pl, wl} \sum_{pm, wm} p(M_{cls} \equiv M_{ilu} | pl, wl) p(M_{cmt} \equiv M_{imv} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm)$$

Previous definitions are extended to pm chain: to pm corresponds ancestor a_{pm} , vectors γ and δ are counterparts of α and β and events $J_{pm_c, wm}^m$ and $J_{pm_i, wm}^m$ of events $J_{pl_c, wl}^l$ and $J_{pl_i, wl}^l$. We have

$$M_{cmt} \equiv M_{imv} \text{ if}$$

$$\begin{cases} t = \mathcal{G}(\gamma_1) \text{ and } v = \mathcal{G}(\delta_1) \\ M_{cmt} \equiv M_{\gamma_1 m \mathcal{G}(\gamma_2)} \equiv M_{\gamma_2 m \mathcal{G}(\gamma_3)} \cdots \equiv M_{a_{pm}mw} \text{ and } M_{imv} \equiv M_{\delta_1 m \mathcal{G}(\delta_2)} \equiv M_{\delta_2 m \mathcal{G}(\delta_3)} \cdots \equiv M_{a_{pm}mw} \text{ with } w = f \text{ or } d \end{cases}$$

$$p(M_{cmt} \equiv M_{imv} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm) = p(J_{pm_c, wm}^m \cap J_{pm_i, wm}^m | J_{pl_c, wl}^l \cap J_{pl_i, wl}^l)$$

The pm chain may be partially confounded with pl chain: an edge (γ_k, γ_{k+1}) from pm may or not be present in pl . If not, $prob(M_{\gamma_k m \mathcal{G}(\gamma_{k+1})} \equiv M_{\gamma_{k+1} m \mathcal{G}(\gamma_{k+2})} | J_{pl_c, wl}^l \cap J_{pl_i, wl}^l) = 1/2$. If $(\gamma_k, \gamma_{k+1}) = (\alpha_h, \alpha_{h+1})$ is both in pm and pl , the probability simplifies to

$prob(M_{\gamma_k m \mathcal{G}(\gamma_{k+1})} \equiv M_{\gamma_{k+1} m \mathcal{G}(\gamma_{k+2})} | M_{\alpha_h l \mathcal{G}(\alpha_{h+1})} \equiv M_{\alpha_{h+1} l \mathcal{G}(\alpha_{h+2})})$. In this case, either $\gamma_{k+2} = \alpha_{h+2}$ (alleles at locus l and m transmitted by $\gamma_{k+1} = \alpha_{h+1}$ to $\gamma_k = \alpha_h$ were on the same grand parental chromosome), giving

$prob(M_{\gamma_k m \mathcal{G}(\gamma_{k+1})} \equiv M_{\gamma_{k+1} m \mathcal{G}(\gamma_{k+2})} | M_{\alpha_h l \mathcal{G}(\alpha_{h+1})} \equiv M_{\alpha_{h+1} l \mathcal{G}(\alpha_{h+2})}) = 1 - r_{ml}$, or $\gamma_{k+2} \neq \alpha_{h+2}$, giving $prob(M_{\gamma_k m \mathcal{G}(\gamma_{k+1})} \equiv M_{\gamma_{k+1} m \mathcal{G}(\gamma_{k+2})} | M_{\alpha_h l \mathcal{G}(\alpha_{h+1})} \equiv M_{\alpha_{h+1} l \mathcal{G}(\alpha_{h+2})}) = r_{ml}$.

Computing φ_{ml}^{stuv} , all types of $g = (g_{cls}, g_{cmt}, g_{ilu}, g_{imv})$ vectors must be considered ($\mathcal{J}_{s=t}^{u=v}, \mathcal{J}_{s=t}^{u \neq v}, \mathcal{J}_{s \neq t}^{u=v}, \mathcal{J}_{s \neq t}^{u \neq v}$) for all combination of pl and pm chains.

Examples

A. i and c are unrelated

In this case, probability $\varphi_{\cancel{ml}}^{ssuu} = 1$ is the only non null, and the expectation turns to be:

$$E[x_{cl}x_{cm}x_{il}x_{im}] = 4\Delta_{lm}^2(1 - 2p_m)(1 - 2p_l)$$

B. i is one of c 's parents

Two chains are possible: $p_1 = a_{p_1} \rightarrow i \rightarrow c$ and $p_2 = a_{p_2} \rightarrow i \rightarrow c$ with a_{p_1} and a_{p_2} , parents of i as ancestors. Here $n_{a_{p_1}c} = 2, n_{a_{p_1}i} = 1, n_s = 1$.

a. l and m genes on the same chromosome in c and i ($\mathcal{J}_{s=t}^{u=v}$)

Non null probabilities are

$$p(M_{cls} \equiv M_{ilu} | pl, wl) = \frac{1}{4} \text{ if } \mathcal{G}(i) = s \text{ and } \mathcal{G}(a_{pl}) = u; \forall wl = f \text{ or } d \text{ and } \forall pl = p_1 \text{ or } p_2$$

$$p(M_{cms} \equiv M_{imu} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm) = (1 - r_{ml})^2 \text{ if } \mathcal{G}(i) = s, \mathcal{G}(a_{pl}) = u; pl = pm, wl = wm$$

$$p(M_{cms} \equiv M_{imu} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm) = r_{ml}(1 - r_{ml}) \text{ if } \mathcal{G}(i) = s, \mathcal{G}(a_{pl}) = u; pl = pm, wl \neq wm$$

$$\text{Finally, } \varphi_{ml}^{ssuu} = \sum_{pl} \frac{1}{4} ((1 - r_{ml})^2 + r_{ml}(1 - r_{ml})) = \sum_{pl} \frac{1}{4} (1 - r_{ml}) = \frac{1 - r_{ml}}{2}$$

$$\varphi_{\cancel{ml}}^{ssuu} = \frac{r_{ml}}{2}; \varphi_{\cancel{ml}}^{ssuu} = \frac{r_{ml}}{2}; \varphi_{\cancel{ml}}^{ssuu} = \frac{1 - r_{ml}}{2}$$

b. l and m genes on the same chromosome in c , on two chromosomes in i ($\mathcal{J}_{s=t}^{u \neq v}$)

$$p(M_{cls} \equiv M_{ilu} | pl, wl) = \frac{1}{4} \text{ if } \mathcal{G}(i) = s \text{ and } \mathcal{G}(a_{pl}) = u; \forall wl = f \text{ or } d \text{ and } \forall pl = p_1 \text{ or } p_2$$

$$p(M_{cms} \equiv M_{imv} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm) = p(M_{cms} \equiv M_{im\mathcal{G}(a_{pm})} | M_{cls} \equiv M_{i\mathcal{G}(a_{pl})}) p(M_{im\mathcal{G}(a_{pm})} \equiv M_{a_{pm}mwm}) = \frac{1}{2} r_{ml} \text{ if } \mathcal{G}(i) = s, \mathcal{G}(a_{pl}) = u, \mathcal{G}(a_{pm}) = v; pl \neq pm, wl \neq wm. \text{ Indeed, as } u \neq v, (1) \text{ either } pl = p_1 \text{ and } pm = p_2, \text{ or } pl = p_2 \text{ and } pm = p_1 \text{ and } (2) \text{ either } wl = f \text{ and } wm = d, \text{ or } wl = d \text{ and } wm = f$$

$$\text{Finally, } \varphi_{ml}^{ssuv} = \sum_{pl, wl} \frac{1}{8} r_{ml} = \frac{r_{ml}}{2}$$

$$\varphi_{\cancel{ml}}^{ssuv} = \frac{1 - r_{ml}}{2}; \varphi_{\cancel{ml}}^{ssuv} = \frac{1 - r_{ml}}{2}; \varphi_{\cancel{ml}}^{ssuv} = \frac{r_{ml}}{2}$$

c. l and m genes are on two chromosomes in c ($\mathcal{J}_{s \neq t}^{u=v}$ or $\mathcal{J}_{s \neq t}^{u \neq v}$)

In these cases $p(M_{cmt} \equiv M_{imu} | M_{cls} \equiv M_{ilu}, pl, wl, pm, wm) = 0$ since parent i cannot be both c sire and dam.

$$\varphi_{ml}^{stuu} = 0; \varphi_{m\bar{l}}^{stuu} = \frac{1}{2}; \varphi_{\bar{m}l}^{stuu} = \frac{1}{2}; \varphi_{\bar{m}\bar{l}}^{stuu} = 0$$

$$\varphi_{ml}^{stuv} = 0; \varphi_{m\bar{l}}^{stuv} = \frac{1}{2}; \varphi_{\bar{m}l}^{stuv} = \frac{1}{2}; \varphi_{\bar{m}\bar{l}}^{stuv} = 0$$

Using these probabilities, the expectation turns to be:

$$E[x_{cl}x_{cm}X_{il}X_{im}] = (1-p_m)(1-p_l)p_m p_l + \Delta_{lm}(1-2p_l)(1-2p_m) + 2\Delta_{lm}^2 \left[r_{ml} \left(\frac{p_m^3 + (1-p_m)^3}{p_m(1-p_m)} + \frac{p_l^3 + (1-p_l)^3}{p_l(1-p_l)} \right) + (1-r_{ml})(1-2p_m)(1-2p_l) \right]$$

When there is no linkage disequilibrium, $E[x_{cl}x_{cm}X_{il}X_{im}] = (1-p_m)(1-p_l)p_m p_l = 4\sigma_m^2 \sigma_l^2 a_{ci}^2$ as expected.

C. i and c had the same sire

Only one genealogical chain links i and c with subchains $p_c = a_p \rightarrow c$ and $p_i = a_p \rightarrow i$ with a_p their sire. Here $n_{a_p c} = n_{a_p i} = 1, n_s = 0$. Non null probabilities are obtained only when l and m genes on the same chromosome in c and $(\mathcal{J}_{s=t}^{u=v})$, and when $s = u = f = \mathcal{G}(a_p)$. In this case we have:

$$p(M_{cls} \equiv M_{ilu} | p, wl) = p(M_{cls} \equiv M_{a_p lwl} \& M_{ilu} \equiv M_{a_p lwl} | p, wl) = \frac{1}{4} \quad \forall wl = f \text{ or } d$$

$$p(M_{cms} \equiv M_{imu} | M_{cls} \equiv M_{ilu}, p, wl, wm) = p(M_{cms} \equiv M_{a_p mwm} | M_{cls} \equiv M_{a_p lwl}, p, wl, wm) p(M_{imu} \equiv M_{a_p mwm} | M_{ilu} \equiv M_{a_p lwl}, p, wl, wm) = (1-r_{ml})^2 \quad \text{if } wl = wm, r_{ml}^2 \quad \text{if } wl \neq wm$$

$$\text{Finally, } \varphi_{ml}^{ssuu} = \frac{1}{2} \left((1-r_{ml})^2 + r_{ml}^2 \right) = \varphi_{\bar{m}\bar{l}}^{ssuu} \text{ and } \varphi_{m\bar{l}}^{ssuu} = \varphi_{\bar{m}l}^{ssuu} = r_{ml}(1-r_{ml})$$

$$E[x_{cl}x_{cm}X_{il}X_{im}] = \frac{(1-r_{ml})^2 + r_{ml}^2}{2} [(1-p_m)(1-p_l)p_m p_l + (\Delta_{lm} + \Delta_{lm}^2)(1-2p_l)(1-2p_m)] + r_{ml}(1-r_{ml})\Delta_{lm}^2 \left(\frac{p_m^3 + (1-p_m)^3}{p_m(1-p_m)} + \frac{p_l^3 + (1-p_l)^3}{p_l(1-p_l)} \right)$$