



HAL
open science

Impacts of climate-related dieback on regeneration in a drought-prone European beech population

Sylvie Muratorio, Matthieu Lingrand, Elodie Magnanou, Joseph Guarigue,
Jean-André Magdalou

► **To cite this version:**

Sylvie Muratorio, Matthieu Lingrand, Elodie Magnanou, Joseph Guarigue, Jean-André Magdalou. Impacts of climate-related dieback on regeneration in a drought-prone European beech population. BES-SFE Joint 2014 Annual Meeting, Dec 2014, Lille, France. hal-01344424

HAL Id: hal-01344424

<https://hal.science/hal-01344424>

Submitted on 5 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impacts of climate-related dieback on regeneration in a drought-prone European beech population


The issue: does climate-driven individual tree dieback reduce effective seed output and population size (N_e)?

Interactions between biotic and abiotic environment and tree functional traits determine carbohydrate and water fluxes at tree-level, which in turn affect demographic rates and fitness at tree- and population-level. The total number of reproductive trees (N_{obs}) and the among-trees variations of seed production determine the effective population size (N_e).



Climate-induced tree dieback may result in a reduced quantity and/or quality of seed output at population level, even without or before tree mortality. We investigate the relationship between individual tree dieback and fecundity in the forest tree *Fagus sylvatica* (the European beech) in Southern France (La Massane, Pyrénées Orientales). *Fagus sylvatica* is known to be sensitive to summer drought, but shows so far limited mortality; for instance at the study site the mortality rate is 1.75%/year (between 2003 and 2005) while the rate of tree dieback (measured through foliar mortality) is 10.5%.


Results: estimated dispersal and fecundity parameters




1. Pollen and seed dispersal abilities

- Pollen and seed dispersal kernel were modeled with exponential power distribution with parameters (δ_p, b_p) and (δ_s, b_s).
- Pollen is dispersed on average at $\delta_p = 62$ m, with non negligible events of long distance dispersal ($b_p = 0.46$)
- Seeds are dispersed on average at $\delta_s = 12$ m, with fewer long distance dispersal events than for pollen ($b_p = 0.83$)
- 13% of the seedlings have their mother-tree out of the plot, and 67% of the offspring with local mother have their father-tree out of the plot.


2. Distribution of female and male fecundities on each plot




Methods



2- Estimation of realized fecundities and dispersal with the SEMM



Results: relationship between dieback and fecundity



Conclusions

- Individual tree dieback (through foliar mortality) is associated to lower relative female fecundity, consistently with costly nut-seeds in *Fagus sylvatica*.
- Non negligible female fecundity can however be achieved for intermediate leaf mortality, suggesting adaptive plastic response aiming at reducing functioning costs.
- Perspective: investigating whether there is a change in allocation to reproduction versus growth for trees showing evidence of dieback.