



**HAL**  
open science

# Monitoring spatial accuracy of oil palm cultivation mapping in southern Cameroon from Landsat series images

Prune Christobelle Komba Mayossa, Sébastien Gadal

► **To cite this version:**

Prune Christobelle Komba Mayossa, Sébastien Gadal. Monitoring spatial accuracy of oil palm cultivation mapping in southern Cameroon from Landsat series images. 12th International Symposium "Spatial Accuracy Assessment in Natural Resources and Environmental Sciences, Jul 2016, Montpellier, France. hal-01348467

**HAL Id: hal-01348467**

**<https://hal.science/hal-01348467>**

Submitted on 24 Jul 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Monitoring Spatial accuracy of oil palm cultivation mapping in southern Cameroon from Landsat series images

Prune Christobelle KOMBA MAYOSSA and Sébastien GADAL



Aix-Marseille Université, CNRS ESPACE UMR 7300  
 Europôle Méditerranéen de l'Arbois, avenue Louis Philibert Bâtiment Laennec, hall C  
 BP 80, 13545 Aix en Provence Cedex 4 Tel: +33(0)1 39 2557 00  
 Email: [prune.komba-mayossa@etu.univ-amu.fr](mailto:prune.komba-mayossa@etu.univ-amu.fr) / [sebastien.gadal@univ-amu.fr](mailto:sebastien.gadal@univ-amu.fr)



## 1 INTRODUCTION

### Oil palm in Cameroon

- \*Great economic importance, with an industrialisation dating from colonial period.
- \*Incomes generated by its cultivation have generated agroindustries such as the Cameroonian Society of Palm grove (SOCAPALM).
- \*This activity is behind of socio-environmental damages (deforestation, loss of biodiversity and pollution).


### Context

- \*Long-term management of oil palm resources in Congo basin.
- \*Studying and mapping palm grove evolution to understand the impact related to its cultivation.

### Objectives

- \*Map SOCAPALM industrial palm grove using Landsat series images.
- \* Measure produced maps accuracy.

## 2 STUDY AREA



### Data

- \* Landsat ETM+ (2001)
- \* Landsat OLI-TIRS (2015)

Table 1. Data specification

Sensor	Acquisition date	Mode	Resolution	Cloud cover
ETM+	26/04/2001	MS/PAN	30m/15m	30%
OLI-TIRS	25/04/2015	MS/PAN	30m/15m	30%


## 3 METHOD

### Palm grove mapping

- \*Per-pixel classification (maximum likelihood algorithm)

### Accuracy assessment

- \*Confusion matrix method



## 4 RESULTS




Figure 3. Land cover mapping (2001), from Landsat 7 ETM+




Figure 4. Land cover mapping (2015), from Landsat 8 OLI-TIRS

- \* Overall accuracy of 92% (Kappa=0,92) in 2001 and 80% (kappa=0,89) in 2015.
- \* An improvement of palm grove surfaces of 18% (from 2001 to 2015)

## 5 CONCLUSION

- \* High correlated maps were obtained ( Kappa =0,92 in 2001 Vs 0,89 in 2015). However, because of the presence of mixed pixels, some confusions were observed between vegetation and oil palm classes. These confusions resulting from the spatial and spectral characteristics of palm groves, the method used to map and validate the map, and the uncertainty related to data, affect maps accuracy.
- \* To increase the accuracy we suggest, (1) use another mapping method such as super-resolution; (2) develop a classification system of cartographic products.