



## Modulations of olfactory responses by GAP junctions uncoupling in the OM.

Mikael Le Bourhis, Bertrand Bryche, Denise Grebert, Edith Pajot, Patrice Congar, Nicolas Meunier

### ► To cite this version:

Mikael Le Bourhis, Bertrand Bryche, Denise Grebert, Edith Pajot, Patrice Congar, et al.. Modulations of olfactory responses by GAP junctions uncoupling in the OM.. XXII. Congress of the European Chemoreception Research Organization (ECRO 2013), Aug 2013, Louvain, Belgium. Chemical Senses, 39, 2013, Abstracts from the XXIIInd Congress of the European Chemoreception Research Organization, ECRO 2013. 10.1093/chemse/bjt055 . hal-01349543

HAL Id: hal-01349543

<https://hal.science/hal-01349543>

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Modulations of olfactory responses by GAP junctions uncoupling in the OM.


Mikaël LE BOURHIS, Bertrand BRYCHE, Denise GREBERT, Edith PAJOT, Patrice CONGAR and Nicolas MEUNIER.

## Introduction

## Results

### I - ET activates SCs ex vivo


Method : Olfactory Mucosa (OM) slices 200 $\mu$ m (Wistar P0-P4), loaded with Oregon Green (488nm), Calcium Imaging.



I - ET evokes both tonic and phasic calcium responses in the SCs.


### II - Plasticity of GAP junctions ex vivo and in vitro.

Method : Olfactory Mucosa explant (Wistar P7), loaded with Oregon Green, Electric stimulation and Calcium Imaging. n > 20 for each group.



CBX and ET decrease spreading of the calcium wave.

Method : Olfactory Mucosa Primary Culture, Whole cell Voltage Clamp. n > 15 for each group.



CBX and ET increase membrane resistance of SCs.


\*\* : P<0,01

\*\*\* : P<0,001


II - ET and CBX uncouple GAP junctions of SCs.

### III - ET and CBX modify olfactory responses.

Method : Hemi-Head (P30), Electro-Olfactogram (EOG), Isoamyl Acetate 1:10 in mineral oil, n = 6 in each group.



Repolarization kinetics of the olfactory response, but not amplitude, are affected by GAP junctions uncoupling.



III - GAP junctions of SCs are involved in olfactory responses.

## Conclusion

SCs respond in vitro (Gouadon et al., 2010) and ex vivo to ET. This peptide acts as an autocrine and paracrine factor. ET induces uncoupling of SC's GAP junctions both in vitro and ex vivo, like the classically used GAP junction uncoupling agent CBX. Such modulations affect olfactory responses. Both ET and CBX treatments increase EOG signal repolarization kinetics. Our results are consistent with an implication of glial-like cells in the peripheral modulation of the olfactory information. This has to be confirmed using patch clamp on the knobs of olfactory sensory neurons.

- Blomstrand F et al., Endothelins regulate astrocyte gap junctions in rat hippocampal slices. *Eur J Neurosci.* (2004)
- Gouadon et al., Endothelin evokes distinct calcium transients in neuronal and non-neuronal cells of rat olfactory mucosa primary cultures. *Neuroscience.* (2010)
- Fivos Vogalis et al., Electrical Coupling in Sustentacular Cells of the Mouse Olfactory Epithelium. *J. Neurophysiol.* (2005)

Mikael Le Bourhis, PhD student  
mlebourhis@jouy.inra.fr  
06.50.98.50.42 / 01.34.65.25.55