



HAL
open science

K-Stability of Fano spherical varieties

Thibaut Delcroix

► **To cite this version:**

| Thibaut Delcroix. K-Stability of Fano spherical varieties. 2016. hal-01352060v1

HAL Id: hal-01352060

<https://hal.science/hal-01352060v1>

Preprint submitted on 5 Aug 2016 (v1), last revised 26 Nov 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

K-STABILITY OF FANO SPHERICAL VARIETIES

THIBAUT DELCROIX

ABSTRACT. We prove a criterion for K-stability of a \mathbb{Q} -Fano spherical variety with respect to equivariant special test configurations, in terms of its moment polytope and some combinatorial data associated to the open orbit. Combined with the equivariant version of the Yau-Tian-Donaldson conjecture for Fano manifolds proved by Datar and Székelyhidi, it yields a criterion for the existence of a Kähler-Einstein metric on a spherical Fano manifold. The results hold also for modified K-stability and existence of Kähler-Ricci solitons.

INTRODUCTION

Kähler-Einstein metrics on a Kähler manifold are the solutions (if they exist) of a highly non linear second order partial derivative equation on the manifold. It is not clear at the moment under which conditions the equation admits solutions on a Fano manifold. In the recent years a major advance in this direction has been made through the resolution of the Yau-Tian-Donaldson conjecture in the Fano case, by Chen, Donaldson, and Sun [CDS15a, CDS15b, CDS15c]. This conjecture states in its general form that the existence of some canonical metrics on a Kähler manifold should be related to an algebro-geometric condition of K-stability on the manifold.

The K-stability condition is a condition involving the positivity of numerical invariants associated to polarized one parameter degenerations of the manifold, equipped with an action of \mathbb{C}^* , called *test configurations*. In the Fano case, it was proved by Li and Xu [LX14] (and also Chen, Donaldson and Sun) that it is enough to consider test configurations with normal central fiber, which are called *special* test configurations. Other proofs of the Yau-Tian-Donaldson conjecture were obtained by Tian [Tia15], Datar and Székelyhidi [DS], Chen, Sun and Wang [CSW], Berman, Boucksom and Jonsson [BBJ]. The work of Datar and Székelyhidi is of special interest to us as it allows us to take into account automorphisms of the manifold, by considering only equivariant test configurations, and extends the result to Kähler-Ricci solitons.

The necessity of the K-stability condition with respect to special test configurations was established earlier by Tian who provided an example of Fano manifold with vanishing Futaki invariant but no Kähler-Einstein metrics [Tia97]. It was not clear at first if the K-stability condition could be used to prove the existence of Kähler-Einstein metrics on explicit examples of Fano manifolds. One aim of the present article is to provide an illustration of the power of the approach to the existence of Kähler-Einstein metrics *via* K-stability, on highly symmetric manifolds.

Namely, we obtain a criterion for the existence of Kähler-Einstein metrics on a Fano spherical manifold, involving only the moment polytope and the valuation cone of the spherical manifold. Both are classical and central objects in the theory of spherical varieties. The class of spherical varieties is a very large class of highly

symmetric varieties, which contains toric varieties, generalized flag manifolds, homogeneous toric bundles, biequivariant compactifications of reductive groups. For all of these subclasses for which a criterion was known, our result specializes to the same criterion (compare with [WZ04, PS10, Del15]). Examples of new varieties to which our criterion applies include colored horospherical varieties and symmetric varieties (for examples the varieties constructed in [DP83]). Let us mention here the work of Ilten and Süß on Fano manifolds with an action of a torus with complexity one [IS], and also the discussion in [DS, Section 4], which were to the author's knowledge the first applications of K-stability as a sufficient criterion for explicit examples.

Another aim of this article is to provide a framework for understanding the K-stability condition in this class of spherical varieties, which should lead to a better understanding of K-stability in general. The author obtained in [Del15, Del] an example of group compactification with no Kähler-Einstein metric but vanishing Futaki invariant, which is furthermore not K-semistable unlike the Mukai-Umemura type example from [Tia97]. This is evidence that non trivial K-stability phenomena appear in the class of spherical Fano manifolds, which was not true for toric Fano manifolds.

Before stating the main result of the article, let us introduce some notations, the moment polytope and the valuation cone.

Let G be a complex connected reductive algebraic group. Let B be a Borel subgroup of G and T a maximal torus of B . Let B^- be the Borel subgroup of G opposite with respect to B . Let $\mathfrak{X}(T)$ denote the group of algebraic characters of T . Denote by $\Phi \subset \mathfrak{X}(T)$ the root system of (G, T) and Φ^+ the positive roots defined by B .

Let X be a Fano manifold, spherical under the action of G , which means that B acts on X with an open and dense orbit. The *moment polytope* $\Delta^+ \subset \mathfrak{X}(T) \otimes \mathbb{R}$ of X with respect to B is a polytope encoding the structure of representation of G on the spaces of sections of tensor powers of the anticanonical line bundle. Alternatively, from a symplectic point of view, it can be characterized as the Kirwan moment polytope of (X, ω) with respect to the action of a maximal compact subgroup K of G , where ω is a K -invariant Kähler form in $2\pi c_1(X)$ (see [Bri87]). The moment polytope Δ^+ determines a sub-root system Φ_L of Φ , composed of those roots that are orthogonal to the affine span of Δ^+ with respect to the *Killing form* κ . Let Φ_P^+ be the set $\Phi^+ \setminus \Phi_L$, and $2\rho_P$ be the sum of the elements of Φ_P^+ .

A spherical variety X also has an open and dense orbit O under the action of G . The valuation cone of X depends only on this open orbit O . Let $\mathcal{M}_- \subset \mathfrak{X}(T)$ be the set of characters of B^- -semi-invariant functions in the function field $\mathbb{C}(O)$ of O , and let \mathcal{N}_- be its \mathbb{Z} -dual. The restriction of a \mathbb{Q} -valued valuation on $\mathbb{C}(O)$ to the B^- -semi-invariant functions defines an element of $\mathcal{N}_- \otimes \mathbb{Q}$. The *valuation cone* \mathcal{V}_- with respect to B^- is defined as the set of those elements of $\mathcal{N}_- \otimes \mathbb{Q}$ induced by G -invariant valuations on $\mathbb{C}(O)$.

Remark that the vector space $\mathcal{N}_- \otimes \mathbb{Q}$ is a quotient of the vector space $\mathfrak{Y}(T) \otimes \mathbb{Q}$, where $\mathfrak{Y}(T)$ is the group of algebraic one parameter subgroups of T . Denote by $\pi : \mathfrak{Y}(T) \otimes \mathbb{Q} \rightarrow \mathcal{N}_- \otimes \mathbb{Q}$ the quotient map, so that $\pi^{-1}(\mathcal{V}_-) \subset \mathfrak{Y}(T) \otimes \mathbb{Q}$. Let $\Xi \subset \mathfrak{X}(T) \otimes \mathbb{R}$ be the dual cone to the closure of $\pi^{-1}(\mathcal{V}_-)$ in $\mathfrak{Y}(T) \otimes \mathbb{R}$, with respect to the natural pairing $\langle \cdot, \cdot \rangle : \mathfrak{X}(T) \times \mathfrak{Y}(T) \rightarrow \mathbb{Z}$.

The group of G -equivariant automorphisms $\text{Aut}_G(X)$ of the spherical manifold X is diagonalizable. The real vector space generated by the linear part of \mathcal{V}_- is in fact isomorphic to $\mathfrak{Y}(\text{Aut}_G(X)) \otimes \mathbb{R}$.

Given ζ in $\mathfrak{Y}(\text{Aut}_G(X)) \otimes \mathbb{R}$ identified with an element of $\mathcal{N}_- \otimes \mathbb{R}$, and a choice $\tilde{\zeta} \in \pi^{-1}(\zeta)$ of lift of ζ , we denote by $\text{bar}_{DH, \tilde{\zeta}}(\Delta^+)$ the barycenter of the polytope Δ^+ with respect to the measure with density $p \mapsto e^{\langle 4\rho - 2p, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \kappa(\alpha, p)$ with respect to the Lebesgue measure dp on $\mathfrak{X}(T) \otimes \mathbb{R}$. Our main result is the following.

Theorem A. *Let X be a Fano spherical manifold. The following are equivalent.*

- (1) *There exists a Kähler-Ricci soliton on X with associated complex holomorphic vector field ζ .*
- (2) *The barycenter $\text{bar}_{DH, \tilde{\zeta}}(\Delta^+)$ is in the relative interior of the cone $2\rho_P + \Xi$.*
- (3) *The manifold X is modified K-stable with respect to equivariant special test configurations.*
- (4) *The manifold X is modified K-stable.*

The equivalence between (1) and (4) holds for any Fano manifold, it is the consequence of the work of Chen, Donaldson and Sun recalled earlier for Kähler-Einstein metrics, and of the work of Datar and Székelyhidi for Kähler-Ricci solitons. When X admits an action of a reductive group G , the equivalence between (1) and (3) was shown by Datar and Székelyhidi. What we prove in this article is the equivalence between (2) and (3) in the case of a spherical Fano manifold. Furthermore we prove that the equivalence between (2) and (3) holds for singular \mathbb{Q} -Fano spherical varieties.

The intuition for our main result came from our previous work on group compactifications, which did not involve K-stability. The proof of a Kähler-Einstein criterion for smooth and Fano group compactifications in [Del15, Del] can be adapted to provide another proof of the criterion for Kähler-Ricci solitons on the same manifolds. Similarly, Wang-Zhu type methods (as used in [WZ04] and [PS10]), together with some results proved for horospherical manifolds in the present paper, could be used to obtain the Kähler-Ricci soliton criterion for these manifolds. One advantage of this other approach is that the value of the greatest Ricci lower bound can be explicitly computed. Alternatively, for horospherical varieties at least, this quantity could be computed using *twisted* modified K-stability (see [DS]).

The computation of the K-stability of a manifold requires two ingredients. The first one is a description of all test configurations (rather, using [DS], all special equivariant test configurations), and the second one is a way to compute the Donaldson-Futaki invariant for all of these test configurations.

The description of special equivariant test configurations is obtained using the general theory of spherical varieties. Generalizing the fan of a toric variety, the *colored fan* of a spherical variety consists essentially of a fan subdividing the valuation cone together with additional data called *colors*. The total space of a (normal) test configuration is still a spherical variety, and can thus be described by its colored fan. In the case of a special test configuration, the central fiber itself is a spherical variety and the action of \mathbb{C}^* on the central fiber may be deduced from the colored fan of the test configuration. More precisely, we prove the following.

Theorem B. *Let X be a spherical variety under the action of a reductive algebraic group G . Let \mathcal{X} be a special equivariant test configuration for X , with central fiber X_0 . Then*

- X_0 is spherical under the action of G ,
- to \mathcal{X} is associated an element ξ of the valuation cone such that for an appropriate choice of H_0 such that the open orbit of G in X is identified with G/H_0 , the action of $e^\tau \in \mathbb{C}^*$ induced by the test configuration is given by:

$$e^\tau \cdot gH_0 = g \exp(-\tau \tilde{\xi}) H_0$$

for any lift $\tilde{\xi} \in \mathfrak{Y}(T) \otimes \mathbb{Q}$ considered as an element of the Lie algebra of T .

Furthermore, for any ξ in the valuation cone, there exists such a test configuration, and X is isomorphic to X_0 if and only if ξ is in the linear part of the valuation cone.

Degenerations of spherical varieties, and moduli questions, have been studied by Alexeev and Brion [AB06] (see also [AB04a, AB04b] for the case of reductive varieties). We adopt here a different approach to be better able to keep track of the actions of \mathbb{C}^* on the degenerations. Remark that the central fiber of a normal and equivariant test configuration which is not special would be a stable spherical variety in the sense of [AB06]. We do not study these test configurations here. Our description of the action of \mathbb{C}^* on the central fiber of a special equivariant test configuration relies heavily on the work of Brion and Pauer [BP87] on elementary embeddings of spherical homogeneous spaces.

The Donaldson-Futaki invariant of a test configuration depends only on the central fiber and the induced action of \mathbb{C}^* on the central fiber. Namely, it reduces to the Futaki invariant of the central fiber evaluated at the holomorphic vector field generating the action. The basic idea behind our computation of these numerical invariants is that we may degenerate the central fiber even more in order to acquire more symmetries, then compute the Futaki invariant on the corresponding degeneration.

For spherical varieties, this idea leads to consider only the Futaki invariants of horospherical varieties. Indeed, there always exist a test configuration with horospherical central fiber. The existence of a horospherical degeneration for spherical varieties is a classical result [Bri86, Pop87]. In the notations of Theorem B, the test configuration \mathcal{X} has a horospherical central fiber if and only if ξ is in the interior of the valuation cone. Horospherical varieties are the simplest among spherical varieties, but still form a large class containing toric varieties and homogeneous toric bundles. They should be considered as the "most symmetric" spherical varieties. In particular, even if it is possible to degenerate any spherical variety to a toric variety [AB04c], it is at the expense of some symmetries, as it can be made equivariantly only with respect to a maximal torus.

Our computation of the modified Futaki invariant on \mathbb{Q} -Fano horospherical varieties gives the following statement, where we keep the notations introduced earlier.

Theorem C. *Let X be a \mathbb{Q} -Fano horospherical variety, with moment polytope Δ^+ . Let ζ, ξ be two elements of $\mathfrak{Y}(\text{Aut}_G(X)) \otimes \mathbb{R}$ and let $\tilde{\zeta}$ and $\tilde{\xi}$ be choices of lifts in $\mathfrak{Y}(T) \otimes \mathbb{R}$. Then the modified (with respect to ζ) Futaki invariant of X evaluated at ξ is*

$$\text{Fut}_{X,\zeta}(\xi) = C \left\langle \text{bar}_{DH,\tilde{\zeta}}(\Delta^+) - 2\rho_P, \tilde{\xi} \right\rangle$$

where C is a positive constant independent of ξ .

This statement was first obtained by Mabuchi [Mab87] for smooth toric manifolds. For smooth homogeneous toric bundles, this statement was obtained by Podesta and Spiro in [PS10]. Let us also mention here the work of Alexeev and Katzarkov on K-stability of group compactifications [AK05]. Our computation is based on an expression for the curvature form of a positive hermitian metric on a polarized horospherical homogeneous space which is invariant under the action of a maximal compact subgroup of G , in terms of a convex potential associated to the metric, and a description of the asymptotic behavior of this convex function.

We will use two definitions of the Futaki invariant. First, we will use an analytic definition to obtain its precise value. The original definition of the Futaki invariant by Futaki [Fut83] was generalized by Ding and Tian to normal \mathbb{Q} -Fano varieties [DT92]. The modified Futaki invariant, the analogue for Kähler-Ricci solitons, was introduced by Tian and Zhu [TZ02], and generalized to singular varieties by Berman and Witt Nyström [BW]. To show that we can compute the Futaki invariant on a degeneration though, we will use the algebraic definition of the Futaki invariant. This was first proposed by Donaldson [Don02]. He showed that the two definitions coincide when the variety is non singular. In fact, as remarked by Li and Xu [LX14, Section 7, Remark 1], his proof extends to normal varieties. It is also Berman and Witt Nyström who generalized this algebraic definition to the modified Futaki invariant (see also [WZZ16] and [Xio14] for related recent work on Futaki invariants).

Consider the partial order on G -spherical \mathbb{Q} -Fano varieties, given by $X_0 \preceq X$ if X_0 is the central fiber of a special equivariant test configuration for X . Given a G -spherical \mathbb{Q} -Fano variety X , it is interesting to consider the set of all G -spherical \mathbb{Q} -Fano varieties smaller than X with respect to this order. It follows from [AB04b, AK05] that this poset is very well understood for reductive varieties. In particular there is a single minimum which is the central fiber of any special test configuration with horospherical central fiber, and the other elements are in bijection with the walls of the Weyl chamber, up to a finite number of isomorphisms induced by exterior automorphisms of the root system. It would be interesting to extend this precise description to all spherical \mathbb{Q} -Fano varieties. The fact that the minima are horospherical varieties is stated in the present article. More generally, one may consider similar posets for more general \mathbb{Q} -Fano varieties by considering the partial order $X_0 \preceq X$ if X_0 is the central fiber of a special test configuration for X , equivariant with respect to a Levi subgroup of the group of automorphisms of X . Many questions about these posets could clarify the interpretation of K-stability. As a precise example, let us ask the following question: *What are the minima of these posets?* In other words, if X is a \mathbb{Q} -Fano variety, and G is a Levi subgroup of $\text{Aut}^0(X)$, *under which conditions on X are there no special G -equivariant test configurations for X with central fiber not isomorphic to X ?* The only examples known to the author are horospherical varieties. One interest of knowing these minima is that, thanks to the argument we use in Section 5, they are the only varieties on which we really need to compute Futaki invariants. Of course the poset alone is not enough to recover all the information about K-stability, as one should keep track of the different ways the elements of the poset appear as central fibers of test configurations, more precisely of the different holomorphic vector fields induced by these test configurations.

The ideas and the methods of the present article should have interesting applications to several related problems. The most obvious one is K-stability of spherical varieties for polarizations not given by the anticanonical line bundle. Another direction, to which we will come back in a later article, is in Sasakian geometry, where an analogue of the Yau-Tian-Donaldson conjecture was obtained by Collins and Székelyhidi [CS]. Finally, it would be interesting to combine the ideas from our work with the work of Ilten and Süß [IS] to study varieties with a spherical action of complexity one. In particular, Langlois and Terpereau [LT16, LT] have started a study of horospherical varieties of complexity one that should lead them to a criterion for being Fano, which would be a starting point.

The reader interested in learning more about spherical varieties may consult [Per14, Tim11, Bri].

1. NOTATIONS

We introduce in this section some notions and notations on groups and Lie algebras that will be used throughout the text.

Let G be a complex linear algebraic group. We will denote its Lie algebra by the corresponding fraktur lower case letter \mathfrak{g} , and the exponential map by \exp . An (algebraic) *character* of G is an algebraic group morphism $G \rightarrow \mathbb{C}^*$. We denote the group of algebraic characters of G by $\mathfrak{X}(G)$. An (algebraic) *one parameter subgroup* of G is an algebraic group morphism $\mathbb{C}^* \rightarrow G$. We denote the set of algebraic one parameter subgroups of G by $\mathfrak{Y}(G)$. We denote by G^u the *unipotent radical* of G , and by $[G, G]$ its derived subgroup. If H is a closed subgroup of G , let $N_G(H) = \{g \in G; gHg^{-1} = H\}$ be the *normalizer* of H in G .

In this article, G will denote a connected complex linear reductive group, K will denote a maximal compact subgroup of G , and θ the *Cartan involution* of G such that $K = G^\theta$ is the fixed point set of θ . The involution θ is identified with the complex conjugation under the identification of G with the complexification of K . Let T be a maximal torus of G , stable under θ , let B be a Borel subgroup of G containing T , and let B^- be the unique Borel subgroup of G such that $B \cap B^- = T$, called the *opposite Borel subgroup* with respect to T . Let $\Phi \subset \mathfrak{X}(T)$ be the root system of (G, T) and let $\Phi^+ \subset \Phi$ be the positive roots determined by B .

A parabolic subgroup P of G containing B admits a unique Levi subgroup L containing T . The parabolic subgroup of G *opposite* to P with respect to L is defined as the unique parabolic subgroup of G such that $P \cap Q = L$ and L is also a Levi subgroup of Q . Let Φ_P denote the set of roots of P with respect to T . We denote by Φ_P^\pm the set of roots of the unipotent radical P^u of P . Alternatively, these are the roots of G that are not roots of Q , or the roots of P that are not roots of L . Let

$$2\rho_P = \sum_{\alpha \in \Phi_P^+} \alpha.$$

Consider the root decomposition of \mathfrak{g} :

$$\mathfrak{g} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha$$

where $\mathfrak{g}_\alpha = \{x \in \mathfrak{g}; \text{ad}(t)(x) = \alpha(t)x \forall t \in \mathfrak{t}\}$ is the root space for α . The Lie algebra of P is

$$\mathfrak{p} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Phi_P} \mathfrak{g}_\alpha.$$

The Cartan involution θ descends to an involution of \mathfrak{g} , still denoted by θ . It sends \mathfrak{g}_α to $\mathfrak{g}_{-\alpha}$ for $\alpha \in \Phi$.

Assume now that T is any algebraic complex torus. There is a natural duality pairing $\langle \lambda, \chi \rangle$ between $\lambda \in \mathfrak{Y}(T)$ and $\chi \in \mathfrak{X}(T)$ defined as the integer such that $\chi \circ \lambda(z) = z^{\langle \lambda, \chi \rangle}$ for $z \in \mathbb{C}^*$, which gives an isomorphism between $\mathfrak{Y}(T)$ and $\text{Hom}(\mathfrak{X}(T), \mathbb{Z})$. Given an algebraic complex subtorus $S \subset T$, there is a natural inclusion of $\mathfrak{Y}(S)$ in $\mathfrak{Y}(T)$ as a direct factor, and a natural inclusion of $\mathfrak{X}(T/S)$ in $\mathfrak{X}(T)$ as a direct factor, where a character of T/S is identified with a character of T which is trivial on S . By duality, these inclusions imply that $\mathfrak{Y}(T/S)$ is a quotient of $\mathfrak{Y}(T)$, and that $\mathfrak{X}(S)$ is a quotient of $\mathfrak{X}(T)$.

Recall that since G is reductive, $\mathfrak{g} = \mathfrak{k} \oplus J\mathfrak{k}$, where J denotes the complex structure on \mathfrak{g} . Let $\mathfrak{a} = \mathfrak{t} \cap J\mathfrak{k}$. Given $\lambda : \mathbb{C}^* \rightarrow T$ an algebraic one parameter subgroup, we consider its restriction to \mathbb{R}_+^* and associate to λ the derivative a of the restriction at 1, which lives in the tangent space $\mathfrak{t} = T_e T$. In fact, $a \in \mathfrak{a}$. Conversely any $a \in \mathfrak{a}$ defines a Lie group morphism $\tau \in \mathbb{C} \rightarrow \exp(\tau a) \in T$, which factorizes by $\mathbb{C} \rightarrow \mathbb{C}^* = \mathbb{C}/2\pi\mathbb{Z}$ if and only if a is obtained as the derivative of an algebraic one parameter subgroup as above. This correspondence embeds $\mathfrak{Y}(T)$ in \mathfrak{a} , and we identify $\mathfrak{Y}(T) \otimes \mathbb{R}$ with \mathfrak{a} . The natural duality $\langle \cdot, \cdot \rangle$ between $\mathfrak{X}(T) \otimes \mathbb{R}$ and $\mathfrak{Y}(T) \otimes \mathbb{R}$ translates, for $a \in \mathfrak{a}$ identified with an element of $\mathfrak{Y}(T) \otimes \mathbb{R}$, and $\chi \in \mathfrak{X}(T)$, as $\langle \chi, a \rangle = \ln \chi(\exp(a))$.

We will denote by κ the *Killing form* on \mathfrak{g} . It defines a scalar product on the semisimple part $\mathfrak{a} \cap [\mathfrak{g}, \mathfrak{g}]$ of \mathfrak{a} . We choose a scalar product $\{\cdot, \cdot\}$ on \mathfrak{a} which is invariant under the action of the Weyl group W , and whose restriction to the semisimple part is the Killing form κ . Note that the vector subspace on which W acts trivially, which is the intersection of \mathfrak{a} with the center of \mathfrak{g} , is then orthogonal to the semisimple part of \mathfrak{a} . Let e_α , for $\alpha \in \Phi$, be a generator of the root space \mathfrak{g}_α , such that $e_{-\alpha} = -\theta(e_\alpha)$, and $[e_\alpha, e_{-\alpha}] = t_\alpha$, where t_α is defined as the unique element of \mathfrak{a} such that $\{t_\alpha, a\} = \langle \alpha, a \rangle$ for all $a \in \mathfrak{a}$. More generally, given $\chi \in \mathfrak{X}(T)$, we denote by t_χ the unique element of \mathfrak{a} such that $\{t_\chi, a\} = \langle \chi, a \rangle$ for all $a \in \mathfrak{a}$.

2. CURVATURE FORMS ON HOROSPHERICAL HOMOGENEOUS SPACES

2.1. Horospherical homogeneous spaces. We begin this section by introducing horospherical homogeneous spaces. Our reference for these homogeneous spaces, and the horospherical \mathbb{Q} -Fano varieties that we will discuss in later sections, is [Pas08].

2.1.1. Definition and normalizer fibration.

Definition 2.1. A closed subgroup H of a connected complex reductive group G is called *horospherical* if it contains the unipotent radical U of a Borel subgroup B of G . The homogeneous space G/H is then called a *horospherical homogeneous space*.

Example 2.2. Consider the natural action of $\text{SL}_2(\mathbb{C})$ on \mathbb{C}^2 . It has two orbits : the fixed point 0 and its complement. The stabilizer of $(1, 0)$ is easily seen to be the set of upper triangular matrices with diagonal coefficients equal to one. It is

the unipotent radical U of the Borel subgroup B of $\mathrm{SL}_2(\mathbb{C})$ consisting of the upper triangular matrices, so $\mathbb{C}^2 \setminus \{0\}$ is a horospherical homogeneous space.

A torus $(\mathbb{C}^*)^n$ is a horospherical homogeneous space under its action on itself. A generalized flag manifold G/P where G is a semisimple group and P a parabolic of G is a horospherical homogeneous space. Combining the two examples gives that products of tori with generalized flag manifolds are examples of horospherical homogeneous spaces. In fact all horospherical homogeneous spaces are torus fibrations over generalized flag manifolds (see for example [Pas08]).

Proposition 2.3. *Assume that H is horospherical, then the normalizer fibration*

$$N_G(H)/H \longrightarrow G/H \longrightarrow G/N_G(H)$$

is a torus fibration over a generalized flag manifold. More precisely, the normalizer $P = N_G(H)$ of H in G is a parabolic subgroup containing B , and the quotient $P/H = T/T \cap H$ is a torus. Conversely, if H is such that $N_G(H)$ is a parabolic subgroup of G and P/H is a torus, then H is horospherical.

Example 2.4. The normalizer fibration associated to $\mathrm{SL}_2(\mathbb{C})/U$ is the one defining the complex projective line : $\mathbb{C}^* \longrightarrow \mathbb{C}^2 \setminus \{0\} \longrightarrow \mathbb{P}^1$. In this case, $P = B$ is the subgroup of upper triangular matrices.

For any closed subgroup H , the normalizer $N_G(H)$ acts on G/H by multiplication on the right by the inverse. The subgroup H in $N_G(H)$ acts trivially. The action of $(g, pH) \in G \times N_G(H)/H$ on the coset $xH \in G/H$ is then given by $(g, pH) \cdot xH = gxp^{-1}H$. The isotropy group of eH under this action is the group $\{(p, pH), p \in N_G(H)\}$. We will mainly use this for horospherical homogeneous spaces, and we denote the corresponding isotropy group by $\mathrm{diag}(P)$ in this case. It is isomorphic to P via the first projection.

2.1.2. *Polar decomposition on a horospherical homogeneous space.* We fix now H a horospherical subgroup of G , and denote by P its normalizer. The inclusion of $\mathfrak{Y}(T \cap H)$ in $\mathfrak{Y}(T)$ gives rise to a subspace $\mathfrak{a}_0 \subset \mathfrak{a}$ under the identification of \mathfrak{a} with $\mathfrak{Y}(T) \otimes \mathbb{R}$. Let \mathfrak{a}_1 denote the orthogonal complement of \mathfrak{a}_0 with respect to $\{\cdot, \cdot\}$.

Proposition 2.5. *The image of \mathfrak{a}_1 in G under the exponential is a fundamental domain for the action of $K \times H$ on G , where K acts by multiplication on the left and H by multiplication on the right by the inverse. As a consequence, the set $\{\exp(a)H; a \in \mathfrak{a}_1\} \subset G/H$ is a fundamental domain for the action of K on G/H .*

Proof. The case when the horospherical subgroup is the unipotent radical U of B is a classical result known as the *Iwasawa decomposition* (see for example [Hel78, Chapter IX, Theorem 1.3]). It states that the map

$$K \times \mathfrak{a} \times U \longrightarrow G, \quad (k, a, u) \longmapsto k \exp(a)u$$

is a diffeomorphism.

Now let H be any horospherical subgroup containing U , and \mathfrak{a}_1 be as defined above. Given $g \in G$, use the Iwasawa decomposition to write $g = k \exp(a)u$ where $a \in \mathfrak{a}$ is uniquely determined. Decompose a as $a_0 + a_1 \in \mathfrak{a}_0 \oplus \mathfrak{a}_1 = \mathfrak{a}$. Then $g = k \exp(a_1) \exp(a_0)u$ where $k \in K$, $\exp(a_0)u \in H$ and $a_1 \in \mathfrak{a}_1$ uniquely determined. \square

2.2. Hermitian metrics on linearized line bundles. We will now associate to a K -invariant hermitian metric on a linearized bundle on a horospherical homogeneous space two functions. One function will be associated to the pull back of the line bundle to G under the quotient map, and the other will be associated to the restriction to $\exp(\mathfrak{a}_1)$. Let us first describe the linearized line bundles obtained by this pull back and this restriction.

2.2.1. *Associated linearized line bundles.*

Definition 2.6. Let X be a G -variety. A G -linearized line bundle L on X is a line bundle on X equipped with an action of G such that the bundle map $L \rightarrow X$ is equivariant and the maps between the fibers induced by the action are linear.

Let G/H be a horospherical homogeneous space. Let L be a $G \times P/H$ -linearized line bundle on G/H . The fiber L_{eH} above $eH \in G/H$ defines a one dimensional representation of the isotropy group $\text{diag}(P)$. Denote by χ the character of P associated to this representation, so that $(p, pH) \cdot \xi = \chi(p)\xi$ for $\xi \in L_{eH}$.

Consider the quotient map $\pi : G \rightarrow G/H, g \mapsto gH$, and the pull back π^*L of the line bundle L to G . We still denote by π the induced map $\pi^*L \rightarrow L$. Since π is equivariant under the action of G by multiplication on the left, π^*L admits a pulled back G -linearization. Let us choose a non zero element $s(e) \in (\pi^*L)_e$. Together with the linearization, it provides a global trivialization

$$s : g \in G \mapsto g \cdot s(e) \in (\pi^*L)_g.$$

Denote by $\iota : P/H \rightarrow G/H$ the inclusion. It is equivariant under the action of $P \times P/H$. The restriction ι^*L of the line bundle L to P/H provides a $P \times P/H$ -linearized line bundle on the torus P/H . In general the linearizations of ι^*L for the action of $P \times \{eH\} \simeq P$ and the action of P through $\{e\} \times P/H$ are different. They are related by:

$$\begin{aligned} (p, H) \cdot \xi &= (e, p^{-1}H)(p, pH) \cdot \xi \\ &= \chi(p)(e, p^{-1}H) \cdot \xi \end{aligned}$$

for $p \in P, \xi \in \iota^*L$.

We still denote by ι the inclusion $\iota^*L \rightarrow L$. We define two trivializations s_l and s_r of ι^*L :

$$\begin{aligned} s_l(pH) &= (p, H) \cdot (\iota^{-1} \circ \pi)(s(e)), \\ s_r(pH) &= (e, p^{-1}H) \cdot (\iota^{-1} \circ \pi)(s(e)). \end{aligned}$$

By the previous calculation, they satisfy

$$s_l(pH) = \chi(p)s_r(pH).$$

The following diagram summarizes the objects introduced in this subsection.

$$\begin{array}{ccccc} \pi^*L & \longrightarrow & L & \longleftarrow & \iota^*L \\ \downarrow & & \downarrow & & \downarrow \\ G & \xrightarrow{\pi} & G/H & \xleftarrow{\iota} & P/H \end{array} \begin{array}{c} \nearrow s \\ \searrow s_l \\ \nearrow s_r \end{array}$$

2.2.2. *Functions associated to hermitian metrics.* Let L be a $G \times P/H$ -linearized line bundle on G/H , and denote by χ the corresponding character of $\text{diag}(P)$. Let q be a smooth hermitian metric on L . Consider the pull-back π^*q of the metric q to π^*L . Denote by $\phi : G \rightarrow \mathbb{R}$ the potential of π^*q with respect to the section s , that is the function on G defined by

$$\phi(g) = -2 \ln |s(g)|_{\pi^*q}.$$

We associate to q another function, this time associated with its restriction to $L|_{P/H}$. Denote by $u : \mathfrak{a}_1 \rightarrow \mathbb{R}$ the function defined by

$$u(x) = -2 \ln |s_r(\exp(x)H)|_{\iota^*q}.$$

Proposition 2.7. *Assume that the metric q is invariant under the action of the compact group K . Then q is uniquely determined by u . Furthermore, we have*

$$\phi(k \exp(x)h) = u(x) - 2 \ln |\chi(\exp(x)h)|.$$

for any $k \in K$, $x \in \mathfrak{a}_1$ and $h \in H$.

Proof. It is clear that π^*q is completely determined by its global potential ϕ on G , thus that q is completely determined by ϕ . We only need to prove the relation between u and ϕ , since any $g \in G$ can be written as $g = k \exp(x)h$ with $k \in K$, $x \in \mathfrak{a}_1$ and $h \in H$ by Proposition 2.5.

First remark that since q is invariant under the action of K , and π is equivariant under the action of G , π^*q is also invariant under the action of K . Then the invariance of the section s yields

$$\phi(k \exp(x)h) = -2 \ln |s(k \exp(x)h)|_{\pi^*q} = -2 \ln |k \cdot s(\exp(x)h)|_{\pi^*q} = \phi(\exp(x)h).$$

We then remark that

$$\begin{aligned} \pi(s(\exp(x)h)) &= \pi((\exp(x)h) \cdot s(e)) \\ &= (\exp(x)h, H) \cdot \pi(s(e)) \\ &= (\exp(x), H)(h, hH) \cdot \pi(s(e)) \\ &= \chi(h)(\exp(x), H) \cdot \pi(s(e)) \end{aligned}$$

by definition of χ .

We then write, since $\exp(x) \in P$,

$$\begin{aligned} \phi(\exp(x)h) &= -2 \ln (|\chi(h)| |(\exp(x), H) \cdot \pi(s(e))|_q) \\ &= -2 \ln |(\exp(x), H) \cdot \iota^{-1} \circ \pi(s(e))|_{\iota^*q} - 2 \ln |\chi(h)| \\ &= -2 \ln |s_l(\exp(x)H)|_{\iota^*q} - 2 \ln |\chi(h)| \end{aligned}$$

we then use the relation between sections described in Section 2.2.1 to obtain

$$\begin{aligned} \phi(\exp(x)h) &= -2 \ln (|\chi(\exp(x))| |s_r(\exp(x)H)|_{\iota^*q}) - 2 \ln |\chi(h)| \\ &= -2 \ln |s_r(\exp(x)H)|_{\iota^*q} - 2 \ln |\chi(\exp(x)h)|. \end{aligned}$$

Recalling the definition of u , we obtain the statement. \square

2.3. Pointwise expression of a curvature form. Let L be a $G \times P/H$ -linearized line bundle on G/H , with associated character of $\text{diag}(P)$ denoted by χ . Let q be a K -invariant smooth hermitian metric on L , with associated functions $\phi : G \rightarrow \mathbb{R}$ and $u : \mathfrak{a}_1 \rightarrow \mathbb{R}$.

Recall that the curvature form ω of q is a global (1,1)-form defined locally as follows. If $s_0 : U \subset G/H \rightarrow L$ is a local trivialization of L , and $\psi(z) := -2 \ln |s_0(z)|_q$, then $\omega = i\partial\bar{\partial}\psi$ on U . If q is invariant under the action of K then ω is also invariant. We want to compute the expression of ω in terms of χ and u .

In general we cannot find a global trivialization of L on G/H , and cannot find a global potential for ω . This is the case in particular for generalized flag manifolds. We will bypass this difficulty by computing ω through its pull back to G under the quotient map. This approach is similar to the use of *quasipotentials* by Azad and Biswas in [AB03] for generalized flag manifolds.

We can identify the tangent space at eH to G/H with

$$\mathfrak{g}/\mathfrak{h} \simeq \bigoplus_{\alpha \in \Phi_P^+} \mathbb{C}e_{-\alpha} \oplus \mathfrak{a}_1 \oplus J\mathfrak{a}_1.$$

Choose a basis l_1, \dots, l_r of the real vector space \mathfrak{a}_1 . Denote by (x_1, \dots, x_r) the corresponding coordinates of a point $x = \sum_i x_i l_i \in \mathfrak{a}_1$. A complex basis of the tangent space $T_{eH}G/H$ is then given by the union of the l_j and the $e_{-\alpha}$ for $\alpha \in \Phi_P^+$.

On $P/H \subset G/H$, given a tangent vector ξ at the coset eH , we can define a smooth real holomorphic vector field $R\xi$, invariant under the action of P/H by multiplication on the right, simply by transporting the given tangent vector by the holomorphic action:

$$R\xi : pH \mapsto (H, p^{-1}H) \cdot \xi \in T_{pH}P/H.$$

Consider the complex basis of holomorphic (1,0)-vector fields composed of the $(Rl_j - iJRl_j)/2$ and $(Re_{-\alpha} - iJRe_{-\alpha})/2$, where J denotes the complex structure on G/H , and i is the complex structure coming from the complexification in $TG/H \otimes \mathbb{C}$. We denote the dual basis of (1,0)-forms by $\{\gamma_j\}_j \cup \{\gamma_\alpha\}_\alpha$. We will compute the curvature (1,1)-form ω pointwise in the basis of (1,1)-forms obtained from these.

Theorem 2.8. *Let ω be the K -invariant curvature form of a K -invariant metric q . Then the form ω is determined by its restriction to P/H , given for $x \in \mathfrak{a}_1$ by*

$$\omega_{\exp(x)H} = \sum_{1 \leq j_1, j_2 \leq r} \frac{1}{4} \frac{\partial^2 u}{\partial x_{j_1} \partial x_{j_2}}(x) i\gamma_{j_1} \wedge \bar{\gamma}_{j_2} + \sum_{\alpha \in \Phi_P^+} \langle \alpha, \nabla u(x)/2 - t_\chi \rangle i\gamma_\alpha \wedge \bar{\gamma}_\alpha$$

where ∇u is the gradient of u with respect to the scalar product $\{\cdot, \cdot\}$.

In order to prove the theorem we need to obtain an infinitesimal decomposition, adapted to the polar decomposition $G = K \exp(\mathfrak{a}_1)H$, for elements of $\mathfrak{g}/\mathfrak{h}$. Recall that U is the unipotent radical of B and is a subgroup of H . It is also the image under the exponential of $\bigoplus_{\alpha \in \Phi^+} \mathbb{C}e_\alpha$. It will be enough to obtain an infinitesimal decomposition adapted to the Iwasawa decomposition $G = K \exp(\mathfrak{a})U$, which is achieved by the following lemma.

Lemma 2.9. *Let $\{z_j\}_j$ and $\{z_\alpha\}_\alpha$ denote complex numbers, and let*

$$f = \sum_{1 \leq j \leq r} z_j l_j + \sum_{\alpha \in \Phi_P^+} z_\alpha e_{-\alpha} \in \mathfrak{g}/\mathfrak{h}$$

Then $\exp(f) = k \exp(y + O)u$ where $k \in K$, $y \in \mathfrak{a}$, $O \in \mathfrak{g}$ is of order strictly higher than two in the z_j and z_α , and $u \in U$. Furthermore, if $z_j = x_j + iy_j$,

$$y = \sum_{1 \leq j \leq r} x_j l_j + \sum_{\alpha \in \Phi_P^+} z_\alpha \bar{z}_\alpha t_\alpha / 2.$$

Proof. Recall that θ denotes the Cartan involution on \mathfrak{g} , with fixed point set \mathfrak{k} . We will denote by O a term in \mathfrak{g} of order strictly higher than two in the z_j and z_α , which may change from line to line.

Write $f = A_1 + A_2 - A_3$, with

$$A_1 = \sum_{1 \leq j \leq r} y_j J l_j + \sum_{\alpha \in \Phi_P^+} (z_\alpha e_{-\alpha} + \theta(z_\alpha e_{-\alpha})) \in \mathfrak{k}$$

$$A_2 = \sum_{1 \leq j \leq r} x_j l_j \in \mathfrak{a}_1$$

$$A_3 = \sum_{\alpha \in \Phi_P^+} \theta(z_\alpha e_{-\alpha}) \in \sum_{\alpha \in \Phi_P^+} \mathbb{C} e_\alpha.$$

Using the Baker-Campbell-Hausdorff formula yields

$$\exp(-A_1) \exp(f) \exp(A_3) = \exp(A_2 + 1/2([A_2, A_1] + [A_1, A_3] + [A_2, A_3]) + O)$$

We now decompose $A_2 + 1/2([A_2, A_1] + [A_1, A_3] + [A_2, A_3]) = B_1 + B_2 + B_3$ with $B_1 \in \mathfrak{k}$, $B_2 \in \mathfrak{a}$ and $B_3 \in \mathfrak{u}$. It is easy to see that B_1 and B_3 are of order two in the z_j , z_α , so that, by the Baker-Campbell-Hausdorff formula again,

$$\exp(-B_1) \exp(-A_1) \exp(f) \exp(A_3) \exp(-B_3) = \exp(B_2 + O).$$

We have $\exp(-B_1) \exp(-A_1) \in K$, and $\exp(A_3) \exp(-B_3) \in U$, so to conclude the proof it remains to compute $y = B_2$. There are two contributions to this term, the first being A_2 and the second, coming from $\frac{1}{2}[A_1, A_3]$, which is

$$\begin{aligned} \frac{1}{2} \sum_{\alpha \in \Phi_P^+} [z_\alpha e_{-\alpha}, \theta(z_\alpha e_{-\alpha})] &= \frac{1}{2} \sum_{\alpha \in \Phi_P^+} z_\alpha \bar{z}_\alpha [e_{-\alpha}, \theta(e_{-\alpha})] \\ &= \frac{1}{2} \sum_{\alpha \in \Phi_P^+} z_\alpha \bar{z}_\alpha t_\alpha \end{aligned}$$

□

We now proceed to prove the theorem.

Proof of Theorem 2.8. Let π denote again the quotient map $G \rightarrow G/H$. Consider $\pi^*\omega$. This is the curvature form of the pulled back metric π^*q on π^*L . Let ϕ be the global potential of π^*q on G . Then ϕ is a global $i\partial\bar{\partial}$ potential for $\pi^*\omega$, which means that $\pi^*\omega = i\partial\bar{\partial}\phi$. Recall from Proposition 2.7 that $\phi(k \exp(x)h) = u(x) - 2 \ln(\chi(\exp(x)h))$.

Consider on $P \subset G$ the right invariant vector fields $\tilde{R}l_j$, respectively $\tilde{R}e_{-\alpha}$ obtained by transporting the elements l_j respectively $e_{-\alpha}$ from $\mathfrak{g} \simeq T_e G$ by the action of P on G by multiplication on the right by the inverse. These vector fields are sent to Rl_j respectively $Re_{-\alpha}$ by π_* . Since for any elements f_1, f_2 of $T_g G \otimes \mathbb{C}$, we have

$$\pi^*\omega_g(f_1, f_2) = \omega_{\pi(g)}(\pi_*(f_1), \pi_*(f_2))$$

it will be enough to compute $\pi^*\omega$ on pairs of holomorphic $(1,0)$ vector fields Z_j or Z_α corresponding to the real holomorphic vector fields just defined.

Let f_1 and f_2 be two elements of \mathfrak{g} , and let Z_1, Z_2 be the corresponding right- P -invariant holomorphic $(1,0)$ vector fields on $P \subset G$ as defined above. Since $\pi^*\omega = i\partial\bar{\partial}\phi$, we have, at $p \in P$,

$$(\pi^*\omega)_p(Z_1, \bar{Z}_2) = i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 \phi(\exp(z_1 f_1 + z_2 f_2)p).$$

We will carry out this computation at $p = \exp(x)$ for f_1 and f_2 two elements in $\mathfrak{g}/\mathfrak{h}$.

Consider now f as in Lemma 2.9 and $x \in \mathfrak{a}_1$. Then

$$\begin{aligned} \exp(f)\exp(x) &= k \exp(y + O)u \exp(x) \\ &= k \exp(y + O)\exp(x)u' \end{aligned}$$

where $u' \in U$, because T normalizes U . Then by the Baker-Campbell-Hausdorff formula and since O is of order strictly higher than two in z_j, z_α , there exists an $O' \in \mathfrak{g}$, still of order strictly higher than two in z_j, z_α , such that

$$\exp(f)\exp(x) = k \exp(x + y + O')u.$$

We deduce that $\phi(\exp(f)\exp(x)) = \phi(\exp(x + y + O'))$, because $u \in U \subset H$, and any character of P vanishes on U . Then we apply this to obtain, given f_1, f_2 in $\mathfrak{g}/\mathfrak{h}$,

$$\begin{aligned} (\pi^*\omega)_{\exp(x)}(Z_1, \bar{Z}_2) &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 \phi(\exp(z_1 f_1 + z_2 f_2)\exp(x)) \\ &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 \phi(\exp(x + y + O')) \end{aligned}$$

where y is given by Lemma 2.9 for $f = z_1 f_1 + z_2 f_2$,

$$\begin{aligned} &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 \phi(\exp(x + y)) \\ &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 \phi(\exp(x + y^1)\exp(y - y^1)) \end{aligned}$$

where y^1 is the projection in \mathfrak{a}_1 of $y \in \mathfrak{a} = \mathfrak{a}_0 \oplus \mathfrak{a}_1$,

$$\begin{aligned} &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 (u(x + y^1) - 2 \ln(\chi(\exp(x + y)))) \\ &= i \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 (u(x + y^1) - 2 \langle \chi, x + y \rangle). \end{aligned}$$

Together with the precise value of y given by Lemma 2.9, this allows us to compute the values of $(\pi^*\omega)_{\exp(x)}(Z_1, \bar{Z}_2)$ for all choices of Z_1, Z_2 in the set of right- P -invariant holomorphic $(1,0)$ -vector fields obtained from elements f_1, f_2 of $\mathfrak{g}/\mathfrak{h}$.

(i) Let us first apply this to $f_1 = l_{j_1}$, $f_2 = l_{j_2}$. We obtain

$$\begin{aligned} (\pi^*\omega)_{\exp(x)}(Z_1, \bar{Z}_2) &= \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 (u(x + x_1 l_{j_1} + x_2 l_{j_2}) - 2 \langle \chi, x \rangle) \\ &= \frac{1}{4} \frac{\partial^2}{\partial x_1 \partial x_2} \Big|_0 u(x + x_1 l_{j_1} + x_2 l_{j_2}) \\ &= \frac{1}{4} \frac{\partial^2 u}{\partial x_{j_1} \partial x_{j_2}}(x). \end{aligned}$$

(ii) If $f_1 = l_j$ and $f_2 = e_{-\alpha}$ then Lemma 2.9 gives

$$y = x_1 l_j + \frac{1}{2} z_2 \bar{z}_2 t_\alpha$$

and it is easy to see that the double derivative vanishes.

(iii) Similarly, if $f_1 = e_{-\alpha_1}$ and $f_2 = e_{-\alpha_2}$, with $\alpha_1 \neq \alpha_2$, then

$$y = \frac{1}{2} (z_1 \bar{z}_1 t_{\alpha_1} + z_2 \bar{z}_2 t_{\alpha_2})$$

and the double derivative vanishes again.

(iv) The remaining case is when $f_1 = f_2 = e_{-\alpha}$. In that case,

$$y = \frac{1}{2} |z_1 + z_2|^2 t_\alpha,$$

so that

$$\begin{aligned} (\pi^*\omega)_{\exp(x)}(Z_1, Z_2) &= \frac{\partial^2}{\partial z_1 \partial \bar{z}_2} \Big|_0 (u(x + |z_1 + z_2|^2 t_\alpha^1 / 2) - 2 \langle \chi, x + |z_1 + z_2|^2 t_\alpha / 2 \rangle) \\ &= \{ \nabla u(x), t_\alpha^1 / 2 \} - \langle \chi, t_\alpha \rangle \\ &= \{ \nabla u(x) / 2 - t_\chi, t_\alpha \}. \end{aligned}$$

We can indeed replace t_α^1 with t_α because $\nabla u(x) \in \mathfrak{a}_1$, and, by definition, \mathfrak{a}_1 is orthogonal to \mathfrak{a}_0 . \square

3. TEST CONFIGURATIONS OF SPHERICAL VARIETIES

3.1. Colored fans and spherical varieties. We first review general results about spherical varieties. We will use [Kno91] as main reference for this section. As mentioned by Knop, the theory was initially developed by Luna and Vust [LV83].

Definition 3.1. A normal variety X equipped with an action of G is called *spherical* if a Borel subgroup B of G acts on X with an open and dense orbit.

A homogeneous space G/H which is a spherical variety under the action of G is a *spherical homogeneous space*. A *spherical subgroup* is a closed subgroup H such that G/H is a spherical homogeneous space.

Let X be a spherical variety and x a point in the open orbit of B . Denote by H the isotropy group of x in G . The pair (X, x) is called a *spherical embedding* of the spherical homogeneous space G/H , and is equipped with a natural inclusion of G/H in X through the G -equivariant map $gH \mapsto g \cdot x$.

Example 3.2. A horospherical homogeneous space is spherical: if H contains the unipotent radical U of B , and B^- is a Borel subgroup opposite to B , then B^-H is open and dense in G , or equivalently, B^-H/H is open and dense in G/H . An

embedding of a horospherical space is called a *horospherical embedding* of G/H , or a *horospherical variety*.

Example 3.3. The group G itself is a spherical homogeneous space under the action of $G \times G$ defined by $(g_1, g_2) \cdot g = g_1 g g_2^{-1}$. Indeed, if B and B^- are opposite Borel subgroups of G , the *Bruhat decomposition* shows that the $B \times B^-$ -orbit BB^- is open and dense in G .

3.1.1. *Valuation cone and colors.* Let O be a spherical homogeneous space under the action of G . Let $k = \mathbb{C}(O)$ be the function field of O . The action of G on k is defined by $(g \cdot f)(x) = f(g^{-1} \cdot x)$ for $g \in G$, $f \in k$, $x \in O$.

Definition 3.4. A *valuation* of k is a map $\nu : k^* = k \setminus \{0\} \rightarrow \mathbb{Q}$ such that:

- $\nu(\mathbb{C}^*) = 0$,
- $\nu(f_1 + f_2) \geq \min\{\nu(f_1), \nu(f_2)\}$ when f_1, f_2 and $f_1 + f_2$ are in k^* ,
- $\nu(f_1 f_2) = \nu(f_1) + \nu(f_2)$ for all $f_1, f_2 \in k^*$.

Let us now choose B a Borel subgroup of G . Define $\mathcal{M}_B(O) \subset \mathfrak{X}(B)$ as the set of characters χ such that there exists a function $f \in k^*$ with $b \cdot f = \chi(b)f$. It is a subgroup of $\mathfrak{X}(B)$, and hence a finitely generated free abelian group.

Define $\mathcal{N}_B(O) = \text{Hom}(\mathcal{M}, \mathbb{Z})$. To any valuation ν of k we can associate an element ρ_ν of $\mathcal{N}_B(O) \otimes \mathbb{Q}$, defined by $\rho_\nu(\chi) = \nu(f)$ where $f \in k^*$ is such that $b \cdot f = \chi(b)f$ for all $b \in B$. This is well defined because B has an open and dense orbit, so two such functions are non-zero scalar multiples of each other.

The map $\nu \mapsto \rho_\nu$ is injective on the set of G -invariant valuations, and we denote by $\mathcal{V}_B(O)$ the image of the set of G -invariant valuations of k under this map. This is a convex cone in $\mathcal{N}_B(O) \otimes \mathbb{Q}$ called the *valuation cone* of O (with respect to B). Although the set of G -invariant valuations of k does not depend on the choice of Borel subgroup, we will use in the following its image $\mathcal{V}_B(O)$, which does depend on the choice of B .

As an example, let us record the following characterization of horospherical varieties. Other examples will be described in Section 5.4.2.

Proposition 3.5. [BP87, Corollaire 5.4] *A spherical homogeneous space O is a horospherical homogeneous space if and only if its valuation cone $\mathcal{V}_B(O)$ is $\mathcal{N}_B(O) \otimes \mathbb{Q}$.*

Denote the set of B -stable prime divisors in O by $\mathcal{D}_B(O)$. An element of $\mathcal{D}_B(O)$ is called a *color* of O . A color $D \in \mathcal{D}_B(O)$ defines a valuation on G/H and thus an element $\rho(D)$ in $\mathcal{N}_B(O)_\mathbb{Q}$. However, the map $\rho : \mathcal{D}_B(O) \rightarrow \mathcal{N}_B(O) \otimes \mathbb{Q}$ is not injective in general.

We will in general drop the mention of B and O in the notations, as no confusion should be possible. When we will use two opposite Borel subgroups B and B^- , we will use a minus sign in subscript to denote objects defined with respect to B^- .

3.1.2. *Colored fans.* Let G/H be a spherical homogeneous space, and choose B a Borel subgroup of G . Let $\mathcal{V} \subset \mathcal{N} \otimes \mathbb{Q}$ be the valuation cone of G/H with respect to B and let \mathcal{D} be its set of colors.

Definition 3.6.

- A *colored cone* is a pair $(\mathcal{C}, \mathcal{R})$, where $\mathcal{R} \subset \mathcal{D}$ with $0 \notin \rho(\mathcal{R})$, and $\mathcal{C} \subset \mathcal{N} \otimes \mathbb{Q}$ is a strictly convex cone generated by $\rho(\mathcal{R})$ and finitely many elements of \mathcal{V} such that the intersection of the relative interior of \mathcal{C} with \mathcal{V} is not empty.

- Given two colored cones $(\mathcal{C}, \mathcal{R})$ and $(\mathcal{C}_0, \mathcal{R}_0)$, we say that $(\mathcal{C}_0, \mathcal{R}_0)$ is a *face* of $(\mathcal{C}, \mathcal{R})$ if \mathcal{C}_0 is a face of \mathcal{C} and $\mathcal{R}_0 = \mathcal{R} \cap \rho^{-1}(\mathcal{C}_0)$.
- A *colored fan* is a non-empty finite set \mathcal{F} of colored cones such that the face of any colored cone in \mathcal{F} is still in \mathcal{F} , and any $v \in \mathcal{V}$ is in the relative interior of at most one cone.

Theorem 3.7. [Kno91, Theorem 3.3] *There is a bijection $(X, x) \mapsto \mathcal{F}_X$ between embeddings of G/H up to G -equivariant isomorphism and colored fans. There is a bijection $Y \mapsto (\mathcal{C}_Y, \mathcal{R}_Y)$ between the orbits of G in X , and the colored cones in \mathcal{F}_X . An orbit Y is in the closure of another orbit Z in X if and only if the colored cone $(\mathcal{C}_Z, \mathcal{R}_Z)$ is a face of $(\mathcal{C}_Y, \mathcal{R}_Y)$.*

The *support* of the colored fan \mathcal{F}_X is defined as $|\mathcal{F}_X| = \bigcup \{\mathcal{C}; (\mathcal{C}, \mathcal{R}) \in \mathcal{F}_X\}$.

Proposition 3.8. [Kno91, Theorem 4.2] *A spherical variety X is complete if and only if the support $|\mathcal{F}_X|$ of its colored fan contains \mathcal{V} .*

Given X a spherical embedding of G/H , we denote by \mathcal{D}_X its set of colors, which is the union of all sets $\mathcal{R} \subset \mathcal{D}$ for $(\mathcal{C}, \mathcal{R}) \in \mathcal{F}_X$.

Definition 3.9. A spherical variety X is called *toroidal* if \mathcal{D}_X is empty.

Example 3.10. [Pas08, Exemple 1.10] A horospherical variety is toroidal if the fibration structure of the horospherical homogeneous space given in Proposition 2.3 extends to the embedding. In other words, toroidal horospherical varieties are precisely the homogeneous fibrations over generalized flag manifolds, with fibers toric varieties.

In the case of the horospherical homogeneous space $\mathbb{C}^2 \setminus \{0\} \simeq \mathrm{SL}_2(\mathbb{C})/U$, there are two complete embeddings: \mathbb{P}^2 and the blow up of \mathbb{P}^2 at one point. The latter is toroidal, and the fibration structure is obvious, while the former is not toroidal.

3.1.3. Equivariant automorphisms. The classification of spherical embeddings up to G -equivariant automorphisms shows that the group of G -equivariant automorphisms of a spherical variety X is isomorphic to the group of G -equivariant automorphisms of its open G -orbit, through the restriction. Indeed, the construction in [Kno91] of the colored fan \mathcal{F}_X of a spherical embedding (X, x) of G/H does not depend on the choice of base point x such that its isotropy group is H .

Choose x a base point in the open G -orbit in X , and let σ be an equivariant automorphism of the open G -orbit. Then the stabilizer H of x in G is also the stabilizer of its image $\sigma(x)$ by σ , since σ commutes with G . Then (X, x) , and $(X, \sigma(x))$ are two embeddings of G/H with the same fan \mathcal{F}_X , so they are G -equivariantly isomorphic by Theorem 3.7, which means that the equivariant automorphism of G/H sending x to $\sigma(x)$ extends to X .

If we fix a base point (or rather the stabilizer H of a base point), then we get an explicit description of these equivariant automorphisms. Indeed, the group of G -equivariant automorphisms of G/H is isomorphic to the quotient $N_G(H)/H$, whose action on G/H is induced by the action of $N_G(H)$ by multiplication on the right by the inverse: $p \cdot gH = gHp^{-1} = gp^{-1}H$ (see for example [Tim11, Proposition 1.2]).

3.1.4. Morphisms between spherical varieties. Let $H < H'$ be two spherical subgroups of G . Denote by $\phi : G/H \rightarrow G/H'$ the corresponding G -equivariant surjective map of homogeneous spaces. It induces a surjective morphism

$$\phi_* : \mathcal{N}(G/H) \otimes \mathbb{Q} \rightarrow \mathcal{N}(G/H') \otimes \mathbb{Q}.$$

Let $\mathcal{D}_\phi \subset \mathcal{D}(G/H)$ be the set of all $D \in \mathcal{D}(G/H)$ such that $\phi(D) = G/H$.

Theorem 3.11. [Kno91, Theorem 4.1] *Let (X, x) , respectively (X', x') , be a spherical embedding of G/H , respectively G/H' , then $\phi : G/H \rightarrow G/H'$ extends to a G -equivariant morphism $\phi : X \rightarrow X'$ sending x to x' if and only if for every colored cone $(\mathcal{C}, \mathcal{R}) \in \mathcal{F}_X$, there exists a colored cone $(\mathcal{C}', \mathcal{R}') \in \mathcal{F}_{X'}$ such that $\phi_*(\mathcal{C}) \subset \mathcal{C}'$ and $\phi_*(\mathcal{R} \setminus \mathcal{D}_\phi) \subset \mathcal{R}'$.*

3.2. Line bundles on spherical varieties. Let us recall some results from [Bri89] about line bundles on spherical varieties.

3.2.1. Cartier divisors. Let X be a complete spherical variety, embedding of some spherical homogeneous space O . Let \mathcal{I}_X^G denote the finite set of G -stable irreducible divisors of X . Any divisor $Y \in \mathcal{I}_X^G$ corresponds to a ray $(\mathcal{C}, \emptyset) \in \mathcal{F}_X$, and we denote by u_Y the indivisible generator of this ray in \mathcal{N} . The set \mathcal{D} of colors of O is in bijection with the set of irreducible B -stable but not G -stable divisors in X , by associating to a color of O its closure in X .

Any B -stable Weil divisor d on X writes

$$d = \sum_{Y \in \mathcal{I}_X^G} n_Y Y + \sum_{D \in \mathcal{D}} n_D \bar{D}$$

for some integers n_Y, n_D . In fact, any Weil divisor is linearly equivalent to a B -invariant divisor and Brion proved the following criterion to characterize Cartier divisors.

Proposition 3.12. [Bri89, Proposition 3.1] *A B -stable Weil divisor in X is Cartier if and only if there exists an integral piecewise linear function l_d on the fan \mathcal{F}_X such that*

$$d = \sum_{Y \in \mathcal{I}_X^G} l_d(u_Y) Y + \sum_{D \in \mathcal{D}_X} l_d(\rho(D)) \bar{D} + \sum_{D \in \mathcal{D} \setminus \mathcal{D}_X} n_D \bar{D}$$

for some integers n_D .

3.2.2. Ample Cartier divisors and polytopes. Let \mathcal{F}_X^{\max} denote the set of cones $\mathcal{C} \subset \mathcal{N} \otimes \mathbb{Q}$ of maximal dimension, such that there exists $\mathcal{R} \subset \mathcal{D}$ with $(\mathcal{C}, \mathcal{R}) \in \mathcal{F}_X$. If d is a Cartier divisor and $\sigma \in \mathcal{F}_X^{\max}$, let m_σ be the element of \mathcal{M} such that $l_d(x) = m_\sigma(x)$ for $x \in \sigma$. Since we assumed X complete, l_d is uniquely determined by the m_σ .

Proposition 3.13. [Bri89, Théorème 3.3] *Assume X is complete and*

$$d = \sum_{Y \in \mathcal{W}_G} l_d(v_Y) Y + \sum_{D \in \mathcal{D}_X} l_d(\rho(D)) \bar{D} + \sum_{D \in \mathcal{D} \setminus \mathcal{D}_X} n_D \bar{D}$$

is a Cartier divisor on X . It is ample if and only if the following conditions are satisfied:

- the function l_d is convex,
- $m_{\sigma_1} \neq m_{\sigma_2}$ if $\sigma_1 \neq \sigma_2 \in \mathcal{F}_X(n)$,
- $n_D > l_d(\rho(D))$ for all $D \in \mathcal{D} \setminus \mathcal{D}_X$.

To a Cartier divisor d , we associate a polytope $\Delta_d \subset \mathcal{M} \otimes \mathbb{R}$ defined as the set of $m \in \mathcal{M} \otimes \mathbb{R}$ such that $m \in -m_\sigma + \sigma^\vee$ for all $\sigma \in \mathcal{F}_X(n)$, and $m(\rho_D) > n_D$ for all $D \in \mathcal{D} \setminus \mathcal{D}_X$.

The support function $v_\Delta : \mathcal{N} \otimes \mathbb{R} \rightarrow \mathbb{R}$ of a polytope $\Delta \subset \mathcal{M} \otimes \mathbb{R}$ is defined by

$$v_\Delta(x) = \sup\{m(x); m \in \Delta\}.$$

If d is ample then $l_d(x) = v_{\Delta_d}(-x)$ for $x \in |\mathcal{F}_X|$.

3.2.3. Linearized line bundles and moment polytopes. Let L be a G -linearized ample line bundle on a spherical variety X . Let B be a Borel subgroup of G and T a maximal torus of B . Denote by V_λ an irreducible representation of G of highest weight $\lambda \in \mathfrak{X}(T)$ with respect to B . Since X is spherical, for all $r \in \mathbb{N}$, there exists a finite set $\Delta_r \subset \mathfrak{X}(T)$ such that

$$H^0(X, L^r) = \bigoplus_{\lambda \in \Delta_r} V_\lambda.$$

Definition 3.14. The *moment polytope* Δ_L of L with respect to B is defined as the closure of $\bigcup_{r \in \mathbb{N}^*} \Delta_r/r$ in $\mathfrak{X}(T) \otimes \mathbb{R}$.

Even though it is not clear from the definition, Δ_L is a polytope. More precisely, we may recall the explicit relation between Δ_L and the polytope associated to a Cartier divisor whose associated sheaf is L . Choose s a global B -semi-invariant section of L , so that the zero divisor d of s is an ample B -invariant Cartier divisor. Let $\chi(s)$ be the character of B defined by s .

Proposition 3.15. ([Bri89, Proposition 3.3], see also [Bri, Section 5.3]) *The moment polytope Δ_L of L and the polytope Δ_d associated to d are related by $\Delta_L = \chi(s) + \Delta_d$.*

3.2.4. Anticanonical line bundle. Let us now recall some results from [GH15]. In this article, Gagliardi and Hofscheier study the anticanonical line bundle on a spherical variety, in particular on \mathbb{Q} -Fano spherical varieties. It is based on the work of Brion [Bri97], and the analogue for \mathbb{Q} -Fano horospherical varieties by Pasquier [Pas08]. We consider the anticanonical divisor on a \mathbb{Q} -Fano spherical variety X . It is clear that the discussion of Cartier divisors and moment polytopes above extends to \mathbb{Q} -Cartier divisors and linearized \mathbb{Q} -line bundles. Let K_X^{-1} denote the (naturally linearized) \mathbb{Q} -line bundle on X .

Let P be the stabilizer of the open orbit of B in X . There exists a B -semi-invariant section of K_X^{-1} with weight $2\rho_P$ and divisor

$$d = \sum_{Y \in \mathcal{I}_X^G} Y + \sum_{D \in \mathcal{D}} n_D \bar{D}$$

where the n_D are explicitly obtained in terms of $2\rho_P$ and the types of the roots (see [GH15] for a precise description of these coefficients).

The moment polytope Δ^+ is then $2\rho_P + \Delta_d$ by Proposition 3.15, and furthermore the dual polytope Δ_d^* of Δ_d is a \mathbb{Q} - G/H -reflexive polytope in the sense of [GH15], which can be defined as the convex hull:

$$\Delta_d^* = \text{conv}(\{\rho_D/n_D, D \in \mathcal{D}\} \cup \{u_Y, Y \in \mathcal{I}_X^G\}).$$

The \mathbb{Q} -Fano variety X can further be recovered from its \mathbb{Q} - G/H -reflexive polytope Δ_d^* by the following procedure, detailed in [GH15]. The colored fan of X is obtained from Δ_d as the union of the colored cones $(\text{Cone}(F), \rho^{-1}(F))$ for all faces F of Δ_d^* such that the intersection of the relative interior of $\text{Cone}(F)$ with the valuation cone \mathcal{V} is not empty.

3.3. Equivariant degenerations of spherical spaces.

3.3.1. *Adapted parabolic and Levi subgroups.* Let X be a spherical variety under the action of G and B a Borel subgroup of G . The stabilizer in G of the open orbit of B is a parabolic subgroup of G containing B called the *adapted parabolic*.

Definition 3.16. Let H be a spherical subgroup of G . An *elementary embedding* of G/H is a spherical embedding (E, x) of G/H such that the boundary $E_0 = E \setminus (G/H)$ is a single codimension one G -orbit (necessarily closed).

The colored fan of an elementary embedding is a single ray in the valuation cone, with no colors [BP87, 2.2].

Choose B a Borel subgroup of G such that BH is open in G . The adapted parabolic P is also the stabilizer in G of BH .

Definition 3.17. A Levi subgroup L of P with connected center C is called *adapted to H* if the following conditions hold:

- $P \cap H = L \cap H$,
- $L \cap H$ contains the derived subgroup $[L, L]$,
- for any elementary embedding (E, x) of G/H with closed orbit E_0 , the closure $\overline{C \cdot x}$ of the C -orbit of x meets the open orbit of B in E_0 .

The choice of an adapted Levi subgroup L together with a maximal torus $T \subset L$ allows us to identify \mathcal{M} with an explicit subgroup of $\mathfrak{X}(T)$ [BP87, 2.9]. More precisely, the group \mathcal{M} is identified with $\mathfrak{X}(T/T \cap H)$, the group \mathcal{N} is identified with the group $\mathfrak{Y}(T/T \cap H)$ of one parameter subgroups of $T/T \cap H$, and so \mathcal{V} is identified with a cone in the \mathbb{Q} -vector space $\mathfrak{Y}(T/T \cap H) \otimes \mathbb{Q}$. Denote by π the quotient map $\mathfrak{Y}(T) \otimes \mathbb{Q} \rightarrow \mathfrak{Y}(T/T \cap H) \otimes \mathbb{Q}$.

Definition 3.18. Let (E, x) be an elementary embedding of G/H , and \mathcal{C}_E be the ray in \mathcal{V} associated to E . We say that a one parameter subgroup $\lambda \in \mathfrak{Y}(T)$ is *adapted to E* if it projects to an element of $\mathcal{C}_E \cap \mathfrak{Y}(T/T \cap H)$ under the quotient map π .

Proposition 3.19. [BP87, 2.10] *Let (E, x) be an elementary embedding of G/H , and $\lambda \in \mathfrak{Y}(T)$ adapted to E , then $\lim_{z \rightarrow 0} \lambda(z) \cdot x$ exists and is a point in the open orbit of B in E_0 .*

3.3.2. *Choice of an adapted Levi subgroup.* We will need to use some properties of the adapted Levi subgroups, essentially proved in [BLV86, BP87]. However, since they are sometimes proved for an adapted Levi subgroup of a particular form, we need to show that we can choose a good Levi subgroup, up to changing the base point. Let us first prove the following elementary fact.

Proposition 3.20. *Let L be a Levi subgroup adapted to H , and $u \in P^u$ the unipotent radical of P , then uLu^{-1} is a Levi subgroup of P adapted to uHu^{-1} .*

Remark that all Levi subgroups of P are conjugate under an element of P^u . Remark also that if H is the stabilizer of x , then uHu^{-1} is the stabilizer of the point $u \cdot x$, which is still in the open B -orbit by definition of P .

Proof. We can first see that

$$P \cap uHu^{-1} = u(P \cap H)u^{-1} = u(L \cap H)u^{-1} = uLu^{-1} \cap uHu^{-1},$$

and

$$[uLu^{-1}, uLu^{-1}] = u[L, L]u^{-1} \subset uHu^{-1}.$$

Assume now that (E, x) is an elementary embedding of G/uHu^{-1} with closed orbit E_0 . Then $(E, u^{-1} \cdot x)$ is an elementary embedding of G/H with closed orbit E_0 . Since L is adapted to H , if C is the connected center of L , $\overline{C \cdot u^{-1} \cdot x}$ meets the open orbit of B in E_0 . The connected center of uLu^{-1} is uCu^{-1} , and we get $\overline{uCu^{-1} \cdot x} = u \cdot \overline{C \cdot u^{-1} \cdot x}$. This meets the open orbit of B in E_0 since $u \in P^u \subset U$, so uLu^{-1} is adapted to uHu^{-1} . \square

Proposition 3.21. *Let T be a maximal torus of B . Then up to changing the base point in the open orbit of B and its stabilizer H , we can choose an adapted Levi subgroup L containing the torus T and such that $N_G(H) = H(C \cap N_G(H))$.*

Proof. First choose any base point \tilde{x} in the open orbit of B , and let \tilde{H} be its stabilizer. Let $\tilde{L} = \text{Stab}_G(df_e)$ where $f \in \mathbb{C}[G]$ is a regular function on G which vanishes everywhere on $G \setminus BH$ and df_e is the differential of f at the neutral element e , considered as an element of the coadjoint representation. Let also \tilde{C} denote the connected center of \tilde{L} . Then \tilde{L} is adapted to \tilde{H} [BLV86, Section 3].

Furthermore, it is shown in [BP87, Section 5] that $B\tilde{H} = BN_G(\tilde{H})$, so $N_G(\tilde{H})$ is spherical and P and \tilde{L} are also adapted to $N_G(\tilde{H})$. In particular, we have $P \cap N_G(\tilde{H}) = \tilde{L} \cap N_G(\tilde{H})$.

The Levi subgroup \tilde{L} might not contain the maximal torus T , but there is a conjugate L under an element u of P^u that does contain T . By Proposition 3.20, $L = u\tilde{L}u^{-1}$ is adapted to the subgroup $H = u\tilde{H}u^{-1}$.

Furthermore, we have $P \cap N_G(H) = L \cap N_G(H)$. From this we deduce that the inclusion of CH in $CN_G(H)$ is an equality. Indeed, $P^u \cap N_G(H) \subset P \cap N_G(H) = L \cap N_G(H)$, and $L \cap P^u = \{e\}$, so the following surjective maps are isomorphisms.

$$P^u \times CN_G(H) \longrightarrow PN_G(H) = BH = PH \longleftarrow P^u \times CH.$$

These isomorphisms imply that $CH = CN_G(H)$, and this equality implies the last conclusion: $N_G(H) = H(C \cap N_G(H))$. \square

3.3.3. Equivariant degenerations. We fix now a spherical homogeneous space O under the action of G , a Borel subgroup B of G and a maximal torus T in B . Using Proposition 3.21, we choose a base point $x \in O$ with isotropy group H so that there exists a Levi subgroup L of P adapted to H and containing the fixed maximal torus T of B . Note that $G/H \times \mathbb{C}^*$ is a spherical homogeneous space under the action of $G \times \mathbb{C}^*$.

We use the work of Brion and Pauer on isotropy subgroups of elementary embeddings [BP87, section 3] to obtain information about the equivariant degenerations of a spherical homogeneous space. In this article, equivariant degenerations of spherical homogeneous spaces are defined as follows.

Definition 3.22. An *equivariant degeneration* of G/H is an elementary embedding (E, \hat{x}) of $G/H \times \mathbb{C}^*$, equipped with a surjective $G \times \mathbb{C}^*$ -equivariant morphism $p: E \longrightarrow \mathbb{C}$, where G acts trivially on \mathbb{C} , and \mathbb{C}^* acts by multiplication on \mathbb{C} , with $p^{-1}(0) = E_0$ the closed orbit of E .

Let us remark that $P \times \mathbb{C}^*$ is an adapted parabolic for $G \times \mathbb{C}^*/H \times \{1\}$, and $L \times \mathbb{C}^*$ is an adapted Levi subgroup with connected center $C \times \mathbb{C}^*$. We identify

$\mathfrak{Y}(T \times \mathbb{C}^*)$ with $\mathfrak{Y}(T) \oplus \mathbb{Z}$. Furthermore, the valuation cone of $G \times \mathbb{C}^*/H \times \{1\}$ can be identified with $\mathcal{V} \times \mathbb{Q} \subset (\mathcal{N} \otimes \mathbb{Q}) \oplus \mathbb{Q}$.

Proposition 3.23. *Let (E, \tilde{x}) be an equivariant degeneration of G/H . Let $(\lambda, m) \in \mathfrak{Y}(T) \oplus \mathbb{Z}$ be a one parameter subgroup adapted to E . Then $m > 0$, the action of G on E_0 is transitive, and if H_0 denotes the isotropy subgroup in G of $\tilde{x}_0 = \lim_{z \rightarrow 0} (\lambda(z), z^m) \cdot \tilde{x}$, then the action of $e^\tau \in \mathbb{C}^*$ on G/H_0 is given by multiplication on the right by $\lambda(e^{-\tau/m})$.*

Proof. Let $p : E \rightarrow \mathbb{C}$ denote the $G \times \mathbb{C}^*$ -equivariant morphism associated to the degeneration. Then since $\tilde{x}_0 \in E_0 = p^{-1}(0)$, we have $m > 0$. Denote by \tilde{H}_0 the isotropy subgroup of \tilde{x}_0 in $G \times \mathbb{C}^*$. Obviously, $(\lambda(z), z^m) \in \tilde{H}_0$ for all $z \in \mathbb{C}^*$.

Let us first show that G acts transitively on $G \times \mathbb{C}^*/\tilde{H}_0$. Consider (g_1, z_1) and (g_2, z_2) in $G \times \mathbb{C}^*$. Let $s \in \mathbb{C}^*$ be such that $s^m = z_2/z_1$. Then

$$(g_1, z_1) = (g_1, z_2/s^m) = (g_1 \lambda(s), z_2)(\lambda(1/s), 1/s^m),$$

so since $(\lambda(1/s), 1/s^m) \in \tilde{H}_0$,

$$(g_1, z_1)\tilde{H}_0 = (g_1 \lambda(s)g_2^{-1}, 1)(g_2, z_2)\tilde{H}_0,$$

which shows the transitivity of the action of G . In particular if H_0 is the isotropy group of \tilde{x}_0 in G , we can identify E_0 with G/H_0 .

The action of \mathbb{C}^* is obtained similarly. Given $z = e^\tau \in \mathbb{C}^*$, we have

$$\begin{aligned} (e, z)(g, 1) \cdot \tilde{x}_0 &= (e, z)(g, 1)\tilde{H}_0 \\ &= (g, e^\tau)\tilde{H}_0 \\ &= (g\lambda(e^{-\tau/m}), 1)(\lambda(e^{\tau/m}), e^\tau)\tilde{H}_0 \\ &= (g\lambda(e^{-\tau/m}), 1)\tilde{H}_0 \\ &= (g\lambda(e^{-\tau/m}), 1) \cdot \tilde{x}_0. \end{aligned}$$

This finishes the proof of the proposition. \square

Conversely, any elementary embedding whose ray is generated by some $(\lambda, m) \in \mathfrak{Y}(T/T \cap H) \otimes \mathbb{Q} \times \mathbb{Q}$ with $m > 0$ provides an equivariant degeneration of G/H thanks to Theorem 3.11.

Proposition 3.24. *Keeping the same notations as in Proposition 3.23, we have: BH_0 is open in G (in particular H_0 is spherical), $P = \text{Stab}_G(BH_0)$, and L is adapted to H_0 .*

Proof. By [BP87, Théorème 3.6], $(B \times \mathbb{C}^*)\tilde{H}_0$ is open in $G \times \mathbb{C}^*$, $P \times \mathbb{C}^* = \text{Stab}_{G \times \mathbb{C}^*}((B \times \mathbb{C}^*)\tilde{H}_0)$, and $L \times \mathbb{C}^*$ is adapted to \tilde{H}_0 , where \tilde{H}_0 denotes the isotropy group of \tilde{x}_0 in $G \times \mathbb{C}^*$ as in the proof of Proposition 3.23.

First remark that for any $b \in B$ and $z \in \mathbb{C}^*$, we have $(b, z^m) \cdot \tilde{x}_0 = (b\lambda(1/z), 1) \cdot \tilde{x}_0$. Since λ is a one parameter subgroup of $T \subset B$, we obtain that $B \times \mathbb{C}^* \cdot \tilde{x}_0 = B \times \{1\} \cdot \tilde{x}_0$, thus the orbit of $\tilde{x}_0 = eH_0$ in G/H_0 is open, which implies that BH_0 is open in G .

Now let us show that $\text{Stab}_G(BH_0) = P$. We have proved above that $B \cdot \tilde{x}_0 = (B \times \mathbb{C}^*) \cdot \tilde{x}_0$, and furthermore we have $\{e\} \times \mathbb{C}^* \subset \text{Stab}_{G \times \mathbb{C}^*}((B \times \mathbb{C}^*) \cdot \tilde{x}_0)$, so it

follows that

$$\begin{aligned} P \times \mathbb{C}^* &= \text{Stab}_{G \times \mathbb{C}^*}((B \times \mathbb{C}^*) \cdot \tilde{x}_0) \\ &= \text{Stab}_G((B \times \mathbb{C}^*) \cdot \tilde{x}_0) \times \mathbb{C}^* \\ &= \text{Stab}_G(B \cdot \tilde{x}_0) \times \mathbb{C}^* \end{aligned}$$

Hence $P = \text{Stab}_G(B \cdot \tilde{x}_0) = \text{Stab}_G(BH_0)$.

Finally we have to show that L is adapted to H_0 .

First let us describe the subgroup \tilde{H}_0 more explicitly in terms of H_0 , λ and m . By the description of the action of $G \times \mathbb{C}^*$ on G/H_0 we easily check that

$$\tilde{H}_0 = \bigcup_{\tau \in \mathbb{C}} H_0 \lambda(e^{\tau/m}) \times \{e^\tau\}.$$

Since $L \times \mathbb{C}^*$ is adapted to \tilde{H}_0 we obtain

$$\begin{aligned} (P \cap H_0) \times \{1\} &= ((P \times \mathbb{C}^*) \cap \tilde{H}_0) \cap (G \times \{1\}) \\ &= ((L \times \mathbb{C}^*) \cap \tilde{H}_0) \cap (G \times \{1\}) \\ &= (L \cap H_0) \times \{1\} \end{aligned}$$

hence $P \cap H_0 = L \cap H_0$. Similarly, $[L, L] \subset H_0$ follows from:

$$[L, L] \times \{1\} = [L \times \mathbb{C}^*, L \times \mathbb{C}^*] \subset \tilde{H}_0 \cap (G \times \{1\}) = H_0 \times \{1\}.$$

Consider (Z, z) an elementary G -embedding of $G/H_0 = G \times \mathbb{C}^*/\tilde{H}_0$, with closed orbit Z_0 . Since the action of \mathbb{C}^* commutes with the action of G on G/H_0 , this is also an elementary embedding for the action of $G \times \mathbb{C}^*$. Since $L \times \mathbb{C}^*$ is adapted to $G \times \mathbb{C}^*/\tilde{H}_0$, there exists a one parameter subgroup $t \mapsto (\mu(t), t^k)$ of $T \times \mathbb{C}^*$ such that $z_0 := \lim_{t \rightarrow 0} (\mu(t), t^k) \cdot z$ is in the open $B \times \mathbb{C}^*$ -orbit in Z_0 . This z_0 obviously lies in the closure of $C \cdot z$ in Z , since $(\mu(t), t^k) \cdot z = (\mu(t)\lambda(s^k), 1) \cdot z$ if $s^m = t$. As previously, $(b, t) \in B \times \mathbb{C}^*$ acts on z_0 as $(b, t) \cdot z_0 = (b, t)(\mu(1/s), 1/t) \cdot z_0 = (b\mu(1/s), 1) \cdot z_0$ where $s^k = t$, so $B \cdot z_0 = B \times \mathbb{C}^* \cdot z_0$ is open in Z_0 . We have thus shown that L is adapted to H_0 . \square

3.3.4. Elementary embeddings and equivariant automorphisms. Let (E, \tilde{x}) be an equivariant degeneration of G/H , and $(\lambda, m) \in \mathfrak{J}(T) \times \mathbb{Z}$ a one parameter subgroup adapted to E . Let $x_0 = \lim_{z \rightarrow 0} (\lambda(z), z^m) \cdot \tilde{x}$ and denote by H_0 the isotropy subgroup of x_0 in G .

Proposition 3.25. *We have $T \cap H \subset T \cap H_0$ and $T \cap N_G(H) \subset T \cap N_G(H_0)$.*

Proof. It $t \in T \cap H$ then

$$\begin{aligned} t \cdot x_0 &= \lim_{z \rightarrow 0} (t, 1) \cdot (\lambda(z), z^m) \cdot \tilde{x} \\ &= \lim_{z \rightarrow 0} (\lambda(z), z^m) \cdot (t, 1) \cdot \tilde{x} \\ &= x_0. \end{aligned}$$

Recall from Section 3.1.3 that the action of

$$N_G(H)/H = \text{Aut}_G(G/H) \subset \text{Aut}_{G \times \mathbb{C}^*}(G/H \times \mathbb{C}^*)$$

extends to the elementary embedding E . It is *a priori* no longer explicit on G/H_0 . We denote by $y \curvearrowright nH$ the action of $nH \in N_G(H)/H$ on $y \in E$. We have $(n, 1) \cdot \tilde{x} \curvearrowright nH = \tilde{x}$.

Assume that $t \in T \cap N_G(H)$. We then have

$$\begin{aligned} (t, 1) \cdot x_0 &= \lim_{z \rightarrow 0} (t, 1) \cdot (\lambda(z), z^m) \cdot \tilde{x} \\ &= \lim_{z \rightarrow 0} (\lambda(z), z^m) \cdot (t, 1) \cdot \tilde{x} \\ &= \lim_{z \rightarrow 0} (\lambda(z), z^m) \tilde{x} \frown t^{-1}H \\ &= x_0 \frown t^{-1}H. \end{aligned}$$

Since the action of $N_G(H)/H$ commutes with the action of G , the isotropy group of $x_0 \frown t^{-1}H$ in G is the same as the isotropy group of x_0 , which is H_0 . On the other hand, the isotropy subgroup of $(t, 1) \cdot x_0$ in G is tH_0t^{-1} , so we obtain that $H_0 = tH_0t^{-1}$. In other words, $t \in N_G(H_0)$. \square

Let us also highlight the relation between the linear part of the valuation cone, equivariant automorphisms, and equivariant degenerations of G/H whose closed $G \times \mathbb{C}^*$ -orbit G/H_0 is isomorphic to G/H . We assume here that H and L are as in Proposition 3.21.

Assume $(\lambda, m) \in \mathfrak{Y}(T \cap N_G(H)) \cap \pi^{-1}(\mathcal{V}) \oplus \mathbb{N}^*$. In this case, the equivariant degeneration of G/H associated to the ray generated by (λ, m) may be described explicitly [BP87, Section 2.8]. This is the quotient of $G/H \times \mathbb{C}^* \times \mathbb{C}$ by the action of \mathbb{C}^* given by $t \cdot (gH, z, \theta) = (g\lambda(1/t)H, z/t^m, t\theta)$. The open $G \times \mathbb{C}^*$ -orbit is the image of the $G \times \mathbb{C}^* \times \mathbb{C}^*$ -orbit of $(eH, 1, 1)$, isomorphic to $G/H \times \mathbb{C}^*$, and the closed $G \times \mathbb{C}^*$ -orbit is the image of the $G \times \mathbb{C}^* \times \mathbb{C}^*$ -orbit of $(eH, 1, 0)$, whose stabilizer in $G \times \mathbb{C}^*$ is $(\lambda(z), z^m)H \times \{1\}$, and stabilizer in G is H . In particular, the closed orbit is in this case isomorphic to G/H .

Remark that the construction above may be carried out whenever $\lambda \in \mathfrak{Y}(T \cap N_G(H))$ and $m \in \mathbb{N}^*$, which shows that $\mathfrak{Y}(T \cap N_G(H)) \otimes \mathbb{Q}$ is a \mathbb{Q} -vector space contained in $\pi^{-1}(\mathcal{V})$. In fact, it is the maximal such vector space, also called the linear part of $\pi^{-1}(\mathcal{V})$, by [BP87, Proposition 5.3]. Let us give a statement summarizing this paragraph for future reference.

Proposition 3.26. *Let $\lambda \in \mathfrak{Y}(T) \cap \pi^{-1}(\mathcal{V})$, and $m \in \mathbb{N}^*$. Then the following are equivalent:*

- $-\lambda \in \mathfrak{Y}(T) \cap \pi^{-1}(\mathcal{V})$, which means that λ is in the linear part of \mathcal{V} ,
- $\lambda \in \mathfrak{Y}(T \cap N_G(H))$,
- the equivariant degeneration associated to the ray generated by (λ, m) has closed orbit isomorphic to G/H .

Recall the characterization of horospherical varieties in terms of their valuation cones. We then have in particular:

Corollary 3.27. *If G/H is horospherical then all equivariant degenerations of G/H have closed orbit isomorphic to G/H .*

3.3.5. *Horospherical degenerations.* Let Q denote the parabolic subgroup of G opposite to P with respect to the Levi subgroup L containing the maximal torus T .

Proposition 3.28. *We still use the notations from Proposition 3.23. Assume that λ is in the interior of the valuation cone, then H_0 is horospherical for G , and $H_0 = Q^u(L \cap H_0)$.*

Proof. This is essentially [BP87, Proposition 3.10] except we consider only the action of G and not the full action of $G \times \mathbb{C}^*$. But it follows easily from this case. Indeed, if \tilde{H}_0 contains a maximal unipotent subgroup $U \times \{1\}$ of $G \times \mathbb{C}^*$, then H_0 obviously contains U , so it is horospherical. Furthermore, we have $\tilde{H}_0 = (Q^u \times \{1\})(L \times \mathbb{C}^* \cap \tilde{H}_0)$ and taking the intersection with $G \times \{1\}$ yields $H_0 = Q^u(L \cap H_0)$. \square

3.4. Equivariant test configurations for spherical varieties. We will in this section apply the general theory of spherical varieties to construct special equivariant test configurations for \mathbb{Q} -Fano spherical varieties. Combined with the description of the action of \mathbb{C}^* on equivariant degenerations obtained in Proposition 3.23, we will have enough information about spherical test configurations for the proof of our main result.

3.4.1. Equivariant test configurations. Let us first recall the definition of special test configurations, introduced by Tian [Tia97].

Definition 3.29. Let (X, L) be a polarized normal projective variety.

A *special test configuration* for (X, L) is a normal variety \mathcal{X} with an action of \mathbb{C}^* , equipped with a \mathbb{C}^* -linearized line bundle \mathcal{L} and a \mathbb{C}^* -equivariant flat morphism $\pi : \mathcal{X} \rightarrow \mathbb{C}$ such that \mathcal{L} is π -ample, the fiber $(\pi^{-1}(1), \mathcal{L}|_{\pi^{-1}(1)})$ is isomorphic to (X, L^r) for some fixed integer $r > 0$, and $\pi^{-1}(0)$ is a *normal* variety.

If a reductive group G acts on (X, L) , we say that a special test configuration is *G -equivariant* if $(\mathcal{X}, \mathcal{L})$ admits an action of G which commutes with the \mathbb{C}^* action and such that the isomorphism between (X, L^r) and $(\pi^{-1}(1), \mathcal{L}|_{\pi^{-1}(1)})$ is G -equivariant.

The definition extends *verbatim* to normal varieties equipped with an ample \mathbb{Q} -line bundle. In the rest of the paper, we always consider \mathbb{Q} -Fano varieties, and the ample \mathbb{Q} -line bundle L will always be the anticanonical \mathbb{Q} -line bundle K_X^{-1} , so we will omit it in the notations. Furthermore, the induced \mathbb{Q} -line bundle L_0 on the central fiber X_0 is the anticanonical line bundle $K_{X_0}^{-1}$, so the central fiber X_0 is also a \mathbb{Q} -Fano variety (see for example [Ber16, Lemma 2.2]).

Let (X, L) be a polarized spherical variety under G . Then if $(\mathcal{X}, \mathcal{L})$ is a special G -equivariant test configuration, the normal variety \mathcal{X} is a spherical variety under the action of $G \times \mathbb{C}^*$. More precisely, if (X, x) is a spherical embedding of G/H , then (\mathcal{X}, \tilde{x}) is a spherical embedding of $G \times \mathbb{C}^*/H \times \{1\}$, where $\tilde{x} \in \pi^{-1}(1)$ corresponds to x in the isomorphism between $\pi^{-1}(1)$ and X .

The central fiber $\pi^{-1}(0)$ is a normal $G \times \mathbb{C}^*$ -stable subvariety of the $G \times \mathbb{C}^*$ -spherical variety \mathcal{X} , hence it is also a $G \times \mathbb{C}^*$ -spherical variety. In particular, if we consider only the union of the open $G \times \mathbb{C}^*$ -orbits in \mathcal{X} and in $\pi^{-1}(0)$, we obtain an elementary embedding of $G \times \mathbb{C}^*/H \times \{1\}$, which can be studied using Section 3.3.

Finally remark that since the action of $\text{Aut}_{G \times \mathbb{C}^*}(G/H \times \mathbb{C}^*)$ extends to any $G/H \times \mathbb{C}^*$ -embedding by Section 3.1.3, any G -equivariant test configuration for a spherical embedding X of G/H is also $G \times \text{Aut}_G(X)$ -equivariant.

3.4.2. Construction.

Theorem 3.30. *Let X be a \mathbb{Q} -Fano spherical embedding of G/H . Let $m \in \mathbb{N}^*$, and let $\lambda \in \mathfrak{J}(T) \cap \pi^{-1}(\mathcal{V})$. Then there exists a G -equivariant test configuration $(\mathcal{X}, \mathcal{L})$ for X , with central fiber X_0 , such that X_0 is a spherical embedding of G/H_0 and the action of $e^\tau \in \mathbb{C}^*$ on G/H_0 is given by $e^\tau \cdot gH_0 = g\lambda(e^{-\tau/m})H_0$.*

Proof. We may assume that (λ, m) is primitive. The construction uses results of [GH15] as recalled in Section 3.2.4. Let d be the \mathbb{Q} -Cartier divisor representing K_X^{-1} constructed there, and n_D the coefficients of the colors. Let Δ_d^* be the \mathbb{Q} - G/H -reflexive polytope associated to X , considered in $(\mathcal{N} \oplus \mathbb{Z}) \otimes \mathbb{R}$.

We build from this polytope the colored fan $\mathcal{F}_{\mathcal{X}}$ of \mathcal{X} . Remark that the colors of G/H are in bijection with the colors of $G/H \times \mathbb{C}^*$ by sending $D \in \mathcal{D}$ to $D \times \mathbb{C}^*$. For every face F of Δ_d^* , consider the three colored cones

$$\begin{aligned} & (\text{Cone}(F), \rho^{-1}(\text{Cone}(F))), \\ & (\text{Cone}(F \cup \{(0, -1)\}), \rho^{-1}(\text{Cone}(F \cup \{(0, -1)\}))) \\ & (\text{Cone}(F \cup \{(\lambda, m)\}), \rho^{-1}(\text{Cone}(F \cup \{(\lambda, m)\}))). \end{aligned}$$

Among these cones, keep only the cones such that the intersection of the relative interior of their support with $\mathcal{V} \times \mathbb{Q}$ is non empty. Then $\mathcal{F}_{\mathcal{X}}$ is the set of these cones.

It is clear that the corresponding spherical embedding of $G/H \times \mathbb{C}^*$ is complete. Furthermore, by the description of equivariant morphisms between spherical varieties, it admits a $G \times \mathbb{C}^*$ -equivariant inclusion of $X \times \mathbb{C}^*$, and a $G \times \mathbb{C}^*$ -equivariant surjective morphism p to the spherical embedding \mathbb{P}^1 of the $G \times \mathbb{C}^*$ homogeneous space $G \times \mathbb{C}^*/G \times \{1\} \simeq \mathbb{C}^*$, such that $p^{-1}(\mathbb{C}^*) = X \times \mathbb{C}^*$.

We then build the line bundle. We chose to build \mathcal{X} complete, and will choose a line bundle \mathcal{L} that is ample. Denote by $X_\infty = p^{-1}(\infty)$, respectively $X_0 = p^{-1}(0)$, the fibers of p , corresponding as irreducible $G \times \mathbb{C}^*$ -invariant divisors of \mathcal{X} to the rays generated by $(0, -1)$, respectively (λ, m) . Remark that these are spherical varieties, and that they are the closures in \mathcal{X} of equivariant degenerations of G/H . Consider the $B \times \mathbb{C}^*$ -invariant Weil divisor

$$\delta = a \left(\sum_{Y \in W_G} \overline{Y \times \mathbb{C}^*} + \sum_{D \in \mathcal{D}} n_D \overline{D \times \mathbb{C}^*} \right) + b(X_\infty + X_0)$$

where $a, b \in \mathbb{N}$.

Let v denote the support function of the dilated polytope $a\Delta_d = \Delta_{ad}$ and consider the function l_δ defined for $(x, n) \in \mathcal{N} \otimes \mathbb{Q} \oplus \mathbb{Q}$ by

- $l_\delta(x, n) = v(-x + n/m\lambda) + bn/m$ if $n \geq 0$ and
- $l_\delta(x, n) = v(x) - bn$ if $n \leq 0$.

This is clearly a piecewise linear function on the support of the fan $\mathcal{F}_{\mathcal{X}}$. It furthermore satisfies the three conditions of Proposition 3.13, provided $b > 0$ is large enough. It is however not integral in general. By definition, it satisfies $l_\delta(0, -1) = b$ and $l_\delta(\lambda, m) = b$. By the relation between l_d and the support function of the polytope Δ_d recalled in Section 3.2.2, we thus have

$$\begin{aligned} \delta &= \sum_{Y \in \mathcal{I}_{\mathcal{X}}^c} l_\delta(u_Y, 0) \overline{Y \times \mathbb{C}^*} + \sum_{D \in \mathcal{D}_X} l_\delta(\rho(D), 0) \overline{D \times \mathbb{C}^*} \\ &+ \sum_{D \in \mathcal{D} \setminus \mathcal{D}_X} a n_D \overline{D \times \mathbb{C}^*} + l_\delta(0, -1) X_\infty + l_\delta(\lambda, m) X_0. \end{aligned}$$

Let us now check that l_δ is integral piecewise linear for a and b divisible enough. Denote by $m_F \in \mathcal{M} \otimes \mathbb{Q}$ the vertex of Δ_d corresponding to the maximal face F of Δ_d^* . Any maximal cone of $\mathcal{F}_{\mathcal{X}}$ is either the cone over the union of some maximal face F of Δ_d^* and $(0, -1)$, in which case l_δ is given by the linear function $(-am_F, -b)$ on this cone, or it is the union of some maximal face F of Δ_d^* and (λ, m) , in which

case l_δ is given by the linear function $(-am_F, -am_F(\lambda)/m + b/m)$ on this cone. Up to choosing a and b divisible enough, we can thus assume that l_δ is given by an element of $\mathcal{M} \oplus \mathbb{Z}$ on all maximal cones.

This finally shows that δ is an ample Cartier divisor. The corresponding ample line bundle \mathcal{L} obviously restricts to $p_1^* K_X^{-a}$ on $X \times \mathbb{C}^*$, where p_1 is the first projection. It remains to show that it may be equipped with a $G \times \mathbb{C}^*$ -linearization such that its restriction to $X \times \mathbb{C}^*$ respects the natural linearization of $p_1^* K_X^{-a}$. Since X is normal, as a spherical variety, a tensor power \mathcal{L}^n of \mathcal{L} admits a $G \times \mathbb{C}^*$ -linearization [KKV89]. Furthermore, two such linearizations differ by a character of $G \times \mathbb{C}^*$. As a consequence, we may change the linearization of \mathcal{L}^n to obtain the one we want.

The test configuration is obtained by considering $p^{-1}(\mathbb{C}) = X \times \mathbb{C}^* \cup X_0$, for $\mathbb{C} = \mathbb{C}^* \cup \{0\} \in \mathbb{P}^1$, equipped with the map p and the restriction of \mathcal{L} . If we consider the spherical embedding of $G/H \times \mathbb{C}^*$ given by the ray generated by (λ, m) , we obtain an equivariant degeneration of G/H , and the action of \mathbb{C}^* on the closed orbit G/H_0 , which is the open orbit of G in the central fiber X_0 , is as expected by Proposition 3.23. \square

Taking any λ projecting to the interior of the valuation cone yields, thanks to Proposition 3.28,

Corollary 3.31. *Any \mathbb{Q} -Fano spherical variety admits a test configuration with horospherical central fiber.*

4. MODIFIED FUTAKI INVARIANT ON \mathbb{Q} -FANO HOROSPHERICAL VARIETIES

4.1. Modified Futaki invariant on singular varieties. Let X be a normal \mathbb{Q} -Fano variety. Let A be an ample line bundle on X such that there exists $m \in \mathbb{N}^*$ such that the restriction of A to the regular part X_{reg} of X is $K_{X_{\text{reg}}}^{-m}$. The cohomology class $c_1(X)$ is the class $c_1(A)/m$. We will always assume that X has log terminal singularities. This is harmless in our setting since \mathbb{Q} -Fano (horo)spherical varieties have log terminal singularities [AB04b, Section 5.1]. See [Pas] for a survey on singularities of spherical varieties.

Let $\text{Aut}^0(X)$ denote the connected component of the neutral element in the automorphism group of X . Since X is a \mathbb{Q} -Fano variety, this is a linear algebraic group. Choose a maximal compact subgroup K in $\text{Aut}^0(X)$. Its complexification G is a Levi subgroup of $\text{Aut}^0(X)$. Choose a maximal torus T in G such that $K \cap T$ is the maximal compact torus of T , and denote by \mathfrak{a} the subspace $i\mathfrak{k} \cap \mathfrak{t}$ in the Lie algebra of G . Any element $\xi \in \mathfrak{a}$ is such that $J\xi \in \mathfrak{k}$ generates a compact torus in K .

Let q_A be a *smooth, positive* K -invariant hermitian metric on A . On a normal variety, this means that the potentials of q_A with respect to local trivializations of A are the restrictions of smooth and strictly plurisubharmonic functions to X , given a local embedding of X in \mathbb{C}^N . Let $\omega_A \in 2\pi c_1(A)$ denote the curvature current of q_A , and $\omega = \omega_A/m \in 2\pi c_1(X)$. The current ω is still K -invariant, and defines a Kähler form on the regular part X_{reg} of X .

Remark that on X_{reg} , the anticanonical line bundle K_X^{-1} is a well defined line bundle, and the metric q_A induces a smooth positive metric q on $K_{X_{\text{reg}}}^{-1}$. The form ω is the curvature form of q on X_{reg} . Let dV_q denote the volume form on X_{reg} associated to q , defined by

$$dV_q(x) = |s^{-1}|_q^2 s \wedge \bar{s}$$

where s is any non zero element of the fiber over x of the canonical line bundle of X_{reg} , and s^{-1} is the corresponding non zero element of the dual line bundle $K_{X_{\text{reg}}}^{-1}$.

We call *Hamiltonian function* of $\xi \in \mathfrak{a}$ with respect to ω the smooth function θ_ξ on X_{reg} defined by $L_\xi \omega = i\partial\bar{\partial}\theta_\xi$, where L_ξ is the Lie derivative with respect to ξ . It is shown to exist and to be bounded in [BW]. This function is well defined up to a constant and we fix this constant by the normalization condition :

$$\int_{X_{\text{reg}}} \theta_\xi dV_q = 0.$$

Definition 4.1. The *modified Futaki invariant* of X with respect to $\zeta \in \mathfrak{a}$ is defined for $\xi \in \mathfrak{a}$ by

$$\text{Fut}_{X,\zeta}(\xi) = - \int_{X_{\text{reg}}} \theta_\xi e^{\theta_\zeta} \omega^n.$$

It is proved in [BW] that it is well defined and independent of the choice of the metric q_A .

If we want to consider the K-stability of X with respect to G -equivariant test configurations (for any reductive subgroup G of $\text{Aut}^0(X)$), the only ξ at which we will need to evaluate the Futaki invariant generate automorphisms commuting with G . Recall that we described these G -equivariant automorphisms when X is spherical under the action of G in Section 3.1.3.

Assume that X is horospherical under G , and that $x \in X$ is such that its isotropy subgroup H contains the unipotent radical U of a Borel subgroup B . Then the group of equivariant automorphisms of X is isomorphic to P/H , where we recall that $P = N_G(H)$ is a parabolic containing B , and the action of P/H on the open orbit $G \cdot x = G/H$ is by multiplication on the right by the inverse. Recall that P/H is a torus. Its maximal compact subtorus is thus well defined and we denote by

$$\mathfrak{b}_1 = \mathfrak{J}(P/H) \otimes \mathbb{R}$$

the subalgebra of the Lie algebra of P/H obtained as J times the Lie algebra of the maximal compact subtorus, where J denotes the complex structure on P/H . By Proposition 2.5, we can identify \mathfrak{b}_1 with \mathfrak{a}_1 . We keep a different notation to emphasize that an element of \mathfrak{b}_1 acts on the right and not on the left.

4.2. Hermitian metrics on polarized horospherical varieties. Let X be a projective embedding of a horospherical homogeneous space G/H , equipped with a $G \times P/H$ -linearized ample line bundle L , where P is the normalizer of H in G .

Denote by χ the character of the isotropy subgroup $\text{diag}(P) \subset G \times P/H$ associated to L . Let Δ^+ be the moment polytope associated to the G -linearized line bundle L with respect to B . Consider the translated polytope $\Delta = \chi + \Delta^+$, and let $v_{2\Delta}$ denote the support function of the dilated polytope 2Δ .

Proposition 4.2. *Let q be a K -invariant locally bounded metric on L , and let $u : \mathfrak{a}_1 \rightarrow \mathbb{R}$ be the function associated to its restriction to G/H . Assume that q is smooth and positive over G/H . Then u is a smooth and strictly convex function, and the function $u - v_{2\Delta}$ is bounded on \mathfrak{a}_1 .*

Proof. The fact that u is smooth and strictly convex is obvious thanks to the expression of the curvature of the restriction of q over G/H obtained in Theorem 2.8.

Let us now consider the *decoloration* \tilde{X} of X . This is the spherical embedding of G/H corresponding to the fan $\mathcal{F}_{\tilde{X}}$ obtained as the union of the colored cones (\mathcal{C}, \emptyset) ,

for all colored cones $(\mathcal{C}, \mathcal{R}) \in \mathcal{F}_X$. It is a complete toroidal embedding of G/H which admits a natural $G \times P/H$ -equivariant surjective morphism $\pi : \tilde{X} \rightarrow X$, thanks to the description of spherical morphisms. The map π is further an isomorphism between the open orbits, both identified with G/H .

Consider the pull back $\tilde{L} := \pi^*L$ of L on \tilde{X} , with the $G \times P/H$ -linearization induced by the linearization of L . Its restriction to G/H corresponds to the same character χ of P . Let s be a $B \times P/H$ -semi-equivariant section of L , let μ denote the character of B associated to s , and let d be the B -invariant Cartier divisor defined by s .

Consider the pulled back section \tilde{s} of \tilde{L} . It is still $B \times P/H$ -semi-invariant with the same B -character μ , and its divisor \tilde{d} is the pull back of the Cartier divisor d defined by s . The function $l_{\tilde{d}}$ associated to \tilde{d} coincide with the function l_d (see [Pas, Proof of Lemma 5.3]). On a *toroidal* horospherical manifold, the function $l_{\tilde{d}}$ coincides with the function associated to the restriction of the Cartier divisor \tilde{d} to the toric subvariety $\tilde{Z} := \overline{P/H} \subset \tilde{X}$ under the action of the torus P/H , whose fan is precisely the fan underlying the colored fan $\mathcal{F}_{\tilde{X}}$ [Bri89, Section 3.2].

Since L is ample, the polytope Δ^+ is equal to $\mu + \Delta$, where Δ is the polytope with support function $x \mapsto l_d(-x)$, associated to the ample Cartier divisor d . On the other hand, since $l_{\tilde{d}} = l_d$, the moment polytope of $\tilde{L}|_{\tilde{Z}}$ is $\eta + \Delta$ where η is the character of T associated to the restriction of \tilde{s} to \tilde{Z} . Remark that even if \tilde{d} is not ample on \tilde{X} , its restriction to \tilde{Z} is. It is clear that $\eta = \mu$, so the moment polytope of $\tilde{L}|_{\tilde{Z}}$ is Δ^+ .

Consider now the metric q on L . The pulled back metric π^*q on \tilde{L} is still K -invariant and locally bounded. Furthermore, the associated function \tilde{u} on \mathfrak{a}_1 is still u since π is an equality on G/H . Recall that with the notations of Section 2,

$$\begin{aligned} u(x) &= -2 \ln |s_r(\exp(x)H)|_{\tilde{q}} \\ &= \langle 2\chi, x \rangle - 2 \ln |s_l(\exp(x)H)|_{\tilde{q}}. \end{aligned}$$

Consider the Batyrev-Tschinkel metric q_0 on $\tilde{L}|_{P/H}$ [Mai00, Section 3.3]. This is a continuous hermitian metric on $\tilde{L}|_{P/H}$, invariant under the action of the compact torus, with potential $u_0 : \mathfrak{a}_1 \rightarrow \mathbb{R}$ such that

$$u_0(x) = -2 \ln |s_l(\exp(x)H)|_{q_0} = v_{2\Delta^+}(x).$$

Then since \tilde{q} is locally bounded and q_0 is continuous, the function

$$u(x) - \langle 2\chi, x \rangle - v_{2\Delta^+}(x) = -2 \ln \frac{|s_l(\exp(x)H)|_{\tilde{q}}}{|s_l(\exp(x)H)|_{q_0}}$$

is bounded. Since we denoted $\chi + \Delta^+$ by Δ , we obtain the statement: $u - v_{2\Delta}$ is bounded. \square

Remark 4.3. The idea to use the decoloration of X was suggested to me by Boris Pasquier. If the closure of P/H in X is a normal toric variety, we can also deduce the moment polytope of the restriction of L to $\overline{P/H}$ from the proof.

4.3. Computation of the modified Futaki invariant. We fix G a connected reductive complex group, B a Borel subgroup of G , T a maximal torus of B , and K a maximal compact subgroup of G such that $K \cap T$ is a maximal compact torus of T . Let X be a horospherical \mathbb{Q} -Fano variety, and choose a base point in X such

that its isotropy group H contains the unipotent radical of B . Denote by P the normalizer of H in G .

Recall from Section 3.2.4 that the moment polytope of the ample \mathbb{Q} -line bundle K_X^{-1} is the polytope obtained as the dual of the \mathbb{Q} - G/H -reflexive polytope associated to X , translated by $2\rho_P$.

The weight associated to the anticanonical line bundle on G/H is obtained as the weight of the action of $\text{diag}(P)$ on the one dimensional representation $\det(\mathfrak{g}/\mathfrak{h})$. It follows that this character is precisely

$$-2\rho_P = \sum_{\alpha \in \Phi_P^+} -\alpha.$$

Let $m \in \mathbb{N}^*$ be such that $A = K_X^{-m}$ is an ample line bundle. Choose q_A a smooth positive K -invariant metric on A . Denote by $q = (q_A|_{G/H})^{1/m}$ the induced metric on $K_{G/H}^{-1}$. Let ω be the curvature form of q and u be the convex potential of q . Then $u - v_{2\Delta}$ is bounded, where $\Delta = \Delta^+ - 2\rho_P$ and Δ^+ is the moment polytope of the anticanonical \mathbb{Q} -line bundle.

We will compute the Futaki invariant for all $\xi \in \mathfrak{b}_1$ for X .

4.3.1. Computation of Hamiltonian functions.

Proposition 4.4. *Let $\xi \in \mathfrak{b}_1 \simeq \mathfrak{Y}(T/T \cap H) \otimes \mathbb{R}$ and θ_ξ be the Hamiltonian function of ξ with respect to ω . Let $\tilde{\xi} \in \mathfrak{Y}(T) \otimes \mathbb{R} \simeq \mathfrak{a}$ be any lift of ξ . Then θ_ξ is the K -invariant smooth function on X_{reg} defined for $x \in \mathfrak{a}_1$ by*

$$\theta_\xi(\exp(x)H) = - \left\{ \nabla u(x), \tilde{\xi} \right\}.$$

Proof. First remark that since the function θ_ξ is smooth on X_{reg} , and $G/H \subset X_{\text{reg}}$ is dense, it is enough to work on G/H . By K -invariance it will be enough to obtain $\theta_\xi(\exp(x)H)$ for $x \in \mathfrak{a}_1$.

We will work on G using the pullback by $\pi : G \rightarrow G/H$. We defined in Section 2 a function ϕ on G as the potential of π^*q with respect to a global left invariant section. This function is a global $i\partial\bar{\partial}$ -potential for $\pi^*\omega$ on G satisfying $\phi(k \exp(x)h) = u(x) - 2 \ln(\chi(\exp(x)h))$ for $k \in K$, $x \in \mathfrak{a}_1$ and $h \in H$.

Let $\tilde{\xi}$ be any lift of ξ , and denote by $L_{\tilde{\xi}}$ the Lie derivative with respect to the right- G -invariant vector field defined by $\tilde{\xi}$ on G . We have on the one hand

$$\begin{aligned} L_{\tilde{\xi}}\pi^*\omega &= L_{\tilde{\xi}}i\partial\bar{\partial}\phi \\ &= i\partial\bar{\partial}L_{\tilde{\xi}}\phi \end{aligned}$$

and on the other hand,

$$\begin{aligned} L_{\tilde{\xi}}\pi^*\omega &= \pi^*L_{\xi}\omega \\ &= \pi^*i\partial\bar{\partial}\theta_\xi \\ &= i\partial\bar{\partial}(\theta_\xi \circ \pi). \end{aligned}$$

Let $g = k \exp(x)h \in G$, where $k \in K$, $x \in \mathfrak{a}_1$ and $h \in H$. Recall that $\exp(t\xi)$ acts here by multiplication on the right by the inverse. We compute

$$\begin{aligned} (\exp(t\xi)^*\phi)(g) &= \phi(g \exp(-t\xi)) \\ &= \phi(k \exp(x - t\xi)h') \end{aligned}$$

where $h' \in H$ since P normalizes H

$$\begin{aligned} &= u(x - t\xi^1) - 2 \ln(\chi(\exp(x - t\xi)h')) \\ &= u(x - t\xi^1) - 2 \ln(\chi(\exp(x)h)) + 2 \langle \chi, t\xi \rangle, \end{aligned}$$

where ξ^1 is the \mathfrak{a}_1 component of $\xi \in \mathfrak{a} = \mathfrak{a}_0 \oplus \mathfrak{a}_1$.

Using this we obtain

$$\begin{aligned} (L_\xi \phi)(g) &= \left. \frac{d}{dt} \right|_0 (\exp(t\xi)^* \phi)(g) \\ &= - \{ \nabla u(x), \xi^1 \} + 2 \langle \chi, \xi \rangle \end{aligned}$$

In particular, this function is $K \times H$ invariant (where H acts by multiplication on the right), just as $\pi^* \theta_\xi$.

The difference between the $L_\xi \phi$ and $\pi^* \theta_\xi$ is thus a $K \times H$ invariant smooth function on G which is pluriharmonic on G . The only such functions are of the form $\psi(k \exp(x)h) = v(x)$ where v is an affine function on \mathfrak{a}_1 , but both $\pi^* \theta_\xi$ and $- \{ \nabla u(x), \xi^1 \}$ are bounded so the difference between the two is a constant.

Remark that $\nabla u(x) \in \mathfrak{a}_1$ is orthogonal to \mathfrak{a}_0 , so we can replace ξ^1 with ξ . It remains to check the normalization condition. We use the notation

$$\tilde{\theta}_\xi(\exp(x)H) = - \{ \nabla u(x), \xi \}.$$

Note that we can restrict to G/H since $X_{\text{reg}} \setminus (G/H)$ is of codimension at least one in X_{reg} . Recall that G/H is a fiber bundle with fiber P/H over the generalized flag manifold $G/P = K/(K \cap P)$. We use fiber integration and K -invariance to obtain that :

$$\int_{G/H} \tilde{\theta}_\xi dV_q = C \int_{P/H} \tilde{\theta}_\xi |s_r|_q^2 s_r^{-1} \wedge \overline{s_r}^{-1},$$

where C is a positive constant, and s_r is the right- P/H -invariant section of the restriction $K_{G/H}^{-1}|_{P/H}$. Using invariance under the action of the maximal compact subtorus of P/H on itself, we obtain that this is up to a positive constant, equal to

$$\int_{\mathfrak{a}_1} - \{ \nabla u(x), \xi \} e^{-u(x)} dx$$

and the last expression vanishes. Indeed, we have

$$- \{ \nabla u(x), \xi \} e^{-u(x)} = \left\{ \nabla e^{-u(x)}, \xi \right\}$$

and the Stokes formula yields the vanishing, because by boundedness of $u - v_{2\Delta}$, $e^{-u} \leq C e^{-v_{2\Delta}}$ and Δ contains 0 in its interior, so $e^{-v_{2\Delta}}$ has exponential decay. \square

4.3.2. Computation of the Futaki invariant.

Theorem 4.5. *Let X be a \mathbb{Q} -Fano horospherical embedding of G/H , let Δ^+ be the moment polytope of X , and let $\zeta, \xi \in \mathfrak{b}_1$. Let $\tilde{\zeta}$ and $\tilde{\xi}$ be lifts in \mathfrak{a} of ζ and ξ . Then*

$$\text{Fut}_{X,\zeta}(\xi) = C \left\langle \text{bar}_{DH,\tilde{\zeta}}(\Delta^+) - 2\rho_P, \tilde{\xi} \right\rangle$$

where $\text{bar}_{DH,\tilde{\zeta}}$ is the barycenter with respect to the measure

$$e^{-\langle 2p-4\rho_P, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi^+ \setminus \Phi_I^+} \kappa(\alpha, p) dp$$

with dp the Lebesgue measure, and C is a positive constant independent of ξ and depending on ζ only via the volume of Δ^+ with respect to the measure above.

Proof. We have to compute

$$\text{Fut}_{X,\zeta}(\xi) = - \int_{X_{\text{reg}}} \theta_\xi e^{\theta_\zeta} \omega^n.$$

Let us first use Theorem 2.8 to express the volume form ω^n on P/H . The theorem gives that, for $x \in \mathfrak{a}_1$,

$$\omega_{\text{exp}(x)H} = \sum_{1 \leq j_1, j_2 \leq r} \frac{1}{4} \frac{\partial^2 u}{\partial x_{j_1} \partial x_{j_2}}(x) \gamma_{j_1} \wedge \bar{\gamma}_{j_2} + \sum_{\alpha \in \Phi_P^+} \langle \alpha, \nabla u(x)/2 - t_\chi \rangle \gamma_\alpha \wedge \bar{\gamma}_\alpha.$$

We then obtain, recalling that $\chi = -2\rho_P$,

$$\omega_{\text{exp}(x)H}^n = \frac{\text{MA}_{\mathbb{R}}(u)(x)}{4^r 2^{\text{Card}(\Phi_P^+)}} \prod_{\alpha \in \Phi_P^+} \langle \alpha, \nabla u(x) + 4t_{\rho_P} \rangle \Omega$$

where

$$\Omega := \bigwedge_{1 \leq j \leq r} \gamma_j \wedge \bar{\gamma}_j \bigwedge_{\alpha \in \Phi_P^+} \gamma_\alpha \wedge \bar{\gamma}_\alpha.$$

We now use the same techniques as in the previous proof, that is, K -invariance, fiber integration on the fiber bundle $G/H \rightarrow G/P$, and integration of a compact torus-invariant function on a torus, to compute :

$$\begin{aligned} \text{Fut}_{X,\zeta}(\xi) &= - \int_{G/H} \theta_\xi e^{\theta_\zeta} \omega^n \\ &= C \int_{\mathfrak{a}_1} \left\{ \nabla u(x), \tilde{\xi} \right\} e^{-\langle \nabla u(x), \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \langle \alpha, \nabla u(x) + 4t_{\rho_P} \rangle \text{MA}_{\mathbb{R}}(u)(x) dx \end{aligned}$$

where $C > 0$ is a constant independent of ξ and ζ . Since u is smooth and strictly convex, we can use the change of variables $p = d_x u + 4\rho_P$, where $d_x u \in \mathfrak{a}^*$ is the derivative of u at x . Remark that this equivalently means $t_p = \nabla u(x) + 4t_{\rho_P}$, and that the domain of integration after this change of variables becomes $2\Delta + 4\rho_P$ by Proposition 4.2. Hence we obtain that

$$\text{Fut}_{X,\zeta}(\xi) = C \int_{2\Delta + 4\rho_P} \left\langle p - 4\rho_P, \tilde{\xi} \right\rangle e^{-\langle p - 4\rho_P, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \langle \alpha, p \rangle dp$$

By Section 3.2.4, the polytope $\Delta + 2\rho_P$ coincides with the moment polytope Δ^+ , so we obtain

$$\begin{aligned} \text{Fut}_{X,\zeta}(\xi) &= 2^{\text{Card}(\Phi_P^+)+1} C \int_{\Delta^+} \left\langle p - 2\rho_P, \tilde{\xi} \right\rangle e^{-\langle 2p - 4\rho_P, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \langle \alpha, p \rangle dp \\ &= 2^{\text{Card}(\Phi_P^+)+1} C \int_{\Delta^+} \left\langle p - 2\rho_P, \tilde{\xi} \right\rangle e^{-\langle 2p - 4\rho_P, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \kappa(\alpha, p) dp \end{aligned}$$

since α is in the semisimple part,

$$= C' \langle \text{bar}_{DH,\zeta}(\Delta^+) - 2\rho_I, \xi \rangle.$$

The final constant $C' > 0$ depends on $\tilde{\zeta}$ only at the last step, where we multiplied $2^{\text{Card}(\Phi_P^+)+1}C$ by the volume of Δ^+ with respect to the measure

$$e^{-\langle 2p-4\rho_P, \tilde{\zeta} \rangle} \prod_{\alpha \in \Phi_P^+} \kappa(\alpha, p) dp.$$

□

5. MODIFIED K-STABILITY CRITERION FOR \mathbb{Q} -FANO SPHERICAL VARIETIES

5.1. Statement of the criterion. We will prove in this section our main theorem. The proof combines the results from Section 3 and Section 4, together with an argument allowing us to compute the Futaki invariant of a \mathbb{Q} -Fano variety on an equivariant degeneration.

5.1.1. Definition of modified K-stability. Let X be a \mathbb{Q} -Fano variety. Given $\zeta \in \text{Lie}(\text{Aut}(X))$, let

$$T_\zeta = \{\exp(z\zeta); z \in \mathbb{C}\}.$$

It defines a subgroup of $\text{Aut}^0(X)$ which may very well not be closed. It is equipped with a privileged map $\mathbb{C} \rightarrow T_\zeta, z \mapsto \exp(z\zeta)$, and ζ may be recovered as the derivative at 1 of the restriction of this map to \mathbb{R} .

Definition 5.1. Let $\zeta \in \text{Lie}(\text{Aut}(X))$ be a semisimple element such that $J\zeta$ generates a compact real subgroup. Let $(\mathcal{X}, \mathcal{L})$ be a \overline{T}_ζ -equivariant special test configuration for X with central fiber X_0 . Denote by ζ_0 the element of $\text{Lie}(\text{Aut}(X_0))$ obtained as derivative at 1 of the map obtained by composition:

$$\mathbb{R} \subset \mathbb{C} \rightarrow T_\zeta \rightarrow \text{Aut}(X_0).$$

Denote by ξ the element of $\text{Lie}(\text{Aut}(X_0))$ generating the action of \mathbb{C}^* on X_0 induced by the test configuration. The *modified Donaldson-Futaki invariant* of the test configuration is

$$\text{DF}_\zeta(\mathcal{X}, \mathcal{L}) = \text{Fut}_{X_0, \zeta_0}(\xi).$$

We assume now that X is equipped with an action of a reductive group G , and fix $\zeta \in \text{Lie}(\text{Aut}(X))$ a semisimple element such that $J\zeta$ generates a compact real subgroup. We further assume that the action of G commutes with \overline{T}_ζ .

Definition 5.2. The \mathbb{Q} -Fano variety X is *modified K-semistable (with respect to G-equivariant special test configurations)* for the vector field ζ if for any $G \times \overline{T}_\zeta$ -equivariant special test configuration $(\mathcal{X}, \mathcal{L})$ for X , $\text{DF}_\zeta(\mathcal{X}, \mathcal{L}) \geq 0$. It is *K-stable (with respect to G-equivariant special test configurations)* for the vector field ζ if furthermore $\text{DF}_\zeta(\mathcal{X}, \mathcal{L}) = 0$ only if the central fiber X_0 of \mathcal{X} is isomorphic to X .

5.1.2. *Statement.* Our main result is a criterion for modified K-stability in terms of combinatorial data describing a \mathbb{Q} -Fano spherical variety.

Let X be a \mathbb{Q} -Fano variety, spherical under the action of a reductive group G . Let B be a Borel subgroup of G , T a maximal torus of B , B^- the Borel opposite to B with respect to T .

Let $\Delta^+ \subset \mathfrak{X}(T) \otimes \mathbb{R}$ be the moment polytope of X with respect to B . Let \mathcal{M}_- be the subgroup of $\mathfrak{X}(T)$ as defined in Section 3.1 with respect to the Borel subgroup B^- , let $\mathcal{N}_- = \text{Hom}(\mathcal{M}_-, \mathbb{Z})$ be the dual, and let $\mathcal{V}_- \subset \mathcal{N}_- \otimes \mathbb{R}$ be the valuation cone of X with respect to B^- . Denote by $\pi : \mathfrak{Y}(T) \otimes \mathbb{R} \rightarrow \mathcal{N}_- \otimes \mathbb{R}$ the quotient map induced by the inclusion of \mathcal{M}_- in $\mathfrak{X}(T)$. Let us denote by $\Xi \subset \mathfrak{X}(T) \otimes \mathbb{R}$ the dual cone of $\overline{\pi^{-1}(\mathcal{V}_-)}$, and by $\text{Relint}(\Xi)$ its relative interior.

Recall that we denote by Φ the root system of (G, T) and Φ^+ the positive roots defined by B . Let Φ_P^+ denote the roots in Φ^+ which are not orthogonal to Δ^+ , and let $2\rho_P$ be the sum of the elements of Φ_P^+ .

Let ζ be an element of the linear part of $\overline{\mathcal{V}_-} \subset \mathcal{N}_- \otimes \mathbb{R}$. Choose $\tilde{\zeta} \in \pi^{-1}(\zeta)$ any lift of ζ in $\mathfrak{Y}(T) \otimes \mathbb{R}$.

Theorem 5.3. *The variety X is modified K-semistable (with respect to special G -equivariant test configurations) for ζ if and only if*

$$\text{bar}_{DH, \tilde{\zeta}}(\Delta^+) \in 2\rho_P + \Xi.$$

It is modified K-stable (with respect to special G -equivariant test configurations) for ζ if and only if

$$\text{bar}_{DH, \zeta}(\Delta^+) \in 2\rho_P + \text{Relint}(\Xi).$$

5.2. Computing the Futaki invariant on a degeneration.

5.2.1. *Algebraic definition of the modified Futaki invariant.* Let us recall the algebraic definition of the modified Futaki invariant, given by Berman and Witt Nyström [BW].

Let X be a \mathbb{Q} -Fano variety, and A an ample line bundle on X so that $A|_{X_{\text{reg}}} = K_{X_{\text{reg}}}^{-m}$. Consider ξ, ζ two semisimple elements in the Lie algebra of $\text{Aut}^0(X)$ such that $J\xi$ and $J\zeta$ generate commuting compact real subgroups.

The complex torus $\overline{T}_\xi \times \overline{T}_\zeta$ acts on $H^0(X, A^k)$ for all $k \in \mathbb{N}$. This action is diagonalizable. Choose a basis $(e_j)_{j=1}^{N_k}$ of $H^0(X, A^k)$, such that $\overline{T}_\xi \times \overline{T}_\zeta$ acts on $\mathbb{C}e_j$ through the character χ_j . Define $s_j = \langle \chi_j, (\xi, 0) \rangle$ and $r_j = \langle \chi_j, (0, \zeta) \rangle$, where $\langle \cdot, \cdot \rangle$ denotes the duality pairing $\mathfrak{X}(\overline{T}_\xi \times \overline{T}_\zeta) \otimes \mathbb{R} \times \mathfrak{Y}(\overline{T}_\xi \times \overline{T}_\zeta) \otimes \mathbb{R} \rightarrow \mathbb{R}$.

The *quantized modified Futaki invariant at level k* is defined by:

$$\text{Fut}_{X, \zeta}^{(k)}(\xi) = - \sum_{j=1}^{N_k} \exp\left(\frac{r_j}{mk}\right) s_j.$$

The algebraic definition of the modified Futaki invariant is obtained thanks to the following result.

Proposition 5.4. [BW, Proposition 4.7] *The modified Futaki invariant of X is obtained as the following limit:*

$$\text{Fut}_{X, \zeta}(\xi) = \lim_{k \rightarrow \infty} \frac{1}{mkN_k} \text{Fut}_{X, \zeta}^{(k)}(\xi).$$

5.2.2. Equivariant degenerations and space of sections of linearized line bundles.

The fact that we can compute the modified Futaki invariant of a \mathbb{Q} -Fano variety on an equivariant degeneration will be a consequence of the following result. Our proof of this statement consists of a little twist in the method used by Donaldson to compute the Futaki invariant on toric test configurations in [Don02], also used by Li and Xu to prove an intersection formula for the Donaldson-Futaki invariant [LX14].

Proposition 5.5. *Let \mathcal{X} be a projective G -variety, where G is a reductive group, and let \mathcal{L} be a G -linearized line bundle on \mathcal{X} . Let $p : \mathcal{X} \rightarrow \mathbb{P}^1$ be a surjective G -invariant morphism, and assume that \mathcal{L} is relatively ample with respect to p . For $t \in \mathbb{P}^1$, let $X_t = p^{-1}(t)$ and $L_t = \mathcal{L}|_{X_t}$. Let $t_1, t_2 \in \mathbb{P}^1$. Then for large enough $k \in \mathbb{N}$, the representation of G given by $H^0(X_{t_1}, L_{t_1}^k)$ is isomorphic, as a representation of G , to $H^0(X_{t_2}, L_{t_2}^k)$.*

Proof. Let us first show that we can assume that \mathcal{L} is ample on \mathcal{X} . By [Har77, II.7.10], there exists an $N \in \mathbb{N}$ such that $\mathcal{L} \otimes p^*(\mathcal{O}_{\mathbb{P}^1}(N))$ is ample on \mathcal{X} . Furthermore, we have $H^0(X_t, L_t) \simeq H^0(X_t, (\mathcal{L} \otimes p^*(\mathcal{O}_{\mathbb{P}^1}(N)))|_{X_t})$ equivariantly for the action of G .

From now on we assume that \mathcal{L} is ample.

Consider on \mathbb{P}^1 the divisor given by a point $t \in \mathbb{P}^1$. Choose a global section s_t of $\mathcal{O}(1)$ with zero divisor $\{t\}$. Consider the pull back $p^*\mathcal{O}(1)$ of this line bundle, and the pulled back global section p^*s_t of $p^*\mathcal{O}(1)$. The zero divisor of p^*s_t is the irreducible divisor X_t , and since p is G -invariant, the section p^*s_t is G -invariant.

The divisor X_t provides, for all $k \in \mathbb{N}$, an exact sequence:

$$1 \rightarrow \mathcal{L}^k \otimes \mathcal{O}(-X_t) \rightarrow \mathcal{L}^k \rightarrow \mathcal{L}^k|_{X_t} \rightarrow 1$$

If we fix t , then for large k , we have by Serre vanishing Theorem,

$$H^1(X, \mathcal{L}^k \otimes \mathcal{O}(-X_t)) = 0,$$

so the short exact sequence of line bundles gives the following exact sequence in cohomology:

$$0 \rightarrow H^0(\mathcal{X}, \mathcal{L}^k \otimes \mathcal{O}(-X_t)) \rightarrow H^0(\mathcal{X}, \mathcal{L}^k) \rightarrow H^0(X_t, L_t^k) \rightarrow 0$$

where the first map is given by multiplication by the global section p^*s_t of $\mathcal{O}(X_t)$.

Remark that $H^0(\mathcal{X}, \mathcal{L}^k \otimes \mathcal{O}(-X_t)) = H^0(\mathcal{X}, \mathcal{L}^k \otimes p^*\mathcal{O}(-1))$ is independent of t . We can thus rewrite the exact sequence as :

$$0 \rightarrow H^0(\mathcal{X}, \mathcal{L}^k \otimes p^*\mathcal{O}(-1)) \rightarrow H^0(\mathcal{X}, \mathcal{L}^k) \rightarrow H^0(X_t, L_t^k) \rightarrow 0$$

It is then clear from the exact sequence that the dimension of $H^0(X_t, L_t^k)$ is independent of t for k large enough. Let us now consider the structure of G -representation. The second map is given by restriction to X_t , which is G -stable, so we obtain that $H^0(X_t, L_t^k)$ is the quotient representation of $H^0(\mathcal{X}, \mathcal{L}^k)$ by the image of $H^0(\mathcal{X}, \mathcal{L}^k \otimes p^*\mathcal{O}(-1))$. But for any $t \in \mathbb{P}^1$, our choice of a section p^*s_t so that p^*s_t is G -invariant, ensures that the inclusion of $H^0(\mathcal{X}, \mathcal{L}^k \otimes p^*\mathcal{O}(-1))$ in $H^0(\mathcal{X}, \mathcal{L}^k)$ is G -equivariant. In particular, the images $H^0(\mathcal{X}, \mathcal{L}^k \otimes p^*\mathcal{O}(-1))$ in $H^0(\mathcal{X}, \mathcal{L}^k)$ for different values of t are isomorphic as G -representations.

Given $t_1, t_2 \in \mathbb{P}^1$, we thus obtain that for large enough k , the structure of G -representation of $H^0(X_t, L_t^k)$ is independent of $t \in \{t_1, t_2\}$. \square

Let X be a normal \mathbb{Q} -Fano variety, and $(\mathcal{X}, \mathcal{L})$ be a special test configuration for X . Let X_t denote the fiber $\pi^{-1}(t)$. Assume that $(\mathcal{X}, \mathcal{L})$ is equipped with actions of two commuting complex tori \overline{T}_ζ and \overline{T}_ξ , generated by the holomorphic vector field ζ , respectively ξ , such that the map $\pi : \mathcal{X} \rightarrow \mathbb{C}$ is *invariant* under both actions. Let ζ_t , respectively ξ_t , denote the holomorphic vector fields on X_t induced by the restriction of the action of T_ζ respectively T_ξ , identified with elements of $\text{Lie}(\text{Aut}^0(X_t))$.

Corollary 5.6. *The function $\mathbb{C} \rightarrow \mathbb{R}; t \mapsto \text{Fut}_{X_t, \zeta_t}(\xi_t)$ is constant.*

Proof. By assumption, $(\mathcal{X}, \mathcal{L})$ is a special test configuration for X . It is a standard procedure (see for example [LX14, Section 8.1]) to extend this test configuration to a family $\pi : \mathcal{X} \rightarrow \mathbb{P}^1$ that we denote by the same letter, still equipped with a π -ample line bundle \mathcal{L} . We can further do this procedure in a way that is invariant under $\overline{T}_\zeta \times \overline{T}_\xi$.

Apply Proposition 5.5 to this family. Given $t_1, t_2 \in \mathbb{P}^1$, we obtain that the $\overline{T}_\zeta \times \overline{T}_\xi$ -representations $H^0(X_{t_1}, A_{t_1}^k)$ and $H^0(X_{t_2}, A_{t_2}^k)$ are isomorphic for k large enough. In particular, the quantized modified Futaki invariants are equal for k large enough:

$$\text{Fut}_{X_{t_1}, \zeta_{t_1}}^{(k)}(\xi_{t_1}) = \text{Fut}_{X_{t_2}, \zeta_{t_2}}^{(k)}(\xi_{t_2}).$$

By Proposition 5.4, this implies that the modified Futaki invariants of X_{t_1} and X_{t_2} are equal, hence the result. \square

5.3. Proof of Theorem 5.3. Let us denote by Q the stabilizer of the open orbit of B^- in X . By Proposition 3.21, we can choose a base point x , whose isotropy group is denoted by H , and a Levi subgroup L of Q adapted to H , containing T , such that $N_G(H) = H(T \cap N_G(H))$. In particular, the following are isomorphic

$$\text{Aut}_G(X) \simeq N_G(H)/H \simeq T \cap N_G(H)/T \cap H.$$

The lift $\tilde{\zeta}$ of ζ in $\mathfrak{Y}(T) \otimes \mathbb{R}$ is thus in $T \cap N_G(H)$, and the action of T_ζ on the open orbit identified with the coset space G/H is given by:

$$\exp(\tau\zeta) \cdot gH = g \exp(-\tau\tilde{\zeta})H.$$

Let \mathcal{X}' be a special G -equivariant test configuration for the \mathbb{Q} -Fano variety X . Recall that it is automatically equivariant under the action of $\text{Aut}_G(X)$ which contains T_ζ . Its central fiber, denoted by X'_0 , is a spherical \mathbb{Q} -Fano variety under the action of G by Proposition 3.23. Let $(\lambda, m) \in \mathfrak{Y}(T) \otimes \mathbb{N}^*$ be such that $y = \lim_{z \rightarrow 0} (\lambda(z), z^m) \cdot (x, 1)$ is in the open orbit of B^- in X'_0 , and denote the isotropy group of y by H'_0 .

Then, by Proposition 3.23 again, the action of $\mathbb{C}^* = T_\xi$ on X'_0 induced by the test configuration is described on G/H'_0 by

$$e^\tau \cdot gH'_0 = g \exp(-\tau\tilde{\xi})H'_0,$$

where $\tilde{\xi} = \lambda/m$. Furthermore, by Proposition 3.25, the action of T_ζ on X'_0 is still given, on G/H'_0 , by

$$\exp(\tau\zeta) \cdot gH'_0 = g \exp(-\tau\tilde{\zeta})H'_0.$$

Let us now choose \mathcal{X} any G -equivariant test configuration for the \mathbb{Q} -Fano spherical variety X'_0 with horospherical central fiber, using Corollary 3.31. Let y_0 be the base point obtained from $y \in X'_0$ through the choice of an adapted one parameter

subgroup, and let H_0 be its isotropy group. Using Proposition 3.24 and Proposition 3.25, we see that the action of T_ζ and T_ξ are still given by multiplication on the right by $\exp(-\tau\tilde{\zeta})$, respectively $\exp(-\tau\tilde{\xi})$, on G/H_0 . Furthermore, by Proposition 3.28, $H_0 = P^u(H_0 \cap T)$, where P is the parabolic subgroup of G , opposite to Q with respect to $T \subset L$. Remark that P is also the normalizer of H_0 in G . Remark also that the roots of the Levi subgroup $L \subset Q$ are determined by the moment polytope Δ^+ , as the roots that are orthogonal to Δ^+ (see [Bri89, Section 4.2]). Thus Φ_P^\perp and $2\rho_P$ as defined before the statement of Theorem 5.3 coincide with the data associated to P .

The moment polytope of X_0 with respect to B is the same as the moment polytope of X and X'_0 . This is a known fact for equivariant polarized degenerations of spherical varieties, and can be recovered using Proposition 5.5. We can now apply our computation of the Futaki invariant on \mathbb{Q} -Fano horospherical varieties to obtain:

$$\begin{aligned} \mathrm{DF}_\zeta(\mathcal{X}, \mathcal{L}) &= \mathrm{Fut}_{X'_0, \zeta}(\xi) \\ &= \mathrm{Fut}_{X_0, \zeta}(\xi) \end{aligned}$$

by Corollary 5.6, then by Theorem 4.5, this is

$$= C \left\langle \mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P, \tilde{\xi} \right\rangle$$

where C is some positive constant. Furthermore, we have seen that for horospherical \mathbb{Q} -Fano varieties, $\mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+)$ and the quantity above do not depend on the choice of lifts $\tilde{\zeta}$ and $\tilde{\xi}$ of ζ , $\xi \in \mathfrak{Y}(T/T \cap H_0) \otimes \mathbb{R}$. Since $T \cap H \subset T \cap H'_0 \subset T \cap H_0$, we obtain that the condition obtained does not depend on the choice of a lift $\tilde{\zeta}$ of $\zeta \in \mathcal{N}_- \otimes \mathbb{R} = \mathfrak{Y}(T/T \cap H) \otimes \mathbb{R}$.

By Theorem 3.30, for any choice of $(\lambda, m) \in \mathfrak{Y}(T) \oplus \mathbb{N}^*$ such that λ projects to the valuation cone \mathcal{V}_- , there exists a G -equivariant test configuration for X with $\tilde{\xi} = \lambda/m$. We then obtain that X is K -semistable (with respect to special equivariant test configurations) if and only if

$$\left\langle \mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P, \tilde{\xi} \right\rangle \geq 0$$

for all $\tilde{\xi} \in \pi^{-1}(\mathcal{V}_-)$ where $\pi : \mathfrak{Y}(T) \otimes \mathbb{R} \rightarrow \mathcal{N}_- \otimes \mathbb{R}$. This means precisely that $\mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P$ is in the dual cone Ξ to $\pi^{-1}(\mathcal{V}_-)$.

Furthermore, X'_0 is isomorphic to X if and only if ξ projects to an element of the linear part of the valuation cone. Indeed, if X'_0 is isomorphic to X , then the action induced by ξ on X is G -equivariant, so ξ projects to an element of the linear part of the valuation cone. Conversely, given such a ξ , the corresponding equivariant degeneration of G/H satisfies $H = H'_0$ by Proposition 3.26. Additionally, the moment polytope of X'_0 is the same as the moment polytope of X , and they are both \mathbb{Q} -Fano. Hence by the results of Gagliardi and Hofscheier recalled in Section 3.2.4, X and X'_0 are equivariantly isomorphic.

It is clear that $\left\langle \mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P, \tilde{\xi} \right\rangle = 0$ if and only if $\mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P$ lies in one of the hyperplanes defining the dual cone $(\pi^{-1}(\mathcal{V}_-))^\vee$. It is always the case if ξ projects to the linear part of \mathcal{V}_- . On the other hand, if $-\xi \notin \pi^{-1}(\mathcal{V}_-)$, then it means that $\mathrm{bar}_{DH, \tilde{\zeta}}(\Delta^+) - 2\rho_P$ is on the boundary of the cone $(\pi^{-1}(\mathcal{V}_-))^\vee$. This finishes the proof of the Theorem.

5.4. **Examples.** Let us give some illustrations of new situations where our main result can be applied. We restrict to smooth varieties for simplicity.

5.4.1. *Horospherical varieties.* Since the valuation cone is the whole space for horospherical varieties, the K-stability criterion becomes very simple in that case. We thus obtain a generalization of the main result of [PS10].

Corollary 5.7. *Let X be a smooth and Fano horospherical variety, with moment polytope Δ^+ . Then X admits a Kähler-Ricci soliton. Furthermore, the following are equivalent:*

- X is Kähler-Einstein,
- X is K-stable,
- X is K-semistable,
- the Futaki invariant of X vanishes,
- $\text{bar}_{DH}(\Delta^+) = 2\rho_P$.

Examples of interesting smooth, Fano, colored horospherical varieties, (thus not treated by [PS10]), are given by Pasquier in [Pas09]. Indeed, he classifies in this article horospherical manifolds with Picard number one. These are necessarily Fano, and colored when they are not homogeneous under a larger group. There exists in fact two infinite families (and three additional examples) of such manifolds [Pas09, Theorem 0.1]. Pasquier shows that their automorphism group is non reductive, which implies that they admit no Kähler-Einstein metrics and are not K-stable.

Our result shows that they are not K-semistable, and admit Kähler-Ricci solitons. The first conclusion is especially interesting in the following context. A conjecture by Odaka and Odaka [OO13, Conjecture 5.1], saying that a Fano manifold with Picard number one should be K-semistable, has been disproved by Fujita [Fuj], who provided two counterexamples. The examples of Pasquier provide infinite families of counterexamples, showing that Fujita's examples are not exceptional.

5.4.2. *Symmetric varieties.* An important class of spherical varieties is given by symmetric varieties. The combinatorial data from Section 3.1 have a nice expression in the case of symmetric spaces, that we recall here. The reference for this point of view is [Vus90], which is summarized in [Per14, Section 3.4.3]. Ruzzi [Ruz12] obtained a classification of smooth and Fano symmetric varieties of small rank. We use this, together with [GH15], to obtain explicit examples of moment polytopes of smooth and Fano symmetric varieties.

Let σ be a group involution of G , Let G^σ be the fixed point set of σ , and H a closed subgroup such that $G^\theta \subset H \subset N_G(G^\theta)$. The subgroup H is called a *symmetric subgroup* and G/H is called a *symmetric space*. A spherical embedding of G/H is called a *symmetric variety*.

A torus S in G is *split* if $\sigma(s) = s^{-1}$ for $s \in S$. Let S be a split torus in G , maximal for this property, and let T be a maximal torus of G containing S . Then we have $\sigma(T) = T$, and σ descends to an involution of $\mathfrak{X}(T)$, still denoted by σ .

Recall that Φ denotes the root system of G with respect to T . There exists a Borel subgroup B of G containing T such that for all $\alpha \in \Phi^+$, either $\sigma(\alpha) = \alpha$ or $\sigma(\alpha)$ is a negative root. Observe that the Borel subgroup B^- opposite to B with respect to T also satisfies this property. Both BH and B^-H are dense in G , in particular H is spherical.

Let Φ^σ denote the set of $\alpha \in \Phi$ such that $\alpha = \sigma(\alpha)$. It is a sub root system of Φ . Let $\Psi^+ = \Phi^+ \setminus \Phi^\sigma$, and

$$2\rho_\sigma = \sum_{\alpha \in \Psi^+} \alpha.$$

The set

$$\Phi_\sigma = \{\alpha - \sigma(\alpha); \alpha \in \Phi \setminus \Phi^\sigma\}$$

is a (possibly non-reduced) root system in $\mathfrak{X}(T/T \cap H) \otimes \mathbb{R}$.

We have $\mathcal{M} = \mathcal{M}_- = \mathfrak{X}(T/T \cap H)$ and the valuation cone \mathcal{V}_- with respect to B^- is the positive Weyl chamber C_σ^+ of the root system Φ_σ in $\mathfrak{Y}(T/T \cap H) \otimes \mathbb{Q}$.

Example 5.8. Consider the involution σ of $G = \mathrm{SL}_n(\mathbb{C})$ defined by sending a matrix g to the inverse of its transpose matrix $\sigma(g) = (g^t)^{-1}$. By definition, the set of fixed point is $\mathrm{SO}_n(\mathbb{C})$. As a consequence, any subgroup of G between $\mathrm{SO}_n(\mathbb{C})$ and $N_G(\mathrm{SO}_n(\mathbb{C}))$ is a symmetric subgroup.

The maximal torus T in $\mathrm{SL}_n(\mathbb{C})$ formed by diagonal matrices is split. We choose as Borel subgroup B the subgroup of upper triangular matrices, B^- is then the subgroup of lower triangular matrices. Then Φ^σ is empty and Φ_σ is the root system formed by the roots 2α for $\alpha \in \Phi$. In particular, the positive Weyl chamber C_σ^+ of the root system Φ_σ is also the positive Weyl chamber of Φ . Hence the valuation cone with respect to B^- is the positive Weyl chamber of Φ and its dual is the cone generated by Φ^+ .

For $H = G^\sigma = \mathrm{SO}_n(\mathbb{C})$, the lattice $\mathcal{M} = \mathcal{M}_- = \mathfrak{X}(T/T \cap H)$ is the lattice formed by the set of characters 2χ , where χ is a character of T (or equivalently χ is an element of the weight lattice of Φ).

For $H = N_G(G^\sigma) = N_{\mathrm{SL}_n(\mathbb{C})}(\mathrm{SO}_n(\mathbb{C}))$, the lattice $\mathcal{M} = \mathcal{M}_- = \mathfrak{X}(T/T \cap H)$ is the lattice formed by the set of characters 2χ , where χ is an element of the lattice generated by roots in Φ .

Corollary 5.9. *Let X be a smooth and Fano spherical embedding of G/H , with moment polytope Δ^+ . Then with the notations introduced above, X is Kähler-Einstein if and only if the barycenter of Δ^+ with respect to the measure $\prod_{\alpha \in \Psi^+} \kappa(\alpha, p) dp$ is in the relative interior of the translated cone $2\rho_\sigma + (C_\sigma^+)^{\vee}$.*

Example 5.10. The classification of spherical projective varieties under the action of $\mathrm{SL}_2(\mathbb{C})$ is explained in [AB06, Example 1.4.3]. Two of them are symmetric varieties, and the other are horospherical. Let us consider the symmetric ones.

The first one is $\mathbb{P}^1 \times \mathbb{P}^1$, where $\mathrm{SL}_2(\mathbb{C})$ acts diagonally. There are only two orbits. The closed orbit is the diagonal and the open orbit is the complement of the diagonal. The open orbit is isomorphic to the symmetric space $\mathrm{SL}_2(\mathbb{C})/\mathrm{SO}_2(\mathbb{C})$. The moment polytope of this Fano variety with respect to the Borel subgroup B of upper triangular matrices is the following.

$$\begin{array}{c} \text{---} \xrightarrow{\quad} \Delta^+ \\ | \qquad \qquad | \\ 0 \qquad \qquad 2\rho \qquad 8\rho/3 \end{array}$$


The barycenter $\mathrm{bar}_{DH}(\Delta^+)$ is computed as

$$\mathrm{bar}_{DH}(\Delta^+) = \frac{\int_0^4 x^2 dx}{\int_0^4 x dx} \rho = \frac{8}{3} \rho.$$

It of course satisfies the K-stability condition: $8\rho/3 \in \mathrm{Relint}(2\rho + \mathbb{R}_+\rho)$. Remark that in this case, there is only one possible central fiber different from $\mathbb{P}^1 \times \mathbb{P}^1$, it

is the horospherical variety S_2 in the notations of [AB06, Example 1.4.3]. More explicitly, consider the rational ruled surface $\mathbb{F}_2 = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(2))$. This is an $\mathrm{SL}_2(\mathbb{C})$ -horospherical variety with two closed orbits. One of these orbits is a curve with self-intersection -2 , which may be contracted, and the image of the contraction is denoted by S_2 . The anticanonically polarized equivariant horospherical degeneration of $\mathbb{P}^1 \times \mathbb{P}^1 \supset \mathrm{SL}_2(\mathbb{C})/\mathrm{SO}_2(\mathbb{C})$ is S_2 .

The second one is \mathbb{P}^2 , where $\mathrm{SL}_2(\mathbb{C})$ acts by the projectivization of its linear action on the quadratic forms in two variables. The open orbit is isomorphic to $\mathrm{SL}_2(\mathbb{C})/N_{\mathrm{SL}_2(\mathbb{C})}(\mathrm{SO}_2(\mathbb{C}))$. Again, this manifold obviously satisfies the K-stability criterion. Its moment polytope, together with the barycenter $\mathrm{bar}_{DH}(\Delta^+) = 4\rho$ are represented as follows.




The polarized equivariant horospherical degeneration of the symmetric variety \mathbb{P}^2 under this action of $\mathrm{SL}_2(\mathbb{C})$ is the variety S_4 , constructed similarly as S_2 from $\mathbb{F}_4 = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(4))$ by contracting the curve with self intersection equal to -4 .

Example 5.11. Consider the space of all conics in \mathbb{P}^2 . It may be identified with \mathbb{P}^5 by identifying a conic with its equation, which has six coefficients. Consider the action of $\mathrm{SL}_3(\mathbb{C})$ on this space. We see that it is a spherical variety with open orbit the orbit of nondegenerate conics, isomorphic to the symmetric space $\mathrm{SL}_3(\mathbb{C})/N_{\mathrm{SL}_3(\mathbb{C})}(\mathrm{SO}_3(\mathbb{C}))$.

There is another smooth and Fano spherical embedding of the symmetric space $\mathrm{SL}_3(\mathbb{C})/N_{\mathrm{SL}_3(\mathbb{C})}(\mathrm{SO}_3(\mathbb{C}))$, called the *variety of complete conics*. It may be constructed as the closure in $\mathbb{P}^5 \times (\mathbb{P}^5)^*$ of the set of couples (C, C^*) where C is a nondegenerate conic in \mathbb{P}^2 and C^* is the dual conic, defined as the set of tangents to C . This construction was generalized by De Concini and Procesi [DP83], to define *wonderful* compactifications of symmetric spaces under an adjoint semisimple group, which are often Fano so provide numerous potential examples.

The moment polytope corresponding to the variety of complete conics is the following.



It is easy to check our K-stability criterion for this manifold, and we obtain that the variety of complete conics admits Kähler-Einstein metrics.

Example 5.12. A reductive group $\hat{G} = \hat{G} \times \hat{G} / \mathrm{diag}(\hat{G})$ is a symmetric homogeneous space under the action of the group $G = \hat{G} \times \hat{G}$ for the involution $\sigma(g_1, g_2) = (g_2, g_1)$. If $\hat{B} \supset \hat{T}$ is a Borel subgroup of \hat{G} containing a maximal torus \hat{T} , then


$B = \hat{B} \times \hat{B}^-$ is an adapted Borel subgroup, Φ_G^σ is empty, $\mathfrak{X}(\hat{T} \times \hat{T}/\text{diag}(\hat{T}))$ is the anti diagonal embedding of $\mathfrak{X}(\hat{T})$ and can thus be identified with $\mathfrak{X}(\hat{T})$ by projection to the first coordinate. Under this identification, $2\rho_\sigma = 2\rho_B \in \mathfrak{X}(\hat{T} \times \hat{T}/\text{diag}(\hat{T}))$ is identified with $2\rho_{\hat{B}}$, and the restricted root system Φ_σ is identified with $\hat{\Phi}$ the root system of \hat{G} with respect to \hat{T} . We may identify, still under the projection to the first coordinate, the moment polytope Δ^+ with the polytope in $\mathfrak{X}(T) \otimes \mathbb{R}$ as defined in [Del15, Del], and the barycenter becomes

$$\text{bar}_{DH}(\Delta^+) = \int_{\Delta^+} p \prod_{\alpha \in \hat{\Phi}^+} \kappa(\alpha, p)^2 dp / \int_{\Delta^+} \prod_{\alpha \in \hat{\Phi}^+} \kappa(\alpha, p)^2 dp,$$

as used in [Del15, Del].

We then recover our previous result: a smooth and Fano group compactification X with moment polytope $\Delta^+ \subset \mathfrak{X}(T) \otimes \mathbb{R}$ is Kähler-Einstein if and only if $\text{bar}_{DH}(\Delta^+) \in 2\rho + \Xi$ where Ξ is the relative interior of the cone generated by Φ^+ . The similar statement for Kähler-Ricci solitons is new, even though it may be obtained using the techniques of [Del15, Del].

Let us take the opportunity to mention other interesting examples of group compactifications which were not mentioned in [Del15, Del], where the only examples were toroidal compactifications of simple groups. First consider the smooth and Fano compactifications of the reductive but not semisimple group $\text{GL}_2(\mathbb{C})$. Using [Ruz12], we obtain the following list of six moment polytopes.




It is obvious that the last three do not admit Kähler-Einstein metrics. Computations show that the first three admit Kähler-Einstein metrics. We may also

approximate, numerically, the unique holomorphic vector field such that the modified Futaki invariant vanishes (see [TZ02]), then check numerically that the modified K-stability condition is satisfied.


As an example, the precise coordinates of the barycenter $\text{bar}_{DH}(\Delta^+)$ in the case of the third polytope are $(2343/1750, -2343/1750)$. Remark that only the third and the fifth polytopes correspond to toroidal compactifications of $\text{GL}_2(\mathbb{C})$.

Considering colored compactifications also allows to obtain smaller dimensional examples of group compactifications with vanishing Futaki invariant which are not K-semistable. Namely, the three smooth and Fano compactifications of the (semisimple but not simple) group $\text{SO}_4(\mathbb{C})$, of dimension six, are colored, with the following moment polytopes:



While the first example satisfies the K-stability criterion, both the two others do not satisfy the K-semistability criterion, while having a trivially zero Futaki invariant since their group of equivariant automorphisms are finite. Hence they do not admit any Kähler-Ricci solitons. It would be interesting to know their full automorphism group and in particular if it is reductive.

Example 5.13. There are examples of smooth and Fano symmetric varieties with Picard number one, which are not homogeneous under a larger group [Ruz10, Ruz11]. Unlike the similar horospherical examples of Pasquier, these may very well be K-stable. For example, the smooth and Fano spherical embedding of $\text{SL}_3(\mathbb{C})/\text{SO}_3(\mathbb{C})$ with Picard number one, the smooth and Fano biequivariant compactification of the group G_2 with Picard number one, and the smooth and Fano biequivariant compactification of the group $\text{SL}_3(\mathbb{C})$ admit Kähler-Einstein metrics. Their respective moment polytopes are as follows.



REFERENCES

- [AB03] Hassan Azad and Indranil Biswas. Quasi-potentials and Kähler-Einstein metrics on flag manifolds. II. *J. Algebra*, 269(2):480–491, 2003.
- [AB04a] Valery Alexeev and Michel Brion. Stable reductive varieties. I. Affine varieties. *Invent. Math.*, 157(2):227–274, 2004.
- [AB04b] Valery Alexeev and Michel Brion. Stable reductive varieties. II. Projective case. *Adv. Math.*, 184(2):380–408, 2004.
- [AB04c] Valery Alexeev and Michel Brion. Toric degenerations of spherical varieties. *Sel. Math., New Ser.*, 10(4):453–478, 2004.
- [AB06] Valery Alexeev and Michel Brion. Stable spherical varieties and their moduli. *IMRP, Int. Math. Res. Pap.*, 2006(16):57, 2006.
- [AK05] Valery Alexeev and Ludmil Katzarkov. On K -stability of reductive varieties. *Geom. Funct. Anal.*, 15(2):297–310, 2005.
- [BBJ] Robert Berman, Sébastien Boucksom, and Mattias Jonsson. A variational approach to the Yau-Tian-Donaldson conjecture. arxiv:1509.04561v1.
- [Ber16] Robert J. Berman. K -polystability of \mathbb{Q} -Fano varieties admitting Kähler-Einstein metrics. *Invent. Math.*, 203(3):973–1025, 2016.
- [BLV86] Michel Brion, Dominique Luna, and Thierry Vust. Espaces homogènes sphériques. (Spherical homogeneous spaces). *Invent. Math.*, 84:617–632, 1986.
- [BP87] Michel Brion and Franz Pauer. Valuations des espaces homogènes sphériques. (Valuations of spherical homogeneous spaces). *Comment. Math. Helv.*, 62:265–285, 1987.
- [Bri] Michel Brion. Variétés sphériques. Notes de la session de la S. M. F. "Opérations hamiltoniennes et opérations de groupes algébriques".
- [Bri86] Michel Brion. Quelques propriétés des espaces homogènes sphériques. (Some properties of spherical homogeneous spaces). *Manuscr. Math.*, 55:191–198, 1986.
- [Bri87] Michel Brion. Sur l'image de l'application moment. In *Séminaire d'algèbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986)*, volume 1296 of *Lecture Notes in Math.*, pages 177–192. Springer, Berlin, 1987.
- [Bri89] Michel Brion. Groupe de Picard et nombres caractéristiques des variétés sphériques. *Duke Math. J.*, 58(2):397–424, 1989.

- [Bri97] Michel Brion. Curves and divisors in spherical varieties. In *Algebraic groups and Lie groups. A volume of papers in honour of the late R. W. Richardson*, pages 21–34. Cambridge: Cambridge University Press, 1997.
- [BW] Robert J. Berman and David Witt Nyström. Complex optimal transport and the pluripotential theory of Kähler-ricci solitons. arxiv:1401.8264v1.
- [CDS15a] Xiuxiong Chen, Simon K. Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities. *J. Amer. Math. Soc.*, 28(1):183–197, 2015.
- [CDS15b] Xiuxiong Chen, Simon K. Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π . *J. Amer. Math. Soc.*, 28(1):199–234, 2015.
- [CDS15c] Xiuxiong Chen, Simon K. Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof. *J. Amer. Math. Soc.*, 28(1):235–278, 2015.
- [CS] Tristan Collins and Gábor Székelyhidi. Sasaki-Einstein metrics and K-stability. arxiv:1512.07213v1.
- [CSW] Xiuxiong Chen, Song Sun, and Bing Wang. Kähler-Ricci flow, Kähler-Einstein metric, and K-stability. arxiv:1508.04397v1.
- [Del] Thibaut Delcroix. Kähler-Einstein metrics on group compactifications. arxiv:1510.07384v1.
- [Del15] Thibaut Delcroix. *Métriques de Kähler-Einstein sur les compactifications de groupes*. Theses, Université Grenoble Alpes, October 2015.
- [Don02] Simon K. Donaldson. Scalar curvature and stability of toric varieties. *J. Differ. Geom.*, 62(2):289–349, 2002.
- [DP83] Corrado De Concini and Claudio Procesi. Complete symmetric varieties. In *Invariant theory (Montecatini, 1982)*, volume 996 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 1983.
- [DS] Ved Datar and Gábor Székelyhidi. Kähler-Einstein metrics along the smooth continuity method. arxiv:1506.07495v1.
- [DT92] Weiyue Ding and Gang Tian. Kähler-Einstein metrics and the generalized Futaki invariant. *Invent. Math.*, 110(2):315–335, 1992.
- [Fuj] Kento Fujita. Examples of K-unstable Fano manifolds with the Picard number one. arxiv:1508.04290v2.
- [Fut83] Akito Futaki. An obstruction to the existence of Einstein Kähler metrics. *Invent. Math.*, 73(3):437–443, 1983.
- [GH15] Giuliano Gagliardi and Johannes Hofscheier. Gorenstein spherical Fano varieties. *Geom. Dedicata*, 178:111–133, 2015.
- [Har77] Robin Hartshorne. Algebraic geometry. Graduate Texts in Mathematics. 52. New York-Heidelberg-Berlin: Springer-Verlag. XVI, 496 p. DM 53.30; \$ 26.65 (1977)., 1977.
- [Hel78] Sigurdur Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 80 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978.
- [IS] Nathan Ilten and Hendrick Süß. K-stability for Fano manifolds with torus action of complexity one. arxiv:1507.04442v2.
- [KKV89] Friedrich Knop, Hanspeter Kraft, and Thierry Vust. The Picard group of a G -variety. In *Algebraische Transformationsgruppen und Invariantentheorie*, volume 13 of *DMV Sem.*, pages 77–87. Birkhäuser, Basel, 1989.
- [Kno91] Friedrich Knop. The Luna-Vust theory of spherical embeddings. In *Proceedings of the Hyderabad conference on algebraic groups held at the School of Mathematics and Computer/Information Sciences of the University of Hyderabad, India, December 1989*, pages 225–249. Madras: Manoj Prakashan, 1991.
- [LT] Kevin Langlois and Ronan Terpereau. The Cox ring of a complexity-one horospherical variety. arxiv:1603.06919v2.
- [LT16] Kevin Langlois and Ronan Terpereau. On the geometry of normal horospherical G -varieties of complexity one. *Journal of Lie Theory*, 26(1):049–078, 2016.
- [LV83] Dominique Luna and Thierry Vust. Plongements d’espaces homogènes. *Comment. Math. Helv.*, 58:186–245, 1983.

- [LX14] Chi Li and Chenyang Xu. Special test configuration and K -stability of Fano varieties. *Ann. Math. (2)*, 180(1):197–232, 2014.
- [Mab87] Toshiki Mabuchi. Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties. *Osaka J. Math.*, 24(4):705–737, 1987.
- [Mai00] Vincent Maillot. Géométrie d’Arakelov des variétés toriques et fibrés en droites intégrables. *Mém. Soc. Math. Fr. (N.S.)*, (80):vi+129, 2000.
- [OO13] Yuji Odaka and Takuzo Okada. Birational superrigidity and slope stability of Fano manifolds. *Math. Z.*, 275(3-4):1109–1119, 2013.
- [Pas] Boris Pasquier. A survey on the singularities of spherical varieties. arxiv:1510.03995v1.
- [Pas08] Boris Pasquier. Variétés horosphériques de Fano. *Bull. Soc. Math. Fr.*, 136(2):195–225, 2008.
- [Pas09] Boris Pasquier. On some smooth projective two-orbit varieties with Picard number 1. *Math. Ann.*, 344(4):963–987, 2009.
- [Per14] Nicolas Perrin. On the geometry of spherical varieties. *Transform. Groups*, 19(1):171–223, 2014.
- [Pop87] Vladimir L. Popov. Contraction of the actions of reductive algebraic groups. *Math. USSR, Sb.*, 58:311–335, 1987.
- [PS10] Fabio Podestà and Andrea Spiro. Kähler-Ricci solitons on homogeneous toric bundles. *J. Reine Angew. Math.*, 642:109–127, 2010.
- [Ruz10] Alessandro Ruzzi. Geometrical description of smooth projective symmetric varieties with Picard number one. *Transform. Groups*, 15(1):201–226, 2010.
- [Ruz11] Alessandro Ruzzi. Smooth projective symmetric varieties with Picard number one. *Int. J. Math.*, 22(2):145–177, 2011.
- [Ruz12] Alessandro Ruzzi. Fano symmetric varieties with low rank. *Publ. Res. Inst. Math. Sci.*, 48(2):235–278, 2012.
- [Tia97] Gang Tian. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.*, 130(1):1–37, 1997.
- [Tia15] Gang Tian. K -stability and Kähler-Einstein metrics. *Comm. Pure Appl. Math.*, 68(7):1085–1156, 2015.
- [Tim11] Dmitri A. Timashev. *Homogeneous spaces and equivariant embeddings*, volume 138 of *Encyclopaedia of Mathematical Sciences*. Springer, Heidelberg, 2011. Invariant Theory and Algebraic Transformation Groups, 8.
- [TZ02] Gang Tian and Xiaohua Zhu. A new holomorphic invariant and uniqueness of Kähler-Ricci solitons. *Comment. Math. Helv.*, 77(2):297–325, 2002.
- [Vus90] Thierry Vust. Plongements d’espaces symétriques algébriques: Une classification. (Embeddings of algebraic symmetric spaces: a classification). *Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser.*, 17(2):165–195, 1990.
- [WZ04] Xu-Jia Wang and Xiaohua Zhu. Kähler-Ricci solitons on toric manifolds with positive first Chern class. *Adv. Math.*, 188(1):87–103, 2004.
- [WZZ16] Feng Wang, Bin Zhou, and Xiaohua Zhu. Modified Futaki invariant and equivariant Riemann-Roch formula. *Adv. Math.*, 289:1205–1235, 2016.
- [Xio14] Meikui Xiong. Kähler-Ricci solitons and generalized Tian-Zhu’s invariant. *Int. J. Math.*, 25(7):13, 2014.

UNIV. GRENOBLE ALPES, IF, F-38000 GRENOBLE, FRANCE, CNRS, IF, F-38000 GRENOBLE, FRANCE

E-mail address: thibaut.delcroix@ujf-grenoble.fr