



HAL
open science

On the Distinguishing Number of Cyclic Tournaments: Towards the Albertson-Collins Conjecture

Kahina Meslem, Eric Sopena

► **To cite this version:**

Kahina Meslem, Eric Sopena. On the Distinguishing Number of Cyclic Tournaments: Towards the Albertson-Collins Conjecture. 2016. hal-01353772v3

HAL Id: hal-01353772

<https://hal.science/hal-01353772v3>

Preprint submitted on 17 Aug 2017 (v3), last revised 8 Oct 2018 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Distinguishing Number of Cyclic Tournaments: Towards the Albertson-Collins Conjecture

Kahina MESLEM¹ Éric SOPENA^{2,3}

August 10, 2017

Abstract

A distinguishing r -labeling of a digraph G is a mapping λ from the set of vertices of G to the set of labels $\{1, \dots, r\}$ such that no nontrivial automorphism of G preserves all the labels. The distinguishing number $D(G)$ of G is then the smallest r for which G admits a distinguishing r -labeling. From a result of Gluck (David Gluck, Trivial set-stabilizers in finite permutation groups, *Can. J. Math.* 35(1) (1983), 59–67), it follows that $D(T) = 2$ for every cyclic tournament T of (odd) order $2p+1 \geq 3$. Let $V(T) = \{0, \dots, 2p\}$ for every such tournament. Albertson and Collins conjectured in 1999 that the canonical 2-labeling λ^* given by $\lambda^*(i) = 1$ if and only if $i \leq p$ is distinguishing.

We prove that whenever one of the sub-tournaments of T induced by the set of vertices $\{0, \dots, p\}$ or $\{p+1, \dots, 2p\}$ is rigid, T satisfies the Albertson-Collins Conjecture. Using this property, we prove that several classes of cyclic tournaments satisfy the Albertson-Collins Conjecture. Moreover, we also prove that every Paley tournament satisfies the Albertson-Collins Conjecture.

Keywords: Distinguishing number; Automorphism group; Cyclic tournament; Albertson-Collins Conjecture.

MSC 2010: 05C20, 20B25.

1 Introduction

An r -labeling of a graph or a digraph G is a mapping λ from the set of vertices $V(G)$ of G to the set of labels $\{1, \dots, r\}$. An automorphism ϕ of G is λ -preserving if $\lambda(\phi(u)) = \lambda(u)$ for every vertex u of G . An r -labeling λ of G is *distinguishing* if the only λ -preserving automorphism of G is the identity, that is, the labeling λ breaks all the symmetries of G . In [4], Albertson and Collins introduce the *distinguishing number* of G , denoted $D(G)$, defined as the smallest r for which G admits a distinguishing r -labeling.

A digraph G is *rigid* (or *asymmetric*) if the only automorphism of G is the identity. Therefore, $D(G) = 1$ if and only if G is rigid.

In the last decade, distinguishing numbers have been studied for several families of graphs, such as trees [11, 26], interval graphs [12], planar graphs [5], hypercubes [6], Cartesian products

¹LaROMaD, Faculty of mathematics, University of Sciences and Technology Houari Boumediene (USTHB), B.P. 32 El-Alia, Bab-Ezzouar, 16111 Algiers, Algeria.

²Univ. Bordeaux, Bordeaux INP, CNRS, LaBRI, UMR5800, F-33400 Talence, France.

³Corresponding author. Eric.Sopena@labri.fr

of graphs [1, 9, 14, 15, 17, 18, 20], Kneser graphs [2, 10], infinite graphs [7, 19, 25], for instance. However, distinguishing numbers of digraphs have been less frequently considered since the paper of Albertson and Collins [3] (see [21, 22, 23]).

Symmetry breaking in tournaments is studied in [23] by Lozano who considers other ways of distinguishing vertices, namely by means of determining sets [8] (sometimes called fixing sets) or resolving sets.

We denote by \mathbb{Z}_n the group of residues modulo n . Let p be an integer, $p \geq 1$, and S be a subset of $\mathbb{Z}_{2p+1} \setminus \{0\}$ such that for every $k \in \mathbb{Z}_{2p+1} \setminus \{0\}$, $|S \cap \{k, 2p+1-k\}| = 1$. The *cyclic tournament* $T = T(2p+1; S)$ is the tournament of order $2p+1$ defined by $V(T) = \mathbb{Z}_{2p+1}$ and ij is an arc in T if and only if $j - i \in S$. Cyclic tournaments are sometimes called *circulant* or *rotational* tournaments in the literature.

Let Γ be a permutation group on a finite set Ω . A subset R of Ω is said to be Γ -*regular* if its stabilizer $S(R) = \{g \in \Gamma \mid gR = R\}$ is trivial, that is, $S(R) = \{Id_\Omega\}$, where $Id_\Omega \in \Gamma$ stands for the identity permutation acting on Ω . In [16], Gluck proves the following theorem (see also [24] for a simpler proof of this result).

Theorem 1 (Gluck [16]) *Let Γ be a permutation group of odd order on a finite set Ω . Then Γ has a regular subset in Ω .*

Theorem 1 implies that $D(T) = 2$ for every cyclic tournament $T = T(2p+1; S)$, $p \geq 1$. Indeed, it follows from Gluck's Theorem that there exists a subset R_T of $V(T)$ such that $\gamma(R_T) \neq R_T$ for every nontrivial automorphism γ of T . Therefore, the 2-labeling λ_{R_T} defined by $\lambda_{R_T}(i) = 1$ if and only if $i \in R_T$ is clearly distinguishing. On the other hand, $D(T) > 1$ since T is a cyclic tournament, and thus not rigid.

In [3], Albertson and Collins study distinguishing 2-labelings of cyclic tournaments and propose the following conjecture.

Conjecture 2 (Albertson-Collins [3]) *For every cyclic tournament $T = T(2p+1; S)$, $p \geq 1$, the 2-labeling λ^* , given by $\lambda^*(i) = 1$ if $0 \leq i \leq p$ and $\lambda^*(i) = 2$ otherwise, is distinguishing.*

In this paper, we prove that several classes of cyclic tournaments satisfy the Albertson-Collins Conjecture. We first give some definitions, notation and basic results in Section 2. We then consider the so-called pseudo-cyclic tournaments in Section 3 and prove our main results in Section 4. We finally propose some directions for future work in Section 5.

2 Preliminaries

We denote by $V(G)$ and $E(G)$ the set of vertices and the set of arcs of a digraph G , respectively. Let G be a digraph and u a vertex of G . The *outdegree* of u in G , denoted $d_G^+(u)$, is the number of arcs in $E(G)$ of the form uv , and the *indegree* of u in G , denoted $d_G^-(u)$, is the number of arcs in $E(G)$ of the form vu . The *degree* of u , denoted $d_G(u)$, is then defined by $d_G(u) = d_G^+(u) + d_G^-(u)$. If uv is an arc in G , u is an *in-neighbour* of v and v is an *out-neighbour* of u . We denote by $N_G^+(u)$ and $N_G^-(u)$ the set of out-neighbours and the set of in-neighbours of u in G , respectively. Hence, $d_G^+(u) = |N_G^+(u)|$ and $d_G^-(u) = |N_G^-(u)|$. Let v and w be two neighbours of u . We say that v and w *agree on u* if either both v and w are in-neighbours of u , or both v and w are out-neighbours of u , and that v and w *disagree on u* otherwise.

A digraph G is *regular* if all its vertices have the same in-degree and the same out-degree, that is, $d_G^+(u) = d_G^+(v)$ and $d_G^-(u) = d_G^-(v)$ for every two vertices u and v in G . For a subset W of $V(G)$, we denote by $G[W]$ the sub-digraph of G induced by W .

For any finite set Ω , Id_Ω denotes the identity permutation acting on Ω . Since the set Ω is always clear from the context, we simply write Id instead of Id_Ω in the following.

An *automorphism* of a digraph G is an arc-preserving permutation of its vertices, that is, a one-to-one mapping $\phi : V(G) \rightarrow V(G)$ such that $\phi(u)\phi(v)$ is an arc in G whenever uv is an arc in G . The set of automorphisms of G is denoted $\text{Aut}(G)$. The *order* of an automorphism ϕ is the smallest integer $k > 0$ for which $\phi^k = Id$. An automorphism ϕ of a digraph G is *nontrivial* if $\phi \neq Id$. A vertex u of G is *fixed* by ϕ if $\phi(u) = u$. The *orbit of u with respect to ϕ* is the set $\{u, \phi(u), \dots, \phi^{q-1}(u)\}$, where q is the *order of u with respect to ϕ* , that is, the smallest positive integer for which $\phi^q(u) = u$ (note that the order of u necessarily divides the order of ϕ). An *orbit of ϕ* is then an orbit of some vertex with respect to ϕ . A tournament cannot admit an automorphism of order 2 (such an automorphism would interchange the ends of some arc) and thus the automorphism group of a tournament has odd order, and every orbit of an automorphism of a tournament contains either one or at least three elements.

The following observations will be useful later.

Observation 3 *Let T be a tournament and u, v two vertices of T . If v is fixed by an automorphism $\phi \in \text{Aut}(T)$ and the orbit of u with respect to ϕ is of size at least 3, then all vertices in the orbit of u agree on v .*

Indeed, if u' belongs to the orbit of u with respect to ϕ , say $u' = \phi^r(u)$ for some integer $r > 0$, the result follows from the fact that $\phi^r(v) = v$ and ϕ^r is an automorphism of T .

Observation 4 *Let T be a tournament and $\phi \in \text{Aut}(T)$. If O_1 and O_2 are two (not necessarily distinct) orbits of ϕ of size at least 3, then, for every two vertices u and u' in O_1 ,*

$$|N_T^+(u) \cap O_2| = |N_T^+(u') \cap O_2| \quad \text{and} \quad |N_T^-(u) \cap O_2| = |N_T^-(u') \cap O_2|.$$

To see that, suppose $\phi^s(u) = u'$ for some integer $s > 0$, and let $N_T^+(u) \cap O_2 = \{v_1, \dots, v_k\}$, $k > 0$ (if $N_T^+(u) \cap O_2 = \emptyset$, the result is obvious). We then have $N_T^+(u') \cap O_2 = \{\phi^s(v_1), \dots, \phi^s(v_k)\}$, which implies $|N_T^+(u) \cap O_2| = |N_T^+(u') \cap O_2|$ since ϕ^s is an automorphism of T , and thus $|N_T^-(u) \cap O_2| = |N_T^-(u') \cap O_2|$.

Observation 5 *Let T be a tournament and $\phi \in \text{Aut}(T)$. Every orbit of ϕ induces a regular sub-tournament of T .*

This directly follows from Observation 4, when considering the case $O_1 = O_2$.

The *transitive tournament* of order n , denoted TT_n , is defined by $V(TT_n) = \{0, \dots, n-1\}$ and, for every $i, j \in V(TT_n)$, ij is an arc whenever $i < j$. Clearly, every transitive tournament is rigid (all its vertices have distinct indegrees) and thus $D(TT_n) = 1$ for every n . The *almost transitive tournament* of order n , denoted TT_n^* , is obtained from TT_n by reversing the arc from 0 to $n-1$. The tournament TT_n^* is thus Hamiltonian. It is also known that every almost transitive tournament of order at least 4 is rigid (note that TT_3^* is the directed 3-cycle, and this is not rigid).

Observation 6 *For every $n > 3$, the tournament TT_n^* is rigid, and thus $D(TT_n^*) = 1$.*

Indeed, every two vertices in TT_n^* have distinct indegrees, and thus distinct outdegrees, except the pairs $\{0, 1\}$ and $\{n-2, n-1\}$, as $d_{TT_n^*}^-(0) = d_{TT_n^*}^-(1) = 1$ and $d_{TT_n^*}^+(n-2) = d_{TT_n^*}^+(n-1) = 1$. Since every automorphism of TT_n^* has odd order, TT_n^* is a rigid tournament.

Let p be an integer, $p \geq 1$, and S be a subset of $\mathbb{Z}_{2p+1} \setminus \{0\}$ such that for every $k \in \mathbb{Z}_{2p+1} \setminus \{0\}$, $|S \cap \{k, 2p+1-k\}| = 1$. Let

$$S^+ = S \cap \{1, \dots, p\}, \quad \text{and} \quad S^- = \{1, \dots, p\} \setminus S^+.$$

We call the elements of S^+ the *positive connectors* of the cyclic tournament $T(2p+1, S)$ and the elements of S^- the *negative connectors* of $T(2p+1, S)$. Note that knowing either S^+ or S^- is enough to determine S since

$$S = S^+ \cup \{-s : s \in \{1, \dots, p\} \setminus S^+\} = \{1, \dots, p\} \setminus S^- \cup \{-s : s \in S^-\}.$$

Therefore, we preferably denote the cyclic tournament $T = T(2p+1; S)$ by $T = T(2p+1; S^-)$ whenever we deal with an explicit set S . In that case, for every $i, j \in \mathbb{Z}_{2p+1}$, $i < j$, ij is an arc in $T = T(2p+1; S^-)$ if and only if either $j - i \leq p$ and $j - i \notin S^-$, or $j - i > p$ and $2p + 1 - j + i \in S^-$.

Note that any cyclic tournament $T = T(2p+1; S)$ is regular. More precisely, $d_T^-(u) = d_T^+(u) = p = |S|$ for every vertex $u \in T$.

The *converse* T^c of a tournament T is obtained from T by reversing all the arcs. Clearly, the converse of any cyclic tournament $T = T(2p+1; S)$ is the cyclic tournament $T^c = T(2p+1; S^c)$ with $S^c = \{1, \dots, 2p\} \setminus S$. Moreover, T and T^c are isomorphic, via the mapping γ defined by $\gamma(0) = 0$ and $\gamma(i) = 2p + 1 - i$ for every $i \in V(T) \setminus \{0\}$.

3 Pseudo-cyclic tournaments

Let $T = T(2p+1; S)$ be a cyclic tournament and i, j two vertices of T with $i < j$. We denote by $T_{i,j}$ the sub-tournament of T induced by the set of vertices $\{i, i+1, \dots, j\}$. Note that the sub-tournament $T_{i,j}$ is not necessarily a cyclic tournament.

In the rest of this paper, we call the 2-labeling λ^* defined in Conjecture 2 the *canonical 2-labeling* of T . The canonical 2-labeling λ^* thus assigns label 1 to vertices of $T_{0,p}$ and label 2 to vertices of $T_{p+1,2p}$.

Since the tournament $T = T(2p+1; S)$ is cyclic, the sub-tournaments $T_{p+1,2p}$ and $T_{0,p-1}$ are isomorphic. Recall that the set S is characterized by the set S^- of its negative connectors. In the following, we thus study (not necessarily cyclic) tournaments of the following form.

Definition 7 Let N be a subset of $\mathbb{Z}_{p+1} \setminus \{0\}$, $p \geq 2$, whose elements are called *negative connectors*. The *pseudo-cyclic* tournament $P = P(p; N)$ is the tournament of order $p+1$ defined by $V(P) = \mathbb{Z}_{p+1}$ and ij , $i > j$, is an arc in P if and only if $i - j \in N$.

Note that if $T = T(2p+1; S)$ is a cyclic tournament, then $T_{0,p} \cong P(p; S^-)$ and $T_{p+1,2p} \cong T_{0,p-1} \cong P(p-1; S^-)$ (if $p \notin S^-$), or $T_{p+1,2p} \cong T_{0,p-1} \cong P(p-1; S^- \setminus \{p\})$ (if $p \in S^-$).

We first prove that $T = T(2p+1; S)$ satisfies the Albertson-Collins Conjecture whenever $T_{0,p}$ or $T_{p+1,2p}$ is rigid.

Proposition 8 *Let $T = T(2p+1; S)$ be a cyclic tournament. If $T_{0,p}$ is rigid or $T_{p+1,2p}$ is rigid, then the canonical 2-labeling λ^* of T is distinguishing.*

Proof. Let ϕ be a λ^* -preserving automorphism of T , that is, $\lambda^* \circ \phi = \lambda^*$, and let ϕ_1 and ϕ_2 denote the restriction of ϕ to $T_{0,p}$ and $T_{p+1,2p}$, respectively. Since $\lambda^* \circ \phi = \lambda^*$, both ϕ_1 and ϕ_2 are automorphisms. Moreover, since $T_{0,p}$ is rigid or $T_{p+1,2p}$ is rigid, we get $\phi_1 = Id$ or $\phi_2 = Id$. We will prove that we necessarily have $\phi_1 = Id$ and $\phi_2 = Id$, which gives $\phi = Id$ so that λ^* is a distinguishing labeling of T .

Suppose first that $\phi_1 = Id$ and assume to the contrary that $\phi_2 \neq Id$. Let $a_1 \in \{p+1, \dots, 2p\}$ be the “smallest” non-fixed vertex of $T_{p+1,2p}$. Since T is cyclic, we can assume without loss of generality $a_1 = p+1$. (If $a_1 \neq p+1$, by using the “shift” automorphism $\alpha : i \mapsto i + a_1 - p - 1$, the sub-tournaments $T_{0,p}$ and $T_{p+1,2p}$ are shifted to T_{a_1-p-1, a_1-1} and T_{a_1, a_1-p-2} , respectively. The two restricted automorphisms ϕ_1 and ϕ_2 become $\phi'_1 = \alpha\phi_1\alpha^{-1}$ and $\phi'_2 = \alpha\phi_2\alpha^{-1}$, respectively,

and we still have $\phi'_1 = Id$.) Let O_1 denote the orbit of a_1 with respect to ϕ_2 . Since the size of O_1 is at least 3 and the sub-tournament $T[O_1]$ induced by O_1 is regular (Observation 5), a_1 necessarily belongs to a cycle in $T[O_1]$ and thus to a 3-cycle in $T[O_1]$, say $a_1 a_2 a_3$.

Since $a_1 a_2 a_3$ is a 3-cycle, we get $\{a_2 - a_1, a_3 - a_2, a_1 - a_3\} \subseteq S$. If $a_2 < a_3$, let $w = a_1 - a_3 + a_2$, so that wa_1 is an arc of T . We then have $w \in V(T_{0,p})$ and thus, since w is fixed by ϕ , wa_1 and wa_2 are both arcs of T , a contradiction since $a_2 - w = a_3 - a_1$ and $a_3 - a_1 \notin S$. If $a_2 > a_3$, we get a similar contradiction by considering the vertex $w = a_1 + a_3 - a_2$: $a_1 w$ is an arc of T , again $w \in V(T_{0,p})$, which implies that $a_1 w$ and $a_3 w$ are both arcs of T , a contradiction since $w - a_3 = a_1 - a_2 \notin S$.

The case $\phi_2 = Id$ is similar. □

Note that Proposition 8 implies that whenever $T_{0,p}$ or $T_{p+1,2p}$ is rigid, both sets $V(T_{0,p})$ and $V(T_{p+1,2p})$ are $\text{Aut}(T)$ -regular. It should also be noticed that the condition in Proposition 8 is sufficient for a cyclic tournament to satisfy the Albertson-Collins Conjecture, but not necessary, as shown by the following example.

Example 9 Consider the cyclic tournament $T = T(13; \{2, 5, 6\})$. The automorphism group of T only contains rotations (that is, mappings $\phi : i \mapsto i + b$, $0 \leq b \leq 12$), so that the canonical 2-labeling λ^* is clearly distinguishing, and thus T satisfies the Albertson-Collins Conjecture. However, none of the sub-tournaments $T_{0,6}$ and $T_{7,12}$ is rigid. The sub-tournament $T_{0,6} \cong P(6; \{2, 5, 6\})$ admits an automorphism of order 3, namely $\phi' = (0, 3, 6)$, and the sub-tournament $T_{7,12} \cong P(5; \{2, 5\})$ also admits an automorphism of order 3, namely $\phi'' = (7, 8, 9)(10, 11, 12)$.

Moreover, we have the following.

Proposition 10 *Let $T = T(2p+1; S)$ be a cyclic tournament and λ^* be the canonical 2-labeling of T . If ϕ is a nontrivial λ^* -preserving automorphism of T , then ϕ has at least two orbits, both in $T_{0,p}$ and in $T_{p+1,2p}$.*

Proof. Suppose first that ϕ has only a single orbit in $T_{0,p}$, which implies that p is even. No vertex a of $T_{p+1,2p}$ can be fixed by ϕ , since otherwise, by Observation 3, all vertices in $T_{0,p}$ would agree on a , which would imply either $d_T^-(a) \geq p+1$ or $d_T^+(a) \geq p+1$, in contradiction with the definition of T . Therefore, ϕ has an even number of orbits in $T_{p+1,2p}$, of respective odd sizes $p_1, \dots, p_{2\ell}$, $\ell \geq 1$, with $p_i \geq 3$ for every i , $1 \leq i \leq 2\ell$, and $\sum_{i=1}^{2\ell} p_i = p$. Since $p_1 < p$, the automorphism $\phi_1 = \phi^{p_1}$ is λ^* -preserving and has a single orbit in $T_{0,p}$. But ϕ_1 fixes p_1 vertices of $T_{p+1,2p}$, contradicting the fact that any λ^* -preserving automorphism for which $T_{0,p}$ has a single orbit fixes no vertices of $T_{p+1,2p}$.

Suppose now that ϕ has only a single orbit in $T_{p+1,2p}$, which implies that p is odd. As before, we claim that no vertex of $T_{0,p}$ is fixed by ϕ . Assume to the contrary that a is such a vertex. As above, by Observation 3, we get $d_{T_{0,p}}^-(a) \geq p$ or $d_{T_{0,p}}^+(a) \geq p$, and thus $d_{T_{0,p}}^-(a) = d_{T_{0,p}}^+(a) = p$ according to the definition of T . Since $V(T_{p+1,2p})$ is either the set of in-neighbours or the set of out-neighbours of a , we get that the elements of S^- or of S^+ , respectively, are consecutive integers, and thus every two vertices in $T_{0,p}$ have distinct indegrees in $T_{0,p}$. Hence, the sub-tournament $T_{0,p}$ is rigid, which implies $\phi = Id$ by Proposition 8, a contradiction. Therefore, ϕ has an even number of orbits in $T_{0,p}$, each of size at least 3, and a contradiction arises as in the previous case. □

Let $P = P(p; N)$ be a pseudo-cyclic tournament. The *indegree sequence* of P is the sequence defined by

$$\sigma(P) = (d_P^-(0), \dots, d_P^-(p)).$$

The value of $d_P^-(i)$ for any vertex i of P is given by the following.

Proposition 11 *The indegree $d_P^-(i)$ of any vertex i of the pseudo-cyclic tournament $P = P(p; N)$, $0 \leq i \leq p$, is given by*

$$d_P^-(i) = \begin{cases} i + |N \cap \{i+1, \dots, p-i\}| & \text{if } 0 \leq i \leq \lfloor \frac{p}{2} \rfloor, \\ p - d_P^-(p-i) & \text{otherwise.} \end{cases}$$

Proof. By definition of P , there is an arc from j to i , $j > i$, if and only if $j - i \in N$. Hence,

$$d_P^-(i) = (i - |N \cap \{1, \dots, i\}|) + |N \cap \{1, \dots, p-i\}|.$$

Therefore, we get

$$d_P^-(i) = i + |N \cap \{i+1, \dots, p-i\}|$$

if $0 \leq i \leq \lfloor \frac{p}{2} \rfloor$ (with, in particular, $d_P^-(\frac{p}{2}) = \frac{p}{2}$ if p is even), and

$$d_P^-(i) = i - |N \cap \{p-i+1, \dots, i\}| = p - d_P^-(p-i)$$

if $\lceil \frac{p}{2} \rceil \leq i \leq p$. □

From Proposition 11, we get that for every pseudo-cyclic tournament $P = P(p; N)$, $d_P^-(0) = |N|$ and $d_P^-(p) = p - |N|$, and that the indegree sequence $\sigma(P)$ admits a central symmetry. Moreover, the difference of the indegrees of any two consecutive vertices is either -1, 0 or 1.

Proposition 12 *For any two consecutive vertices i and $i+1$ of the pseudo-cyclic tournament $P = P(p; N)$, $0 \leq i \leq p-1$, $d_P^-(i+1) - d_P^-(i) \in \{-1, 0, 1\}$. More precisely, we have*

1. $d_P^-(i+1) - d_P^-(i) = 1 - |N \cap \{i+1, p-i\}|$,
2. if p is odd, $d_P^-(\lceil \frac{p}{2} \rceil) - d_P^-(\lfloor \frac{p}{2} \rfloor) = 1 - 2|N \cap \{\lfloor \frac{p}{2} \rfloor + 1\}|$.

Proof. By Proposition 11, if $0 \leq i < \lfloor \frac{p}{2} \rfloor$ we have

$$\begin{aligned} d_P^-(i+1) - d_P^-(i) &= i+1 + |N \cap \{i+2, \dots, p-i-1\}| \\ &\quad - i - |N \cap \{i+1, \dots, p-i\}| \\ &= 1 - |N \cap \{i+1, p-i\}|. \end{aligned}$$

Since $0 \leq |N \cap \{i+1, p-i\}| \leq 2$, we get $d_P^-(i+1) - d_P^-(i) \in \{-1, 0, 1\}$.

If $\lceil \frac{p}{2} \rceil \leq i < p$, then, by Proposition 11, we get

$$\begin{aligned} d_P^-(i+1) - d_P^-(i) &= p - d^-(p-i-1) - p + d^-(p-i) \\ &= d^-(p-i) - d^-(p-i-1) \\ &= 1 - |N \cap \{p-i-1+1, p-p+i+1\}| \\ &= 1 - |N \cap \{i+1, p-i\}|. \end{aligned}$$

Finally, if p is odd, we have

$$\begin{aligned} d_P^-(\lceil \frac{p}{2} \rceil) - d_P^-(\lfloor \frac{p}{2} \rfloor) &= p - (\lfloor \frac{p}{2} \rfloor + |N \cap \{\lfloor \frac{p}{2} \rfloor + 1\}|) - (\lfloor \frac{p}{2} \rfloor + |N \cap \{\lfloor \frac{p}{2} \rfloor + 1\}|) \\ &= 1 - 2|N \cap \{\lfloor \frac{p}{2} \rfloor + 1\}|. \end{aligned}$$

This completes the proof. □

Note that if p is odd, the value of $d_P^-(\lceil \frac{p}{2} \rceil) - d_P^-(\lfloor \frac{p}{2} \rfloor)$ is either 1 or -1.

We can then define three types of vertices as follows.

Definition 13 Let $P = P(p; N)$ be a pseudo-cyclic tournament. A vertex $i \in V(P)$, $0 \leq i \leq p - 1$, is said to be

1. an *ascent-vertex* if $d_P^-(i + 1) = d_P^-(i) + 1$, that is, $N \cap \{i + 1, p - i\} = \emptyset$,
2. a *descent-vertex* if $d_P^-(i + 1) = d_P^-(i) - 1$, that is, $\{i + 1, p - i\} \subseteq N$, or
3. a *plateau-vertex* if $d_P^-(i + 1) = d_P^-(i)$, that is, $|N \cap \{i + 1, p - i\}| = 1$.

A pseudo-cyclic tournament $P = P(p; N)$ is cyclic if and only if all its vertices have the same indegree, which implies p is even and $|N| = \frac{p}{2}$, and the set N is such that $i \in N$ if and only if $p + 1 - i \notin N$. In other words, we have the following.

Observation 14 A pseudo-cyclic tournament $P = P(p; N)$ is cyclic if and only if every vertex $i \in V(P)$, $0 \leq i \leq p - 1$, is a plateau-vertex.

Moreover, we also have the following.

Observation 15 Let $P^c = (p; N^c)$ be the converse pseudo-cyclic tournament of $P = (p; N)$, that is, $N^c = \{1, \dots, p\} \setminus N$. If i is an ascent-vertex (resp. a descent-vertex, a plateau-vertex) in P , $0 \leq i \leq p - 1$, then i is a descent-vertex (resp. an ascent-vertex, a plateau-vertex) in P^c .

This observation directly follows from the fact that $d_{P^c}^-(i) = p - d_P^-(i)$ for every vertex i .

We denote by $\alpha(P)$, $\delta(P)$ and $\pi(P)$ the number of ascent-vertices, descent-vertices and plateau-vertices in $\sigma(P)$, respectively, so that $\alpha(P) + \delta(P) + \pi(P) = p$. Moreover, according to the three above types of vertices, we define three types of subsequences of the indegree sequence $\sigma(P)$.

Definition 16 Let $P = P(p; N)$ be a pseudo-cyclic tournament. A sequence of $k \geq 2$ consecutive vertices $(i, \dots, i + k - 1)$, $i \leq n - k + 1$, of P is called

1. an *ascent of size k* if $d_P^-(i + j + 1) = d_P^-(i + j) + 1$ for every j , $0 \leq j \leq k - 2$,
2. a *descent of size k* if $d_P^-(i + j + 1) = d_P^-(i + j) - 1$ for every j , $0 \leq j \leq k - 2$,
3. a *plateau of size k* if $d_P^-(i + j + 1) = d_P^-(i + j)$ for every j , $0 \leq j \leq k - 2$.

Note here that an ascent, a descent or a plateau of size k contains $k - 1$ ascent-vertices, descent-vertices or plateau-vertices, respectively.

Example 17 The pseudo-cyclic tournament $P = P(8; \{2, 4, 5\})$ is depicted in Figure 1(a) (only arcs corresponding to negative connectors are drawn, every missing arc is thus directed from left to right). The indegree sequence of P is given by


$$\sigma(P) = (3, 4, 4, 5, 4, 3, 4, 4, 5)$$

and is represented by the (centrally symmetric) *indegree path* of P depicted in Figure 1(b). We then have $\pi(P) = 2$, $\delta(P) = 2$ and $\alpha(P) = 4$.


The number of descent-vertices and the number of plateau-vertices are related to the cardinality of the set N as follows.

Proposition 18 For every pseudo-cyclic tournament $P = P(p; N)$,

$$\delta(P) + \frac{1}{2}\pi(P) = |N|.$$



(a) The pseudo-cyclic tournament $P(8; \{2, 4, 5\})$



(b) The indegree path of $P(8; \{2, 4, 5\})$

Figure 1: The pseudo-cyclic tournament $P(8; \{2, 4, 5\})$ and its indegree path

Proof. If $N = \emptyset$, then $\sigma(P) = (0, 1, \dots, p)$ (P is transitive), so that $\delta(P) = \pi(P) = 0$ and we are done.

Otherwise, let $N = \{a_1, \dots, a_q\}$, so that $|N| = q \geq 1$. We claim that each a_i generates either a descent-vertex or two plateau-vertices. Indeed, for each $a_i \in N$, either $p + 1 - a_i \notin N$, which implies that both vertices $a_i - 1$ and $p - a_i$ are plateau-vertices, or $p + 1 - a_i \in N$, which implies that $a_i - 1$ is a descent-vertex. We thus get $|N| = \delta(P) + \frac{1}{2}\pi(P)$, as required. \square

The following proposition shows that vertices with same indegree in a pseudo-cyclic tournament are necessarily “not too far” from each other.

Proposition 19 *Let $P = P(p; N)$ be a pseudo-cyclic tournament. If i and j are two vertices of P with $i < j$ and $d_P^-(i) = d_P^-(j)$, then*

1. *if $0 \leq i < j \leq \lfloor \frac{p}{2} \rfloor$ or $\lceil \frac{p}{2} \rceil \leq i < j \leq p$, then $j \leq i + |N|$,*
2. *if $0 \leq i \leq \lfloor \frac{p}{2} \rfloor < j \leq p$, then $j \leq i + 2|N|$.*

Proof. Let i and j be such that $0 \leq i < j \leq \lfloor \frac{p}{2} \rfloor$ and $j > i + |N|$. By Proposition 11, we have

$$\begin{aligned} d_P^-(j) - d_P^-(i) &= j + |N \cap \{j + 1, \dots, p - j\}| - i - |N \cap \{i + 1, \dots, p - i\}| \\ &= j - i - |N \cap (\{i + 1, j\} \cup \{p - j + 1, p - i\})| \\ &> |N| - |N| = 0. \end{aligned}$$

Therefore, $d_P^-(i) \neq d_P^-(j)$. Since the indegree path of P is centrally symmetric, the result also holds when $\lceil \frac{p}{2} \rceil \leq i < j \leq p$.

Suppose now that $0 \leq i \leq \lfloor \frac{p}{2} \rfloor < j \leq p$ and $j > i + 2|N|$. By Proposition 11, we have

$$\begin{aligned} d_P^-(j) - d_P^-(i) &= j - |N \cap \{p - j + 1, \dots, j\}| - i - |N \cap \{i + 1, \dots, p - i\}| \\ &= j - i - (|N \cap \{p - j + 1, \dots, j\}| + |N \cap \{i + 1, \dots, p - i\}|) \\ &> 2|N| - (|N| + |N|) = 0. \end{aligned}$$

Again, we get $d_P^-(i) \neq d_P^-(j)$. □

Finally, the following property will be useful in the sequel.

Proposition 20 *Let $P = P(p; N)$ be a pseudo-cyclic tournament. If a vertex i is fixed by every automorphism of P , then the vertex $p - i$ is also fixed by every automorphism of P .*

Proof. Suppose to the contrary that i is fixed by every automorphism of P and that there exists an automorphism ϕ of P such that $\phi(p - i) \neq p - i$.

Let $\phi^* : V(P) \rightarrow V(P)$ be the mapping defined by $\phi^*(i) = p - \phi(p - i)$ for every $i \in V(P)$.

We claim that $\phi^* \in \text{Aut}(P)$. Let ij be an arc in P with $i > j$, that is, $i - j \in N$. We then have

$$\phi^*(i) - \phi^*(j) = p - \phi(p - i) - p + \phi(p - j) = \phi(p - j) - \phi(p - i).$$

Since $p - j > p - i$ and $p - j - p + i = i - j \in N$, $(p - j)(p - i)$ is an arc in P . Since $\phi \in \text{Aut}(P)$, $\phi(p - j)\phi(p - i)$ is an arc in P .

Suppose first that $\phi(p - j) > \phi(p - i)$, which implies $\phi(p - j) - \phi(p - i) \in N$. In that case $\phi^*(j) = p - \phi(p - j) < p - \phi(p - i) = \phi^*(i)$ and, since $\phi^*(i) - \phi^*(j) = \phi(p - j) - \phi(p - i) \in N$, $\phi^*(i)\phi^*(j)$ is an arc in P .

Suppose now that $\phi(p - j) < \phi(p - i)$, which implies $\phi(p - i) - \phi(p - j) \notin N$. In that case $\phi^*(i) = p - \phi(p - i) < p - \phi(p - j) = \phi^*(j)$ and, since $\phi^*(j) - \phi^*(i) = \phi(p - i) - \phi(p - j) \notin N$, $\phi^*(i)\phi^*(j)$ is an arc in P .

The case of an arc ij with $i < j$ is similar, and thus $\phi^* \in \text{Aut}(P)$.

We then have $\phi^*(i) = p - \phi(p - i) \neq i$, contradicting the fact that i is fixed by every automorphism of P . □

This immediately gives the following.

Corollary 21 *Let $P = P(p; N)$ be a pseudo-cyclic tournament. If all vertices $i \leq \lfloor \frac{p}{2} \rfloor$ of P are fixed by every automorphism of P , then P is rigid.*

4 Cyclic tournaments satisfying the Albertson-Collins Conjecture

In this section, we prove that several classes of cyclic tournaments $T(2p + 1; S)$ satisfy the Albertson-Collins Conjecture. We first propose a few sufficient conditions for a cyclic tournament to satisfy the Albertson-Collins Conjecture and then consider several specific classes of cyclic tournaments, depending on the structure of the set S of connectors.

4.1 Simple sufficient conditions

By Proposition 10, for every nontrivial λ^* -preserving automorphism ϕ of $T = T(2p + 1; S)$, none of $T_{0,p}$ and $T_{p+1,2p}$ has a single orbit with respect to ϕ . Using this property, we can prove the following result.

Lemma 22 *The cyclic tournament $T = T(2p + 1; S)$ satisfies the Albertson-Collins Conjecture whenever one of the following conditions is satisfied.*

1. $\text{Aut}(T)$ is of order $2p + 1$,
2. p is even, $|S^-| = \frac{p}{2}$, $p + 1 - s \notin S^-$ for every $s \in S^-$, and $|\text{Aut}(T_{0,p})| = p + 1$,
3. p is odd, $|S^- \setminus \{p\}| = \frac{p-1}{2}$, $p - s \notin S^-$ for every $s \in S^-$, and $|\text{Aut}(T_{p+1,2p})| = p$.

Proof. Recall that for every cyclic tournament T_n of order n , the n rotations of T_n form a subgroup of $\text{Aut}(T_n)$.

In Case 1, $\text{Aut}(T)$ contains only rotations, so that λ^* is clearly distinguishing. In Case 2, $T_{0,p}$ is a cyclic tournament and every automorphism of $T_{0,p}$ is a rotation. Therefore, the restriction ϕ_1 to $T_{0,p}$ of any λ^* -preserving automorphism ϕ of T is a rotation and thus $\phi_1 = \text{Id}$. The result then follows by Proposition 10. Case 3 is similar, using $T_{p+1,2p}$ instead of $T_{0,p}$. \square

Let T be a tournament. We denote by $V_d(T)$, $d \geq 0$, the set of vertices of T with indegree d . Since any automorphism maps every vertex to a vertex with same degree, we have the following result.

Lemma 23 *Let $P = P(n; N)$ be a pseudo-cyclic tournament. If the sub-tournament $P[V_d(P)]$ of P , induced by $V_d(T)$, is rigid for every d , $d \geq 0$, then P is rigid.*

This lemma directly gives the following results.

Theorem 24 *Let $T = T(2p + 1; S)$ be a cyclic tournament, $P_1 = T_{0,p}$ and $P_2 = T_{p+1,2p}$. If $P_1[V_d(P_1)]$ is rigid for every d , or $P_2[V_d(P_2)]$ is rigid for every d , $d \geq 0$, then T satisfies the Albertson-Collins Conjecture.*

Proof. The result directly follows from Lemma 23 and Proposition 8. \square

Theorem 25 *Let $T = T(2p + 1; S)$ be a cyclic tournament with $S^- \neq \emptyset$ and let $\min(S^-)$ denote the smallest negative connector of S . If $\min(S^-) > 2|S^-|$, then T satisfies the Albertson-Collins Conjecture.*

Proof. Consider the pseudo-cyclic sub-tournament $P = T_{0,p}$. By Proposition 19, we know that if two vertices i and j , $i < j$, have the same indegree in P , then $j - i \leq 2|S^-|$. Since $\min(S^-) > 2|S^-|$, we get that the sub-tournament $P[V_d(P)]$ is transitive, and thus rigid, for every d , $d \geq 0$. The result then follows by Theorem 24. \square

4.2 Cyclic tournaments with few negative connectors

We consider in this subsection the case of cyclic tournaments with at most two negative connectors. We prove that every such tournament satisfies the Albertson-Collins Conjecture.

Theorem 26 *If $T = T(2p + 1; S)$ is a cyclic tournament with $p \geq 1$ and $|S^-| \leq 2$, then T satisfies the Albertson-Collins Conjecture.*

Proof. The result is obvious if $p = 1$ since, in that case, T is the directed 3-cycle. We thus assume $p \geq 2$. We will prove that one of the two sub-tournaments $P_1 = T_{0,p}$ or $P_2 = T_{p+1,2p}$ is rigid, so that the result follows by Proposition 8.

If $S^- = \emptyset$, then P_1 is transitive and thus rigid.

If $|S^-| = 1$, say $S^- = \{a\}$, then, by Proposition 11, the indegree of any vertex i of P_1 is as follows.

$$d_{P_1}^-(i) = \begin{cases} i & \text{if } a \notin \{i + 1, \dots, p - i\} \text{ and } 0 \leq i \leq \lfloor \frac{p}{2} \rfloor, \\ i + 1 & \text{if } a \in \{i + 1, \dots, p - i\} \text{ and } 0 \leq i \leq \lfloor \frac{p}{2} \rfloor, \\ p - d_{P_1}^-(p - i) & \text{otherwise.} \end{cases}$$

Therefore, $d_{P_1}^-(i) = d_{P_1}^-(j)$, $i < j \leq \lfloor \frac{p}{2} \rfloor$, if and only if $j = i + 1$ and $j \in \{a, p + 1 - a\}$. If $a \neq \lfloor \frac{p}{2} \rfloor$, then the number of vertices in P_1 with degree d is at most 2 for every d . In that case,

P_1 satisfies the condition of Lemma 23 and is thus rigid. If $a = \lfloor \frac{p}{2} \rfloor$ the same property holds except if p is even (in that case, we have $|V_{\frac{p}{2}}(P_1)| = 3$ and $|V_d(P_1)| = 1$ for every $d \neq \frac{p}{2}$). But then, P_2 satisfies the condition of Lemma 23 (since $p - 1$ is odd) and we are done too.

Assume now that $|S^-| = 2$ and let $S^- = \{a_1, a_2\}$. If $p = 2$, then P_2 has order 2 and is thus rigid. If $p = 3$, then either $S^- = \{1, 2\}$, $S^- = \{1, 3\}$ or $S^- = \{2, 3\}$. It is then easily checked that $|V_1(P_1)| = |V_2(P_1)| = 2$ in all cases, so that P_1 satisfies the condition of Lemma 23 and we are done.

Suppose thus that $p > 3$ and, moreover, that p is even (if p is odd we simply consider P_2 instead of P_1). By Proposition 18, we know that $\delta(P_1) + \frac{1}{2}\pi(P_1) = 2$. We thus have three cases to consider.

1. $\delta(P_1) = 2$ and $\pi(P_1) = 0$.

In that case, the sequence $\sigma(P_1)$ has no plateau and exactly two descent-vertices, say d_1 and d_2 . Since $\sigma(P_1)$ is centrally symmetric and p is even, we necessarily have $d_2 = p - d_1 - 1$ which implies in particular that the difference $d_2 - d_1$ is necessarily odd.

If $d_2 - d_1 = 1$, we necessarily have $d_2 = \frac{p}{2}$ (see Figure 2(a)). Moreover, since $p > 3$, the vertex 0 cannot be a descent-vertex (this would imply $\frac{p}{2} = 1$) and thus $1 \notin S^-$. The only set $V_d(P_1)$ containing more than two elements is thus $V_{\frac{p}{2}}(P_1) = \{\frac{p}{2} - 2, \frac{p}{2}, \frac{p}{2} + 2\}$. But since the vertex $\frac{p}{2} - 1$ is fixed by every automorphism of P_1 and vertices $\frac{p}{2} - 2$ and $\frac{p}{2}$ disagree on $\frac{p}{2} - 1$ (as $1 \notin S^-$), $\frac{p}{2} - 2$ and $\frac{p}{2}$ are also fixed so that P_1 is rigid by Corollary 21.

If $d_2 - d_1 > 3$ (see Figure 2(b)), then every set $V_d(P_1)$ contains at most two elements and Lemma 23 again allows us to conclude.

Suppose finally that $d_2 - d_1 = 3$ (see Figure 2(c)). This case is similar to the case $d_2 - d_1 = 1$: again, the only set $V_d(P_1)$ containing more than two elements is $V_{\frac{p}{2}}(P_1) = \{\frac{p}{2} - 2, \frac{p}{2}, \frac{p}{2} + 2\}$ while the vertex $\frac{p}{2} - 1$ is fixed by every automorphism of P_1 and vertices $\frac{p}{2} - 2$ and $\frac{p}{2}$ disagree on $\frac{p}{2} - 1$. Again, P_1 is rigid thanks to Corollary 21.

2. $\delta(P_1) = 1$ and $\pi(P_1) = 2$.

Since $\delta(P_1) = 1$, we cannot have two consecutive descent-vertices in P_1 . Therefore, since $\sigma(P_1)$ is centrally symmetric, if i is a descent-vertex, then $p - (i + 1)$ is also a descent-vertex. As $\delta(P_1) = 1$, we must have $i = p - (i + 1)$, in contradiction with our assumption on the parity of p . Hence, this case cannot happen.


3. $\delta(P_1) = 0$ and $\pi(P_1) = 4$.

Since p is even and $\sigma(P_1)$ is centrally symmetric, we have four possibilities: (a) P_1 contains four plateaus of size 2, (b) P_1 contains two plateaus of size 3, (c) P_1 contains one plateau of size 3, centered at $\frac{p}{2}$, and two plateaus of size 2, or (d) P_1 contains a single plateau, of size 5 and centered at $\frac{p}{2}$, see Figure 3(a), (b), (c) or (d), respectively.

(a) If P_1 contains four plateaus of size 2, then P_1 satisfies the condition of Lemma 23 and we are done.


(b) If P_1 contains two plateaus of size 3, then the plateau-vertices are necessarily of the form $i, i + 1, p - i - 2$ and $p - i - 1$, with $i \leq \frac{p}{2} - 3$, so that $S^- = \{a_1, a_2\}$, with $a_1 \in \{i + 1, p - i\}$, $a_2 \in \{i + 2, p - i - 1\}$, and $d_{P_1}^-(i) = 2 + i$. We then have $V_{2+i}(P_1) = \{i, i + 1, i + 2\}$ and all other sets $V_d(P_1)$, $2 \leq d \leq \frac{p}{2}$, $d \neq 2 + i$, are singletons.

If $i > 0$ and $i + 1 \in S^-$, then, since $i \notin S^-$, vertices i and $i + 1$ disagree on 0. Since vertex 0 is fixed by every automorphism of P_1 , i and $i + 1$ cannot belong to the same orbit of any automorphism and thus, by Corollary 21, P_1 is rigid. If $i > 0$ and $p - i \in S^-$, we get the same conclusion since, in that case, either $i + 2 \in S^-$, which implies that i and $i + 2$ disagree on 0, or $p - i - 1 \in S^-$, which implies, since



(a) $\delta(P_1) = 2, \pi(P_1) = 0, d_2 - d_1 = 1$

(b) $\delta(P_1) = 2, \pi(P_1) = 0, d_2 - d_1 > 3$



(c) $\delta(P_1) = 2, \pi(P_1) = 0, d_2 - d_1 = 3$

Figure 2: Indegree paths for the proof of Theorem 26

$p - i - 2 \notin S^-$, that i and $i + 2$ disagree on p (and p is fixed by every automorphism of P_1). Suppose finally that $i = 0$. Since p is even, the vertex 3 is not a plateau-vertex and thus is fixed by every automorphism of P_1 . If 0, 1 and 2 disagree on 3, these three vertices cannot belong to the same orbit of any automorphism and P_1 is rigid thanks to Corollary 21. Otherwise, since 3 cannot be an in-neighbour of 0, 1 and 2 (this would imply $|S^-| \geq 3$), 3 is an out-neighbour of 0, 1 and 2, so that $1, 2 \notin S^-$, which implies that $P_1[\{0, 1, 2\}]$ is transitive, and P_1 is rigid by Lemma 23.

(c) If P_1 contains one plateau of size 3 and two plateaus of size 2, then the plateau-vertices are necessarily of the form $i, \frac{p}{2} - 1, \frac{p}{2}$ and $p - i - 1$, with $i \leq \frac{p}{2} - 3$, so that $S^- = \{a_1, a_2\}$, with $a_1 \in \{i + 1, p - i\}$, $a_2 \in \{\frac{p}{2}, \frac{p}{2} + 1\}$, and $d_{P_1}^-(\frac{p}{2}) = \frac{p}{2}$. Therefore, the only set $V_d(P_1)$ containing more than two elements is $V_{\frac{p}{2}}(P_1) = \{\frac{p}{2} - 1, \frac{p}{2}, \frac{p}{2} + 1\}$. Since S^- contains either $\frac{p}{2}$ or $\frac{p}{2} + 1$, vertices $\frac{p}{2}$ and $\frac{p}{2} + 1$ disagree on 0. Since 0 is fixed by every automorphism of P_1 , $\frac{p}{2} - 1, \frac{p}{2}$ and $\frac{p}{2} + 1$ are also fixed and thus, by Corollary 21, P_1 is rigid.

(d) If P_1 contains a single plateau of size 5, then the plateau-vertices are necessarily $\frac{p}{2} - 2, \frac{p}{2} - 1, \dots, \frac{p}{2} + 1$, so that $S^- = \{a_1, a_2\}$, with $a_1 \in \{\frac{p}{2} - 1, \frac{p}{2} + 2\}$, $a_2 \in \{\frac{p}{2}, \frac{p}{2} + 1\}$, and $d_{P_1}^-(\frac{p}{2}) = \frac{p}{2}$. Moreover, if $p > 4$, then every set $V_d(P_1)$ except $V_{\frac{p}{2}}(P_1)$ is a singleton. Suppose first that $p > 4$, so that $1 \notin S^-$ (otherwise, this would imply $1 = \frac{p}{2} - 1$). If $\{2, 3\} \cap S^- = \emptyset$ (which happens in particular if $p \geq 10$), the sub-tournament induced by $V_{\frac{p}{2}}(P_1)$ is either transitive or almost transitive, and thus rigid, so that P_1 is rigid by Lemma 23 (this also follows from Theorem 25).

Suppose now that $p \in \{6, 8\}$, so that vertices 0 and p are fixed by every automorphism of P_1 . If all vertices belonging to the plateau are fixed by every automorphism of P_1 , then P_1 is rigid by Lemma 23. Otherwise, some of these vertices may form an orbit, say O , of some automorphism, of size 3 or 5. Considering the structure of the set S^- , we get that the vertex 0 has three out-neighbours and two in-neighbours in $\{\frac{p}{2} - 2, \frac{p}{2} - 1, \frac{p}{2}, \frac{p}{2} + 1, \frac{p}{2} + 2\}$. Therefore, the size of the orbit O must be 3. Moreover, since $0(\frac{p}{2} - x)$ is an arc if and only if $(\frac{p}{2} + x)p$ is an arc for every $x \in \{0, 1, 2\}$, we get that this orbit, if it exists, is necessarily $O = \{\frac{p}{2} - 2, \frac{p}{2}, \frac{p}{2} + 2\}$, and thus $S^- = \{\frac{p}{2} - 1, \frac{p}{2} + 1\}$. Moreover, O must induce a 3-cycle in P_1 , which implies either $2 \in S^-$ and $4 \notin S^-$, or $2 \notin S^-$ and $4 \in S^-$. This forces $S^- = \{4, 6\}$ and thus $\{2, 3\} \cap S^- = \emptyset$ for which we have seen before that P_1 is rigid.

Suppose finally that $p = 4$. Without loss of generality, we can suppose that $1 \notin S^-$ (otherwise, we consider P^c instead of P and we have $|(S^-)^c| = |S^-|$, so that either $S^- = \{2, 4\}$ or $S^- = \{3, 4\}$ and, in both cases, P_1 is a cyclic tournament). Due to our initial assumption on the parity of p , this case occurs either if $T = T(9; S^-)$, in which case $P(4; S^-) = P_1$, or $T = T(11; S^-)$, in which case $P(4; S^-) = P_2$, so that $P_1 = P(5; S^-)$.

In the former case, the pseudo-cyclic tournament P_2 is then either $P(3; \{2\})$ or $P(3; \{3\})$, respectively. In both cases, P_2 is rigid since $|S^-| = 1$.

In the latter case, we will prove that $P = P(5; S^-)$ is rigid. We consider two subcases, according to the set S^- .


i. $S^- = \{2, 4\}$.

In that case, we have $V_2(P) = \{0, 2, 4\}$ and $V_3(P) = \{1, 3, 5\}$. Since both sub-tournaments $P[V_2(P)]$ and $P[V_3(P)]$ are transitive, P is rigid by Lemma 23.

ii. $S^- = \{3, 4\}$.


In that case, we have $V_2(P) = \{0, 3, 4\}$ and $V_3(P) = \{1, 2, 5\}$. Again, both sub-tournaments $P[V_2(P)]$ and $P[V_3(P)]$ are transitive, P is rigid by Lemma 23.

This completes the proof. \square




(a) Four plateaus of size 2

(b) Two plateaus of size 3



(c) Two plateaus of size 2



(d) One plateau of size 5

Figure 3: Indegree paths for the proof of Theorem 26 (Cont.)

For proving Theorem 26, we showed that whenever $|S^-| \leq 2$, at least one of the two subtournaments $T_{0,p}$ or $T_{p+1,2p}$ of the cyclic tournament $T = T(2p+1; S)$ is rigid. It should be noticed that this property does not hold in general when $|S^-| \geq 3$, as shown by the cyclic tournament $T = T(13; \{2, 5, 6\})$ given in Example 9.

Since any tournament T is isomorphic to its converse, Theorem 26 gives the following.

Corollary 27 *If $T = T(2p+1; S)$ is a cyclic tournament, such that $|S^-| \geq p-1$, then T satisfies the Albertson-Collins Conjecture.*

4.3 The set S^- forms an interval

Observe first the following easy result.

Theorem 28 *For every $p \geq 1$, if $S = \{1, \dots, p\}$ or $S = \{1, \dots, p-1, p+1\}$, then the cyclic tournament $T(2p+1; S)$ satisfies the Albertson-Collins Conjecture.*

Proof. This directly follows from Theorem 26 since we have either $|S^-| = 0$ or $|S^-| = 1$. \square

In fact, we can prove that whenever the set S^- of negative connectors forms an interval of integers, the corresponding cyclic tournament $T(2p+1; S)$ satisfies the Albertson-Collins Conjecture.

We first prove two lemmas. The first one says that for every pseudo-cyclic tournament P , if vertices $0, \dots, i-1$ are fixed by every automorphism of P , i is the first vertex of a plateau whose corresponding negative connectors form an interval and every vertex outside the plateau having the same indegree as the vertices of the plateau is fixed by every automorphism of P , then all vertices of the plateau are also fixed by every automorphism of P .

Lemma 29 *Let $P = P(p; N)$ be a pseudo-cyclic tournament, and i a vertex of P with $i < \frac{p}{2}$, satisfying all the following conditions.*

- (i) *every vertex j , $0 \leq j < i$, is fixed by every automorphism of P ,*
- (ii) *$i-1$ is an ascent vertex,*
- (iii) *$(i, i+1, \dots, i+k-1)$ is a plateau of size $k \geq 2$ such that either*
 - (a) *$i+k-1 < \frac{p}{2}$ and either $[i+1, i+k-1] \subseteq N$, or $[p-i-k+2, p-i] \subseteq N$, or*
 - (b) *$i+k-1 > \frac{p}{2}$ and either $[i+1, \frac{p}{2}] \subseteq N$, or $[\frac{p}{2}+1, p-i] \subseteq N$,*
- (iv) *$|V_d(P)| \geq k$, for $d = d_P^-(i)$, and, if $|V_d(P)| > k$, then every vertex in $V_d(P) \setminus \{i, i+1, \dots, i+k-1\}$ is fixed by every automorphism of P .*

Then, every vertex in $\{i, i+1, \dots, i+k-1\}$ is fixed by every automorphism of P .

Proof. Suppose that P and i satisfy the conditions of the lemma. According to condition (iv), it is enough to prove that no two vertices in $\{i, i+1, \dots, i+k-1\}$ can belong to the same orbit of any automorphism of P , since this implies that every such vertex is fixed by every automorphism of P . Observe that $i \notin N$ since $i-1$ is an ascent-vertex.

We consider two cases, according to the position of the plateau.

1. $i+k-1 < \frac{p}{2}$.

We consider two subcases, depending on the corresponding interval of negative connectors.

- (a) $[i+1, i+k-1] \subseteq N$.

In that case, P contains arcs $0i, (i+1)0, \dots, (i+k-1)0$, so that i and all other vertices of the plateau disagree on 0 , and thus, by Observation 3, i and any other vertex of the plateau cannot belong to the same orbit of any automorphism of P . Similarly, for every $j = i+1, \dots, i+k-1$ (in that order), j and vertices $j+1, \dots, i+k-1$ disagree on $j-i$ (which is fixed by every automorphism of P), so that, by Observation 3, j and any other vertex of the plateau cannot belong to the same orbit of any automorphism of P . Therefore, all vertices of the plateau are fixed by every automorphism of P .

- (b) $[p-i-k+2, p-i] \subseteq N$.

If N contains no negative connector $a < i$, then the sub-tournament induced by the plateau is transitive (since we then have $\min(N) > i+k-1$), so that all vertices of the plateau are fixed by every automorphism of P .

Otherwise, let a be the largest negative connector such that $a < i$. We then have $[a+1, i+k-1] \cap N = \emptyset$, so that i and all other vertices of the plateau disagree on $i-a$, and thus, by Observation 3, i and any other vertex of the plateau cannot belong to the same orbit of any automorphism of P . Similarly, for every $j = i+1, \dots, i+k-1$ (in that order), j and vertices $j+1, \dots, i+k-1$ disagree on $j-a$ (which is fixed by every automorphism of P), so that, by Observation 3, j and any other vertex of the plateau cannot belong to the same orbit of any automorphism of P . Therefore, all vertices of the plateau are fixed by every automorphism of P .

2. $i+k-1 > \frac{p}{2}$.

In that case, p is necessarily even. Since $\sigma(P)$ is centrally symmetric, we have $(i, i+1, \dots, i+k-1) = (\frac{p}{2}-q, \dots, \frac{p}{2}+q)$, for some $q \geq 1$.

If $[\frac{p}{2} - q + 1, \frac{p}{2}] \subseteq N$, then, using the same proof as in Case 1a above, we get that no two vertices in $\{\frac{p}{2} - q, \dots, \frac{p}{2}\}$ can belong to the same orbit of any automorphism of P . Symmetrically, starting with vertex p instead of vertex 0 (recall that $p - j$, $j \geq 0$, is fixed whenever j is fixed, by Proposition 20), we can prove similarly that no two vertices in $\{\frac{p}{2}, \dots, \frac{p}{2} + q\}$ can belong to the same orbit of any automorphism of P . Therefore, no vertex of the plateau can belong to an orbit of size at least 3 of any automorphism of P , which implies that every vertex in $\{\frac{p}{2} - q, \dots, \frac{p}{2} + q\}$ is fixed by every automorphism of P and we are done.

If $[\frac{p}{2} + 1, \frac{p}{2} + q] \subseteq N$, then we proceed as in Case 1b above: if the vertices of the plateau do not induce a transitive tournament, then (i) using the largest negative connector $a < \frac{p}{2} - q$, we get that no two vertices in $\{\frac{p}{2} - q, \dots, \frac{p}{2}\}$ can belong to the same orbit of any automorphism of P , and (ii) symmetrically, using the smallest negative connector $a' > \frac{p}{2} + q$, we get that no two vertices in $\{\frac{p}{2}, \dots, \frac{p}{2} + q\}$ can belong to the same orbit of any automorphism of P and thus, again, every vertex in $\{\frac{p}{2} - q, \dots, \frac{p}{2} + q\}$ is fixed by every automorphism of P .

This concludes the proof. \square

The second lemma says that every pseudo-cyclic tournament containing either ascent-vertices and no descent-vertices, or descent-vertices and no ascent-vertices, is rigid whenever every plateau, if any, is produced by an interval of negative connectors.

Lemma 30 *If $P = P(p; N)$ is a pseudo-cyclic tournament such that*

- (i) $\alpha(P) + \delta(P) > 0$,
- (ii) $\alpha(P)\delta(P) = 0$,
- (iii) *for every plateau $(i, i+1, \dots, i+k-1)$ of size $k \geq 3$, with $i+k-1 < \frac{p}{2}$, either $[i+1, i+k-1] \subseteq N$, or $[p-i-k+2, p-i] \subseteq N$, and*
- (iv) *for every plateau $(\frac{p}{2} - q, \dots, \frac{p}{2} + q)$, p even, $q \geq 1$, of size $2q+1$, either $[\frac{p}{2} - q + 1, \frac{p}{2}] \subseteq N$, or $[\frac{p}{2} + 1, \frac{p}{2} + q] \subseteq N$,*

then P is rigid.

Proof. If $\pi(P) = 0$, then P is a transitive tournament and is thus rigid.

Assume now that $\pi(P) > 0$ and suppose first that P has no descent-vertices, that is, $\delta(P) = 0$ and $\alpha(P) > 0$. We will prove by induction on i that every vertex $i \leq \lfloor \frac{p}{2} \rfloor$ is fixed by every automorphism of P , so that the result follows by Corollary 21.

If 0 is an ascent-vertex or the first vertex of a plateau of size 2, then 0 is fixed by every automorphism of P . If 0 is the first vertex of a plateau $(0, \dots, k-1)$ of size $k \geq 3$, then the sub-tournament induced by this plateau is transitive since we have either $[1, k-1] \in N$ or $[p-k+2, p] \in N$. Therefore, all vertices $0, \dots, k-1$ are fixed by every automorphism of P .

Suppose now that all vertices $0, \dots, i-1 < \lfloor \frac{p}{2} \rfloor$ are fixed by every automorphism of P . If i is an ascent-vertex, then i is fixed by every automorphism of P , otherwise i is the first vertex of a plateau and the result follows from Lemma 29.

The case $\alpha(P) = 0$ and $\delta(P) > 0$ follows from Observation 15, considering P^c instead of P (we then have $\delta(P^c) = 0$ and $\alpha(P^c) > 0$). \square

Note that conditions (iii) and (iv) in Lemma 30 are both necessary. For instance, the pseudo-cyclic tournament $P = P(5; \{2, 5\})$ has only ascent-vertices and plateau-vertices (we have $\sigma(P) = (2, 2, 2, 3, 3, 3)$) while $(0, 1, 2)(3, 4, 5)$ is an automorphism of P so that P is not rigid, and thus condition (iii) is necessary. On the other hand, the pseudo-cyclic tournament $P = P(8; \{2, 3, 5\})$ has only ascent-vertices and plateau-vertices, but a single central plateau




Figure 4: Indegree paths for the proof of Theorem 31, case $b - a + 1 > \frac{p}{2}$

(we have $\sigma(P) = (3, 4, \dots, 4, 5)$) while $(1, 4, 7)$ is an automorphism of P so that P is not rigid, and thus condition (iv) is necessary.

We are now able to prove the following result.

Theorem 31 *If $T = T(2p + 1; S)$ is a cyclic tournament, such that $S^- = [a, b]$, $1 \leq a \leq b \leq p$, then T satisfies the Albertson-Collins Conjecture.*

Proof. We will prove that $P = P(p; [a, b])$ is rigid whenever p is even. Therefore, considering either $T_{0,p}$ or $T_{p+1,2p}$ depending on the parity of p , the result will follow by Proposition 8.

If $b \leq a + 1$, the result follows from Theorem 26, so that we can assume $b \geq a + 2$. Suppose now that $\frac{p}{2} \geq b$ or $\frac{p}{2} < a$. If $[a, b] \neq [1, \frac{p}{2}]$ and $[a, b] \neq [\frac{p}{2} + 1, p]$, then $T_{0,p}$ is rigid by Lemma 30. Otherwise, $T_{p+1,2p}$ is rigid by Lemma 30 (since $\sigma(T_{p+1,2p}) = (\frac{p}{2}, \dots, \frac{p}{2}, \frac{p}{2} - 1, \dots, \frac{p}{2} - 1)$ if $[a, b] = [1, \frac{p}{2}]$, and $\sigma(T_{p+1,2p}) = (\frac{p}{2} - 1, \dots, \frac{p}{2} - 1, \frac{p}{2}, \dots, \frac{p}{2})$ if $[a, b] = [\frac{p}{2} + 1, p]$).


From now on, we thus assume that $a \leq \frac{p}{2} < b$. We consider three subcases, depending on the size of $[a, b]$.

1. $b - a + 1 > \frac{p}{2}$.


In that case, $d_P^-(0) = b - a + 1 > p - (b - a + 1) = d_P^-(p)$. We consider three subcases, corresponding to the three possible forms of the indegree path of P .

- (a) $a + b = p + 1$.

In that case, no vertex in P can be a plateau-vertex, and thus the indegree path of P contains an ascent, a descent and an ascent (see Figure 4(a)). Therefore, each set $V_d(P)$ has cardinality at most 2 and P is rigid by Lemma 23.



(a) $a + b = p + 1$



(b) $a + b \neq p + 1$

Figure 5: Indegree paths for the proof of Theorem 31, case $b - a + 1 = \frac{p}{2}$

(b) $a = 1$ or $b = p$.

In that case, vertices $0, \dots, p - b - 1$ (if $a = 1$) or $0, \dots, a - 1$ (if $p = b$) are plateau-vertices, and thus the indegree path of P contains a plateau of size $p - b + 1$ or $a + 1$, a descent and a plateau of size $p - b + 1$ or $a + 1$ (see Figure 4(b) for the case $a = 1$). In both cases, the sub-tournaments induced by these plateaus are transitive, so that every vertex of P is fixed by every automorphism of P by Observation 5, and P is rigid by Corollary 21.


(c) $1 < a$, $b < p$ and $a + b \neq p + 1$.

In that case, the indegree path of P contains an ascent, a plateau, a descent, a plateau and an ascent (see Figure 4(c)). Every vertex not belonging to a plateau belongs to a set $V_d(P)$ with cardinality 2 and is thus fixed by every automorphism of P . By Lemma 29, every vertex belonging to a plateau is also fixed by every automorphism of P . Using now Corollary 21, we get that P is rigid.


2. $b - a + 1 = \frac{p}{2}$.

Note first that since $a \leq \frac{p}{2} < b$, we necessarily have $a > 1$ and $b < p$. The indegree path of P contains an ascent, possibly a plateau, a descent, possibly a plateau and an ascent. More precisely, $\sigma(P)$ contains a plateau if and only if $a + b \neq p + 1$ (see Figure 5(a) and (b)). Moreover, since $d_P^-(0) = d_P^-(p) = \frac{p}{2}$, we get $V_{\frac{p}{2}}(P) = \{0, \frac{p}{2}, p\}$ (and the sub-tournament induced by $V_{\frac{p}{2}}(P)$ is a 3-cycle). Only two other sets $V_d(P)$ may have cardinality at least 3, namely those two sets corresponding to the two plateaus.

We consider three subcases, depending on the values of a and b .



(a) $a + b = p + 1$ and $p - b \geq b - a + 1$



(b) $a + b = p + 1$ and $p - b < b - a + 1$

Figure 6: Indegree paths for the proof of Theorem 31, case $b - a + 1 < \frac{p}{2}$

- (a) $a > 2$ and $b < p - 1$.

We first prove that every vertex in $V_{\frac{p}{2}}(P) = \{0, \frac{p}{2}, p\}$ is fixed by every automorphism of P . Since the vertex 1 is an ascent-vertex with $d_P^-(1) = \frac{p}{2} + 1$ and $|V_{\frac{p}{2}+1}| = 2$, it is fixed by every automorphism of P . Since $p - 1 \notin [a, b]$, vertices 0 and p disagree on 1, so that, by Observation 3, 0 and p cannot belong to the same orbit of any automorphism of P and we are done.

Moreover, by Lemma 29, all vertices of the first plateau (whenever $a + b \neq p + 1$) are fixed by every automorphism of P . Hence, all vertices $i \leq \frac{p}{2}$ are fixed by every automorphism of P and the result follows by Corollary 21.

- (b) $b = p - 1$.

In that case, we thus have $a = \frac{p}{2}$. The first plateau contains vertices 1 to $\frac{p}{2} - 1$ and thus induces a transitive tournament, so that all its vertices are fixed by every automorphism of P . Now, since vertices $\frac{p}{2}$ and p disagree on $\frac{p}{2} - 1$, the three vertices $\{0, \frac{p}{2}, p\}$ are all fixed by every automorphism of P and the result follows by Corollary 21.

- (c) $a = 2$.

In that case, we thus have $b = \frac{p}{2} + 1$. The first plateau contains vertices 1 to $\frac{p}{2} - 1$ and is isomorphic to the pseudo-cyclic tournament $PC = P(\frac{p}{2} - 2; [2, \frac{p}{2} - 2])$. Since PC^c is rigid by Theorem 26, PC is rigid. Now, since vertices 0 and p disagree on 1, the three vertices $\{0, \frac{p}{2}, p\}$ are all fixed by every automorphism of P and, again, the result follows by Corollary 21.

3. $b - a + 1 < \frac{p}{2}$.

The indegree path of P contains an ascent, possibly a plateau (if $a \neq p + 1 - b$), a descent, possibly a plateau and an ascent. If $a \leq p + 1 - b$, the first vertex of the first plateau is $a - 1$ and its last vertex is $p - b$ (so that there is no plateau if $a = p + 1 - b$). If $a > p + 1 - b$, the first vertex of the first plateau is $p - b$ and its last vertex is $a - 1$.

We will prove that every vertex of P is fixed by every automorphism of P . Note that $d_P^-(0) = b - a + 1$, so that $d_P^-(p) = p - b + a - 1 > d_P^-(0)$ since $b - a + 1 < \frac{p}{2}$.

We consider two subcases, depending on whether $\sigma(P)$ contains a plateau or not.

(a) $a = p + 1 - b$.

In that case, $\sigma(P)$ does not contain any plateau, the first vertex of the descent is $a - 1$ and its last vertex is b . Moreover, $d_P^-(a - 1) = b$ and $d_P^-(b) = p - b$. Let us consider the vertex b . If $d_P^-(b) = p - b \geq b - a + 1 = d_P^-(0)$ (see Figure 6(a)), then $|V_{p-b}(P)| = 2$ and thus b is fixed by every automorphism of P . If $p - b > b - a + 1$ (see Figure 6(b)), then $|V_{p-b}(P)| = 1$ and thus, again, b is fixed by every automorphism of P .

Now, we claim that every set $V_d(P)$ with $|V_d(P)| = 3$ contains three fixed vertices. Each such set is of the form $\{i, j, k\}$, $i < j < k$, with $j + k = 2b$. Then, j and k disagree on b (since $b - j = k - b < p - b$, so that jb and bk are arcs in P) and, as b is fixed by every automorphism of P , by Observation 3, i, j and k are also fixed by every automorphism of P . The result then follows by Corollary 21.

(b) $a \neq p + 1 - b$.

In that case, the vertex $a - 1$ is either the first or the last vertex of the first plateau and $d_P^-(a - 1) = b$ or $d_P^-(a - 1) = p - a + 1$, respectively. Similarly, the vertex b is either the first or the last vertex of the last plateau and $d_P^-(b) = p - b$ or $d_P^-(b) = a - 1$, respectively.

We first claim that whenever 0 is fixed by every automorphism of P , all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P . Indeed, for any such set $\{i, j, k\}$ with $i < j < k$, we have $i < a - 1$ and $a - 1 < j < b$, so that i and j disagree on 0. Therefore, i, j and k are all fixed by every automorphism of P .

We now consider three subcases, depending on the values of $d_P^-(b)$ and $d_P^-(0)$.

i. $d_P^-(b) > d_P^-(0)$.

In that case (see Figure 7(a)), the vertex 0 is fixed by every automorphism of P , so that all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P .

If $a - 1$ is the first vertex of the first plateau, let $x > \frac{p}{2}$ be the unique vertex with $d_P^-(x) = d_P^-(a - 1)$. We then have $V_b(P) = \{a - 1, a, \dots, p - b, x\}$. Since $a - 1$ and x are the only in-neighbours of 0 in $V_b(P)$, both of them are fixed by every automorphism of P . Therefore, by Lemma 29, every vertex belonging to the first plateau is fixed by every automorphism of P .

Suppose now that $a - 1$ is the last vertex of the first plateau. Similarly, let $x < \frac{p}{2}$ be the unique vertex with $d_P^-(x) = d_P^-(b)$, so that $V_{a-1}(P) = \{x, p - a + 1, \dots, b\}$. Since x is the unique out-neighbour of 0 in $V_{a-1}(P)$ (as $x < a$ and $p - a + 1 > a$, which gives $V_{a-1}(P) \cap [a, b] = \{p - a + 1, \dots, b\}$), x is fixed by every automorphism of P and, by Proposition 20, $p - x$ is also fixed by every automorphism of P . Therefore, by Lemma 29, every vertex belonging to the first plateau is fixed by every automorphism of P .

The result then follows by Corollary 21.

ii. $d_P^-(b) = d_P^-(0)$.

Suppose first that $a - 1$ is the last vertex of the first plateau (that is, $a > p + 1 - b$), so that b is the last vertex of the second plateau. In that case, the

vertices of the second plateau induce a transitive tournament and they all agree on 0. Therefore, the sub-tournament induced by $V_{b-a+1}(P)$ is rigid, so that all its vertices, and in particular 0, are fixed by every automorphism of P . Hence, all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P . By Proposition 20, all vertices of the first plateau are also fixed by every automorphism of P and, again, the result follows by Corollary 21.

Suppose now that $a-1$ is the first vertex of the first plateau (that is, $a < p+1-b$), so that b is the first vertex of the second plateau (see Figure 7(b)).

We first prove that the vertex 0 is fixed by every automorphism of P . Suppose to the contrary that 0 belongs to an orbit of size at least 3 of some automorphism ϕ of P . Among the vertices of the second plateau, b is the only in-neighbour of 0. Therefore, this orbit must be of size 3 (recall that, by Observation 5, the sub-tournament induced by an orbit must be regular) and thus a 3-cycle $b0x$. Note that since xb is an arc of P , we then have $x-b \geq a$. Consider now the set $V_{b-a+2} = \{1, b-1, p-a+2\}$. The vertex 0 has only one in-neighbour in V_{b-a+2} , namely $b-1$, since $b-1 \in [a, b]$, $1 \notin [a, b]$ and $p-a+2 \notin [a, b]$ (since $p-a+2 > b$). On the other hand, b has at least two in-neighbours in V_{b-a+2} , namely 1 (since $b-1 \in [a, b]$) and $p-a+2$ (since $p-a+2-b > x-b \geq a$ and $p-a+2-b < \frac{p}{2} < b$). By Observation 4, this implies that V_{b-a+2} cannot be an orbit of ϕ , so that, in particular, 1 is fixed by ϕ . Now, note that 1 is an in-neighbour of x (since $x-1 \geq b+a-1 > b$) and an out-neighbour of 0, a contradiction since, by Observation 3, all vertices in $\{0, b, x\}$ should agree on every vertex fixed by ϕ .

Therefore, 0 is fixed by every automorphism of P , so that all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P . By Proposition 20, p is also fixed by every automorphism of P and thus, by Lemma 29, every vertex of the first plateau is fixed by every automorphism of P . The result then follows by Corollary 21.


iii. $d_P^-(b) < d_P^-(0)$.

Suppose first that $a-1$ is the last vertex of the first plateau (that is, $a > p+1-b$), so that b is the last vertex of the second plateau (see Figure 7(c)). In that case, the vertices of each plateau induce a transitive tournament and thus all vertices of these plateaus are fixed by every automorphism of P . Consider the set $V_{b-a+1} = \{0, x, y\} = \{0, p-b+a-1, 2b-2a+2\}$. Vertices 0 and $p-b+a-1$ disagree on $p-a+1$ (since $p-a+1 \in [a, b]$ and $q = p-a+1 - (p-b+a-1) < p-b < a-1$ so that $q \notin [a, b]$), which is fixed by every automorphism of P , and thus 0, $p-b+a-1$ and $2b-2a+2$ are all fixed by every automorphism of P .


Therefore, all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P , and the result follows by Corollary 21.

Suppose now that $a-1$ is the first vertex of the first plateau (that is, $a < p+1-b$), so that b is the first vertex of the second plateau.


We first prove that the vertex 0 is fixed by every automorphism of P . Suppose to the contrary that 0 belongs to an orbit of size 3 of some automorphism ϕ of P , say a 3-cycle $0xy$, with $a < x < b$ and $y > b+1$, so that $x0$, $0y$ and yx are arcs in P , implying in particular $a \leq y-i \leq b$. Consider now the set $V_{b-a+2} = \{1, x-1, y+1\}$. Since $1 < a$, $a \leq x-2$ (as $a-1 < \frac{p}{2} < x$) and $x < b$, we get that 01 , $x1$, $(x-1)0$ and $(x-1)x$ are all arcs in P , so that 1 and $x-1$ disagree on the orbit of 0 with respect to ϕ . This implies that V_{b-a+2} cannot be an orbit of ϕ , so that, in particular, 1 is fixed by ϕ . Now, note that 1 is an



(a) $a + b \neq p + 1$ and $p - b > b - a + 1$



(b) $a + b \neq p + 1$, $p - b = b - a + 1$ and $a < p + 1 - b$



(c) $a + b \neq p + 1$, $a - 1 < b - a + 1$ and $a > p + 1 - b$

Figure 7: Indegree paths for the proof of Theorem 31, case $b - a + 1 < \frac{p}{2}$ (Cont.)

in-neighbour of y (since $y - 1 > b$) and an out-neighbour of 0, a contradiction since all vertices in $\{0, x, y\}$ should agree on every vertex fixed by ϕ .

Therefore, 0 is fixed by every automorphism of P , so that all vertices belonging to a set $V_d(P)$ with $|V_d(P)| = 3$ are also fixed by every automorphism of P . By Lemma 29, every vertex of the first plateau is fixed by every automorphism of P and the result follows by Corollary 21.

This concludes the proof. \square

Again, since any tournament T is isomorphic to its converse, Theorem 31 gives the following.

Corollary 32 *If $T = T(2p + 1; S)$ is a cyclic tournament, such that $S^- = [1, a] \cup [b, p]$, $1 \leq a \leq b \leq p$, then T satisfies the Albertson-Collins Conjecture.*

4.4 Paley tournaments

Let n be a prime, $n \equiv 3 \pmod{4}$, so that $n = 2p + 1$ with p odd. The *Paley tournament* is the cyclic tournament $QR_n = T(n; S_n)$ where S_n is the set of non-zero quadratic residues of n . That is, for every i, j , $i < j$, ij is an arc in QR_n if and only if $j - i$ is a non-zero square in \mathbb{Z}_n . Paley tournaments are arc-transitive and the set $\text{Aut}(QR_n)$ is the set of mappings $i \mapsto ai + b$, where $a, b \in \mathbb{Z}_n$ and a is a non-zero square. Since the number of non-zero squares in \mathbb{Z}_n is $p = \lfloor \frac{n}{2} \rfloor$, we thus have $|\text{Aut}(QR_n)| = np$.

In [3], Albertson and Collins show that the Paley tournament QR_7 satisfies their conjecture. We prove here that all Paley tournaments satisfy the Albertson-Collins Conjecture.

Theorem 33 *For every prime n , $n \equiv 3 \pmod{4}$, QR_n satisfies the Albertson-Collins Conjecture.*

Proof. Suppose to the contrary that there exists a nontrivial λ^* -preserving automorphism $\phi : i \mapsto ai + b$, with $a, b \in \mathbb{Z}_n$ and a is a non-zero quadratic residue and let ℓ denote the order of ϕ .

Consider the action on $V(QR_n)$ of the group $H = \langle \phi \rangle = \{\phi^k : 1 \leq k \leq \ell\}$. Note that $|H|$ divides $|\text{Aut}(QR_n)| = np$ and, since ϕ is λ^* -preserving, $|H|$ divides p . Moreover, p cannot be prime, since otherwise the sub-tournament T_2 of QR_n induced by the set of vertices $\{p+1, \dots, 2p\}$ would have a single orbit with respect to ϕ , in contradiction with Proposition 10.

We claim that we necessarily have $b = 0$. Assume to the contrary that $b \neq 0$ and let r be the size of the orbit of 0 with respect to ϕ . We have $\phi(0) = b$, $\phi^2(0) = ab + b$ and so on, so that

$$\phi^r(0) = a^{r-1}b + a^{r-2}b + \dots + ab + b = 0$$

and $\phi^{r+1}(0) = \phi(0) = b$. On one other hand, we have

$$\phi^{r+1}(0) = a^r b + a^{r-1}b + \dots + ab + b = a^r b.$$

Therefore, $a^r b = b$ and thus $a^r = 1$. Since a is a non-zero quadratic residue in \mathbb{Z}_n , its order in the multiplicative group \mathbb{Z}_n^* is p and thus $p | r$. Moreover, since ϕ is λ^* -preserving, we have $r = p$. Hence, the sub-tournament T_1 of QR_n induced by the set of vertices $\{0, \dots, p\}$ contains a fixed vertex x such that all vertices of the orbit of 0 with respect to ϕ agree on x , which implies either $d_{T_1}^+(x) = 0$ or $d_{T_1}^+(x) = p$ in contradiction with the definition of QR_n since both sets $\{1, \dots, p\}$ and $\{p+1, \dots, 2p\}$ contain non-zero quadratic residues.

We thus have $b = 0$, so that 0 is fixed by ϕ . Moreover, the orbit of 1 with respect to ϕ is $\{1, a, a^2, \dots, a^{p-1}\}$ (recall that the order of a in the multiplicative group \mathbb{Z}_n^* is p) and its size is p . Since ϕ is λ^* -preserving, the orbit of 1 with respect to ϕ is thus $\{1, \dots, p\}$. Therefore, all vertices in $\{1, \dots, p\}$ agree on 0 and we obtain a contradiction as above. \square

5 Discussion

In this paper, we proposed several sufficient conditions for a cyclic tournament to satisfy the Albertson-Collins Conjecture. Proposition 8 says that the cyclic tournament $T = T(2p + 1; S)$ satisfies the Albertson-Collins Conjecture whenever any of the sub-tournaments $T_{0,p}$ or $T_{p+1,2p}$ is rigid (however, this condition is not necessary, as shown by Example 9). Using this property, we then got several sufficient conditions on the structure of the set S^- of negative connectors for at least one of these two sub-tournaments to be rigid.

We finally propose some directions for future research.

1. The main question following our work is obviously to prove, or disprove, the Albertson-Collins Conjecture.
2. In [8], Boutin introduces the notion of a *determining set* of a graph G , that is, a subset X of $V(G)$ such that, for every two automorphisms $\phi_1, \phi_2 \in \text{Aut}(G)$, $\phi_1 = \phi_2$ whenever $\phi_1(x) = \phi_2(x)$ for every vertex $x \in X$ (this notion was independently introduced by Erwin and Harary in [13], under the name of *fixing set*). Albertson and Boutin prove in [2] that $D(G) \leq d$ if and only if G has a determining set X with $D(X) \leq d - 1$. Therefore $D(G) = 2$ if and only if the subgraph of G induced by X is rigid.

For each cyclic tournament T of order $2p + 1$ for which we proved the Albertson-Collins Conjecture, we exhibited a rigid determining set of size at most $\lceil \frac{p}{2} \rceil$. It is not difficult to see that every Paley tournament has a rigid determining set of size 2 (indeed, every pair of vertices is a rigid determining set). In [23], Lozano proves that every tournament contains a (not necessarily rigid) determining set of size $\lfloor \frac{p}{3} \rfloor$.

It would thus be interesting to determine the minimal size of a rigid determining set in a cyclic tournament.

3. In [9, 10], Boutin also defines the *cost of distinguishing a graph* G such that $D(G) = 2$, denoted $\rho(G)$, as the minimum size of a label class in a distinguishing 2-labeling of G . It is easy to prove that $\rho(QR_n) = 2$ for every Paley tournament QR_n (indeed, for every two vertices i and j in QR_n , the 2-labeling $\lambda_{i,j}$ defined by $\lambda_{i,j}(v) = 1$ if and only if $v = i$ or $v = j$, is distinguishing). It would be interesting to determine $\rho(T)$ for every cyclic tournament $T = T(2p + 1; S)$, or to characterize cyclic tournaments T with $\rho(T) = 2$.

References

- [1] Michael O. Albertson. Distinguishing Cartesian powers of graphs. *Electron. J. Combin.* 12 (2005), #N17
- [2] Michael O. Albertson and Debra L. Boutin. Using Determining Sets to Distinguish Kneser Graphs. *Electron. J. Combin.* 14 (2007), #R20.
- [3] Michael O. Albertson and Karen L. Collins. A Note on Breaking the Symmetries of Tournaments. *Proc. 13th Southeastern Int. Conf. on Combinatorics, Graph Theory, and Computing. Congr. Numer.* 136 (1999), 129–131.
- [4] Michael O. Albertson and Karen L. Collins. Symmetry Breaking in Graphs. *Electron. J. Combin.* 3(1) (1996), #R18.
- [5] V. Arvind, Christine T. Cheng and Nikhil R. Devanur. On computing the distinguishing numbers of planar graphs and beyond: a counting approach. *SIAM J. Discrete Math.* 22(4) (2008), 1297–1324.
- [6] Bill Bogstad and Lenore J. Cowen. The distinguishing number of the hypercube. *Discrete Math.* 283(1–3) (2004), 29–35.

- [7] Anthony Bonato and Dejan Delić. Distinguishing number and adjacency properties. *J. Combin.* 1(2) (2010), 141–148.
- [8] Debra L. Boutin. Identifying graph automorphisms using determining sets. *Electron. J. Combin.* 13 (2006), #R78.
- [9] Debra L. Boutin. The Cost of 2-Distinguishing Cartesian Powers. *Electron. J. Combin.* 20(1) (2013), #P74.
- [10] Debra L. Boutin. The Cost of 2-Distinguishing Selected Kneser Graphs and Hypercubes. *J. Combinatorial Math. and Combinatorial Comput.* 85 (2013), 161–171.
- [11] Christine T. Cheng. On computing the distinguishing numbers of trees and forests. *Electron. J. Combin.* 13 (2006), #R11.
- [12] Christine T. Cheng. On computing the distinguishing and distinguishing chromatic numbers of interval graphs and other results. *Discrete Math.* 309(16) (2009), 5169–5182.
- [13] David Erwin and Frank Harary. Destroying automorphisms by fixing nodes. *Discrete Math.* 306(24) (2006), 3244–3252.
- [14] Ehsan Estaji, Wilfried Imrich, Rafal Kalinowski, Monika Piłśniak and Thomas Tucker. Distinguishing Cartesian products of countable graphs. *Discuss. Math. Graph Theory* (2016), in press.
- [15] Michael Fisher and Garth Isaak. Distinguishing colorings of Cartesian products of complete graphs. *Discrete Math.* 308 (2008), 2240–2246.
- [16] David Gluck. Trivial set-stabilizers in finite permutation groups. *Can. J. Math.* 35(1) (1983), 59–67.
- [17] Wilfried Imrich, Janja Jerebic and Sandi Klavžar. The distinguishing number of Cartesian products of complete graphs. *European J. Combin.* 29(4) (2008), 922–929.
- [18] Wilfried Imrich and Sandi Klavžar. Distinguishing Cartesian powers of graphs. *J. Graph Theory* 53(3) (2006), 250–260.
- [19] Wilfried Imrich, Sandi Klavžar and Vladimir Trofimov. Distinguishing infinite graphs. *Electron. J. Combin.* 14 (2007), #R36.
- [20] Sandi Klavžar and Xuding Zhu. Cartesian powers of graphs can be distinguished by two labels. *European J. Combin.* 28(1) (2007), 303–310.
- [21] Claude Laflamme, L. Nguyen Van Thé and Norbert Sauer. Distinguishing Number of Countable Homogeneous Relational Structures. *Electron. J. Combin.* 17 (2010), #R20.
- [22] Andrew Lazowski and Stephen M. Shea. Finite Factors of Bernoulli Schemes and Distinguishing Labelings of Directed Graphs. *Electron. J. Combin.* 19 (2012), #P1.
- [23] Antoni Lozano. Symmetry Breaking in Tournaments. *Electron. J. Combin.* 20(1) (2013), #P69.
- [24] Hiroshi Matsuyama. Another Proof of Gluck’s Theorem. *J. Algebra* 247 (2002), 703–706.
- [25] Simon M. Smith and Mark E. Watkins. Bounding the distinguishing number of infinite graphs and permutation groups. *Electron. J. Combin.* 21(3) (2014), #P3.40.
- [26] Mark E. Watkins and Xiangqian Zhou. Distinguishability of Locally Finite Trees. *Electron. J. Combin.* 14 (2007), #R29.