



HAL
open science

SUB-EXPONENTIAL LOWER BOUNDS FOR EMBEDDED EIGENFUNCTIONS OF SOME DISCRETE SCHRÖDINGER OPERATORS

Marc-Adrien Mandich

► **To cite this version:**

Marc-Adrien Mandich. SUB-EXPONENTIAL LOWER BOUNDS FOR EMBEDDED EIGENFUNCTIONS OF SOME DISCRETE SCHRÖDINGER OPERATORS. 2016. hal-01353783v2

HAL Id: hal-01353783

<https://hal.science/hal-01353783v2>

Preprint submitted on 19 Sep 2016 (v2), last revised 24 Apr 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SUB-EXPONENTIAL LOWER BOUNDS FOR EMBEDDED EIGENFUNCTIONS OF SOME DISCRETE SCHRÖDINGER OPERATORS

MANDICH, MARC-ADRIEN

ABSTRACT. Following the method of Froese and Herbst, we show for a class of potentials V that an embedded eigenfunction ψ with eigenvalue E of the multi-dimensional discrete Schrödinger operator $H = \Delta + V$ on \mathbb{Z}^d decays sub-exponentially whenever the Mourre estimate holds at E . In the one-dimensional case we further show that this eigenfunction decays exponentially with a rate at least of $\cosh^{-1}((E - 2)/(\theta_E - 2))$, where θ_E is the nearest threshold of H located between E and 2. A consequence of the latter result is the absence of eigenvalues between 2 and the nearest threshold to 2 on either side.

1. INTRODUCTION

The analysis of the decay rate of eigenfunctions of Schrödinger operators goes back to the famous works of Slaggie and Wichmann [SW], Agmon [A], and Combes and Thomas [CT]. Their results showed that eigenfunctions corresponding to eigenvalues located outside the essential spectrum decay exponentially. Subsequently, Froese and Herbst [FH], but also [FHHO1] and [FHHO2], investigated what happened for eigenfunctions corresponding to eigenvalues located in the essential spectrum of Schrödinger operators. They showed that eigenfunctions of the continuous Schrödinger operator on \mathbb{R}^n decay exponentially at non-threshold energies for a large class of potentials. Since their pioneering work a solid literature has grown using these ideas. For example, these ideas have been applied to Schrödinger operators on manifolds [V], Schrödinger operators in PDE's [HS], and self-adjoint operators in Mourre Theory [FMS]. This short list of examples is by no means complete. The question however does not seem to have been investigated for the discrete Schrödinger operator and constitutes the subject of this paper.

We now describe the mathematical setup of the article. The configuration space is the multi-dimensional lattice \mathbb{Z}^d for some integer $d \geq 1$. For a multi-index $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$, we set $|n|^2 := n_1^2 + \dots + n_d^2$. Consider the complex Hilbert space $\mathcal{H} := \ell^2(\mathbb{Z}^d)$ of square summable sequences $(u(n))_{n \in \mathbb{Z}^d}$. The discrete Schrödinger acting on \mathcal{H} is

$$(1.1) \quad H := \Delta + V,$$

where Δ is the non-negative discrete Laplacian defined by

$$(\Delta u)(n) := \sum_{\substack{m \in \mathbb{Z}^d, \\ |n-m|=1}} (u(n) - u(m)), \quad \text{for all } n \in \mathbb{Z}^d, u \in \mathcal{H},$$

and V is a multiplication operator by a bounded real-valued sequence $(V(n))_{n \in \mathbb{Z}^d}$. Let us denote by $\tau_i V$ and $\tau_i^* V$ the operators of multiplication by the shifted sequence $(V(n))_{n \in \mathbb{Z}^d}$ to the right and to left respectively on the i^{th} coordinate, namely

$$[(\tau_i V)u](n) := V(n_1, \dots, n_i - 1, \dots, n_d)u(n), \quad \text{for all } n \in \mathbb{Z}^d, u \in \mathcal{H}, \text{ and } i = 1, \dots, d,$$

2010 *Mathematics Subject Classification.* 39A70, 81Q10, 47B25, 47A10, 35P99.

Key words and phrases. discrete Schrödinger operator, decay eigenfunctions, embedded eigenvalues, absence eigenvalues, Mourre theory, Mourre estimate.

and correspondingly for τ_i^*V . In addition to V bounded, two hypotheses on the potential appear in this manuscript :

- Hypothesis 1: V satisfies

$$(1.2) \quad \max_{1 \leq i \leq d} \sup_{n \in \mathbb{Z}^d} |n_i(V - \tau_i V)(n)| < \infty.$$

- Hypothesis 2: The potential V is compact, i.e.

$$(1.3) \quad V(n) \rightarrow 0, \quad \text{as } |n| \rightarrow \infty.$$

The method of Froese and Herbst that we employ works in conjunction with the Mourre estimate, which we now briefly recall in order to present the results. The Mourre estimate is the key relation in the theory developed by Mourre [Mo]. We refer to [ABG] and references therein for a thorough overview of the improved theory. The position operator $N = (N_1, \dots, N_d)$ is defined by

$$(1.4) \quad (N_i u)(n) := n_i u(n), \quad \mathcal{D}(N_i) := \left\{ u \in \ell^2(\mathbb{Z}^d) : \sum_{n \in \mathbb{Z}^d} |n_i u(n)|^2 < \infty \right\},$$

and the shift operators S_i and S_i^* to the right and to left respectively act on \mathcal{H} by

$$(1.5) \quad (S_i u)(n) := u(n_1, \dots, n_i - 1, \dots, n_d), \quad \text{for all } n \in \mathbb{Z}^d \text{ and } u \in \mathcal{H},$$

and correspondingly for S_i^* . The shift operators allow us to rewrite the Laplacian as $\Delta = \sum_{i=1}^d (2 - S_i^* - S_i)$. The generator of dilations A is the closure of the operator A_0 given by

$$(1.6) \quad A_0 := i \sum_{i=1}^d 2^{-1}(S_i^* + S_i) - (S_i^* - S_i)N_i = -i \sum_{i=1}^d 2^{-1}(S_i^* + S_i) + N_i(S_i^* - S_i)$$

with domain $\mathcal{D}(A_0) = \ell_0(\mathbb{Z}^d)$, the collection of sequences with compact support. It is well-known that A is a self-adjoint operator, see e.g. [GG0]. Let T be an arbitrary bounded self-adjoint operator on \mathcal{H} . If the form

$$(u, v) \mapsto \langle u, [T, A]v \rangle := \langle Tu, Av \rangle - \langle Au, Tv \rangle$$

defined on $\mathcal{D}(A) \times \mathcal{D}(A)$ extends to a bounded form on $\mathcal{H} \times \mathcal{H}$, we denote by $[T, A]_\circ$ the bounded operator extending the form, and say that T is of class $C^1(A)$ (cf. [ABG][Lemma 6.2.9]). We refer the reader to [ABG][Theorem 6.2.10] for equivalent definitions of this class. We have that

$$(1.7) \quad [\Delta, iA]_\circ = \sum_{i=1}^d \Delta_i(4 - \Delta_i) = \sum_{i=1}^d (2 - (S_i^*)^2 - (S_i)^2)$$

is a non-negative operator. As for the commutator between V and A , we have for $u, v \in \ell_0(\mathbb{Z}^d)$,

$$\langle u, [V, iA]v \rangle = \sum_{i=1}^d \langle u, [(N_i - 2^{-1})(V - \tau_i V)S_i + (N_i - 2^{-1})(V - \tau_i^*)S_i^*]v \rangle.$$

Assuming V to be bounded, note that $[V, iA]_\circ$ exists if and only if Hypothesis 1 holds. Assuming $[H, iA]_\circ$ to exist, we say that the Mourre estimate holds at $\lambda \in \mathbb{R}$ if there exists an open interval Σ containing λ , a constant $c > 0$ and a compact operator K such that

$$(1.8) \quad E_\Sigma(H)[H, iA]_\circ E_\Sigma(H) \geq cE_\Sigma(H) + K,$$

in the form sense on $\mathcal{H} \times \mathcal{H}$. Here $E_\Sigma(H)$ is the spectral projector of H onto the interval Σ . Denote $\Theta(H)$ the set of points where a Mourre estimate (1.8) holds for H with respect to A . In other words, $\mathbb{R} \setminus \Theta(H)$ is the set of *thresholds* of H . Finally, define for $(\alpha, \gamma) \in [0, \infty) \times [0, 1]$, the following operator of multiplication on \mathcal{H} :

$$(1.9) \quad \vartheta_{\alpha, \gamma} := \exp\left(\alpha(1 + |N|^2)^{\gamma/2}\right), \quad \text{with domain}$$

$$\mathcal{D}(\vartheta_{\alpha,\gamma}) := \left\{ u \in \mathcal{H} : \sum_{n \in \mathbb{Z}^d} \exp\left(2\alpha(1+|n|^2)^{\gamma/2}\right) |u(n)|^2 < \infty \right\}.$$

In this manuscript, we will say that $\psi \in \mathcal{H}$ decays sub-exponentially (resp. exponentially) if $\psi \in \mathcal{D}(\vartheta_{\alpha,\gamma})$ for some $\gamma < 1$ (resp. for $\gamma = 1$) and some $\alpha > 0$. Write $\vartheta_\alpha := \vartheta_{\alpha,1}$. The main result of the paper concerning the one-dimensional operator H is :

Theorem 1.1. *Assume Hypotheses 1 and 2, and $d = 1$. If $H\psi = E\psi$ with $\psi \in \ell^2(\mathbb{Z})$, then if*

$$(1.10) \quad \theta_E := \begin{cases} \sup \{2 + (E - 2)/\cosh \alpha : \alpha \geq 0 \text{ and } \psi \in \mathcal{D}(\vartheta_\alpha)\}, & \text{for } E < 2 \\ \inf \{2 + (E - 2)/\cosh \alpha : \alpha \geq 0 \text{ and } \psi \in \mathcal{D}(\vartheta_\alpha)\}, & \text{for } E > 2, \end{cases}$$

one has that either $\theta_E \in \mathbb{R} \setminus \Theta(H)$ or $\theta_E = 2$. If $E = 2$, the statement is that either $\psi \in \mathcal{D}(\vartheta_\alpha)$ for all $\alpha \geq 0$ or $2 \in \mathbb{R} \setminus \Theta(H)$. Moreover, if $\psi \in \mathcal{D}(\vartheta_\alpha)$ for all $\alpha \geq 0$, then $\psi = 0$.

Remark 1.1. *The function $\mathbb{R}^+ \ni \alpha \mapsto \theta_E(\alpha) := 2 + (E - 2)/\cosh(\alpha) \in [E, 2)$ is strictly increasing to two when $E < 2$ so that $E \leq \theta_E \leq 2$, whereas the function is strictly decreasing to two when $E > 2$ in which case $E \geq \theta_E \geq 2$.*

If E is both an eigenvalue and a threshold, the above result does not give any information about the rate of decay of the corresponding eigenfunction, whereas if E is not a threshold, the corresponding eigenfunction decays at a rate at least of $\cosh^{-1}((E - 2)/(\theta_E - 2))$. As in the continuous operator setting, the possibility of $\psi \in \mathcal{D}(\vartheta_\alpha)$ for all $\alpha \geq 0$ can be eliminated. The last part of the Theorem implies the absence of eigenvalues in the middle of the band $\sigma(\Delta) = [0, 4]$, i.e. between 2 and the nearest threshold to 2 on either side this value. The absence of eigenvalues has already been shown in similar contexts. For example, in [S] a one-dimensional discrete version of Weidmann's Theorem is proven, namely if V is compact and of bounded variation, then the spectrum of $\Delta + V$ is purely absolutely continuous on $(0, 4)$. Another closely related result is that of [JS] where it is shown that the spectrum of the half-line discrete Schrödinger operator $\Delta + W + V$ is purely absolutely continuous on $(0, 4) \setminus \{2 \pm 2 \cos(k/2)\}$, where $W(n) = q \sin(kn)/n^\beta$ with $q, k \in \mathbb{R}, \beta \in (1/2, 1]$ and $(V(n)) \in \ell^1(\mathbb{Z})$. Note that an application of Theorem 1.1 is in conformity with their example when $\beta = 1$ and $V \equiv 0$.

The main result concerning the multidimensional discrete Schrödinger operator H is:

Theorem 1.2. *Let $d \geq 1$. Suppose that Hypothesis 1 holds for the potential V . If $H\psi = E\psi$ with $\psi \in \ell^2(\mathbb{Z}^d)$ and $E \in \Theta(H)$, then $\psi \in \mathcal{D}(\vartheta_{\alpha,\gamma})$ for all $(\alpha, \gamma) \in [0, \infty) \times [0, 2/3)$.*

Although Theorem 1.2 does not yield exponential decay of eigenfunctions, the result is useful for applications in Mourre theory. It appears that the method of Froese and Herbst adapts quite well for the one-dimensional discrete operator; however, there seems to be a non-trivial difference between the dimensions $d \geq 2$ and $d = 1$ in the discrete setting as far as the method is concerned. The exponential decay of eigenfunctions in higher dimensions remains an open question because our proof does not attain it.

From a perspective of Mourre theory and in an abstract setting, a related question is to show that the eigenfunction $\psi \in \mathcal{D}(A^n)$ for some $n \geq 1$. The first results of this kind were obtained in [Ca] and [CGH], where it was shown that if $H\psi = E\psi$ with E embedded in the continuous spectrum of H , and the iterated commutators $\text{ad}_A^k(H)$ are bounded for $k = 1, \dots, \nu$ together with appropriate domain conditions being satisfied by H and A , then $\psi \in \mathcal{D}(A^n)$ for all $n \geq 0$ satisfying $n + 2 \leq \nu$, whenever the Mourre estimate holds at E . Here A is the conjugate operator to the Hamiltonian H in the abstract framework, and the iterated commutators are defined by $\text{ad}_A^1(H) := [H, A]_\circ$ and $\text{ad}_A^k(H) := [\text{ad}_A^{k-1}(H), A]_\circ$. So in the simplest case, one would obtain $\psi \in \mathcal{D}(A)$ provided $\text{ad}_A^3(H)$ exists. Then in [FMS], the authors reduce by one, from $n + 2$ to $n + 1$ the number of commutators that need to be bounded in order to obtain $\psi \in \mathcal{D}(A^n)$, and

show that the result is optimal. In counterpart of these abstract results, we should point out that in the framework of Schrödinger operators, minimal hypotheses yield much stronger results. Indeed, a direct consequence of Theorem 1.2 is that $\psi \in \mathcal{D}(A^n)$ for all $n \geq 0$ assuming only $[H, A]_\circ$ bounded.

The notion of the $C^1(A)$ class of operators also exists for unbounded operators. It appears to us that the results of this paper could also apply to Schrödinger operators with unbounded potentials satisfying the $C^1(A)$ condition. A simple criterion to check if the potential belongs to this class is given in [GM][Lemma A.2]. This criterion is straightforward to verify in the setting of this paper. It is however doubtful to us if the generalization of the result to unbounded potentials is significant.

To further motivate Theorem 1.2, we give an application to discrete Schrödinger operators with Wigner-von Neumann potentials.

Example. (from [Ma]) Let W be the discrete Wigner-von Neumann potential given by

$$(Wu)(n) = W(n)u(n) := \frac{q \sin(k(n_1 + \dots + n_d))}{|n|} u(n), \quad \text{for all } n \in \mathbb{Z}^d, u \in \mathcal{H},$$

for some $(q, k) \in \mathbb{R} \times (-\pi, \pi)$, and let V be a multiplication operator satisfying for some $\rho > 0$,

$$\sup_{n \in \mathbb{Z}^d} \langle n \rangle^\rho |V(n)| < \infty, \quad \text{and} \quad \max_{1 \leq i \leq d} \sup_{n \in \mathbb{Z}^d} \langle n \rangle^\rho |n_i| |(V - \tau_i V)(n)| < \infty.$$

Here $\langle n \rangle := \sqrt{1 + |n|^2}$. Now let $H := \Delta + W + V$ be the Schrödinger operator on \mathcal{H} , and let P and P^\perp be the spectral projectors onto the pure point subspace of H and its complement respectively. Let $E(k) := 4 - 4\text{sign}(k) \cos(k/2)$, and consider the sets

$$\begin{aligned} \mu(H) &:= (0, 4) \setminus \{2 \pm 2 \cos(k/2)\}, \quad \text{for } d = 1, \\ \mu(H) &:= (0, E(k)) \cup (4d - E(k), 4d), \quad \text{for } d \geq 2. \end{aligned}$$

By combining Theorem 1.2 with [Ma, Theorem 1.1], one can remove the abstract assumption $\ker(H - E) \subset \mathcal{D}(A)$ that appears in the latter Theorem. We get the following improved result:

Theorem 1.3. *We have that $\mu(H) \subset \Theta(H)$. For all $E \in \mu(H)$ there is an open interval Σ containing E such that for all $s > 1/2$ and all compact intervals $\Sigma' \subset \Sigma$, the reduced limiting absorption principle for H holds for with respect to (Σ', s, A) , that is,*

$$\sup_{x \in \Sigma', y \neq 0} \|\langle A \rangle^{-s} (H - x - iy)^{-1} P^\perp \langle A \rangle^{-s}\| < \infty.$$

In particular, the spectrum of H is purely absolutely continuous on Σ' whenever $P = 0$ on Σ' , and for $d = 1$, H does not have any eigenvalues on $(2 - 2 \cos(k/2), 2 + 2 \cos(k/2))$.

The plan of the paper is as follows: in Section 2, we prove the main result for the multidimensional Schrödinger operator, namely Theorem 1.2. In section 3, we further develop the method of Section 2 in the case of the one-dimensional operator, and prove Theorem 1.1. Finally Section 4 is the appendix and contains a long technical calculation proving a key relation required for both Sections 2 and 3.

Acknowledgments: It is a pleasure to thank my thesis director Sylvain Golénia for his numerous useful comments and advice, and also Thierry Jecko and Milivoje Lukic for enlightening conversations. I am grateful to the University of Bordeaux for funding my studies.

2. THE MULTIDIMENSIONAL CASE : SUB-EXPONENTIAL LOWER BOUNDS

We begin this section by fixing more notation, and build on the one introduced above. Let

$$\Delta_i := 2 - S_i^* - S_i \quad \text{and}$$

$$A_{0,i} := -i \left(2^{-1}(S_i^* + S_i) + N_i(S_i^* - S_i) \right) = i \left(2^{-1}(S_i^* + S_i) - (S_i^* - S_i)N_i \right).$$

Let

$$(2.1) \quad A'_i := iA_{0,i}, \quad A' := \sum_{i=1}^d A'_i = iA_0 \quad \text{with} \quad \mathcal{D}(A') = \mathcal{D}(A_0).$$

Then the following is a non-negative operator on \mathcal{H} :

$$[\Delta_i, A'_i]_{\circ} = \Delta_i(4 - \Delta_i) = 2 - (S_i^*)^2 - (S_i)^2$$

A useful identity relating the shift operators and the potential is:

$$(2.2) \quad S_i V = (\tau_i V) S_i \quad \text{and} \quad S_i^* V = (\tau_i^* V) S_i^*.$$

Consider an increasing function $F \in C^3([0, \infty))$ with bounded derivative away from the origin. Examples to keep in mind for a later application are $F_{s,\alpha,\gamma} : [0, \infty) \mapsto [0, \infty)$, where $(s, \alpha, \gamma) \in [0, \infty) \times [0, \infty) \times [0, 2/3)$ and

$$(2.3) \quad F_{s,\alpha,\gamma}(x) := \Upsilon_s(\alpha x^\gamma).$$

Here Υ_s is an interpolating function defined for $s \geq 0$ by

$$(2.4) \quad \Upsilon_s(x) := \int_0^x \langle st \rangle^{-2} dt.$$

Then $\Upsilon_s(x) \uparrow x$ as $s \downarrow 0$, and

$$(2.5) \quad \Upsilon_s(x) \leq c_s \quad \text{for} \quad s > 0, \quad \text{and} \quad |\Upsilon_s^{(n)}(x)| \leq cx^{-n+1},$$

where the first constant in (2.5) is dependent of s and the second constant is independent of s . It is readily seen that there are constants $C > 0$ independent of s and γ such that

$$(2.6) \quad |F'_{s,\alpha,\gamma}(x)| \leq Cx^{\gamma-1} \quad \text{and} \quad |F''_{s,\alpha,\gamma}(x)| \leq Cx^{\gamma-2}.$$

We also have that for all $x \geq 0$,

$$(2.7) \quad F'_{s,\alpha,\gamma}(x) \geq 0 \quad \text{and} \quad F''_{s,\alpha,\gamma}(x) \leq 0.$$

So $F_{s,\alpha,\gamma}$ is increasing and concave.

For $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$, let $\langle n \rangle := \sqrt{1 + |n|^2}$. The function F induces a radial operator of multiplication on \mathcal{H} , also denoted by F and acting as follows: $(Fu)(n) := F(\langle n \rangle)u(n)$, for all $u \in \mathcal{H}$. For $i = 1, \dots, d$, we introduce the multiplication operators on \mathcal{H} :

$$(2.8) \quad \varphi_{\ell_i} := (\tau_i e^F - e^F)/e^F = e^{\tau_i F - F} - 1 \quad \text{and} \quad \varphi_{r_i} := (\tau_i^* e^F - e^F)/e^F = e^{\tau_i^* F - F} - 1,$$

$$(2.9) \quad g_{\ell_i} := \varphi_{\ell_i}/N_i \quad \text{and} \quad g_{r_i} := \varphi_{r_i}/N_i.$$

In other words, if $U_i : \mathbb{Z}^d \mapsto \mathbb{Z}^d$ denotes the flow $(n_1, \dots, n_d) \mapsto (n_1, \dots, n_i - 1, \dots, n_d)$ and U_i^{-1} its inverse, then φ_{ℓ_i} and φ_{r_i} are multiplication at n respectively by $\varphi_{\ell_i}(n) = e^{F(\langle U_i n \rangle) - F(\langle n \rangle)} - 1$ and $\varphi_{r_i}(n) = e^{F(\langle U_i^{-1} n \rangle) - F(\langle n \rangle)} - 1$, while g_{ℓ_i} and g_{r_i} are multiplication at n respectively by $g_{\ell_i}(n) = \varphi_{\ell_i}(n)/n_i$ and $g_{r_i}(n) = \varphi_{r_i}(n)/n_i$. Since $g_{\ell_i}(n)$ and $g_{r_i}(n)$ are not well-defined when $n_i = 0$, set $g_{\ell_i}(n) = g_{r_i}(n) := 0$ in that case. We will need the operator g on \mathcal{H} given by

$$(2.10) \quad (gu)(n) = g(n)u(n) := \frac{F'(\langle n \rangle)}{\langle n \rangle} u(n).$$

Three remarks are in order. First, by the mean value theorem, F' bounded away from the origin ensures that φ_{ℓ_i} , φ_{r_i} , g_{ℓ_i} and g_{r_i} are bounded operators on \mathcal{H} ; secondly, F increasing implies $\text{sign}(n_i)\varphi_{r_i}(n) \geq 0$, $\text{sign}(n_i)\varphi_{\ell_i}(n) \leq 0$, $g_{r_i}(n) \geq 0$, $g_{\ell_i}(n) \leq 0$ and $g(n) \geq 0$; and thirdly, we remark that $F, \varphi_{\ell_i}, \varphi_{r_i}$ and g are radial potentials on \mathcal{H} .

Proposition 2.1. *Suppose that Hypotheses 1 holds for the potential V . Let F be a general function as described above and suppose that for all $i, j = 1, \dots, d$,*

- †₁ $|g_{r_i}| \in O(1)$ and $|g_{\ell_i}| \in O(1)$,
- †₂ $|\tau_i g - g|N_j \in O(1)$,
- †₃ $|\tau_i \varphi_{r_i} - \varphi_{r_i}|N_j$, $|\tau_i \varphi_{\ell_i} - \varphi_{\ell_i}|N_j$, $|\tau_i \varphi_{r_j} - \varphi_{r_j}|N_i$ and $|\tau_i \varphi_{\ell_j} - \varphi_{\ell_j}|N_i \in O(1)$,
- †₄ $|(g_{r_i} - g) - (g_{\ell_i} + g)|N_i N_j \in O(1)$.

Suppose that $H\psi = E\psi$, with $\psi \in \mathcal{H}$. Let $\psi_F := e^F \psi$, and assume $\psi_F \in \mathcal{H}$. Then $\psi_F \in \mathcal{D}(\sqrt{g}A')$ and there exist bounded operators $(W_i)_{i=1}^d$, \mathcal{L} , \mathcal{M} and \mathcal{G} on \mathcal{H} depending on F such that

$$(2.11) \quad \begin{aligned} \langle \psi_F, [H, A']_o \psi_F \rangle &= -2 \|\sqrt{g}A'\psi_F\|^2 - \sum_{i=1}^d \|\sqrt{\Delta_i(4 - \Delta_i)}W_i\psi_F\|^2 \\ &\quad + 2^{-1} \langle \psi_F, (\mathcal{L} + \mathcal{M} + \mathcal{G})\psi_F \rangle. \end{aligned}$$

The W_i are multiplication operators given by $W_i = W_{F;i} := \sqrt{\cosh(\tau_i F - F) - 1}$. The expressions of \mathcal{L} , \mathcal{M} and \mathcal{G} are involved; they are given by (4.6), (4.7) and (4.8) respectively. The relevant point is that these operators are a finite sum of terms, each one of the form

$$(2.12) \quad P_1(S_1, \dots, S_d, S_1^*, \dots, S_d^*) T P_2(S_1, \dots, S_d, S_1^*, \dots, S_d^*),$$

where P_1 and P_2 are multivariable polynomials in $S_1, \dots, S_d, S_1^*, \dots, S_d^*$ and T are multiplication operators that satisfy one of the assumptions listed in †₁ – †₄.

Remark 2.1. Formula (2.11) has an additional negative term compared to the corresponding formula for the continuous Schrödinger operator, cf. [FH, Lemma 2.2]:

$$\langle \psi_F, [H, A']_o \psi_F \rangle = -4 \|\sqrt{g}A'\psi_F\|^2 + \langle \psi_F, \mathcal{Q}\psi_F \rangle, \quad \text{with } \mathcal{Q} = (x \cdot \nabla)^2 g - x \cdot \nabla(\nabla F)^2.$$

Remark 2.2. As mentioned in [FH], if we consider the Virial Theorem disregarding operator domains, it is reasonable to expect $\langle \psi, [H, e^F A' e^F] \psi \rangle = 0$. This idea underlies (2.11).

Proof. Let $\phi \in \ell_0(\mathbb{Z}^d)$, the sequences with compact support, and $\phi_F := e^F \phi$. The first step of the proof consists in establishing the following identity :

$$(2.13) \quad \begin{aligned} \langle \phi, [e^F A' e^F, \Delta] \phi \rangle &= \langle \phi_F, [A', \Delta] \phi_F \rangle - 2 \|\sqrt{g}A'\phi_F\|^2 \\ &\quad - \sum_{1 \leq i \leq d} \|\sqrt{\Delta_i(4 - \Delta_i)}W_i\phi_F\|^2 + 2^{-1} \langle \phi_F, (\mathcal{L} + \mathcal{M} + \mathcal{G})\phi_F \rangle. \end{aligned}$$

The proof of (2.13) is technical and long, so it is done in the appendix. The assumptions of this Proposition and F' bounded away from the origin imply that the W_i , \mathcal{L} , \mathcal{M} and \mathcal{G} stemming from this calculation are bounded operators. Exactly where these assumptions are applied are indicated in the appendix by (‡). The second step consists in using (2.13) to prove (2.11). For $m \geq 1$, define the cut-off potentials $\chi_m(n) := \chi(\langle n \rangle / m)$ on \mathbb{Z}^d , where $\chi \in C_c^\infty(\mathbb{R})$ and χ equals one in a neighborhood of the origin. Then (2.13) holds with $\phi = \chi_m \psi$ and $\phi_F = e^F \chi_m \psi$. Adding $\langle \chi_m \psi, [e^F A' e^F, V] \chi_m \psi \rangle = \langle e^F \chi_m \psi, [A', V] e^F \chi_m \psi \rangle$ to each side of (2.13), and introducing the constant E in the commutator on the left, we obtain

$$(2.14) \quad \begin{aligned} \langle \chi_m \psi, [e^F A' e^F, H - E] \chi_m \psi \rangle &= \langle e^F \chi_m \psi, [A', H] e^F \chi_m \psi \rangle - 2 \|\sqrt{g}A'e^F \chi_m \psi\|^2 \\ &\quad - \sum_{1 \leq i \leq d} \|\sqrt{\Delta_i(4 - \Delta_i)}W_i e^F \chi_m \psi\|^2 \\ &\quad + 2^{-1} \langle e^F \chi_m \psi, (\mathcal{L} + \mathcal{M} + \mathcal{G}) e^F \chi_m \psi \rangle. \end{aligned}$$

Since $e^F \chi_m \psi \rightarrow \psi_F$ in \mathcal{H} as $m \rightarrow \infty$, the first, third and fourth terms on the right side of (2.14) converge. The left side of (2.14) is handled in the same way as in [CFKS, Proposition 4.16]:

$$\begin{aligned} \langle \chi_m \psi, [e^F A' e^F, H - E] \chi_m \psi \rangle &= -2\Re(\langle e^F A' e^F \chi_m \psi, (H - E) \chi_m \psi \rangle) \\ &= -2\Re(\langle \langle N \rangle^{-1} A' e^F \chi_m \psi, \langle N \rangle e^F (H - E) \chi_m \psi \rangle). \end{aligned}$$

Since $\text{supp}(\chi_m) \subset [-2m, 2m]^d$, $\text{supp}((H - E)\chi_m \psi) \subset K := [-2m - 1, 2m + 1]^d$ and so commuting χ_m with $(H - E)$ gives

$$(2.15) \quad \begin{aligned} \langle N \rangle e^F (H - E) \chi_m \psi &= \langle N \rangle e^F \mathbf{1}_K (H - E) \chi_m \psi \\ &= \sum_{1 \leq i \leq d} \langle N \rangle (\chi_m - \tau_i \chi_m) e^F S_i \psi + \langle N \rangle (\chi_m - \tau_i^* \chi_m) e^F S_i^* \psi. \end{aligned}$$

An application of the Mean Value Theorem shows that $|\langle N \rangle (\chi_m - \tau_i \chi_m)|$ and $|\langle N \rangle (\chi_m - \tau_i^* \chi_m)|$ are bounded by a constant independent of m . Moreover, $\psi_F \in \mathcal{H}$ and F' bounded imply that $e^F S_i \psi = S_i e^{\tau_i^* F - F} \psi_F$ and $e^F S_i^* \psi = S_i^* e^{\tau_i F - F} \psi_F \in \mathcal{H}$. Thus the sequence (2.15) is uniformly bounded in absolute value in \mathcal{H} . Furthermore, it converges pointwise to zero. By Lebesgue's Dominated Convergence Theorem,

$$(2.16) \quad \|\langle N \rangle e^F (H - E) \chi_m \psi\| \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Since $\langle N \rangle^{-1} A'$ is a bounded operator on \mathcal{H} , the left side of (2.14) converges to zero as $m \rightarrow \infty$. The only remaining term in (2.14) is $2\|\sqrt{g} A' e^F \chi_m \psi\|^2$, hence it must also converge as $m \rightarrow \infty$. To finish the proof, it remains to show that $\psi_F \in \mathcal{D}(\sqrt{g} A')$. Let $\phi \in \ell_0(\mathbb{Z})$. Then

$$|\langle \psi_F, A' \sqrt{g} \phi \rangle| = \lim_{m \rightarrow \infty} |\langle e^F \chi_m \psi, A' \sqrt{g} \phi \rangle| \leq \left(\lim_{m \rightarrow \infty} \|\sqrt{g} A' e^F \chi_m \psi\| \right) \|\phi\|.$$

This shows that $\psi_F \in \mathcal{D}((-A' \sqrt{g})^*) = \mathcal{D}(\sqrt{g} A')$. Then it must be that $\|\sqrt{g} A' e^F \chi_m \psi\|^2 \rightarrow \|\sqrt{g} A' \psi_F\|^2$ and the proof is complete after rearranging the terms accordingly in (2.14). \square

As mentioned in the last Proposition, \mathcal{L}, \mathcal{M} and \mathcal{G} are a finite sum of terms of the form

$$P_1(S_1, \dots, S_d, S_1^*, \dots, S_d^*) T P_2(S_1, \dots, S_d, S_1^*, \dots, S_d^*)$$

for some polynomials P_1 and P_2 . Going forward, it is essential that the multiplication operators $T = T(n)$ decay radially at infinity. In other words, for the minimal assumptions $\dagger_1 - \dagger_4$, we will need $o(1)$ instead of $O(1)$. The following Lemma shows that this is the case for $F = F_{s, \alpha, \gamma}$.

Lemma 2.2. *Let $F = F_{s, \alpha, \gamma}$ be the function defined in (2.3). Consider its corresponding functions $\varphi_{r_i}, \varphi_{\ell_i}, g_{r_i}, g_{\ell_i}$ and g . The following estimates hold uniformly with respect to s and γ :*

- $\dagger_1 \quad |g_{r_i}| \quad \text{and} \quad |g_{\ell_i}| \in O_\alpha(\langle n \rangle^{\gamma-2}),$
- $\dagger_2 \quad |\tau_i g - g| \in O_\alpha(\langle n \rangle^{\gamma-3}),$
- $\dagger_3 \quad |\tau_i^* \varphi_{r_j} - \varphi_{r_j}| \quad \text{and} \quad |\tau_i^* \varphi_{\ell_j} - \varphi_{\ell_j}| \in O_\alpha(\langle n \rangle^{\gamma-2}),$
- $\dagger_4 \quad |(g_{r_i} - g) - (g_{\ell_i} + g)| \in O_\alpha(\langle n \rangle^{3\gamma-4}),$
- $\dagger_5 \quad |(\tau_i F - F) - \tau_i(\tau_i F - F)| \in O_\alpha(\langle n \rangle^{\gamma-2}).$

Therefore \dagger_i improve \dagger_i for $i = 1, 2, 3, 4$ respectively.

Proof. These estimates are simple applications of the Mean Value Theorem (MVT). Let $n = (n_1, \dots, n_d)$ and fix $i \in \{1, \dots, d\}$. There is $n' = (n'_1, \dots, n'_d)$ with $n'_i \in (n_i, n_i + 1)$ and $n'_j = n_j$ for $j \neq i$ such that

$$g_{r_i}(n) = \frac{n'_i}{\langle n' \rangle} \frac{F'(\langle n' \rangle) e^{F(\langle n' \rangle)}}{n_i e^{F(\langle n \rangle)}}.$$

This, together with (2.6), and an analogous calculation for $g_{\ell_i}(n)$ shows \ddagger_1 . Define $g : \mathbb{R}^d \rightarrow \mathbb{R}$, $g(x) := F'(\langle x \rangle) \langle x \rangle^{-1}$. Then \ddagger_2 follows from

$$\frac{\partial g}{\partial x_i}(x) = \frac{x_i}{\langle x \rangle} \frac{F''(\langle x \rangle) \langle x \rangle - F'(\langle x \rangle)}{\langle x \rangle^2}.$$

Now fix $i, j \in \{1, \dots, d\}$. First there is $n' = (n'_1, \dots, n'_d)$ with $n'_j \in (n_j, n_j + 1)$ and $n'_k = n_k$ for $k \neq j$ such that

$$(\tau_j^* F - F)(n) = \frac{\partial \tilde{F}}{\partial x_j}(n') = \frac{n'_j}{\langle n' \rangle} F'(\langle n' \rangle), \quad \text{with } \tilde{F}(x) = F(\langle x \rangle).$$

Then there is $n'' = (n''_1, \dots, n''_d)$ with $n''_i \in (n'_i, n'_i + 1)$ and $n''_k = n'_k$ for $k \neq i$ such that

$$(\tau_i^* \varphi_{r_j} - \varphi_{r_j})(n) = \frac{\partial^2 \tilde{F}}{\partial x_i \partial x_j}(n'') e^{\frac{\partial \tilde{F}}{\partial x_j}(n'')}.$$

This proves \ddagger_3 since

$$\left| \frac{\partial^2 \tilde{F}}{\partial x_i \partial x_j}(x) \right| \leq \frac{|F'(\langle x \rangle)|}{\langle x \rangle} + |F''(\langle x \rangle)|.$$

The latter estimate on $\partial^2 \tilde{F}/(\partial x_i \partial x_j)$ also implies \ddagger_5 . Finally, for \ddagger_4 , we start with

$$g_{r_i}(n) - g(n) = \frac{1}{n_i e^{F(\langle n \rangle)}} \left[\frac{n'_i}{\langle n' \rangle} F'(\langle n' \rangle) e^{F(\langle n' \rangle)} - \frac{n_i}{\langle n \rangle} F'(\langle n \rangle) e^{F(\langle n \rangle)} \right] = \frac{1}{n_i e^{F(\langle n \rangle)}} \frac{\partial k}{\partial x_i}(n'')$$

where

$$k : \mathbb{R}^d \rightarrow \mathbb{R}, \quad k(x) := \frac{x_i}{\langle x \rangle} F'(\langle x \rangle) e^{F(\langle x \rangle)},$$

and $n'' = (n''_1, \dots, n''_d)$ with $n''_i \in (n_i, n'_i)$ and $n''_j = n_j$ for $j \neq i$. We compute

$$\frac{\partial k}{\partial x_i}(x) = \left(\frac{F'(\langle x \rangle)}{\langle x \rangle} - \frac{x_i^2 F'(\langle x \rangle)}{\langle x \rangle^3} + \frac{x_i^2 F''(\langle x \rangle)}{\langle x \rangle^2} + \frac{x_i^2 (F'(\langle x \rangle))^2}{\langle x \rangle^2} \right) e^{F(\langle x \rangle)}.$$

Thus for some $n''' = (n'''_1, \dots, n'''_d)$ with $n'''_i \in (n_i - 1, n_i + 1)$ and $n'''_j = n_j$ for $j \neq i$, we have

$$(g_{r_i}(n) - g(n)) - (g_{\ell_i}(n) + g(n)) = \frac{1}{n_i e^{F(\langle n \rangle)}} \frac{\partial^2 k}{\partial x_i^2}(n''').$$

A calculation of $\partial^2 k / \partial x_i^2$ yields the required estimate. \square

We are now ready to prove the main result concerning the multidimensional operator H :

Proof of Theorem 1.2. Let $\psi_{F_s, \alpha, \gamma} := e^{F_s, \alpha, \gamma} \psi$, and let $\Psi_s := \psi_{F_s, \alpha, \gamma} / \|\psi_{F_s, \alpha, \gamma}\|$. We suppose that for some $(\alpha, \gamma) \in [0, \infty) \times [0, 2/3)$, $\psi \notin \mathcal{D}(\vartheta_{\alpha, \gamma})$ and derive a contradiction. Of course, $\psi_{F_s, \alpha, \gamma} \in \mathcal{H}$ for all $s > 0$, but by the Monotone Convergence Theorem, $\|\psi_{F_s, \alpha, \gamma}\| \rightarrow +\infty$ as $s \downarrow 0$. Thus, for any bounded set $B \subset \mathbb{Z}^d$,

$$(2.17) \quad \lim_{s \downarrow 0} \sum_{n \in B} |\Psi_s(n)|^2 = 0.$$

In particular, Ψ_s converges weakly to zero. As α and γ are fixed, we shall write F_s instead of $F_{s, \alpha, \gamma}$ for simplicity. Introduce the operator $H_{F_s} := e^{F_s} H e^{-F_s}$. Then H_{F_s} is a bounded operator and $H_{F_s} \Psi_s = E \Psi_s$. We claim that

$$(2.18) \quad \lim_{s \downarrow 0} \|(H - E) \Psi_s\| = 0.$$

To see this, write H_{F_s} as follows:

$$H_{F_s} = H + \sum_{1 \leq i \leq d} S_i(1 - e^{\tau_i^* F_s - F_s}) + S_i^*(1 - e^{\tau_i F_s - F_s}).$$

To show (2.18), it is therefore enough to show that

$$(2.19) \quad \lim_{s \downarrow 0} \|(1 - e^{\tau_i^* F_s - F_s})\Psi_s\| = \lim_{s \downarrow 0} \|(1 - e^{\tau_i F_s - F_s})\Psi_s\| = 0.$$

Let $B(N) = \{n \in \mathbb{Z}^d : \langle n \rangle \leq N\}$, and $B(N)^c$ the complement set. For all $\epsilon > 0$, there is $N > 0$ such that

$$\sup_{\substack{n \in B(N)^c \\ s > 0}} \left| 1 - e^{(\tau_i^* F_s - F_s)(n)} \right| = \sup_{\substack{n \in B(N)^c \\ s > 0}} \left| 1 - e^{\alpha \gamma \langle n' \rangle^{\gamma-1} \Upsilon'_s(\alpha \langle n' \rangle^\gamma)} \right| \leq \epsilon$$

(here $n' = (n'_1, \dots, n'_d)$ with $n'_i \in (n_i, n_i + 1)$ and $n'_j = n_j$ for $j \neq i$). Combining this with (2.17) proves the first limit in (2.19), and the second one is shown in the same way. Thus the claim is proven. Because $E \in \Theta(H)$, there exists an interval $\Sigma := (E - \delta, E + \delta)$ with $\delta > 0$, $\eta > 0$ and a compact K such that

$$(2.20) \quad E_\Sigma(H)[H, A']_\circ E_\Sigma(H) \geq \eta E_\Sigma(H) + K.$$

By functional calculus,

$$(2.21) \quad \lim_{s \downarrow 0} \|E_{\mathbb{R} \setminus \Sigma}(H)\Psi_s\| \leq \lim_{s \downarrow 0} \delta^{-1} \|E_{\mathbb{R} \setminus \Sigma}(H)(H - E)\Psi_s\| = 0.$$

It follows by the Mourre estimate (2.20) and (2.21) that

$$(2.22) \quad \liminf_{s \downarrow 0} \langle \Psi_s, E_\Sigma(H)[H, A']_\circ E_\Sigma(H)\Psi_s \rangle \geq \eta \liminf_{s \downarrow 0} \|E_\Sigma(H)\Psi_s\|^2 = \eta > 0.$$

We now look to contradict this equation. We start with

$$(2.23) \quad \langle \Psi_s, E_\Sigma(H)[H, A']_\circ E_\Sigma(H)\Psi_s \rangle = \langle \Psi_s, [H, A']_\circ \Psi_s \rangle - f_1(s) - f_2(s), \quad \text{where} \\ f_1(s) = \langle \Psi_s, E_{\mathbb{R} \setminus \Sigma}(H)[H, A']_\circ E_\Sigma(H)\Psi_s \rangle \quad \text{and} \quad f_2(s) = \langle \Psi_s, [H, A']_\circ E_{\mathbb{R} \setminus \Sigma}(H)\Psi_s \rangle.$$

Applying (2.21) gives

$$\lim_{s \downarrow 0} |f_1(s)| = \lim_{s \downarrow 0} |f_2(s)| = 0.$$

Now apply (2.11) with $F = F_{s, \alpha, \gamma}$, and after dividing this equation by $\|\Psi_s\|^2$, we have

$$\limsup_{s \downarrow 0} \langle \Psi_s, [H, A']_\circ \Psi_s \rangle \leq 0.$$

Here we took advantage of the negativity of the first two terms on the right side of (2.11), and used the uniform decay of $\mathcal{L} + \mathcal{M} + \mathcal{G}$ together with the weak convergence of Ψ_s to get $\langle \Psi_s, (\mathcal{L} + \mathcal{M} + \mathcal{G})\Psi_s \rangle \rightarrow 0$ as $s \downarrow 0$. To check this thoroughly, one needs to apply the estimates of Lemma 2.2 to where indicated in the appendix by a (‡). Note that \mathcal{L} given by (4.6) is the most constraining term; it has the necessary decay provided $3\gamma - 4 < -2$, i.e. $\gamma < 2/3$. Note also that ‡₅ allows to conclude, by continuity of the map $x \mapsto \sqrt{\cosh(x) - 1}$, that $\langle \Psi_s, (W_{F_s; i} - \tau_i W_{F_s; i})\Psi_s \rangle$ and like terms converge to zero. Thus by (2.23),

$$\limsup_{s \downarrow 0} \langle \Psi_s, E_\Sigma(H)[H, A']_\circ E_\Sigma(H)\Psi_s \rangle \leq 0.$$

This is in contradiction with (2.22), so the proof is complete. \square

3. THE ONE-DIMENSIONAL CASE: EXPONENTIAL LOWER BOUNDS

In this section we deal with the one-dimensional Schrödinger operator H on $\mathcal{H} = \ell^2(\mathbb{Z})$. We follow the same definitions as in the introduction and the previous section, but since $i = 1$, we will drop this subscript. We shall write S and S^* instead of S_i and S_i^* , N instead of N_i , etc... Consider

an increasing function $F \in C^2([0, \infty))$ with bounded derivative away from the origin. This function induces a radial operator on \mathcal{H} as in the previous section : $(Fu)(n) := F(\langle n \rangle)u(n)$ for all $u \in \mathcal{H}$.

Proposition 3.1. *Suppose that Hypothesis 1 holds for the potential V . Let F be as above, and suppose additionally that*

$$(3.1) \quad |xF''(x)| \leq C, \quad \text{for } x \text{ away from the origin.}$$

Suppose that $H\psi = E\psi$, with $\psi \in \mathcal{H}$. Let $\psi_F := e^F\psi$, and assume that $\psi_F \in \mathcal{H}$. Then $\psi_F \in \mathcal{D}(\sqrt{g_r - g_\ell}A')$ and there exist bounded operators W , M and G depending on F such that

$$(3.2) \quad \langle \psi_F, [H, A']\psi_F \rangle = -\|\sqrt{g_r - g_\ell}A'\psi_F\|^2 - \|\sqrt{\Delta(4 - \Delta)}W\psi_F\|^2 + 2^{-1}\langle \psi_F, (M + G)\psi_F \rangle.$$

The exact expressions of W, M and G are given by (4.10), (4.11) and (4.12) respectively.

Proof. The proof is done in two steps. The first step consists in proving that

$$(3.3) \quad \begin{aligned} \langle \phi, [e^F A' e^F, \Delta]\phi \rangle &= \langle \phi_F, [A', \Delta]\phi_F \rangle - \|\sqrt{g_r - g_\ell}A'\phi_F\|^2 \\ &\quad - \|\sqrt{\Delta(4 - \Delta)}W\phi_F\|^2 + 2^{-1}\langle \phi_F, (M + G)\phi_F \rangle. \end{aligned}$$

The proof of this is in the appendix starting from (4.9). That F' is bounded away from the origin ensures that W and $(g_r - g_\ell)$ are bounded. The additional assumption (3.1) ensures that $(\tau^*\varphi_r - \varphi_r)N$ and like terms are bounded. The second step is the same as that of Proposition 2.1, and the proof is identical. \square

Lemma 3.2. *Suppose that $H\psi = E\psi$ with $\psi \in \ell^2(\mathbb{Z})$. Let F be a general function as above, and assume that $\psi_F := e^F\psi \in \ell^2(\mathbb{Z})$. Define the operator*

$$(3.4) \quad H_F := e^F H e^{-F}.$$

Then H_F is bounded, $H_F\psi_F = E\psi_F$ and there exist bounded operators C_F and R_F such that

$$(3.5) \quad H_F = C_F H + (2 - 2C_F) + 2^{-1}R_F, \quad \text{where}$$

$$(3.6) \quad C_F := 2^{-1} \left(e^{F-\tau F} + e^{F-\tau^* F} \right) \quad \text{and}$$

$$(3.7) \quad \begin{aligned} R_F &:= V(2 - 2C_F) + (\tau\varphi_r - \varphi_r)(S^* - S) + (\varphi_\ell - \tau^*\varphi_\ell)(S^* - S) \\ &\quad + (g_r - g_\ell)A' - 2^{-1}(g_r - g_\ell)(S^* + S). \end{aligned}$$

Proof. Because F' is bounded away from the origin, both $e^F S e^{-F} \phi = S e^{\tau^* F - F} \phi$ and $e^F S^* e^{-F} \phi = S^* e^{\tau F - F} \phi$ belong to $\ell^2(\mathbb{Z})$ whenever $\phi \in \ell^2(\mathbb{Z})$. Thus H_F is bounded, and $H_F\psi_F = E\psi_F$ follows immediately. Now

$$H_F = 2 + V - e^{F-\tau F} S - e^{F-\tau^* F} S^*.$$

Rewriting this relation in two different ways, we have

$$\begin{aligned} H_F &= e^{F-\tau F} H + (2 + V)(1 - e^{F-\tau F}) + (e^{F-\tau F} - e^{F-\tau^* F})S^*, \\ H_F &= e^{F-\tau^* F} H + (2 + V)(1 - e^{F-\tau^* F}) + (e^{F-\tau^* F} - e^{F-\tau F})S. \end{aligned}$$

Adding these two relations gives

$$(3.8) \quad 2H_F = 2C_F H + (2 + V)(2 - 2C_F) + (e^{F-\tau F} - e^{F-\tau^* F})(S^* - S).$$

We further develop the third term on the right side:

$$\begin{aligned}
 (e^{F-\tau F} - e^{F-\tau^* F})(S^* - S) &= (\tau\varphi_r - \tau^*\varphi_\ell)(S^* - S) \\
 &= (\tau\varphi_r - \varphi_r)(S^* - S) + (\varphi_\ell - \tau^*\varphi_\ell)(S^* - S) + (\varphi_r - \varphi_\ell)(S^* - S) \\
 &= (\tau\varphi_r - \varphi_r)(S^* - S) + (\varphi_\ell - \tau^*\varphi_\ell)(S^* - S) \\
 &\quad + (g_r - g_\ell)A' - 2^{-1}(g_r - g_\ell)(S^* + S) + (\varphi_r - \varphi_\ell)\delta_{\{n=0\}}(S^* - S).
 \end{aligned}$$

Here, δ_B is the projector onto $B \subset \mathbb{Z}$. Note that $(\varphi_r - \varphi_\ell)\delta_{\{n=0\}} = 0$, and thus (3.5) is shown. \square

We are now ready to prove the main result concerning the one-dimensional operator H :

Proof of Theorem 1.1, the first part. We first handle the case $E \neq 2$. Suppose that the statement of the theorem is false. Then $\theta_E = \theta_E(\alpha_0) = (E - 2)/\cosh(\alpha_0) + 2 \in \Theta(H) \setminus \{+2\}$ for some $\alpha_0 \in [0, \infty)$, and there is an interval

$$(3.9) \quad \Sigma_0 := (\theta_E(\alpha_0) - 2\delta, \theta_E(\alpha_0) + 2\delta)$$

such that the Mourre estimate holds there, i.e.

$$(3.10) \quad E_{\Sigma_0}(H)[H, A'] \circ E_{\Sigma_0}(H) \geq \eta E_{\Sigma_0}(H) + K$$

for some $\eta > 0$ and some compact operator K . For the remainder of the proof, δ , η and K are fixed. If $\alpha_0 > 0$, choose $\alpha_1 > 0$ and $\gamma > 0$ such that

$$(3.11) \quad \alpha_1 < \alpha_0 < \alpha_1 + \gamma.$$

If however $\alpha_0 = 0$, let $\alpha_1 = 0$ and $\gamma > 0$. By continuity of the map $\theta_E(\alpha) = (E - 2)/\cosh(\alpha) + 2$, $\theta_E(\alpha_1) \rightarrow \theta_E(\alpha_0)$ as $\alpha_1 \rightarrow \alpha_0$, so taking α_1 close enough to α_0 we obtain intervals

$$\Sigma_1 := (\theta_E(\alpha_1) - \delta, \theta_E(\alpha_1) + \delta) \subset \Sigma_0$$

with the inclusion remaining valid as $\alpha_1 \rightarrow \alpha_0$. Multiplying to the right and left of (3.10) by $E_{\Sigma_1}(H)$, we obtain

$$(3.12) \quad E_{\Sigma_1}(H)[H, A'] \circ E_{\Sigma_1}(H) \geq \eta E_{\Sigma_1}(H) + E_{\Sigma_1}(H)KE_{\Sigma_1}(H).$$

Later in the proof α_1 will be taken even closer to α_0 allowing γ to be as small as necessary in order to lead to a contradiction (in this limiting process, δ , η and K are fixed). Before delving into the details of the proof, we expose the strategy. For a suitable sequence of functions $\{F_s(x)\}_{s>0}$, let

$$(3.13) \quad \Psi_s := e^{F_s}\psi / \|e^{F_s}\psi\|.$$

With F_s and Ψ_s instead of F and ψ_F respectively, we apply Proposition 3.1 to conclude that

$$(3.14) \quad \limsup_{s \downarrow 0} \langle \Psi_s, [H, A'] \circ \Psi_s \rangle \leq \limsup_{s \downarrow 0} |\langle \Psi_s, 2^{-1}(M_{F_s} + G_{F_s})\Psi_s \rangle|.$$

Notice how the the negativity of the first two terms on the right side of (3.2) was crucial. We have also written M_{F_s} and G_{F_s} instead of M and G to show the dependence on F_s . The first part of the proof consists in showing that

$$(3.15) \quad \limsup_{s \downarrow 0} \langle \Psi_s, [H, A'] \circ \Psi_s \rangle \leq \limsup_{s \downarrow 0} |\langle \Psi_s, 2^{-1}(M_{F_s} + G_{F_s})\Psi_s \rangle| \leq c\epsilon_\gamma$$

for some $\epsilon_\gamma > 0$ satisfying $\epsilon_\gamma \rightarrow 0$ when $\gamma \rightarrow 0$. Here and thereafter, $c > 0$ denotes a constant independent of s , α_1 and γ . The second part of the proof consists in showing that

$$(3.16) \quad \limsup_{s \downarrow 0} \|(H - \theta_E(\alpha_1))\Psi_s\| \leq c\epsilon_\gamma.$$

Roughly speaking (3.16) says that Ψ_s has energy concentrated about $\theta_E(\alpha_1)$ and so localizing (3.15) about this energy will lead to

$$(3.17) \quad \limsup_{s \downarrow 0} \langle \Psi_s, E_{\Sigma_1}(H)[H, A'] \circ E_{\Sigma_1}(H) \Psi_s \rangle \leq c\epsilon_\gamma.$$

However, the Mourre estimate (3.12) holds on Σ_1 . In the end, the contradiction will come from the fact that the Mourre estimate asserts that the left side of (3.17) is not that small.

We now begin in earnest the proof. Notice that $\psi \in \mathcal{D}(\vartheta_{\alpha_1})$ but $\psi \notin \mathcal{D}(\vartheta_{\alpha_1+\gamma})$. Let Υ_s be the interpolating function defined in (2.4), and for $s > 0$ let

$$(3.18) \quad F_s(x) := \alpha_1 x + \gamma \Upsilon_s(x).$$

As explained in the multidimensional case, F_s induces a radial potential as follows : $(F_s u)(n) := F_s(\langle n \rangle)u(n)$, for all $u \in \ell^2(\mathbb{Z})$. By (2.5), $e^{F_s} \psi \in \ell^2(\mathbb{Z})$ for all $s > 0$, but $\|e^{F_s} \psi\| \rightarrow \infty$ as $s \downarrow 0$. To ease the notation, we will be bounding various quantities by the same constant $c > 0$, a constant that is independent of α_1 , γ , s and of position x (or n).

Part 1. We use Proposition 3.1 with F_s replacing F , and so we verify that F_s satisfies the hypotheses of that proposition. Since

$$F'_s(x) = \alpha_1 + \gamma \Upsilon'_s(x) \quad \text{and} \quad F''_s(x) = \gamma \Upsilon''_s(x),$$

indeed $|F'_s(x)| \leq c$, $|x F''_s(x)| \leq c$. Dividing (3.2) by $\|e^{F_s} \psi\|^2$ throughout we obtain (3.14) as claimed. To prove (3.15), we need two ingredients. First, for any bounded set $B \subset \mathbb{Z}$,

$$(3.19) \quad \lim_{s \downarrow 0} \sum_{n \in B} |\Psi_s(n)|^2 = 0.$$

In particular, Ψ_s converges weakly to zero. What's more, we also have for any $k \in \mathbb{N}$

$$(3.20) \quad \lim_{s \downarrow 0} \sum_{n \in B} |(S^k \Psi_s)(n)|^2 = 0, \quad \text{and} \quad \lim_{s \downarrow 0} \sum_{n \in B} |((S^*)^k \Psi_s)(n)|^2 = 0.$$

Now M_{F_s} and G_{F_s} are a finite sum of terms of the form $P_1(S, S^*) T P_2(S, S^*)$, where P_1 and P_2 are polynomials and the $T = T(n)$ are sequences. The second item to show is that,

$$(3.21) \quad |T(n)| \leq c \langle n \rangle^{-1} + \epsilon_\gamma.$$

In other words we want smallness coming from decay in position n or from γ . Outside a sufficiently large bounded set, decay in position can be converted into smallness in γ by using (3.19) while $P_1(S, S^*)$ and $P_2(S, S^*)$ get absorbed in the process thanks to (3.20). Consider first $M = M_{F_s}$ given by (4.11). Applying the Mean Value Theorem (MVT) gives the uniform estimates in s

$$(3.22) \quad |\tau F_s - F_s| \quad \text{and} \quad |\tau^* F_s - F_s| \in O(1).$$

It follows that

$$|\varphi_\ell| \quad \text{and} \quad |\varphi_r| \in O(1), \quad \text{and} \quad |g_r - g_\ell| \in O(\langle n \rangle^{-1}).$$

To handle the term $(\tau^* \varphi_\ell - \varphi_\ell)$, define the function $f(x) := e^{F_s(\langle x-1 \rangle) - F_s(\langle x \rangle)}$. Then $(\tau^* \varphi_\ell - \varphi_\ell)(n) = f(n+1) - f(n)$. Applying twice the MVT gives

$$|(\tau^* \varphi_\ell - \varphi_\ell)(n)| \leq c \langle n \rangle^{-3} + \gamma \langle n \rangle^{-1}.$$

The same estimate holds for the similar terms like $(\varphi_r - \tau \varphi_r)$, $(\tau^* \varphi_r - \varphi_r)$ and so forth. We turn our attention to $G = G_{F_s}$ given by (4.12). By (3.22), $|W_{F_s}| \in O(1)$. To estimate $(W_{F_s} - W_{\tau^* F_s})$, let $g(x) := \sqrt{\cosh(F_s(\langle x-1 \rangle) - F_s(\langle x \rangle)) - 1}$, so that $(W_{F_s} - W_{\tau^* F_s})(n) = g(n) - g(n+1)$. Moreover,

$$g'(x) = \frac{(F'_s(\langle x-1 \rangle) - F'_s(\langle x \rangle)) \sinh(F_s(\langle x-1 \rangle) - F_s(\langle x \rangle))}{2\sqrt{\cosh(F_s(\langle x-1 \rangle) - F_s(\langle x \rangle)) - 1}}.$$

If $\alpha_1 > 0$, then $|F_s(\langle x-1 \rangle) - F_s(\langle x \rangle)| \geq c'\alpha_1$ for some constant $c' > 0$ independent of x and s , and so $\cosh(F_s(\langle x-1 \rangle) - F_s(\langle x \rangle)) - 1$ is uniformly bounded from below by a positive number. Applying the MVT to $(F'_s(\langle x-1 \rangle) - F'_s(\langle x \rangle))$ yields the estimate

$$|(W_{F_s} - W_{\tau^*F_s})(n)| \leq c(\langle n \rangle^{-3} + \gamma \langle n \rangle^{-1}).$$

If however $\alpha_1 = 0$, then

$$(3.23) \quad |(\tau F_s - F_s)(n) - (F_s - \tau^*F_s)(n)| \leq c\gamma \langle n \rangle^{-1}.$$

By continuity of the function $x \mapsto \sqrt{\cosh(x) - 1}$ we have that for any $\epsilon_\gamma > 0$,

$$|W_{F_s} - W_{\tau^*F_s}| = |\sqrt{\cosh(\tau F_s - F_s) - 1} - \sqrt{\cosh(F_s - \tau^*F_s) - 1}| \leq \epsilon_\gamma$$

whenever (3.23) holds. A similar argument works for $(W_{F_s} - W_{\tau F_s})$. Thus (3.21) is proven, and this shows (3.15) when combined with the fact that Ψ_s converges weakly to zero.

Part 2. We now prove (3.16). Consider Lemma 3.2 with F_s instead of F . We claim that

$$(3.24) \quad \lim_{s \downarrow 0} \left\| (C_{F_s}H + 2 - E - 2C_{F_s}) \Psi_s \right\| = 0.$$

By (3.5) of Lemma 3.2, this is equivalent to showing that

$$\lim_{s \downarrow 0} \|R_{F_s} \Psi_s\| = 0.$$

Dividing each term in (3.2) by $\|e^{F_s} \psi\|^2$, we see that $\|\sqrt{g_r - g_\ell} A' \Psi_s\| \leq c$. Let χ_N denote the characteristic function of the set $\{n \in \mathbb{Z} : (g_r - g_\ell) < N^{-1}\}$. Then

$$\limsup_{s \downarrow 0} \|(g_r - g_\ell) A' \Psi_s\| \leq \limsup_{s \downarrow 0} N^{-\frac{1}{2}} \|\chi_N \sqrt{g_r - g_\ell} A' \Psi_s\| + \|(1 - \chi_N)(g_r - g_\ell) A' \Psi_s\| \leq cN^{-\frac{1}{2}}.$$

Here we used the fact that $1 - \chi_N$ has support in a fixed, bounded set as $s \downarrow 0$. Since N is arbitrary, this shows that $\|(g_r - g_\ell) A' \Psi_s\| \rightarrow 0$ as $s \downarrow 0$. The other terms of R_{F_s} are handled similarly. Note that for the term containing V we use the fact it goes to zero at infinity, and from Part 1, $(\tau \varphi_r - \varphi_r)$, $(\varphi_\ell - \tau^* \varphi_\ell)$ and $(g_r - g_\ell)$ also go to zero at infinity. Hence (3.24) is proved. Let $\kappa := \kappa(n) = \text{sign}(n)$. From the expression of F'_s , we have the estimates :

$$|(F_s - \tau F_s)(n) - \kappa(n)\alpha_1| \leq c(\alpha_1 \langle n \rangle^{-1} + \gamma) \quad \text{and} \quad |(F_s - \tau^* F_s)(n) - (-\kappa(n)\alpha_1)| \leq c(\alpha_1 \langle n \rangle^{-1} + \gamma).$$

Therefore, outside a fixed bounded set we have

$$(3.25) \quad |(F_s - \tau F_s) - \kappa\alpha_1| \leq c\gamma \quad \text{and} \quad |(F_s - \tau^* F_s) - (-\kappa\alpha_1)| \leq c\gamma.$$

By continuity of the exponential function, we have for any $\epsilon_\gamma > 0$ that

$$|e^{F_s - \tau F_s} - e^{\kappa\alpha_1}| \leq \epsilon_\gamma \quad \text{and} \quad |e^{F_s - \tau^* F_s} - e^{-\kappa\alpha_1}| \leq \epsilon_\gamma$$

whenever the respective terms of (3.25) hold. It follows from (3.24) that

$$\limsup_{s \downarrow 0} \left\| [2^{-1} (e^{\alpha_1} + e^{-\alpha_1}) H + 2 - E - (e^{\alpha_1} + e^{-\alpha_1})] \Psi_s \right\| \leq c\epsilon_\gamma.$$

Dividing this expression by $\cosh(\alpha_1)$ proves (3.16).

Part 3. By functional calculus and (3.16), we have

$$(3.26) \quad \limsup_{s \downarrow 0} \|E_{\mathbb{R} \setminus \Sigma_1}(H) \Psi_s\| \leq \limsup_{s \downarrow 0} \delta^{-1} \|E_{\mathbb{R} \setminus \Sigma_1}(H) (H - \theta_E(\alpha_1)) \Psi_s\| \leq c\epsilon_\gamma.$$

We have

$$(3.27) \quad \langle \Psi_s, E_{\Sigma_1}(H) [H, A'] \circ E_{\Sigma_1}(H) \Psi_s \rangle = \langle \Psi_s, [H, A'] \circ \Psi_s \rangle - f_1(s) - f_2(s), \quad \text{where} \\ f_1(s) = \langle \Psi_s, E_{\mathbb{R} \setminus \Sigma_1}(H) [H, A'] \circ E_{\Sigma_1}(H) \Psi_s \rangle, \quad \text{and} \quad f_2(s) = \langle \Psi_s, [H, A'] \circ E_{\mathbb{R} \setminus \Sigma_1}(H) \Psi_s \rangle.$$

By (3.26),

$$\max_{i=1,2} \limsup_{s \downarrow 0} |f_i(s)| \leq c\epsilon_\gamma.$$

This together with (3.15) and (3.27) implies

$$(3.28) \quad \limsup_{s \downarrow 0} \langle \Psi_s, E_{\Sigma_1}(H)[H, A'] \circ E_{\Sigma_1}(H)\Psi_s \rangle \leq c\epsilon_\gamma.$$

On the other hand, by the Mourre estimate (3.12), we have that

$$(3.29) \quad \langle \Psi_s, E_{\Sigma_1}(H)[H, A'] \circ E_{\Sigma_1}(H)\Psi_s \rangle \geq \eta \|E_{\Sigma_1}(H)\Psi_s\|^2 + \langle \Psi_s, E_{\Sigma_1}(H)KE_{\Sigma_1}(H)\Psi_s \rangle.$$

Thus, since Ψ_s converges weakly to zero and $E_{\Sigma_1}(H)KE_{\Sigma_1}(H)$ is compact, we have, using (3.26)

$$(3.30) \quad \liminf_{s \downarrow 0} \langle \Psi_s, E_{\Sigma_1}(H)[H, A'] \circ E_{\Sigma_1}(H)\Psi_s \rangle \geq \eta(1 - c\epsilon_\gamma^2).$$

Recall that $\epsilon_\gamma \rightarrow 0$ as $\gamma \rightarrow 0$. Taking first α_1 sufficiently close to α_0 , we can then take γ small enough to see that (3.30) contradicts (3.28). The proof is complete for the case $E \neq 2$.

Part 4. Case $E = 2$: the proof is almost the same as before but a bit simpler. We briefly go over the proof to point out the small adjustments. Assuming the statement of the theorem to be false, we have that $2 \in \Theta(H)$, and also that $\psi \notin \mathcal{D}(\vartheta_\alpha)$ for some $\alpha \in (0, \infty)$. Since $\Theta(H)$ is open, there is an interval

$$\Sigma := (2 - \delta, 2 + \delta)$$

such that the Mourre estimate holds there, i.e.

$$(3.31) \quad E_\Sigma(H)[H, A'] \circ E_\Sigma(H) \geq \eta E_\Sigma(H) + K$$

for some $\eta > 0$ and some compact operator K . Let $\alpha_0 := \inf\{\alpha \geq 0 : \psi \notin \mathcal{D}(\vartheta_\alpha)\}$. As before, let α_1 and γ be such that $\alpha_1 < \alpha_0 < \alpha_1 + \gamma$ if $\alpha_0 > 0$; if $\alpha_0 = 0$, let $\alpha_1 = 0$. Let F_s and Ψ_s be defined as before (see (3.18) and (3.13)), so that Ψ_s has norm one but converges weakly to zero. The calculation of Part 1 shows that

$$\limsup_{s \downarrow 0} \langle \Psi_s, [H, A'] \circ \Psi_s \rangle \leq c\epsilon_\gamma,$$

whereas the calculation of Part 2 shows that

$$\lim_{s \downarrow 0} \|(H - 2)\Psi_s\| \leq c\epsilon_\gamma.$$

The functional calculus then gives

$$\limsup_{s \downarrow 0} \|E_{\mathbb{R} \setminus \Sigma}(H)\Psi_s\| \leq \limsup_{s \downarrow 0} \delta^{-1} \|E_{\mathbb{R} \setminus \Sigma}(H)(H - 2)\Psi_s\| \leq c\epsilon_\gamma.$$

As in Part 3, we get inequalities (3.28) and (3.30) with Σ instead of Σ_1 . Taking α_1 very close to α_0 in order to take γ sufficiently small, these two inequalities disagree. The proof is complete. \square

It remains to show however that

$$(3.32) \quad H\psi = E\psi, \quad \text{and} \quad \psi \in \mathcal{D}(\vartheta_\alpha) \quad \text{for all} \quad \alpha \geq 0 \quad \text{implies} \quad \psi = 0.$$

We slightly modify the notation we have been using so far. Let

$$(3.33) \quad F_\alpha(n) := \alpha|n| \quad \text{and} \quad \psi_\alpha(n) := e^{F_\alpha(n)}\psi(n) = e^{\alpha|n|}\psi(n), \quad \text{for all } n \in \mathbb{Z}.$$

Proof of Theorem 1.1, the second part. The proof is by contradiction, and the strategy is as follows: we assume that $\psi \neq 0$ and define $\Psi_\alpha := \psi_\alpha / \|\psi_\alpha\|$. It is not hard to see that Ψ_α converges weakly to zero as $\alpha \rightarrow \infty$ (use the fact that the difference equation $H\psi = E\psi$ implies $\psi(n) \neq 0$ infinitely often). In the first part we apply Proposition 3.1 with F_α replacing F . In this case we

can exactly compute terms to show that

$$(3.34) \quad 0 = \cosh(\alpha)^{-1} \langle \Psi_\alpha, [V, A']_\circ \Psi_\alpha \rangle + 2 \tanh(\alpha) \|\sqrt{|N|}(S^* - S)\Psi_\alpha\|^2 \\ + \|\sqrt{\Delta(4 - \Delta)}\Psi_\alpha\|^2 - \tanh(\alpha) \left(2\Psi_\alpha^2(0) + (\Psi_\alpha(-1) - \Psi_\alpha(1))^2 \right).$$

In the second part, we apply Lemma 3.2 again with F_α replacing F . We show that

$$(3.35) \quad \lim_{\alpha \rightarrow \infty} \|\sqrt{\Delta(4 - \Delta)}\Psi_\alpha\|^2 = \lim_{\alpha \rightarrow \infty} \Re \langle \Psi_\alpha, \Delta(4 - \Delta)\Psi_\alpha \rangle = 2.$$

The conclusion is then imminent: taking the limit $\alpha \rightarrow \infty$ in (3.34), and recalling that $[V, A']_\circ$ exists as a bounded operator and Ψ_α converges weakly to zero leads to a contradiction.

Part 1. From the proof of (4.9),

$$(3.36) \quad \langle \psi_\alpha, [H, A']_\circ \psi_\alpha \rangle = -\|\sqrt{g_r - g_\ell}A'\psi_\alpha\|^2 + \langle \psi_\alpha, 2^{-1}(M + G_r + G_\ell)\psi_\alpha \rangle$$

where $M = M_r + M_\ell$ is given by (4.11) and $G_r + G_\ell$ by (4.15). All terms are computed exactly:

$$(3.37) \quad e^{(\tau F_\alpha - F_\alpha)(n)} = \begin{cases} e^{-\alpha} & \text{if } n \geq 1 \\ e^\alpha & \text{if } n \leq 0 \end{cases} \quad \text{and} \quad e^{(\tau^* F_\alpha - F_\alpha)(n)} = \begin{cases} e^\alpha & \text{if } n \geq 0 \\ e^{-\alpha} & \text{if } n \leq -1, \end{cases}$$

$$(3.38) \quad e^{(F_\alpha - \tau F_\alpha)(n)} = \begin{cases} e^\alpha & \text{if } n \geq 1 \\ e^{-\alpha} & \text{if } n \leq 0 \end{cases} \quad \text{and} \quad e^{(F_\alpha - \tau^* F_\alpha)(n)} = \begin{cases} e^{-\alpha} & \text{if } n \geq 0 \\ e^\alpha & \text{if } n \leq -1. \end{cases}$$

Let δ_B be the projector onto $B \subset \mathbb{Z}$. Therefore

$$\begin{aligned} \tau^* \varphi_\ell - \varphi_\ell &= -2 \sinh(\alpha) \delta_{\{n=0\}}, & \varphi_\ell - \tau \varphi_\ell &= -2 \sinh(\alpha) \delta_{\{n=+1\}}, \\ \tau \varphi_r - \varphi_r &= -2 \sinh(\alpha) \delta_{\{n=0\}}, & \varphi_r - \tau^* \varphi_r &= -2 \sinh(\alpha) \delta_{\{n=-1\}}, \\ \tau^* \varphi_\ell - \tau^{*2} \varphi_\ell &= 2 \sinh(\alpha) \delta_{\{n=-1\}}, & \tau \varphi_r - \tau^2 \varphi_r &= 2 \sinh(\alpha) \delta_{\{n=+1\}}. \end{aligned}$$

We expand the first term on the right side of (3.36), using $A' = 2^{-1}(S^* + S) + N(S^* - S)$:

$$(3.39) \quad -\|\sqrt{g_r - g_\ell}A'\psi_\alpha\|^2 = -\langle A'\psi_\alpha, (g_r - g_\ell)A'\psi_\alpha \rangle = -4^{-1} \langle \psi_\alpha, (S^* + S)(g_r - g_\ell)(S^* + S)\psi_\alpha \rangle \\ - 2^{-1} \langle (S^* + S)\psi_\alpha, (g_r - g_\ell)N(S^* - S)\psi_\alpha \rangle$$

$$(3.40) \quad - 2^{-1} \langle N(S^* - S)\psi_\alpha, (g_r - g_\ell)(S^* + S)\psi_\alpha \rangle$$

$$(3.41) \quad - \langle N(S^* - S)\psi_\alpha, (g_r - g_\ell)N(S^* - S)\psi_\alpha \rangle.$$

First,

$$(g_r - g_\ell)N = 2 \sinh(\alpha) \delta_{\{n \neq 0\}} \text{sign}(N) \quad \text{implies} \quad (3.41) = -2 \sinh(\alpha) \|\sqrt{|N|}(S^* - S)\psi_\alpha\|^2.$$

As for (3.39) and (3.40), we have

$$(3.39) + (3.40) = -\sinh(\alpha) \left[\langle \psi_\alpha, (S^* + S) \delta_{\{n \neq 0\}} \text{sign}(N) (S^* - S) \psi_\alpha \rangle \right. \\ \left. + \langle \psi_\alpha, (S^* - S) \delta_{\{n \neq 0\}} \text{sign}(N) (S^* + S) \psi_\alpha \rangle \right].$$

The following commutation formulas hold

$$(3.42) \quad S^*(\delta_{\{n \neq 0\}} \text{sign}(N)) = [\delta_{\{n \neq 0\}} \text{sign}(N) + \delta_{\{n=0\}} + \delta_{\{n=-1\}}] S^*$$

$$(3.43) \quad S(\delta_{\{n \neq 0\}} \text{sign}(N)) = [\delta_{\{n \neq 0\}} \text{sign}(N) - \delta_{\{n=0\}} - \delta_{\{n=+1\}}] S.$$

Hence we commute S and S^* with $\delta_{\{n \neq 0\}} \text{sign}(N)$ and cancel terms. We get

$$(3.39) + (3.40) = 2 \sinh(\alpha) \langle \psi_\alpha, [2\delta_{\{n=0\}} + \delta_{\{n=-1\}} + \delta_{\{n=+1\}}] \psi_\alpha \rangle.$$

We therefore have

$$(3.44) \quad -\|\sqrt{g_r - g_\ell} A' \psi_\alpha\|^2 = -4^{-1} \langle \psi_\alpha, (S^* + S)(g_r - g_\ell)(S^* + S)\psi_\alpha \rangle$$

$$(3.45) \quad + 2 \sinh(\alpha) \langle \psi_\alpha, [2\delta_{\{n=0\}} + \delta_{\{n=-1\}} + \delta_{\{n=+1\}}] \psi_\alpha \rangle$$

$$(3.46) \quad - 2 \sinh(\alpha) \|\sqrt{|N|}(S^* - S)\psi_\alpha\|^2.$$

We analyze the second term on the right side of (3.36). A straightforward calculation gives

$$(3.47) \quad 2^{-1}M = 4^{-1}(S^* + S)(g_r - g_\ell)(S^* + S) - \sinh(\alpha)[2\delta_{\{n=0\}} + \delta_{\{n=+1\}} + \delta_{\{n=-2\}}], \quad \text{and}$$

$$(3.48) \quad 2^{-1}(G_r + G_\ell) = -\sinh(\alpha)[\delta_{\{n=-1\}}S^{*2} + \delta_{\{n=+1\}}S^2] - (\cosh(\alpha) - 1)\Delta(4 - \Delta).$$

Assembling (3.44)-(3.46), (3.47) and (3.48) to create (3.36) gives

$$\begin{aligned} \langle \psi_\alpha, [H, A']_o \psi_\alpha \rangle &= -2 \sinh(\alpha) \|\sqrt{|N|}(S^* - S)\psi_\alpha\|^2 - \langle \psi_\alpha, (\cosh(\alpha) - 1)\Delta(4 - \Delta)\psi_\alpha \rangle \\ &\quad + \sinh(\alpha) \left(2\psi_\alpha^2(0) + (\psi_\alpha(-1) - \psi_\alpha(1))^2 \right). \end{aligned}$$

Cancelling $\langle \psi_\alpha, [\Delta, A']\psi_\alpha \rangle = \langle \psi_\alpha, \Delta(4 - \Delta)\psi_\alpha \rangle$ on both sides and dividing throughout by $\cosh(\alpha)\|\psi_\alpha\|^2$ yields (3.34) as required.

Part 2. From (3.38),

$$\begin{aligned} 2^{-1}(e^{F_\alpha - \tau F_\alpha} + e^{F_\alpha - \tau^* F_\alpha}) &= \begin{cases} \cosh(\alpha) & \text{if } |n| \geq 1 \\ e^{-\alpha} & \text{if } n = 0, \end{cases} \\ 2^{-1}(e^{F_\alpha - \tau F_\alpha} - e^{F_\alpha - \tau^* F_\alpha}) &= \sinh(\alpha)\delta_{\{n \neq 0\}}\text{sign}(N). \end{aligned}$$

We apply (3.8) of Lemma 3.2:

$$\begin{aligned} H_{F_\alpha} &= \cosh(\alpha)\Delta + \delta_{\{n=0\}}(e^{-\alpha} - \cosh(\alpha))\Delta + V + 2(1 - \cosh(\alpha)) \\ &\quad + 2\delta_{\{n=0\}}(\cosh(\alpha) - e^{-\alpha}) + \sinh(\alpha)\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S). \end{aligned}$$

The goal is to square H_{F_α} . Divide throughout by $\cosh(\alpha)$ and let $c_\alpha := (e^{-\alpha} \cosh(\alpha)^{-1} - 1)$:

$$(3.49) \quad \cosh(\alpha)^{-1}H_{F_\alpha} = \Delta + c_\alpha\delta_{\{n=0\}}\Delta + \cosh(\alpha)^{-1}V + 2(\cosh(\alpha)^{-1} - 1) - 2c_\alpha\delta_{\{n=0\}}$$

$$(3.50) \quad + \tanh(\alpha)\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S).$$

Note that $\sup_{\alpha \geq 0} |c_\alpha| \leq 2$. Since $(S^* - S)$ is antisymmetric, by (3.42) and (3.43), we see that $\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)$ is antisymmetric up to a couple of rank one projectors. The same goes for $\Delta\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)$ and $\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)\Delta$. Therefore

$$\lim_{\alpha \rightarrow \infty} \Re \langle \Psi_\alpha, [\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)]\Psi_\alpha \rangle = 0,$$

$$\lim_{\alpha \rightarrow \infty} \Re \langle \Psi_\alpha, \Delta[\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)]\Psi_\alpha \rangle = 0,$$

$$\lim_{\alpha \rightarrow \infty} \Re \langle \Psi_\alpha, [\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)]\Delta\Psi_\alpha \rangle = 0.$$

We compute $[\tanh(\alpha)\delta_{\{n \neq 0\}}\text{sign}(N)(S^* - S)]^2$ using (3.42) and (3.43):

$$(3.50)^2 = \tanh^2(\alpha) \left[\delta_{\{n \neq 0\}}(S^2 + (S^*)^2 - 2) + \delta_{\{n=-1\}}(1 - (S^*)^2) + \delta_{\{n=+1\}}(S^2 - 1) \right].$$

Thus squaring $\cosh(\alpha)^{-1}H_{F_\alpha}$ given by (3.49)-(3.50) and recalling that $\Delta(4 - \Delta) = 2 - S^2 - (S^*)^2$ we get

$$\cosh(\alpha)^{-2}H_{F_\alpha}^2 = \Delta(\Delta - 4) + 4 - \tanh^2(\alpha)\Delta(4 - \Delta) + P_\alpha,$$

where P_α is a bounded operator satisfying

$$\lim_{\alpha \rightarrow \infty} \Re \langle \Psi_\alpha, P_\alpha \Psi_\alpha \rangle = 0.$$

Rearranging and recalling that $H_{F_\alpha}\Psi_\alpha = E\Psi_\alpha$ yields (3.35) as required. \square

4. APPENDIX : TECHNICAL CALCULATIONS

The appendix is devoted to proving the key relations (2.13) and (3.3) that appear in Propositions 2.1 and 3.1 respectively. Recall that for $B \subset \mathbb{Z}^d$, δ_B denotes the projector onto B . We start with the proof of the multidimensional formula

$$(4.1) \quad \begin{aligned} \langle \phi, [e^F A' e^F, \Delta] \phi \rangle &= \langle \phi_F, [A', \Delta] \phi_F \rangle - 2 \|\sqrt{g} A' \phi_F\|^2 \\ &\quad - \sum_{i=1}^d \|\sqrt{\Delta_i(4 - \Delta_i)} W_i \phi_F\|^2 + 2^{-1} \langle \phi_F, (\mathcal{L} + \mathcal{M} + \mathcal{G}) \phi_F \rangle, \end{aligned}$$

where $\phi \in \ell_0(\mathbb{Z}^d)$ and $\phi_F := e^F \phi$. To jump to the proof of the $1d$ relation, go to (4.9).

Proof. It is understood that the operators are calculated and the commutators developed against $\phi \in \ell_0(\mathbb{Z}^d)$, so we omit the ϕ for ease of notation. Usual commutation relations give

$$[e^F A' e^F, \Delta] = e^F [A', \Delta] e^F + e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F.$$

We now concentrate on the second and third terms on the right side of the latter relation. The goal is to pop out $e^F A' g A' e^F$ and control the remainder. As pointed out in [FH] and [CFKS], this is the key quantity to single out. The following commutators will be used repeatedly:

$$\begin{aligned} [e^F, S_i] &= -(\tau_i e^F - e^F) S_i = S_i (\tau_i^* e^F - e^F) = -e^F \varphi_{\ell_i} S_i = S_i \varphi_{r_i} e^F, \\ [e^F, S_i^*] &= -(\tau_i^* e^F - e^F) S_i^* = S_i^* (\tau_i e^F - e^F) = -e^F \varphi_{r_i} S_i^* = S_i^* \varphi_{\ell_i} e^F. \end{aligned}$$

Part 1 : Creating $e^F A' g A' e^F$ in a first way. We have

$$\begin{aligned} [e^F, \Delta_i] &= \varphi_{r_i} e^F S_i^* + \varphi_{\ell_i} e^F S_i \\ &= g_{r_i} N_i e^F S_i^* + \varphi_{r_i} \delta_{\{n_i=0\}} e^F S_i^* + \varphi_{\ell_i} e^F S_i \\ &= g_{r_i} N_i e^F (S_i^* - S_i) + \varphi_{r_i} \delta_{\{n_i=0\}} e^F (S_i^* - S_i) + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i \\ &= g_{r_i} N_i (S_i^* - S_i) e^F + g_{r_i} N_i [e^F, (S_i^* - S_i)] + \varphi_{r_i} \delta_{\{n_i=0\}} e^F (S_i^* - S_i) + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i \\ &= g N_i (S_i^* - S_i) e^F + (g_{r_i} - g) N_i (S_i^* - S_i) e^F + \varphi_{r_i} \delta_{\{n_i=0\}} (S_i^* - S_i) e^F \\ &\quad + \varphi_{r_i} [e^F, (S_i^* - S_i)] + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i \\ &= g A'_i e^F - 2^{-1} g (S_i^* + S_i) e^F + (g_{r_i} - g) N_i (S_i^* - S_i) e^F + \varphi_{r_i} \delta_{\{n_i=0\}} (S_i^* - S_i) e^F \\ &\quad + \varphi_{r_i} [e^F, (S_i^* - S_i)] + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i. \end{aligned}$$

$$\begin{aligned} [e^F, \Delta_i] &= -S_i e^F \varphi_{r_i} - S_i^* e^F \varphi_{\ell_i} \\ &= -S_i e^F N_i g_{r_i} - S_i e^F \varphi_{r_i} \delta_{\{n_i=0\}} - S_i^* e^F \varphi_{\ell_i} \\ &= (S_i^* - S_i) e^F N_i g_{r_i} + (S_i^* - S_i) e^F \varphi_{r_i} \delta_{\{n_i=0\}} - S_i^* e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\ &= e^F (S_i^* - S_i) N_i g_{r_i} + [(S_i^* - S_i), e^F] N_i g_{r_i} + (S_i^* - S_i) e^F \varphi_{r_i} \delta_{\{n_i=0\}} - S_i^* e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\ &= e^F (S_i^* - S_i) N_i g + e^F (S_i^* - S_i) N_i (g_{r_i} - g) + e^F (S_i^* - S_i) \varphi_{r_i} \delta_{\{n_i=0\}} \\ &\quad - [e^F, (S_i^* - S_i)] \varphi_{r_i} - S_i^* e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\ &= e^F A'_i g + 2^{-1} e^F (S_i^* + S_i) g + e^F (S_i^* - S_i) N_i (g_{r_i} - g) + e^F (S_i^* - S_i) \varphi_{r_i} \delta_{\{n_i=0\}} \\ &\quad - [e^F, (S_i^* - S_i)] \varphi_{r_i} - S_i^* e^F (\varphi_{r_i} + \varphi_{\ell_i}). \end{aligned}$$

Therefore we have obtained

$$(4.2) \quad \begin{aligned} e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F &= 2e^F A' g A' e^F + e^F (L_r + M_r + G_r + H_r) e^F, \quad \text{where} \\ L_r &:= \sum_{i,j} A'_i (g_{r_j} - g) N_j (S_j^* - S_j) + (S_i^* - S_i) N_i (g_{r_i} - g) A'_j, \end{aligned}$$

$$\begin{aligned}
M_r &:= 2^{-1} \sum_{i,j} -A'_i g(S_j^* + S_j) + (S_i^* + S_i) g A'_j, \\
G_r &:= \sum_{i,j} A'_i (\varphi_{r_j} [e^F, (S_j^* - S_j)] e^{-F} + (\varphi_{r_j} + \varphi_{l_j}) e^F S_j e^{-F}) \\
&\quad - \sum_{i,j} (e^{-F} [e^F, (S_i^* - S_i)] \varphi_{r_i} + e^{-F} S_i^* e^F (\varphi_{r_i} + \varphi_{l_i})) A'_j, \quad \text{and} \\
H_r &:= \sum_{i,j} A'_i \varphi_{r_j} \delta_{\{n_j=0\}} (S_j^* - S_j) + (S_i^* - S_i) \varphi_{r_i} \delta_{\{n_i=0\}} A_j.
\end{aligned}$$

We split M_r as follows: $M_r = M_{r;1} + M_{r;2}$, where

$$\begin{aligned}
M_{r;1} &:= 2^{-1} \sum_{i \neq j} -A'_i g(S_j^* + S_j) + (S_i^* + S_i) g A'_j, \\
M_{r;2} &:= 2^{-1} \sum_i -A'_i g(S_i^* + S_i) + (S_i^* + S_i) g A'_i = M_{r;2;1} + M_{r;2;2}, \quad \text{with} \\
M_{r;2;1} &:= 2^{-1} \sum_i -A'_i g_{r_i} (S_i^* + S_i) + (S_i^* + S_i) g_{r_i} A'_i, \\
M_{r;2;2} &:= 2^{-1} \sum_i -A'_i (g - g_{r_i}) (S_i^* + S_i) + (S_i^* + S_i) (g - g_{r_i}) A'_i.
\end{aligned}$$

We calculate $M_{r;1}$ by expanding A'_i and A'_j :

$$\begin{aligned}
M_{r;1} &:= 2^{-1} \sum_{i \neq j} -N_i (S_i^* - S_i) g (S_j^* + S_j) + (S_i^* + S_i) g N_j (S_j^* - S_j) \\
&= 2^{-1} \sum_{i \neq j} -N_i [(\tau_i^* g) S_i^* - (\tau_i g) S_i] (S_j^* + S_j) + [(\tau_i^* (g N_j)) S_i^* + (\tau_i (g N_j)) S_i] (S_j^* - S_j) \\
&= \frac{1}{2} \sum_{i \neq j} N_i [\tau_i g - \tau_j g] S_i S_j + N_i [\tau_j^* g - \tau_i^* g] S_i^* S_j^* + [N_i (\tau_j g - \tau_i^* g) + N_j (\tau_j g - \tau_i^* g)] S_i^* S_j. \quad (\ddagger 2)
\end{aligned}$$

Again expanding A'_i :

$$\begin{aligned}
M_{r;2;1} &= 2^{-1} \sum_i (S_i^* + S_i) g_{r_i} (S_i^* + S_i) - (S_i^* - S_i) \varphi_{r_i} \delta_{\{n_i \neq 0\}} (S_i^* + S_i) \\
&\quad + 2^{-1} \sum_i (S_i^* + S_i) \varphi_{r_i} \delta_{\{n_i \neq 0\}} (S_i^* - S_i) \\
&= \sum_i 2^{-1} (S_i^* + S_i) g_{r_i} (S_i^* + S_i) + S_i \varphi_{r_i} S_i^* - S_i^* \varphi_{r_i} S_i + (S_i^* \varphi_{r_i} \delta_{\{n_i=0\}} S_i - S_i \varphi_{r_i} \delta_{\{n_i=0\}} S_i^*) \\
&= M_{r;2;1;1} + M_{r;2;1;2}, \quad \text{where} \\
M_{r;2;1;1} &:= \sum_i 2^{-1} (S_i^* + S_i) g_{r_i} (S_i^* + S_i) + (\tau_i \varphi_{r_i} - \tau_i^* \varphi_{r_i}), \quad (\ddagger 1, \ddagger 3) \\
M_{r;2;1;2} &:= \sum_i (\tau_i^* \varphi_{r_i}) \delta_{\{n_i=-1\}} - (\tau_i \varphi_{r_i}) \delta_{\{n_i=+1\}}.
\end{aligned}$$

We calculate G_r . We note that

$$(4.3) \quad (\tau_i \varphi_{r_i}) \varphi_{l_i} = \varphi_{l_i} (\tau_i \varphi_{r_i}) = -(\tau_i \varphi_{r_i} + \varphi_{l_i}), \quad \text{and} \quad (\tau_i^* \varphi_{l_i}) \varphi_{r_i} = \varphi_{r_i} (\tau_i^* \varphi_{l_i}) = -(\tau_i^* \varphi_{l_i} + \varphi_{r_i}).$$

$$\begin{aligned}
G_r &:= \sum_{i,j} A'_i (\varphi_{r_j} (S_j^* \varphi_{\ell_j} - S_j \varphi_{r_j}) + (\varphi_{r_j} + \varphi_{\ell_j}) S_j (\varphi_{r_j} + 1)) \\
&\quad - \sum_{i,j} ((-\varphi_{r_i} S_i^* + \varphi_{\ell_i} S_i) \varphi_{r_i} + (\varphi_{r_i} + 1) S_i^* (\varphi_{r_i} + \varphi_{\ell_i})) A'_j \\
&= \sum_{i,j} A'_i (S_j^* (\tau_j \varphi_{r_j}) \varphi_{\ell_j} - S_j (\tau_j^* \varphi_{r_j}) \varphi_{r_j} + S_j (\tau_j^* \varphi_{r_j}) (\varphi_{r_j} + 1) + S_j (\tau_j^* \varphi_{\ell_j}) (\varphi_{r_j} + 1)) \\
&\quad + \sum_{i,j} (\varphi_{r_i} (\tau_i^* \varphi_{r_i}) S_i^* - \varphi_{\ell_i} (\tau_i \varphi_{r_i}) S_i - (\varphi_{r_i} + 1) (\tau_i^* \varphi_{r_i}) S_i^* - (\varphi_{r_i} + 1) (\tau_i^* \varphi_{\ell_i}) S_i^*) A'_j \\
&= \sum_{i,j} A'_i (-S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + S_j (\tau_j^* \varphi_{r_j} - \varphi_{r_j})) + \sum_{i,j} ((\varphi_{r_i} - \tau_i^* \varphi_{r_i}) S_i^* + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i) A'_j \\
&= G_{r;1} + G_{r;2}, \quad \text{where}
\end{aligned}$$

$$G_{r;1} := \sum_{i,j} A'_i S_j (\tau_j^* \varphi_{r_j} - \varphi_{r_j}) + (\varphi_{r_i} - \tau_i^* \varphi_{r_i}) S_i^* A'_j, \quad (\dagger 3)$$

$$G_{r;2} := \sum_{i,j} -A'_i S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i A'_j.$$

To end this section we note that we are left to deal with $L_r + M_{r;2;1;2} + M_{r;2;2} + G_{r;2} + H_r$.

Part 2 : Creating $e^F A' g A' e^F$ a second way. We repeat the calculation with a variation.

$$\begin{aligned}
[e^F, \Delta_i] &= \varphi_{\ell_i} e^F S_i + \varphi_{r_i} e^F S_i^* \\
&= g_{\ell_i} N_i e^F S_i + \varphi_{\ell_i} \delta_{\{n_i=0\}} e^F S_i + \varphi_{r_i} e^F S_i^* \\
&= -g_{\ell_i} N_i e^F (S_i^* - S_i) + \varphi_{\ell_i} \delta_{\{n_i=0\}} e^F (S_i - S_i^*) + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i^* \\
&= -g_{\ell_i} N_i (S_i^* - S_i) e^F - g_{\ell_i} N_i [e^F, (S_i^* - S_i)] + \varphi_{\ell_i} \delta_{\{n_i=0\}} e^F (S_i - S_i^*) + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i^* \\
&= g N_i (S_i^* - S_i) e^F - (g_{\ell_i} + g) N_i (S_i^* - S_i) e^F + \varphi_{\ell_i} \delta_{\{n_i=0\}} (S_i - S_i^*) e^F \\
&\quad - \varphi_{\ell_i} [e^F, (S_i^* - S_i)] + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i^* \\
&= g A'_i e^F - 2^{-1} g (S_i^* + S_i) e^F - (g_{\ell_i} + g) N_i (S_i^* - S_i) e^F + \varphi_{\ell_i} \delta_{\{n_i=0\}} (S_i - S_i^*) e^F \\
&\quad - \varphi_{\ell_i} [e^F, (S_i^* - S_i)] + (\varphi_{r_i} + \varphi_{\ell_i}) e^F S_i^*.
\end{aligned}$$

$$\begin{aligned}
[e^F, \Delta_i] &= -S_i^* e^F \varphi_{\ell_i} - S_i e^F \varphi_{r_i} \\
&= -S_i^* e^F N_i g_{\ell_i} - S_i^* e^F \varphi_{\ell_i} \delta_{\{n_i=0\}} - S_i e^F \varphi_{r_i} \\
&= -(S_i^* - S_i) e^F N_i g_{\ell_i} + (S_i - S_i^*) e^F \varphi_{\ell_i} \delta_{\{n_i=0\}} - S_i e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\
&= -e^F (S_i^* - S_i) N_i g_{\ell_i} - [(S_i^* - S_i), e^F] N_i g_{\ell_i} + (S_i - S_i^*) e^F \varphi_{\ell_i} \delta_{\{n_i=0\}} - S_i e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\
&= e^F (S_i^* - S_i) N_i g - e^F (S_i^* - S_i) N_i (g_{\ell_i} + g) + e^F (S_i - S_i^*) \varphi_{\ell_i} \delta_{\{n_i=0\}} \\
&\quad + [e^F, (S_i^* - S_i)] \varphi_{\ell_i} - S_i e^F (\varphi_{r_i} + \varphi_{\ell_i}) \\
&= e^F A'_i g + 2^{-1} e^F (S_i^* + S_i) g - e^F (S_i^* - S_i) N_i (g_{\ell_i} + g) + e^F (S_i - S_i^*) \varphi_{\ell_i} \delta_{\{n_i=0\}} \\
&\quad + [e^F, (S_i^* - S_i)] \varphi_{\ell_i} - S_i e^F (\varphi_{r_i} + \varphi_{\ell_i})
\end{aligned}$$

Therefore we have obtained

$$(4.4) \quad e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F = 2e^F A' g A' e^F + e^F (L_\ell + M_\ell + G_\ell + H_\ell) e^F, \quad \text{where}$$

$$L_\ell := - \sum_{i,j} A'_i (g_{\ell_j} + g) N_j (S_j^* - S_j) + (S_i^* - S_i) N_i (g_{\ell_i} + g) A'_j,$$

$$\begin{aligned}
M_\ell &:= 2^{-1} \sum_{i,j} -A'_i g(S_j^* + S_j) + (S_i^* + S_i) g A'_j, \\
G_\ell &:= \sum_{i,j} A'_i (-\varphi_{\ell_j} [e^F, (S_j^* - S_j)] e^{-F} + (\varphi_{r_j} + \varphi_{\ell_j}) e^F S_j^* e^{-F}) \\
&\quad + \sum_{i,j} (e^{-F} [e^F, (S_i^* - S_i)] \varphi_{\ell_i} - e^{-F} S_i e^F (\varphi_{r_i} + \varphi_{\ell_i})) A'_j, \quad \text{and} \\
H_\ell &:= \sum_{i,j} A'_i \varphi_{\ell_j} \delta_{\{n_j=0\}} (S_j - S_j^*) + (S_i - S_i^*) \varphi_{\ell_i} \delta_{\{n_i=0\}} A'_j.
\end{aligned}$$

We split M_ℓ as follows: $M_\ell := M_{\ell;1} + M_{\ell;2}$, where

$$\begin{aligned}
M_{\ell;1} &:= 2^{-1} \sum_{i \neq j} -A'_i g(S_j^* + S_j) + (S_i^* + S_i) g A'_j, \\
M_{\ell;2} &:= 2^{-1} \sum_i -A'_i g(S_i^* + S_i) + (S_i^* + S_i) g A'_i = M_{\ell;2;1} + M_{\ell;2;2}, \quad \text{with} \\
M_{\ell;2;1} &:= 2^{-1} \sum_i A'_i g_{\ell_i} (S_i^* + S_i) - (S_i^* + S_i) g_{\ell_i} A'_i, \\
M_{\ell;2;2} &:= 2^{-1} \sum_i -A'_i (g + g_{\ell_i}) (S_i^* + S_i) + (S_i^* + S_i) (g + g_{\ell_i}) A'_i.
\end{aligned}$$

We calculate $M_{\ell;1}$ by expanding A'_i and A'_j :

$$\begin{aligned}
M_{\ell;1} &= 2^{-1} \sum_{i \neq j} -N_i (S_i^* - S_i) g (S_j^* + S_j) + (S_i^* + S_i) g N_j (S_j^* - S_j) \\
&= 2^{-1} \sum_{i \neq j} -N_i [(\tau_i^* g) S_i^* - (\tau_i g) S_i] (S_j^* + S_j) + [(\tau_i^* (g N_j)) S_i^* + (\tau_i (g N_j)) S_i] (S_j^* - S_j) \\
&= \frac{1}{2} \sum_{i \neq j} N_i [\tau_i g - \tau_j g] S_i S_j + N_i [\tau_j^* g - \tau_i^* g] S_i^* S_j^* + [N_i (\tau_j g - \tau_i^* g) + N_j (\tau_j g - \tau_i^* g)] S_i^* S_j. \quad (\ddagger_2)
\end{aligned}$$

Again expanding A'_i :

$$\begin{aligned}
M_{\ell;2;1} &= 2^{-1} \sum_i -(S_i^* + S_i) g_{\ell_i} (S_i^* + S_i) + (S_i^* - S_i) \varphi_{\ell_i} \delta_{\{n_i \neq 0\}} (S_i^* + S_i) \\
&\quad - 2^{-1} \sum_i (S_i^* + S_i) \varphi_{\ell_i} \delta_{\{n_i \neq 0\}} (S_i^* - S_i) \\
&= \sum_i -2^{-1} (S_i^* + S_i) g_{\ell_i} (S_i^* + S_i) + S_i^* \varphi_{\ell_i} S_i - S_i \varphi_{\ell_i} S_i^* + (S_i \varphi_{\ell_i} \delta_{\{n_i=0\}} S_i^* - S_i^* \varphi_{\ell_i} \delta_{\{n_i=0\}} S_i) \\
&= M_{\ell;2;1;1} + M_{\ell;2;1;2}, \quad \text{where} \\
M_{\ell;2;1;1} &:= \sum_i -2^{-1} (S_i^* + S_i) g_{\ell_i} (S_i^* + S_i) + (\tau_i^* \varphi_{\ell_i} - \tau_i \varphi_{\ell_i}), \quad (\ddagger_1, \ddagger_3) \\
M_{\ell;2;1;2} &:= \sum_i (\tau_i \varphi_{\ell_i}) \delta_{\{n_i=+1\}} - (\tau_i^* \varphi_{\ell_i}) \delta_{\{n_i=-1\}}.
\end{aligned}$$

We calculate G_ℓ :

$$\begin{aligned}
G_\ell &:= \sum_{i,j} A'_i (-\varphi_{\ell_j} (S_j^* \varphi_{\ell_j} - S_j \varphi_{r_j}) + (\varphi_{r_j} + \varphi_{\ell_j}) S_j^* (\varphi_{\ell_j} + 1)) \\
&\quad + \sum_{i,j} ((-\varphi_{r_i} S_i^* + \varphi_{\ell_i} S_i) \varphi_{\ell_i} - (\varphi_{\ell_i} + 1) S_i (\varphi_{r_i} + \varphi_{\ell_i})) A'_j \\
&= \sum_{i,j} A'_i (-S_j^* (\tau_j \varphi_{\ell_j}) \varphi_{\ell_j} + S_j (\tau_j^* \varphi_{\ell_j}) \varphi_{r_j} + S_j^* (\tau_j \varphi_{r_j}) (\varphi_{\ell_j} + 1) + S_j^* (\tau_j \varphi_{\ell_j}) (\varphi_{\ell_j} + 1)) \\
&\quad + \sum_{i,j} (-\varphi_{r_i} (\tau_i^* \varphi_{\ell_i}) S_i^* + \varphi_{\ell_i} (\tau_i \varphi_{\ell_i}) S_i - (\varphi_{\ell_i} + 1) (\tau_i \varphi_{r_i}) S_i - (\varphi_{\ell_i} + 1) (\tau_i \varphi_{\ell_i}) S_i) A'_j \\
&= \sum_{i,j} A'_i (S_j^* (\tau_j \varphi_{\ell_j} - \varphi_{\ell_j}) - S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j})) + \sum_{i,j} ((\varphi_{\ell_i} - \tau_i \varphi_{\ell_i}) S_i + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^*) A'_j \\
&= G_{\ell;1} + G_{\ell;2}, \quad \text{where}
\end{aligned}$$

$$G_{\ell;1} := \sum_{i,j} A'_i S_j^* (\tau_j \varphi_{\ell_j} - \varphi_{\ell_j}) + (\varphi_{\ell_i} - \tau_i \varphi_{\ell_i}) S_i A'_j, \quad (\ddagger_3)$$

$$G_{\ell;2} := \sum_{i,j} -A'_i S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* A'_j.$$

Note that we are left to deal with $L_\ell + M_{\ell;2;1;2} + M_{\ell;2;2} + G_{\ell;2} + H_\ell$.

Part 3 : Adding the terms of Parts 1 and 2. Take the average of (4.2) and (4.4):

$$\begin{aligned}
(4.5) \quad [e^F A' e^F, \Delta] &= e^F [A', \Delta] e^F + 2e^F A' g A' e^F \\
&\quad + 2^{-1} e^F (L_r + L_\ell + M_r + M_\ell + G_r + G_\ell + H_r + H_\ell) e^F.
\end{aligned}$$

Applying $\phi \in \ell_0(\mathbb{Z}^d)$ to this equation and taking inner products leads to (4.1). We go into details. The terms that still have to be dealt with are $L_r + M_{r;2;1;2} + M_{r;2;2} + G_{r;2} + H_r$ from the first part and $L_\ell + M_{\ell;2;1;2} + M_{\ell;2;2} + G_{\ell;2} + H_\ell$ from the second part. Since

$$(\tau_i^* \varphi_{r_i} - \tau_i^* \varphi_{\ell_i}) \delta_{\{n_i=-1\}} \phi = (\tau_i \varphi_{\ell_i} - \tau_i \varphi_{r_i}) \delta_{\{n_i=+1\}} \phi = 0, \quad \text{and} \quad (\varphi_{r_i} - \varphi_{\ell_i}) \delta_{\{n_i=0\}} \phi = 0,$$

it follows that

$$(M_{r;2;1;2} + M_{\ell;2;1;2}) \phi = 0, \quad \text{and} \quad (H_r + H_\ell) \phi = 0.$$

We add L_r and L_ℓ and define this to be \mathcal{L} :

$$\begin{aligned}
(4.6) \quad \mathcal{L} := L_r + L_\ell &= \sum_{i,j} A'_i [(g_{r_j} - g) - (g_{\ell_j} + g)] N_j (S_j^* - S_j) \\
&\quad + \sum_{i,j} (S_i^* - S_i) N_i [(g_{r_i} - g) - (g_{\ell_i} + g)] A'_j. \quad (\ddagger_4)
\end{aligned}$$

We add $M_{r;2;2}$ and $M_{\ell;2;2}$:

$$M_{r;2;2} + M_{\ell;2;2} = 2^{-1} \sum_i A'_i [(g_{r_i} - g) - (g_{\ell_i} + g)] (S_i^* + S_i) - (S_i^* + S_i) [(g_{r_i} - g) - (g_{\ell_i} + g)] A'_i. \quad (\ddagger_4)$$

We can now define \mathcal{M} :

$$(4.7) \quad \mathcal{M} := M_r + M_\ell = M_{r;1} + M_{r;2;1;1} + M_{\ell;1} + M_{\ell;2;1;1} + (M_{r;2;2} + M_{\ell;2;2}).$$

The final step is to add $G_{r;2}$ and $G_{\ell;2}$:

$$\begin{aligned}
G_{r;2} + G_{\ell;2} &= \sum_{i,j} -A_i S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i A_j \\
&\quad + \sum_{i,j} -A_i S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* A_j \\
&= - \sum_{i,j} [2^{-1}(S_i^* + S_i) + N_i(S_i^* - S_i)] S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) \\
&\quad + \sum_{i,j} (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i [-2^{-1}(S_j^* + S_j) + (S_j^* - S_j) N_j] \\
&\quad - \sum_{i,j} [2^{-1}(S_i^* + S_i) + N_i(S_i^* - S_i)] S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) \\
&\quad + \sum_{i,j} (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* [-2^{-1}(S_j^* + S_j) + (S_j^* - S_j) N_j] \\
&= G_1 + G_2 + G_3 + G_4 + G_5 + G_6, \quad \text{where}
\end{aligned}$$

$$G_1 := \sum_{i,j} -N_i S_i^* S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j^* N_j,$$

$$G_2 := \sum_{i,j} N_i S_i S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) - (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j N_j,$$

$$G_3 := \sum_{i,j} N_i S_i S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) - N_i S_i^* S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) - (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j N_j + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j^* N_j,$$

$$G_4 := -2^{-1} \sum_{i,j} S_i^* S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j^*,$$

$$G_5 := -2^{-1} \sum_{i,j} (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j + S_i S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}),$$

$$G_6 := -2^{-1} \sum_{i,j} S_i S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j^* + S_i^* S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j.$$

We calculate G_i for $i = 1 \dots 6$. $G_1 = G_{1;1} + G_{1;2} + G_{1;3}$, with

$$G_{1;1} := \sum_{i,j} [(\tau_j^* \varphi_{\ell_j} - \tau_i^* \tau_j^* \varphi_{\ell_j}) + (\varphi_{r_j} - \tau_i^* \varphi_{r_j})] N_i S_i^* S_j^*, \quad (\dagger_3)$$

$$G_{1;2} := \sum_{i \neq j} (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j^* \quad \text{and} \quad G_{1;3} := 2 \sum_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) (S_i^*)^2.$$

$$G_2 = G_{2;1} + G_{2;2} + G_{2;3}, \quad \text{where}$$

$$G_{2;1} := \sum_{i,j} [(\tau_i \varphi_{\ell_j} - \varphi_{\ell_j}) + (\tau_i \tau_j \varphi_{r_j} - \tau_j \varphi_{r_j})] N_i S_i S_j, \quad (\dagger_3)$$

$$G_{2;2} := \sum_{i \neq j} (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j \quad \text{and} \quad G_{2;3} := 2 \sum_i (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) (S_i)^2.$$

$$\begin{aligned}
G_3 &= \sum_{i \neq j} N_i S_i S_j^* (\varphi_{\ell_j} + \tau_j \varphi_{r_j}) - N_i S_i^* S_j (\tau_j^* \varphi_{\ell_j} + \varphi_{r_j}) - (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j N_j + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j^* N_j \\
&\quad + \sum_i N_i (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) - N_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) - (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) N_i + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) N_i \\
&= \sum_{i \neq j} (\tau_i \tau_j^* \varphi_{\ell_j} + \tau_i \varphi_{r_j}) N_i S_i S_j^* - (\tau_i^* \varphi_{\ell_j} + \tau_i^* \tau_j \varphi_{r_j}) N_i S_i^* S_j \\
&\quad + \sum_{i \neq j} -(\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) N_j S_i^* S_j + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) N_j S_i S_j^* \\
&\quad + \sum_{i \neq j} (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j^* + 2 \sum_i [(\varphi_{\ell_i} - \tau_i^* \varphi_{\ell_i}) + (\tau_i \varphi_{r_i} - \varphi_{r_i})] N_i \\
&= G_{3;1} + G_{3;2} + G_{3;3}, \quad \text{where}
\end{aligned}$$

$$G_{3;1} := \sum_{i \neq j} [(\tau_i \tau_j^* \varphi_{\ell_j} - \tau_j^* \varphi_{\ell_j}) + (\tau_i \varphi_{r_j} - \varphi_{r_j})] N_i S_i S_j^* + [(\varphi_{\ell_j} - \tau_i^* \varphi_{\ell_j}) + (\tau_j \varphi_{r_j} - \tau_i^* \tau_j \varphi_{r_j})] N_i S_i^* S_j, \quad (\dagger 3)$$

$$G_{3;2} := \sum_{i \neq j} (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j^*,$$

$$G_{3;3} := 2 \sum_i [(\varphi_{\ell_i} - \tau_i^* \varphi_{\ell_i}) + (\tau_i \varphi_{r_i} - \varphi_{r_i})] N_i. \quad (\dagger 3)$$

$$G_4 = -2^{-1} \sum_{i,j} [(\tau_i^* \tau_j^* \varphi_{\ell_j} + \tau_i^* \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] S_i^* S_j^* = G_{4;1} + G_{4;2}, \quad \text{with}$$

$$G_{4;1} := -2^{-1} \sum_{i \neq j} [(\tau_i^* \tau_j^* \varphi_{\ell_j} + \tau_i^* \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] S_i^* S_j^*,$$

$$G_{4;2} := -2^{-1} \sum_i [(\tau_i^* \tau_i^* \varphi_{\ell_i} + \tau_i^* \varphi_{r_i}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] (S_i^*)^2.$$

$$G_5 = -2^{-1} \sum_{i,j} [(\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i \varphi_{\ell_j} + \tau_i \tau_j \varphi_{r_j})] S_i S_j = G_{5;1} + G_{5;2}, \quad \text{with}$$

$$G_{5;1} = -2^{-1} \sum_{i \neq j} [(\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i \varphi_{\ell_j} + \tau_i \tau_j \varphi_{r_j})] S_i S_j,$$

$$G_{5;2} = -2^{-1} \sum_i [(\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i \varphi_{\ell_i} + \tau_i \tau_i \varphi_{r_i})] (S_i)^2.$$

$$G_6 = -2^{-1} \sum_{i,j} [(\tau_i \tau_j^* \varphi_{\ell_j} + \tau_i \varphi_{r_j}) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i})] S_i S_j^* + [(\tau_i^* \varphi_{\ell_j} + \tau_i^* \tau_j \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] S_i^* S_j$$

$$= G_{6;1} + G_{6;2}, \quad \text{with}$$

$$G_{6;1} := -2^{-1} \sum_{i \neq j} [(\tau_i \tau_j^* \varphi_{\ell_j} + \tau_i \varphi_{r_j}) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i})] S_i S_j^* + [(\tau_i^* \varphi_{\ell_j} + \tau_i^* \tau_j \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] S_i^* S_j,$$

$$G_{6;2} := - \sum_i (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}).$$

We add $G_{1;2}$ and $G_{4;1}$:

$$\begin{aligned}
G_{1;2} + G_{4;1} &= \sum_{i \neq j} (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_j^* - 2^{-1} \sum_{i \neq j} [(\tau_i^* \tau_j^* \varphi_{\ell_j} + \tau_i^* \varphi_{r_j}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] S_i^* S_j^* \\
&= 2^{-1} \sum_{i \neq j} [(\tau_j^* \varphi_{\ell_j} - \tau_i^* \tau_j^* \varphi_{\ell_j}) + (\varphi_{r_j} - \tau_i^* \varphi_{r_j})] S_i^* S_j^*. \quad (\dagger 3)
\end{aligned}$$

We add $G_{1,3}$ and $G_{4,2}$:

$$\begin{aligned} G_{1,3} + G_{4,2} &= 2 \sum_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) (S_i^*)^2 - 2^{-1} \sum_i [(\tau_i^* \tau_i^* \varphi_{\ell_i} + \tau_i^* \varphi_{r_i}) + (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i})] (S_i^*)^2 \\ &= G_7 + G_8, \quad \text{where} \end{aligned}$$

$$G_7 := 2^{-1} \sum_i [(\tau_i^* \varphi_{\ell_i} - \tau_i^* \tau_i^* \varphi_{\ell_i}) + (\varphi_{r_i} - \tau_i^* \varphi_{r_i})] (S_i^*)^2 \quad (\dagger 3) \quad \text{and} \quad G_8 := \sum_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) (S_i^*)^2.$$

We add $G_{2,2}$ and $G_{5,1}$:

$$\begin{aligned} G_{2,2} + G_{5,1} &= \sum_{i \neq j} (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_j - 2^{-1} \sum_{i \neq j} [(\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i \varphi_{\ell_j} + \tau_i \tau_j \varphi_{r_j})] S_i S_j \\ &= 2^{-1} \sum_{i \neq j} [(\varphi_{\ell_j} - \tau_i \varphi_{\ell_j}) + (\tau_j \varphi_{r_j} - \tau_i \tau_j \varphi_{r_j})] S_i S_j. \quad (\dagger 3) \end{aligned}$$

We add $G_{2,3}$ and $G_{5,2}$:

$$\begin{aligned} G_{2,3} + G_{5,2} &= 2 \sum_i (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) (S_i)^2 - 2^{-1} \sum_i [(\varphi_{\ell_i} + \tau_i \varphi_{r_i}) + (\tau_i \varphi_{\ell_i} + \tau_i \tau_i \varphi_{r_i})] (S_i)^2 \\ &= G_9 + G_{10}, \quad \text{where} \end{aligned}$$

$$G_9 := 2^{-1} \sum_i [(\varphi_{\ell_i} - \tau_i \varphi_{\ell_i}) + (\tau_i \varphi_{r_i} - \tau_i \tau_i \varphi_{r_i})] (S_i)^2 \quad (\dagger 3) \quad \text{and} \quad G_{10} := \sum_i (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) (S_i)^2.$$

We add $G_{3,2}$ and $G_{6,1}$:

$$G_{3,2} + G_{6,1} = -2^{-1} \sum_{i \neq j} [(\tau_i \tau_j^* \varphi_{\ell_j} - \tau_j^* \varphi_{\ell_j}) + (\tau_i \varphi_{r_j} - \varphi_{r_j}) + (\tau_j^* \varphi_{\ell_i} - \varphi_{\ell_i}) + (\tau_j^* \tau_i \varphi_{r_i} - \tau_i \varphi_{r_i})] S_i S_j^*. \quad (\dagger 3)$$

We are left to deal with $G_{6,2}$, G_8 and G_{10} :

$$\begin{aligned} G_8 + G_{10} + G_{6,2} &= \sum_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) S_i^* S_i^* + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) S_i S_i - (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) - (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) \\ &= \sum_i (\tau_i^* \varphi_{\ell_i} + \varphi_{r_i}) ((S_i^*)^2 - 1) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) ((S_i)^2 - 1) \\ &= \sum_i [(\tau_i^* \varphi_{\ell_i} - \varphi_{\ell_i}) + (\varphi_{r_i} - \tau_i \varphi_{r_i})] ((S_i^*)^2 - 1) + (\varphi_{\ell_i} + \tau_i \varphi_{r_i}) ((S_i^*)^2 + S_i^2 - 2) \\ &= G_{11} + G_{12}, \quad \text{where} \end{aligned}$$

$$G_{11} := \sum_i [(\tau_i^* \varphi_{\ell_i} - \varphi_{\ell_i}) + (\varphi_{r_i} - \tau_i \varphi_{r_i})] ((S_i^*)^2 - 1) \quad (\dagger 3) \quad \text{and} \quad G_{12} := -2 \sum_i (\cosh(\tau_i F - F) - 1) \Delta_i (4 - \Delta_i).$$

Let $W_{F;i} := \sqrt{\cosh(\tau_i F - F) - 1}$. Commuting $W_{F;i}$ with Δ_i gives

$$W_{F;i} \Delta_i = \Delta_i W_{F;i} + S_i (W_{F;i} - \tau_i^* W_{F;i}) + S_i^* (W_{F;i} - \tau_i W_{F;i}).$$

Thus

$$\begin{aligned} W_{F;i}^2 \Delta_i (4 - \Delta_i) &= W_{F;i} \Delta_i (4 - \Delta_i) W_{F;i} + R_{F;i}, \quad \text{where} \\ R_{F;i} &:= -W_{F;i} \Delta_i S_i (W_{F;i} - \tau_i^* W_{F;i}) - W_{F;i} \Delta_i S_i^* (W_{F;i} - \tau_i W_{F;i}) \\ &\quad + W_{F;i} S_i (W_{F;i} - \tau_i^* W_{F;i}) (4 - \Delta_i) + W_{F;i} S_i^* (W_{F;i} - \tau_i W_{F;i}) (4 - \Delta_i) \quad \dagger 5. \end{aligned}$$

A final accounting job gives the expression of \mathcal{G} :

$$(4.8) \quad \begin{aligned} \mathcal{G} &:= G_{r,1} + G_{\ell,1} + G_{1,1} + G_{2,1} + G_{3,1} + G_{3,3} + (G_{1,2} + G_{4,1}) \\ &\quad + G_7 + (G_{2,2} + G_{5,1}) + G_9 + (G_{3,2} + G_{6,1}) + G_{11} - 2 \sum_i R_{F;i}, \end{aligned}$$

or equivalently, $\mathcal{G} = G_r + G_\ell + 2 \sum_i W_{F;i} \Delta_i (4 - \Delta_i) W_{F;i}$. \square

* * *

We now turn to the proof of relation (3.3) that is key in Proposition 3.1. Here $d = 1$. For convenience we rewrite the relation we want to show. For $\phi \in \ell_0(\mathbb{Z})$, $\phi_F := e^F \phi$:

$$(4.9) \quad \langle \phi, [e^F A' e^F, \Delta] \phi \rangle = \langle \phi_F, [A', \Delta] \phi_F \rangle - \|\sqrt{g_r - g_\ell} A' \phi_F\|^2 \\ - \|\sqrt{\Delta(4 - \Delta)} W \phi_F\|^2 + 2^{-1} \langle \phi_F, (M + G) \phi_F \rangle, \quad \text{where}$$

$$(4.10) \quad W = W_F := \sqrt{\cosh(\tau F - F) - 1},$$

$$(4.11) \quad M = M_F := 2^{-1}(S^* + S)(g_r - g_\ell)(S^* + S) \\ + [(\tau^* \varphi_\ell - \varphi_\ell) + (\varphi_\ell - \tau \varphi_\ell) + (\tau \varphi_r - \varphi_r) + (\varphi_r - \tau^* \varphi_r)], \quad \text{and}$$

$$(4.12) \quad G = G_F := A' S(\tau^* \varphi_r - \varphi_r) + (\varphi_r - \tau^* \varphi_r) S^* A' + A' S^*(\tau \varphi_\ell - \varphi_\ell) + (\varphi_\ell - \tau \varphi_\ell) S A' \\ + [(\tau^* \varphi_\ell - \tau^{*2} \varphi_\ell) + (\varphi_r - \tau^* \varphi_r)] N S^{*2} + [(\tau^2 \varphi_r - \tau \varphi_r) + (\tau \varphi_\ell - \varphi_\ell)] N S^2 \\ + \frac{1}{2} [(\tau^* \varphi_\ell - \tau^{*2} \varphi_\ell) + (\varphi_r - \tau^* \varphi_r)] (S^*)^2 + \frac{1}{2} [(\tau \varphi_r - \tau^2 \varphi_r) + (\varphi_\ell - \tau \varphi_\ell)] S^2 \\ + 2[(\varphi_\ell - \tau^* \varphi_\ell) + (\tau \varphi_r - \varphi_r)] N + [(\tau^* \varphi_\ell - \varphi_\ell) + (\varphi_r - \tau \varphi_r)] ((S^*)^2 - 1) \\ + 2W_F \Delta S(W_F - W_{\tau^* F}) + 2W_F \Delta S^*(W_F - W_{\tau F}) \\ - 2W_F S(W_F - W_{\tau^* F})(4 - \Delta) - 2W_F S^*(W_F - W_{\tau F})(4 - \Delta).$$

Proof of (4.9). For the most part, the proof of this relation is the same as that of (4.1) when $d \geq 1$. However, the main difference is that here we do not introduce the function $g(n) := F'(\langle n \rangle) / \langle n \rangle$. We go over the proof done just above and point out the differences. As before we start with

$$[e^F A' e^F, \Delta] = e^F [A', \Delta] e^F + e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F$$

and develop the last two terms of this relation.

Part 1 : Creating $e^F A' g_r A' e^F$.

$$[e^F, \Delta] = g_r A' e^F - \frac{1}{2} g_r (S^* + S) e^F + \varphi_r \delta_{\{n=0\}} (S^* - S) e^F + \varphi_r [e^F, (S^* - S)] + (\varphi_r + \varphi_\ell) e^F S.$$

$$[e^F, \Delta] = e^F A' g_r + \frac{1}{2} e^F (S^* + S) g_r + e^F (S^* - S) \varphi_r \delta_{\{n=0\}} - [e^F, (S^* - S)] \varphi_r - S^* e^F (\varphi_r + \varphi_\ell).$$

Therefore we have obtained

$$(4.13) \quad e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F = 2e^F A' g_r A' e^F + e^F (M_r + G_r + H_r) e^F, \quad \text{where}$$

$$M_r := -2^{-1} A' g_r (S^* + S) + 2^{-1} (S^* + S) g_r A',$$

$$G_r := A' \varphi_r [e^F, (S^* - S)] e^{-F} + A' (\varphi_r + \varphi_\ell) e^F S e^{-F} \\ - e^{-F} [e^F, (S^* - S)] \varphi_r A' - e^{-F} S^* e^F (\varphi_r + \varphi_\ell) A', \quad \text{and}$$

$$H_r := A' \varphi_r \delta_{\{n=0\}} (S^* - S) + (S^* - S) \varphi_r \delta_{\{n=0\}} A'.$$

We calculate M_r :

$$M_r = -\frac{1}{2} \left(-\frac{1}{2} (S^* + S) + (S^* - S) N \right) g_r (S^* + S) + \frac{1}{2} (S^* + S) g_r \left(\frac{1}{2} (S^* + S) + N (S^* - S) \right) \\ = 2^{-1} (S^* + S) g_r (S^* + S) - 2^{-1} (S^* - S) \varphi_r \delta_{\{n \neq 0\}} (S^* + S) + 2^{-1} (S^* + S) \varphi_r \delta_{\{n \neq 0\}} (S^* - S) \\ = 2^{-1} (S^* + S) g_r (S^* + S) + (S \varphi_r S^* - S^* \varphi_r S) + (S^* \varphi_r \delta_{\{n=0\}} S - S \varphi_r \delta_{\{n=0\}} S^*) \\ = M_{r;1} + M_{r;2}, \quad \text{where}$$

$M_{r;1} := 2^{-1}(S^* + S)g_r(S^* + S) + [(\tau\varphi_r - \varphi_r) + (\varphi_r - \tau^*\varphi_r)]$ and $M_{r;2} := \varphi_r(0)(\delta_{\{n=-1\}} - \delta_{\{n=1\}})$.

Part 2 : Creating $e^F A' g_\ell A' e^F$.

$$[e^F, \Delta] = -g_\ell A' e^F + \frac{1}{2}g_\ell(S^* + S)e^F - \varphi_\ell \delta_{\{n=0\}}(S^* - S)e^F - \varphi_\ell[e^F, (S^* - S)] + (\varphi_r + \varphi_\ell)e^F S^*.$$

$$[e^F, \Delta] = -e^F A' g_\ell - \frac{1}{2}e^F(S^* + S)g_\ell - e^F(S^* - S)\varphi_\ell \delta_{\{n=0\}} - [(S^* - S), e^F]\varphi_\ell - S e^F(\varphi_r + \varphi_\ell).$$

Therefore we have obtained

$$(4.14) \quad e^F A' [e^F, \Delta] + [e^F, \Delta] A' e^F = -2e^F A' g_\ell A' e^F + e^F(M_\ell + G_\ell + H_\ell)e^F, \quad \text{where}$$

$$M_\ell := 2^{-1}A'g_\ell(S^* + S) - 2^{-1}(S^* + S)g_\ell A',$$

$$G_\ell := -A'\varphi_\ell[e^F, (S^* - S)]e^{-F} + A'(\varphi_r + \varphi_\ell)e^F S^* e^{-F} \\ + e^{-F}[e^F, (S^* - S)]\varphi_\ell A' - e^{-F}S e^F(\varphi_r + \varphi_\ell)A', \quad \text{and}$$

$$H_\ell := -A'\varphi_\ell \delta_{\{n=0\}}(S^* - S) - (S^* - S)\varphi_\ell \delta_{\{n=0\}}A'.$$

We calculate M_ℓ :

$$M_\ell = M_{\ell;1} + M_{\ell;2}, \quad \text{where}$$

$$M_{\ell;1} := -2^{-1}(S^* + S)g_\ell(S^* + S) + [(\tau^*\varphi_\ell - \varphi_\ell) + (\varphi_\ell - \tau\varphi_\ell)] \quad \text{and} \quad M_{\ell;2} := \varphi_\ell(0)(\delta_{\{n=1\}} - \delta_{\{n=-1\}}).$$

Part 3 : Adding the terms of Parts 1 and 2. Take the average of (4.13) and (4.14) to get :

$$[e^F A' e^F, \Delta] = e^F[A', \Delta]e^F + e^F A'(g_r - g_\ell)A' e^F + 2^{-1}e^F(M_r + M_\ell + G_r + G_\ell + H_r + H_\ell)e^F.$$

Applying $\phi \in \ell_0(\mathbb{Z})$ to this equation and taking inner products will yield (4.9). Let us elaborate exactly how this is achieved. First, let

$$M := M_r + M_\ell = M_{r;1} + M_{\ell;1}.$$

The latter equality holds because $(M_{r;2} + M_{\ell;2})\phi = 0$. Second, note that G_r, G_ℓ, H_r and H_ℓ are exactly the same as in the preceding proof when $i = j = 1$, which corresponds to $d = 1$. These terms are handled in the same way. In particular $(H_r + H_\ell)\phi = 0$. Finally, we investigate G . Referring to the preceding proof with $i = j = 1$, let

$$G := G_{r;1} + G_{\ell;1} + G_{1;1} + G_{2;1} + G_{3;3} + G_7 + G_9 + G_{11} - 2R_{F;1}.$$

Terms that do not contribute here are: $G_{3;1}, G_{1;2} + G_{4;1}, G_{2;2} + G_{5;1}, G_{3;2} + G_{6;1}$. We warn the careful reader that G is not simply $G_r + G_\ell$, because somewhere hidden in $G_{r;2} + G_{\ell;2}$ is the term $-2W\Delta(4 - \Delta)W$ which needs to be extracted. After taking inner products, this term ultimately produces $-\|\sqrt{\Delta(4 - \Delta)}W\phi_F\|^2$. Alternatively, $G = G_r + G_\ell + 2W\Delta(4 - \Delta)W$. \square

We also note that

$$(4.15) \quad \begin{aligned} G_r + G_\ell &= G_{r;1} + G_{\ell;1} + G_{1;1} + G_{2;1} + G_{3;3} + G_7 + G_9 + G_{11} + G_{12} \\ &= 2[(\tau^*\varphi_\ell - \tau^{*2}\varphi_\ell) + (\varphi_r - \tau^*\varphi_r)]NS^{*2} - 2[(\tau\varphi_r - \tau^2\varphi_r) + (\varphi_\ell - \tau\varphi_\ell)]NS^2 \\ &\quad + 2[(\varphi_\ell - \tau\varphi_\ell) + (\tau^*\varphi_r - \varphi_r) + (\varphi_\ell - \tau^*\varphi_\ell) + (\tau\varphi_r - \varphi_r)]N \\ &\quad + [(\tau^*\varphi_\ell - \tau^{*2}\varphi_\ell) + 2(\varphi_r - \tau^*\varphi_r)]S^{*2} + [(\tau\varphi_r - \tau^2\varphi_r) + 2(\varphi_\ell - \tau\varphi_\ell)]S^2 \\ &\quad + [(\tau\varphi_\ell - \varphi_\ell) + (\tau^*\varphi_r - \varphi_r)] + [(\tau^*\varphi_\ell - \varphi_\ell) + (\varphi_r - \tau\varphi_r)](S^{*2} - 1) \\ &\quad - 2(\cosh(\tau F - F) - 1)\Delta(4 - \Delta). \end{aligned}$$

REFERENCES

- [A] S. Agmon: *Lectures on exponential decay of solutions of second-order elliptic equations: bounds on eigenfunctions of N-body Schrödinger operators.*, Mathematical Notes, 29. Princeton University Press, Princeton, NJ, (1982).

- [ABG] W. O. Amrein, A. Boutet de Monvel, and V. Georgescu: *C₀-groups, commutator methods and spectral theory of N-body hamiltonians.*, Birkhäuser, (1996).
- [CFKS] H. Cycon, R. Froese, W. Kirsch and B. Simon: *Schrödinger operators with application to quantum mechanics and global geometry.*, Texts and Monographs in Physics. Springer-Verlag, Berlin, (1987).
- [Ca] L. Cattaneo: *Mourre's inequality and embedded bound states.*, Bull. Sci. Math. **129**, no. 7, p. 591–614, (2005).
- [CGH] L. Cattaneo, G. M. Graf, and W. Hunziker: *A general resonance theory based on Mourre's inequality.*, Ann. Henri Poincaré, **7**, p. 583–601, (2006).
- [CT] J. M. Combes, L. Thomas: *Asymptotic behaviour of eigenfunctions for multiparticle Schrödinger operators.*, Commun. Math. Phys. **34**, p. 251–270, (1973).
- [FMS] J. Faupin, J. S. Møller, and E. Skibsted: *Regularity of bound states.*, Rev. Math. Phys. **23**, no. 5, p. 453–530, (2011).
- [FH] R. Froese, I. Herbst: *Exponential bounds and absence of positive eigenvalues for N-body Schrödinger operators.*, Comm. Math. Phys. **87**, no. 3, p. 429–447, (1982/83).
- [FHHO1] R. Froese, I. Herbst, M. Hoffmann-Ostenhof, and T. Hoffmann-Ostenhof: *L²-exponential lower bounds for solutions to the Schrödinger equation.*, Commun. Math. Phys. **87**, p. 265–286, (1982).
- [FHHO2] R. Froese, I. Herbst, M. Hoffmann-Ostenhof, and T. Hoffmann-Ostenhof: *On the absence of positive eigenvalues for one-body Schrödinger operators*, J. Anal. Math. **41**, p. 272–284, (1982).
- [GGol] V. Georgescu, S. Golénia: *Isometries, Fock spaces, and spectral analysis of Schrödinger operators on trees*, J. Funct. Anal. **227**, no. 2, p. 389–429, (2005).
- [GM] S. Golénia, S. Moroianu: *Spectral analysis of magnetic Laplacians on conformally cusp manifolds.*, Ann. Henri Poincaré, **9**, no. 1, p. 131–179, (2008).
- [HS] I. Herbst, E. Skibsted: *Decay of eigenfunctions of elliptic PDE's, I.*, Adv. Math. **270**, p. 138–180, (2015).
- [JS] J. Janas, S. Simonov: *Weyl-Titchmarsh type formula for discrete Schrödinger operator with Wigner-von Neumann potential.*, Studia Math. **201**, no. 2, p. 167–189, (2010).
- [Ma] M. Mandich: *The limiting absorption principle for the discrete Wigner-von Neumann operator.*, preprint.
- [Mo] E. Mourre: *Absence of singular continuous spectrum for certain selfadjoint operators.*, Commun. Math. Phys. **78**, no. 3, p. 391–408, (1980/81).
- [S] B. Simon: *Bounded eigenfunctions and absolutely continuous spectra for one-dimensional Schrödinger operators.*, Proc. Amer. Math. Soc. **124**, no. 11, p. 3361–3369, (1996).
- [SW] E. L. Slaggie, E. H. Wichmann: *Asymptotic properties of the wave function for a bound nonrelativistic three-body system*, J. Math. Phys. **3**, p. 946–968, (1962).
- [V] A. Vasy: *Exponential decay of eigenfunctions in many-body type scattering with second-order perturbations.*, J. Funct. Anal. **209**, no. 2, p. 468–492, (2004).