



**HAL**  
open science

# Optimal functional inequalities for fractional operators on the sphere and applications

Jean Dolbeault, An Zhang

► **To cite this version:**

Jean Dolbeault, An Zhang. Optimal functional inequalities for fractional operators on the sphere and applications. 2016. hal-01358059v1

**HAL Id: hal-01358059**

**<https://hal.science/hal-01358059v1>**

Preprint submitted on 30 Aug 2016 (v1), last revised 6 Oct 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# OPTIMAL FUNCTIONAL INEQUALITIES FOR FRACTIONAL OPERATORS ON THE SPHERE AND APPLICATIONS

JEAN DOLBEAULT AND AN ZHANG

ABSTRACT. This paper is devoted to the family of optimal functional inequalities on the  $n$ -dimensional sphere  $\mathbb{S}^n$

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq C_{q,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu \quad \forall F \in H^{s/2}(\mathbb{S}^n)$$

where  $\mathcal{L}_s$  denotes a fractional Laplace operator of order  $s \in (0, n)$ ,  $q \in [1, 2) \cup (2, q_*]$ ,  $q_* = 2n/(n-s)$  is a critical exponent and  $d\mu$  is the uniform probability measure on  $\mathbb{S}^n$ . These inequalities are established with optimal constants using spectral properties of fractional operators. Their consequences for fractional heat flows are considered. If  $q > 2$ , these inequalities interpolate between fractional Sobolev and subcritical fractional logarithmic Sobolev inequalities, which correspond to the limit case as  $q \rightarrow 2$ . For  $q < 2$ , the inequalities interpolate between fractional logarithmic Sobolev and fractional Poincaré inequalities. In the subcritical range  $q < q_*$ , the method also provides us with remainder terms which can be considered as an improved version of the optimal inequalities. Finally, weighted inequalities of Caffarelli-Kohn-Nirenberg type involving the fractional Laplacian are obtained in the Euclidean space, using a stereographic projection and scaling properties. The case  $s \in (-n, 0)$  is also considered.

## 1. INTRODUCTION AND MAIN RESULTS

Let us consider the unit sphere  $\mathbb{S}^n$  with  $n \geq 1$  and assume that the measure  $d\mu$  is the uniform probability measure, which is also the measure induced on  $\mathbb{S}^n$  by Lebesgue's measure on  $\mathbb{R}^{n+1}$ , up to a normalization constant. With  $\lambda \in (0, n)$ ,  $p = \frac{2n}{2n-\lambda} \in (1, 2)$  or equivalently  $\lambda = \frac{2n}{p'}$  where  $\frac{1}{p} + \frac{1}{p'} = 1$ , according to [38], the sharp *Hardy-Littlewood-Sobolev inequality on  $\mathbb{S}^n$*  reads

$$(1) \quad \iint_{\mathbb{S}^n \times \mathbb{S}^n} F(\zeta) |\zeta - \eta|^{-\lambda} F(\eta) d\mu(\zeta) d\mu(\eta) \leq \frac{\Gamma(n) \Gamma(\frac{n-\lambda}{2})}{2^\lambda \Gamma(\frac{n}{2}) \Gamma(\frac{n}{p})} \|F\|_{L^p(\mathbb{S}^n)}^2.$$

For the convenience of the reader, the definitions of all parameters, their ranges and their relations have been collected in Appendix C.

---

*Date:* August 30, 2016.

*2010 Mathematics Subject Classification.* 26D15; 35A23; 35R11; Secondary: 26D10; 26A33; 35B33.

*Key words and phrases.* Hardy-Littlewood-Sobolev inequality; fractional Sobolev inequality; fractional logarithmic Sobolev inequality; spectral gap; fractional Poincaré inequality; fractional heat flow; subcritical interpolation inequalities on the sphere; stereographic projection; Euclidean fractional Caffarelli-Kohn-Nirenberg inequalities.

By the Funk-Hecke formula, the left side of the inequality can be written as

$$(2) \quad \iint_{\mathbb{S}^n \times \mathbb{S}^n} F(\zeta) |\zeta - \eta|^{-\lambda} F(\eta) d\mu(\zeta) d\mu(\eta) \\ = \frac{\Gamma(n) \Gamma(\frac{n-\lambda}{2})}{2^\lambda \Gamma(\frac{n}{2}) \Gamma(\frac{n}{p})} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{n}{p}) \Gamma(\frac{n}{p'} + k)}{\Gamma(\frac{n}{p'}) \Gamma(\frac{n}{p} + k)} \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

where  $F = \sum_{k=0}^{\infty} F_{(k)}$  is a decomposition on spherical harmonics, so that  $F_{(k)}$  is a spherical harmonic function of degree  $k$ . See [33, Section 4] for details on the computations and, *e.g.*, [42] for further related results. With the above representation, Inequality (1) is equivalent to

$$(3) \quad \sum_{k=0}^{\infty} \frac{\Gamma(\frac{n}{p}) \Gamma(\frac{n}{p'} + k)}{\Gamma(\frac{n}{p'}) \Gamma(\frac{n}{p} + k)} \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu \leq \|F\|_{L^p(\mathbb{S}^n)}^2.$$

By duality, with  $q_\star = q_\star(s)$  defined by

$$(4) \quad q_\star = \frac{2n}{n-s}$$

or equivalently  $s = n(1 - 2/q_\star)$ , we obtain the *fractional Sobolev inequality on  $\mathbb{S}^n$*

$$(5) \quad \|F\|_{L^{q_\star}(\mathbb{S}^n)}^2 \leq \int_{\mathbb{S}^n} F \mathcal{K}_s F d\mu \quad \forall F \in \mathbf{H}^{s/2}(\mathbb{S}^n)$$

for any  $s \in (0, n)$ , where

$$(6) \quad \int_{\mathbb{S}^n} F \mathcal{K}_s F d\mu := \sum_{k=0}^{\infty} \gamma_k\left(\frac{n}{q_\star}\right) \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

and

$$\gamma_k(x) := \frac{\Gamma(x) \Gamma(n-x+k)}{\Gamma(n-x) \Gamma(x+k)}.$$

With  $s \in (0, n)$ , the exponent  $q_\star$  is in the range  $(2, \infty)$ . Inequalities (1) and (5) are related by  $q_\star = p'$  so that

$$p = \frac{2n}{n+s} \quad \text{and} \quad \lambda = n-s.$$

We shall refer to  $q = q_\star(s)$  given by (4) as the *critical case* and our purpose is to study the whole range of the *subcritical interpolation inequalities*

$$(7) \quad \frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq C_{q,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu \quad \forall F \in \mathbf{H}^{s/2}(\mathbb{S}^n)$$

for any  $q \in [1, 2) \cup (2, q_\star]$ , where

$$\mathcal{L}_s := \frac{1}{\kappa_{n,s}} (\mathcal{K}_s - \text{Id}) \quad \text{with} \quad \kappa_{n,s} := \frac{\Gamma(\frac{n}{q_\star})}{\Gamma(n - \frac{n}{q_\star})} = \frac{\Gamma(\frac{n-s}{2})}{\Gamma(\frac{n+s}{2})}.$$

If  $q = q_\star$ , (5) and (7) are identical, the optimal constant in (7) is  $C_{q_\star,s} = \frac{\kappa_{n,s}}{q_\star - 2}$ , and we recall that (5) is equivalent to the fractional Sobolev inequality on the

Euclidean space (see the proof of Theorem 6 in Section 3 for details). The usual conformal fractional Laplacian is defined by

$$\mathcal{A}_s := \frac{1}{\kappa_{n,s}} \mathcal{K}_s = \mathcal{L}_s + \frac{1}{\kappa_{n,s}} \text{Id}.$$

For brevity, we shall say that  $\mathcal{L}_s$  is the *fractional Laplacian* of order  $s$ , or simply the *fractional Laplacian*.

We observe that  $\gamma_0(n/q) - 1 = 0$  and  $\gamma_1(n/q) - 1 = q - 2$ . A straightforward computation gives

$$\int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu := \sum_{k=1}^{\infty} \delta_k\left(\frac{n}{q_*}\right) \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

where the spectrum of  $\mathcal{L}_s$  is given by

$$\delta_k(x) := \frac{\Gamma(n-x+k)}{\Gamma(x+k)} - \frac{\Gamma(n-x)}{\Gamma(x)}.$$

The case corresponding to  $s = 2$  and  $n \geq 3$ , where  $1/\kappa_{n,2} = \frac{1}{4}n(n-2)$ ,  $\mathcal{L}_2 = -\Delta$ ,  $\mathcal{A}_2 = -\Delta + \frac{1}{4}n(n-2)$  and  $\Delta$  stands for the *Laplace-Beltrami operator* on  $\mathbb{S}^n$ , has been considered by W. Beckner: in [5, page 233, (35)] he observed that

$$\delta_k\left(\frac{n}{q}\right) \leq \delta_k\left(\frac{n}{q_*}\right) = k(k+n-1)$$

if  $q \in (2, q_*(2)]$ , where  $q_* = q_*(2) = 2n/(n-2)$  and  $(k(k+n-1))_{k \in \mathbb{N}}$  is the sequence of the eigenvalues of  $-\Delta$  according to, e.g., [7]. This establishes the interpolation inequality

$$(8) \quad \|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2 \leq \frac{q-2}{n} \|\nabla F\|_{L^2(\mathbb{S}^n)}^2 \quad \forall F \in H^1(\mathbb{S}^n)$$

where  $C_{q,2} = 1/n$  is the optimal constant: see [5, (35), Theorem 4] for details. An earlier proof of the inequality with optimal constant can be found in [8, Corollary 6.2], with a proof based on *rigidity* results for elliptic partial differential equations. Our main result generalizes the interpolation inequalities (8) to the case of the fractional operators  $\mathcal{L}_s$ , and relies on W. Beckner's approach. In particular, as in [5], we characterize the optimal constant  $C_{q,s}$  in (7) using a spectral gap property.

After dividing both sides of (8) by  $(q-2)$  we obtain an inequality which, for  $s = 2$ , also makes sense for any  $q \in [1, 2)$ . When  $q = 1$ , this is actually a variant of the Poincaré inequality (or, to be precise, the Poincaré inequality written for  $|F|$ ), and the range  $q > 1$  has been studied using the *carré du champ* method, also known as the  $\Gamma_2$  calculus, by D. Bakry and M. Emery in [3]. Actually their method covers the range corresponding to  $1 \leq q < \infty$  if  $n = 1$  and  $1 \leq q \leq 2^\# := (2n^2+1)/(n-1)^2$  if  $n \geq 2$ . In the special case  $q = 2$ , the l.h.s. has to be replaced by the entropy  $\int_{\mathbb{S}^n} F^2 \log(F^2/\|F\|_{L^2(\mathbb{S}^n)}^2) d\mu$ . Still under the condition that  $s = 2$ , the whole range  $1 \leq q < \infty$  when  $n = 2$ , and  $1 \leq q \leq 2n/(n-2)$  if  $n \geq 3$  can be covered using nonlinear flows as shown in [21, 24, 25].

All these considerations motivate our first result, which generalizes known results for  $\mathcal{L}_2 = -\Delta$  to the case of the *fractional Laplacian*  $\mathcal{L}_s$ .

**Theorem 1.** *Let  $n \geq 1$ ,  $s \in (0, n]$ ,  $q \in [1, 2) \cup (2, q_\star]$ , with  $q_\star$  given by (4), if  $s < n$ , and  $q \in [1, 2) \cup (2, \infty)$  if  $s = n$ . Inequality (7) holds with sharp constant*

$$C_{q,s} = \frac{n-s}{2s} \frac{\Gamma\left(\frac{n-s}{2}\right)}{\Gamma\left(\frac{n+s}{2}\right)}.$$

With our previous notations, this amounts to  $C_{q,s} = \frac{\kappa_{n,s}}{q_\star - 2} = \frac{n-s}{2s} \kappa_{n,s}$ . Remarkably,  $C_{q,s}$  is independent of  $q$ . Equality in (7) is achieved by constant functions. The issue of the optimality of  $C_{q,s}$  is henceforth somewhat subtle. If we define the functional

$$(9) \quad \mathcal{Q}[F] := \frac{(q-2) \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu}{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}$$

on the subset  $\mathcal{H}^{s/2}$  of the functions in  $H^{s/2}(\mathbb{S}^n)$  which are not almost everywhere constant, then  $C_{q,s}$  can be characterized by

$$C_{q,s}^{-1} = \inf_{F \in \mathcal{H}^{s/2}} \mathcal{Q}[F].$$

This minimization problem will be discussed in Section 4.

Our key estimate is a simple convexity observation that is stated in Lemma 9. The optimality in (7) is obtained by performing a linearization, which corresponds to an asymptotic regime as we shall see in Section 2.1. Technically, this is the reason why we are able to identify the optimal constant. The asymptotic regime can be investigated using a flow. Indeed, a first consequence of Theorem 1 is that we may apply entropy methods to the generalized fractional heat flow

$$(10) \quad \frac{\partial u}{\partial t} - q \nabla \cdot \left( u^{1-\frac{1}{q}} \nabla (-\Delta)^{-1} \mathcal{L}_s u^{\frac{1}{q}} \right) = 0.$$

Notice that (10) is a 1-homogeneous equation, but that it is nonlinear when  $q \neq 1$  and  $s \neq 2$ . Let us define a *generalized entropy* by

$$\mathcal{E}_q[u] := \frac{1}{q-2} \left[ \left( \int_{\mathbb{S}^n} u d\mu \right)^{\frac{2}{q}} - \int_{\mathbb{S}^n} u^{\frac{2}{q}} d\mu \right].$$

It is straightforward to check that for any positive solution to (10) which is smooth enough and has sufficient decay properties as  $|x| \rightarrow +\infty$ , we have

$$\frac{d}{dt} \mathcal{E}_q[u(t, \cdot)] = -2 \int_{\mathbb{S}^n} \nabla u^{\frac{1}{q}} \cdot \nabla (-\Delta)^{-1} \mathcal{L}_s u^{\frac{1}{q}} d\mu = -2 \int_{\mathbb{S}^n} u^{\frac{1}{q}} \mathcal{L}_s u^{\frac{1}{q}} d\mu,$$

so that by applying (7) to  $F = u^{1/q}$  we obtain the exponential decay of  $\mathcal{E}_q[u(t, \cdot)]$ .

**Corollary 2.** *Let  $n \geq 1$ ,  $s \in (0, n]$ ,  $q \in [1, 2) \cup (2, q_\star]$  if  $s < n$ , with  $q_\star$  given by (4), and  $q \in [1, 2) \cup (2, \infty)$  if  $s = n$ . If  $u$  is a positive function in  $C^1(\mathbb{R}^+; L^\infty(\mathbb{S}^n))$  such that  $u^{1/q} \in C^1(\mathbb{R}^+; H^{s/2}(\mathbb{S}^n))$  and if  $u$  solves (10) on  $\mathbb{S}^n$  with initial datum  $u_0 > 0$ , then*

$$\mathcal{E}_q[u(t, \cdot)] \leq \mathcal{E}_q[u_0] e^{-2C_{q,s}^{-1}t} \quad \forall t \geq 0.$$

The exponential rate is determined by the asymptotic regime as  $t \rightarrow +\infty$ . The value of the *optimal constant*  $\mathcal{C}_{q,s}$  is indeed determined by the *spectral gap of the linearized problem* around non-zero constant functions. From the expression of (10), which is not even a linear equation whenever  $s \neq 2$ , we observe that the interplay of optimal fractional inequalities and fractional diffusion flows is not straightforward, while for  $s = 2$ , the generalized entropy  $\mathcal{E}_q$  enters in the framework of the so-called  $\varphi$ -entropies and is well understood in terms of gradient flows: see for instance [2, 13, 28]. When  $s = 2$ , it is also known from [3] that heat flows can be used in the framework of the *carré du champ* method to establish the inequalities at least for exponents in the range  $q \leq 2^\#$  if  $n \geq 2$ , and that the whole subcritical range of exponents can be covered using nonlinear diffusions as in [21, 24, 25] (and also the critical exponent if  $n \geq 3$ ). Even better, *rigidity* results, that is, uniqueness of positive solutions (which are therefore constant functions) follows by this technique. So far there is no analogue in the case of fractional operators, except for one example found in [12] when  $n = 1$ .

When  $s = 2$ , the *carré du champ* method provides us with an integral remainder term and, as a consequence, with an improved version of (7). As we shall see, our proof of Theorem 1 establishes another improved inequality, by construction: see Corollary 10. This also suggests another direction, which is more connected with the duality that relates (1) and (5). Let us describe the main idea. The operator  $\mathcal{K}_s$  is positive definite and we can henceforth consider  $\mathcal{K}_s^{1/2}$  and  $\mathcal{K}_s^{-1}$ . Moreover, using (2) and (6), we know that

$$\iint_{\mathbb{S}^n \times \mathbb{S}^n} G(\zeta) |\zeta - \eta|^{-\lambda} G(\eta) d\mu(\zeta) d\mu(\eta) = \frac{\Gamma(n) \Gamma(\frac{s}{2})}{2^\lambda \Gamma(\frac{n}{2}) \Gamma(n + \frac{s}{2})} \int_{\mathbb{S}^n} G \mathcal{K}_s^{-1} G d\mu.$$

By expanding the square  $\int_{\mathbb{S}^n} |\mathcal{K}_s^{1/2} F - \mathcal{K}_s^{-1/2} G|^2 d\mu$  with  $G = F^{q_\star - 1}$  so that  $F G = F^{q_\star} = G^p$  where  $q_\star$  and  $p$  are Hölder conjugates, we get a comparison of the difference of the two terms which show up in (1) and (5) and, as a result, an *improved fractional Sobolev inequality on  $\mathbb{S}^n$* . The reader interested in the details of the proof is invited to refer to [27] for a similar result.

**Proposition 3.** *Let  $n \geq 1$ ,  $s \in (0, n)$ , and consider  $q_\star$  given by (4),  $p = q'_\star = \frac{2n}{n+s}$  and  $\lambda = n - s$ . For any  $F \in H^{s/2}(\mathbb{S}^n)$ , if  $G = F^{q_\star - 1}$ , then*

$$\begin{aligned} \|G\|_{L^p(\mathbb{S}^n)}^2 - 2^\lambda \frac{\Gamma(\frac{n}{2}) \Gamma(n + \frac{s}{2})}{\Gamma(n) \Gamma(\frac{s}{2})} \iint_{\mathbb{S}^n \times \mathbb{S}^n} G(\zeta) |\zeta - \eta|^{-\lambda} G(\eta) d\mu(\zeta) d\mu(\eta) \\ \leq \|F\|_{L^{q_\star}(\mathbb{S}^n)}^{2(q_\star - 2)} \left( \int_{\mathbb{S}^n} F \mathcal{K}_s F d\mu - \|F\|_{L^{q_\star}(\mathbb{S}^n)}^2 \right). \end{aligned}$$

Still in the critical case  $q = q_\star$ , using the *fractional Yamabe flow* and taking inspiration from [23, 27, 37, 36, 40], it is possible to give improvements of the above inequality and in particular improve on the constant which relates the left and the right sides of the inequality in Proposition 3. We will not go further in this direction because of the delicate regularity properties of the fractional Yamabe flow and because, so far the method does not allow to characterize the best constant in the improvement. Let us mention that, in the critical case  $q = q_\star$ , further estimates

of Bianchi-Egnell type have also been obtained in [15, 40] for fractional operators. In this paper, we shall rather focus on the subcritical range. It is however clear that there is still space for further improvements, or alternative proofs of (5) which rely neither on rearrangements as in [38] nor on inversion symmetry as in [31, 32, 33], for the simple reason that our method fails to provide us with a proof of the Bianchi-Egnell estimates in the critical case.

For completeness let us quote a few other related results. Symmetrization techniques and the method of competing symmetries are both very useful to identify the optimal functions: the interested reader is invited to refer respectively to [39] and [11], when  $s = 2$ . In this paper, we shall use notations inspired by [5], but at this point it worth mentioning that in [5] the emphasis is put on logarithmic Hardy-Littlewood-Sobolev inequalities and their dual counterparts, which are  $n$ -dimensional versions of the Moser-Trudinger-Onofri inequalities. Some of these results were obtained simultaneously in [10] with some additional insight on optimal functions gained from rearrangements and from the method of competing symmetries. Concerning observations on duality, we refer to the introduction of [10], which clearly refers the earlier contributions of various authors in this area. For more recent considerations on  $n$ -dimensional Moser-Trudinger-Onofri inequalities, see, *e.g.*, [19].

Section 2 is devoted to the proof of Theorem 1. As already said, we shall take advantage of the subcritical range to obtain remainder terms and improved inequalities. Improvements in the subcritical range have been obtained in the case of non-fractional interpolation inequalities in the context of fast diffusion equations in [29, 30]. In this paper we shall simply take into account the terms which appear by difference in the proof of Theorem 1: see Corollary 10 in Section 2.3. Although this approach does not provide us with an alternative proof of the optimality of the constant  $C_{q,s}$  in (7), variational methods will be applied in Section 4 in order to explain *a posteriori* why the value of the optimal value of  $C_{q,s}$  is determined by the spectral gap of a linearized problem. Some useful information on the spectrum of  $\mathcal{L}_s$  is detailed in Appendix A.

Our next result is devoted to the singular case of Inequality (7) corresponding to the limit as  $q = 2$ : we establish a family of *sharp fractional logarithmic Sobolev inequalities*, in the subcritical range.

**Corollary 4.** *Let  $s \in (0, n]$ . Then we have the sharp logarithmic Sobolev inequality*

$$(11) \quad \int_{\mathbb{S}^n} |F|^2 \log \left( \frac{|F|}{\|F\|_2} \right) d\mu \leq C_{2,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu \quad \forall F \in H^{s/2}(\mathbb{S}^n).$$

*Equality is achieved only by constant functions and  $C_{2,s} = \frac{n-s}{2s} \kappa_{n,s}$  is optimal.*

This result completes the picture of Theorem 1 and shows that, under appropriate precautions, the case  $q = 2$  can be put in a common picture with the cases corresponding to  $q \neq 2$ . By taking the limit as  $s \rightarrow 0_+$ , we recover Beckner's fractional logarithmic Sobolev inequality as stated in [4, 6]. In that case,  $q = 2$  is critical, from the point of view of the fractional operator. The proof of Corollary 4 and further considerations on the  $s = 0$  limit will be given in Section 2.4.

The definition (6) of  $\mathcal{K}_s$  also applies to the range  $s \in (-n, 0)$  and the reader is invited to check that

$$\mathcal{K}_s^{-1} = \mathcal{K}_{-s} \quad \forall s \in (0, n)$$

is defined by the sequence of eigenvalues  $\gamma_k(n/p)$  where  $p = 2n/(n+s)$  is the Hölder conjugate of  $q_*(s)$  given by (4). It is then straightforward to check that the sharp *Hardy-Littlewood-Sobolev inequality on  $\mathbb{S}^n$*  (3) can be written as

$$(12) \quad \frac{\|F\|_{L^p(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{p-2} \leq \frac{\kappa_{n,-s}}{2-p} \int_{\mathbb{S}^n} F \mathcal{L}_{-s} F d\mu \quad \forall F \in L^2(\mathbb{S}^n)$$

where  $p = \frac{2n}{n+s} \in (1, 2)$ ,  $\mathcal{L}_{-s} := \frac{1}{\kappa_{n,-s}}(\text{Id} - \mathcal{K}_{-s})$  and  $\kappa_{n,-s} = \Gamma(\frac{n+s}{2})/\Gamma(\frac{n-s}{2})$ . Notice that  $\kappa_{n,-s} = 1/\kappa_{n,s}$ . A first consequence is that we can rewrite the result of Proposition 3 as

$$\|G\|_{L^p(\mathbb{S}^n)}^2 - \int_{\mathbb{S}^n} G \mathcal{K}_{-s} G d\mu \leq \|F\|_{L^{q_*}(\mathbb{S}^n)}^{2(q_*-2)} \left( \int_{\mathbb{S}^n} F \mathcal{K}_s F d\mu - \|F\|_{L^{q_*}(\mathbb{S}^n)}^2 \right).$$

for any  $F \in H^{s/2}(\mathbb{S}^n)$  and  $G = F^{q_*-1}$ , where  $n \geq 1$ ,  $s \in (0, n)$ ,  $q_*$  is given by (4) and  $p = q'_*$ . A second consequence of the above observations is the extension of Theorem 1 to the range  $(-n, 0)$ .

**Theorem 5.** *Let  $n \geq 1$ ,  $s \in (-n, 0)$ ,  $q \in [1, 2n/(n-s))$ . Inequality (7) holds with  $\mathcal{L}_s := \kappa_{n,-s}(\text{Id} - \mathcal{K}_s)$  and sharp constant  $C_{q,s} = \frac{n-s}{2|s|} \Gamma(\frac{n-s}{2})/\Gamma(\frac{n+s}{2})$ .*

The results of Theorems 1 and 5 are summarized in Fig. 2.

To conclude with the outline of this paper, Section 3 is devoted to the stereographic projection and consequences for functional inequalities on the Euclidean space. By stereographic projection, (5) becomes

$$\|f\|_{L^{q_*}(\mathbb{R}^n)}^2 \leq S_{n,s} \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 \quad \forall f \in \dot{H}^{s/2}(\mathbb{R}^n),$$

where  $\|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 := \int_{\mathbb{R}^n} f(-\Delta)^{s/2} f dx$  and the optimal constant is such that

$$S_{n,s} = \kappa_{n,s} |\mathbb{S}^n|^{\frac{2}{q_*}-1}.$$

The fact that (5) is equivalent to the fractional Sobolev inequality on the Euclidean space is specific to the critical exponent  $q = q_*(s)$ . In the subcritical range, weights appear. However, using scaling properties, it is possible to get rid of these weights except for some power law terms. Altogether, we obtain some *fractional Caffarelli-Kohn-Nirenberg inequalities*, with an explicit estimate of the constant. Let us introduce the two weighted norms

$$\|f\|_{L^{q,\beta}(\mathbb{R}^n)}^q := \int_{\mathbb{R}^n} |f|^q |x|^{-\beta} dx \quad \text{and} \quad \|f\|_{L_*^{q,\beta}(\mathbb{R}^n)}^q := \int_{\mathbb{R}^n} |f|^q (1+|x|^2)^{-\frac{\beta}{2}} dx.$$

The next result is inspired by a non-fractional computation done in [26] and relies on the stereographic projection.

**Theorem 6.** *Let  $n \geq 1$ ,  $s \in (0, n)$ ,  $q \in (2, q_*)$  with  $q_*$  given by (4),  $\beta = 2n(1 - \frac{q}{q_*})$ . Then we have the weighted inequality*

$$(13) \quad \|f\|_{L_*^{q,\beta}(\mathbb{R}^n)}^2 \leq a \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 + b \|f\|_{L_*^{2,2s}(\mathbb{R}^n)}^2 \quad \forall f \in C_0^\infty(\mathbb{R}^n)$$



where  $\mathbf{a} = \frac{q-2}{q_\star-2} \kappa_{n,s} 2^{n(\frac{2}{q_\star}-\frac{2}{q})} |\mathbb{S}^n|^{\frac{2}{q}-1}$  and  $\mathbf{b} = \frac{q_\star-q}{q_\star-2} 2^{n(1-\frac{2}{q})} |\mathbb{S}^n|^{\frac{2}{q}-1}$ . Moreover, if  $q < q_\star$ , equality holds in (13) if and only if  $f$  is proportional to  $f_{s,\star}(x) := (1+|x|^2)^{-\frac{n-s}{2}}$ .

This result is one of the few examples of optimal functional inequalities involving fractional operators on  $\mathbb{R}^n$ . It touches the area of fractional Hardy-Sobolev inequalities and weighted fractional Sobolev inequalities, for which we respectively refer to [34, 14] and [16], and references therein. The wider family of Caffarelli-Kohn-Nirenberg type inequalities raises additional difficulties, for instance related with symmetry and symmetry breaking issues, which are so far essentially untouched in the framework of fractional operators, up to few exceptions like [14]. One has to notice that the specific expression of the weight in  $\|f\|_{L^{q,\beta}(\mathbb{R}^n)}$  introduces a scale, which can be removed using scalings, as shown by the last result of this paper.

**Corollary 7.** *Under the assumptions of Theorem 6, if  $s < n/2$ ,  $q \in (2, q_\star)$  and*

$$\vartheta = \frac{1/2 - 1/q}{1/2 - 1/q_\star} = \frac{n(q-2)}{sq},$$

then for any  $f \in C_0^\infty(\mathbb{R}^n)$  we have

$$(14) \quad \|f\|_{L^{q,\beta}(\mathbb{R}^n)}^2 \leq K_{s,q} \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^{2\vartheta} \|f\|_{L^{2,2s}(\mathbb{R}^n)}^{2(1-\vartheta)}$$

with  $K_{s,q} := \vartheta^{-\vartheta} (1-\vartheta)^{\vartheta-1} |\mathbb{S}^n|^{\frac{2}{q}-1} \kappa_{n,s}^\vartheta \left(\frac{q-2}{q_\star-2}\right)^\vartheta \left(\frac{q_\star-q}{q_\star-2}\right)^{1-\vartheta}$ .

In the limit case as  $q \rightarrow 2$ , the above inequality has to be replaced by the fractional, logarithmic Hardy inequality: for any  $f \in C_0^\infty(\mathbb{R}^n)$  such that  $\|f\|_{L^{2,2s}(\mathbb{R}^n)} = 1$ ,

$$(15) \quad \int_{\mathbb{R}^n} \frac{f^2}{|x|^{2s}} \log(|x|^{n-s} f^2) dx \leq \frac{n}{s} \log \left[ C_s^{\text{FLH}} \int_{\mathbb{R}^n} f (-\Delta)^{s/2} f dx \right]$$

with  $C_s^{\text{FLH}} = \frac{n-s}{n} \left(\frac{e}{|\mathbb{S}^n|}\right)^{s/n} \Gamma\left(\frac{n-s}{2}\right) / \Gamma\left(\frac{n+s}{2}\right)$ .

Inequalities (13) and (14)-(15) hold not only for the space  $C_0^\infty(\mathbb{R}^n)$  of all smooth functions with compact support but also for the much larger spaces of functions obtained by completion of  $C_0^\infty(\mathbb{R}^n)$  with respect to the norms defined respectively by  $\|f\|^2 := \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 + \|f\|_{L^{2,2s}(\mathbb{R}^n)}^2$  and  $\|f\|^2 := \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 + \|f\|_{L^{2,2s}(\mathbb{R}^n)}^2$ .

## 2. SUBCRITICAL INTERPOLATION INEQUALITIES

In this section, our purpose is to prove Theorem 1.

**2.1. A Poincaré inequality.** We start by recalling some basic facts:

- (i) If  $q$  and  $q'$  are Hölder conjugates, then  $n/q' = n - x$  with  $x = n/q$ ,
- (i)  $\gamma_0(x) = 1$  for any  $x > 0$ ,
- (ii)  $\gamma_k(n/2) = 1$  and  $\delta_k(n/2) = 0$  for any  $k \in \mathbb{N}$ ,
- (iii)  $\gamma_1(x) = (n-x)/x$ ,  $\gamma_1(n/q) = q-1$  and  $\delta_1(n/q_\star) = (q_\star-2)/\kappa_{n,s}$ . As a consequence, we know that the first positive eigenvalues of  $\mathcal{K}_s$  and  $\mathcal{L}_s$  are

$$\lambda_1(\mathcal{K}_s) = \gamma_1\left(\frac{n}{q_\star}\right) = q_\star - 1 \quad \text{and} \quad \lambda_1(\mathcal{L}_s) = \delta_1\left(\frac{n}{q_\star}\right) = \frac{q_\star - 2}{\kappa_{n,s}} = \frac{2s}{(n-s)\kappa_{n,s}}.$$

A straightforward consequence is the following sharp Poincaré inequality.

**Lemma 8.** *For any  $F \in \mathbf{H}^{s/2}(\mathbb{S}^n)$ , we have*

$$\|F - F_{(0)}\|_{\mathbf{L}^2(\mathbb{S}^n)}^2 \leq C_{1,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F \, d\mu \quad \text{where} \quad F_{(0)} = \int_{\mathbb{S}^n} F \, d\mu,$$

and  $C_{1,s} = \kappa_{n,s}/(q_\star - 2)$  is the optimal constant. Any function  $F = F_{(0)} + F_{(1)}$ , with  $F_{(1)}$  such that  $\mathcal{L}_s F_{(1)} = \lambda_1(\mathcal{L}_s) F_{(1)}$ , realizes the equality case.

*Proof.* The proof is elementary. With the usual notations, we may write

$$\begin{aligned} \int_{\mathbb{S}^n} F \mathcal{L}_s F \, d\mu &= \int_{\mathbb{S}^n} (F - F_{(0)}) \mathcal{L}_s (F - F_{(0)}) \, d\mu = \sum_{k=1}^{\infty} \delta_k \left(\frac{n}{q_\star}\right) \int_{\mathbb{S}^n} |F_{(k)}|^2 \, d\mu \\ &\geq \delta_1 \left(\frac{n}{q_\star}\right) \|F - F_{(0)}\|_{\mathbf{L}^2(\mathbb{S}^n)}^2 = \lambda_1(\mathcal{L}_s) \|F - F_{(0)}\|_{\mathbf{L}^2(\mathbb{S}^n)}^2 \end{aligned}$$

because  $\delta_k(n/q_\star)$  is increasing with respect to  $k \in \mathbb{N}$ .  $\square$

The sharp Poincaré constant  $C_{1,s}$  is a lower bound for  $C_{q,s}$ , for any  $q \in (1, q_\star]$  if  $s < n$ , or any  $q > 1$  if  $s = n$ . Indeed, if  $q \neq 2$ , by testing Inequality (7) with  $F = 1 + \varepsilon G_1$ , where  $G_1$  is an eigenfunction of  $\mathcal{L}_s$  associated with the eigenvalue  $\lambda_1(\mathcal{L}_s)$ , it is easy to see that

$$\begin{aligned} \varepsilon^2 \|G_1\|_{\mathbf{L}^2(\mathbb{S}^n)}^2 &\sim \frac{\|F\|_{\mathbf{L}^q(\mathbb{S}^n)}^2 - \|F\|_{\mathbf{L}^2(\mathbb{S}^n)}^2}{q - 2} \leq C_{q,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F \, d\mu \\ &= C_{q,s} \varepsilon^2 \int_{\mathbb{S}^n} G_1 \mathcal{L}_s G_1 \, d\mu \end{aligned}$$

as  $\varepsilon \rightarrow 0$ , which means that, at leading order in  $\varepsilon$ ,

$$\|G_1\|_{\mathbf{L}^2(\mathbb{S}^n)}^2 = \lambda_1(\mathcal{L}_s) C_{q,s} \|G_1\|_{\mathbf{L}^2(\mathbb{S}^n)}^2.$$

Altogether, this proves that

$$(16) \quad C_{q,s} \geq \frac{1}{\lambda_1(\mathcal{L}_s)} = \frac{\kappa_{n,s}}{q_\star - 2}.$$

A similar computation, with (7) replaced by (11) and  $F = 1 + \varepsilon G_1$ , shows that

$$\int_{\mathbb{S}^n} |F|^2 \log \left( \frac{|F|}{\|F\|_2} \right) \, d\mu \sim C_{2,s} \varepsilon^2 \int_{\mathbb{S}^n} G_1 \mathcal{L}_s G_1 \, d\mu$$

as  $\varepsilon \rightarrow 0$ , so that (16) also holds if  $q = 2$ . Hence, under the Assumptions of Theorem 1, (16) holds for any  $q \geq 1$ . In order to establish Theorem 1 and Corollary 4, we have now to prove that (16) is actually an equality.

**2.2. Some spectral estimates.** Let us start with some observations on the function  $\gamma_k$  in (6). By expanding its expression, we get that

$$\gamma_k(x) = \frac{(n+k-1-x)(n+k-2-x)\dots(n-x)}{(k-1+x)(k-2+x)\dots x}$$

for any  $k \geq 1$ . After taking the logarithmic derivative, we find that

$$(17) \quad \alpha_k(x) := -\frac{\gamma'_k(x)}{\gamma_k(x)} = \sum_{j=0}^{k-1} \beta_j(x) \quad \text{with} \quad \beta_j(x) := \frac{1}{n+j-x} + \frac{1}{j+x}$$

and observe that  $\alpha_k$  is positive. As a consequence,  $\gamma'_k < 0$  on  $[0, n]$  and, from the expression of  $\gamma_k$ , we read that  $\gamma_k(n) = 0$ . Since  $\gamma_k(n/2) = 1$ , we know that  $\gamma_k(n/q) > 1$  if and only if  $q > 2$ . Using the fact that

$$\frac{\gamma''_k(x)}{\gamma_k(x)} = (\alpha_k(x))^2 - \alpha'_k(x) = \left(\frac{\gamma'_k(x)}{\gamma_k(x)}\right)^2 + \sum_{j=0}^{k-1} \frac{(2j+n)(n-2x)}{(n+j-x)^2(j+x)^2},$$

we have  $\gamma''_k(x) \geq 0$ , which establishes the convexity of  $\gamma_k$  on  $[0, n/2]$ . Moreover, we know that

$$\gamma'_k\left(\frac{n}{2}\right) = -\alpha_k\left(\frac{n}{2}\right) = -\sum_{j=0}^{k-1} \frac{4}{n+2j}.$$

See Fig. 1. Taking these observations into account, we can state the following result.

**Lemma 9.** *Assume that  $n \geq 1$ . With the above notations, the function*

$$q \mapsto \frac{\gamma_k\left(\frac{n}{q}\right) - 1}{q - 2}$$

*is strictly monotone increasing on  $(1, \infty)$  for any  $k \geq 2$ .*

*Proof.* Let us prove that  $q \mapsto \gamma_k(n/q)$  is strictly *convex* w.r.t.  $q$  for any  $k \geq 2$ . Written in terms of  $x = n/q$ , it is sufficient to prove that

$$x \gamma''_k + 2 \gamma'_k > 0 \quad \forall x \in (0, n),$$

which can also be rewritten as

$$\alpha_k^2 - \alpha'_k - \frac{2}{x} \alpha_k > 0.$$

Let us prove this inequality. Using the estimates

$$\alpha_k^2 = \left(\sum_{j=0}^{k-1} \beta_j\right)^2 \geq 2\beta_0 \sum_{j=1}^{k-1} \beta_j + \sum_{j=0}^{k-1} \beta_j^2,$$

$$\beta_0^2 - \beta'_0 - \frac{2}{x} \beta_0 = 0,$$

and

$$2\beta_0 \beta_j + \beta_j^2 - \beta'_j - \frac{2}{x} \beta_j = \frac{2(n+j)(n+2j)}{(n-x)(n+j-x)(j+x)^2}$$

for any  $j \geq 1$ , we actually find that

$$\alpha_k^2 - \alpha'_k - \frac{2}{x} \alpha_k \geq \sum_{j=1}^{k-1} \frac{2(n+j)(n+2j)}{(n-x)(j+n-x)(j+x)^2} \quad \forall k \geq 2,$$

which concludes the proof. Note that as a byproduct, we also proved the strict convexity of  $\gamma_k$  for the whole range  $x \in (0, n)$ .  $\square$

*Proof of Theorem 1.* We deduce from (5) that

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq \sum_{k=1}^{\infty} \frac{\gamma_k\left(\frac{n}{q}\right) - 1}{q-2} \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

because  $\gamma_0(x) = 1$ . It follows from Lemma 9 that

$$\begin{aligned} \frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} &\leq \sum_{k=1}^{\infty} \frac{\gamma_k\left(\frac{n}{q_*}\right) - 1}{q_* - 2} \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu \\ &= \frac{\kappa_{n,s}}{q_* - 2} \sum_{k=1}^{\infty} \delta_k\left(\frac{n}{q_*}\right) \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu = \frac{\kappa_{n,s}}{q_* - 2} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu. \end{aligned}$$

This proves that  $C_{q,s} \leq \frac{\kappa_{n,s}}{q_* - 2}$ . The reverse inequality has already been shown in (16).  $\square$

*Proof of Theorem 5.* With  $s \in (-n, 0)$ , it turns out that  $q_*$  defined by (4) is in the range  $(1, 2)$  and plays the role of  $p$  in (12). According to Lemma 9, the inequality holds with the same constant for any  $q \in (1, q_*)$ , and this constant is optimal because of (16).  $\square$

**2.3. An improved inequality with a remainder term.** What we have shown in Section 2.2 is actually that the fractional Sobolev inequality (5) is equivalent to the following improved subcritical inequality.

**Corollary 10.** *Assume that  $n \geq 1$ ,  $q \in [1, 2) \cup (2, q_*)$  if  $s \in (0, n)$ , and  $q \in [1, 2) \cup (2, \infty)$  if  $s = n$ . For any  $F \in H^{s/2}(\mathbb{S}^n)$  we have*

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} + \int_{\mathbb{S}^n} F \mathcal{R}_{q,s} F d\mu \leq \frac{\kappa_{n,s}}{q_* - 2} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu$$

where  $\mathcal{R}_{q,s}$  is a positive semi-definite operator whose kernel is generated by the spherical harmonics corresponding to  $k = 0$  and  $k = 1$ .

*Proof.* We observe that

$$\int_{\mathbb{S}^n} F \mathcal{R}_{q,s} F d\mu := \sum_{k=2}^{\infty} \epsilon_k \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

where

$$\epsilon_k := \frac{\gamma_k\left(\frac{n}{q_*}\right) - 1}{q_* - 2} - \frac{\gamma_k\left(\frac{n}{q}\right) - 1}{q - 2}$$

is positive for any  $k \geq 2$  according to Lemma 9.  $\square$

Equality in (7) is realized only when  $F$  optimizes the critical fractional Sobolev inequality and, if  $q < q_*$ , when  $F_{(k)} = 0$  for any  $k \geq 2$ , which is impossible unless  $F$  is an optimal function for the Poincaré inequality of Lemma 8. This observation will be further exploited in Section 4.

#### 2.4. Fractional logarithmic Sobolev inequalities.

*Proof of Corollary 4.* According to Theorem 1, we know by (7) that

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq \frac{n-s}{2s} \kappa_{n,s} \int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu$$

for any function  $F \in H^{s/2}(\mathbb{S}^n)$  and any  $q \in [1, 2) \cup (2, q_*)$  with  $q_* = q_*(s)$  given by (4) (and the convention that  $q_* = \infty$  if  $s = n$ ). By taking the limit as  $q \rightarrow 2$  for a given  $s \in (0, n)$ , we obtain that (11) holds with  $C_{2,s} \leq \frac{n-s}{2s} \kappa_{n,s}$ . The reverse inequality has already been shown in (16) written with  $q = 2$ .  $\square$

Let us comment on the results of Corollary 4, in preparation for Section 4. Instead of fixing  $s$  and letting  $q \rightarrow 2$  as in the proof of Corollary 4, we can consider the case  $q = q_*(s)$  and let  $s \rightarrow 0$ , or equivalently rewrite (5) as

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq \sum_{k=0}^{\infty} \frac{\gamma_k(\frac{n}{q}) - 1}{q-2} \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu,$$

and take the limit as  $q \rightarrow 2$ . By an endpoint differentiation argument, we recover the conformally invariant *fractional logarithmic Sobolev* inequality

$$(18) \quad \int_{\mathbb{S}^n} F^2 \log \left( \frac{|F|}{\|F\|_{L^2(\mathbb{S}^n)}} \right) d\mu \leq \frac{n}{2} \int_{\mathbb{S}^n} F \mathcal{K}'_0 F d\mu$$

as in [4, 6], where the differential operator  $\mathcal{K}'_0$  is the endpoint derivative of  $\mathcal{K}_s$  at  $s = 0$ . The equality  $\mathcal{K}'_0 = \mathcal{L}'_0$  holds because  $\kappa_{n,0} = 1$  and  $\mathcal{K}_0 = \text{Id}$ . More specifically the right side of (18) can be written using the identities

$$\int_{\mathbb{S}^n} F \mathcal{K}'_0 F d\mu = \int_{\mathbb{S}^n} F \mathcal{L}'_0 F d\mu = \frac{1}{2} \sum_{k=0}^{\infty} \alpha_k(\frac{n}{2}) \int_{\mathbb{S}^n} |F_{(k)}|^2 d\mu$$

with  $\alpha_k(\frac{n}{2}) = -\gamma'_k(\frac{n}{2}) = \sum_{j=0}^{k-1} \frac{4}{n+2j}$ .

Inequality (18) is sharp, and equality holds if and only if  $F$  is obtained after applying any conformal transformation on  $\mathbb{S}^n$  to constant functions. Finally, let us notice that (18) can be recovered as an endpoint of (11) by letting  $s \rightarrow 0$ . The critical case is then achieved as a limit of the subcritical inequalities (11): the optimal constant can be identified, but the set of optimal functions in the limit is larger than in the subcritical regime, because of the conformal invariance.

Even more interesting is the fact that the *fractional logarithmic Sobolev* inequality is critical for  $s = 0$  and  $q = 2$  but subcritical inequalities corresponding to  $q \in [1, 2)$  still make sense.

**Corollary 11.** *Assume that  $n \geq 1$  and  $q \in [1, 2)$ . For any  $F \in L^2(\mathbb{S}^n)$  such that  $\int_{\mathbb{S}^n} F \mathcal{K}'_0 F d\mu$  is finite, we have*

$$\frac{\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2}{q-2} \leq \frac{n}{2} \int_{\mathbb{S}^n} F \mathcal{K}'_0 F d\mu.$$

As for Corollary 4, the proof relies on Lemma 9. Details are left to the reader.

## 3. STEREOGRAPHIC PROJECTION AND WEIGHTED FRACTIONAL INTERPOLATION INEQUALITIES ON THE EUCLIDEAN SPACE

This section is devoted to the proofs of Theorem 6 and Corollary 7. Various results concerning the extension of the Caffarelli-Kohn-Nirenberg inequalities introduced in [9] (also see [20, Theorem 1] in our context) are scattered in the literature, and one can refer for instance to [18, Theorem 1.8] for a quite general result in this direction. However, very little is known so far on optimal constants or even estimates of such constants, except for some limit cases like fractional Sobolev or fractional Hardy-Sobolev inequalities (see, *e.g.*, [45]). What we prove here is that the interpolation inequalities on the sphere provide inequalities on the Euclidean space with weights based on  $(1 + |x|^2)$ , with optimal constants and, using a scaling, Caffarelli-Kohn-Nirenberg inequalities with more standard power law weights.

*Proof of Theorem 6.* Let us consider the stereographic projection  $\mathcal{S}$ , whose inverse is defined by

$$\mathcal{S}^{-1} : \mathbb{R}^n \rightarrow \mathbb{S}^n, \quad x \mapsto \zeta = \left( \frac{2x}{1 + |x|^2}, \frac{1 - |x|^2}{1 + |x|^2} \right).$$

with Jacobian determinant  $|J| = 2^n (1 + |x|^2)^{-n}$ . Given  $s \in (0, n)$  and  $q \in (2, q_*)$ , Inequality (7) can be written using the conformal Laplacian as

$$\|F\|_{L^q(\mathbb{S}^n)}^2 - \frac{q_* - q}{q_* - 2} \|F\|_{L^2(\mathbb{S}^n)}^2 \leq \frac{q - 2}{q_* - 2} \kappa_{n,s} \int_{\mathbb{S}^n} F \mathcal{A}_s F \, d\mu$$

where  $\mathcal{A}_s$  and the fractional Laplacian on  $\mathbb{R}^n$  are related by

$$|J|^{1 - \frac{1}{q_*}} (\mathcal{A}_s F) \circ \mathcal{S}^{-1} = (-\Delta)^{s/2} \left( |J|^{\frac{1}{q_*}} F \circ \mathcal{S}^{-1} \right).$$

Then the interpolation inequality (7) on the sphere is equivalent to the following *fractional interpolation inequality on the Euclidean space*

$$\begin{aligned} |\mathbb{S}^n|^{1 - \frac{2}{q}} \left( \int_{\mathbb{R}^n} |f|^q |J|^{1 - \frac{q}{q_*}} \, dx \right)^{\frac{2}{q}} - \frac{q_* - q}{q_* - 2} \int_{\mathbb{R}^n} f^2 |J|^{1 - \frac{2}{q_*}} \, dx \\ \leq \frac{q - 2}{q_* - 2} \kappa_{n,s} \int_{\mathbb{R}^n} f (-\Delta)^{s/2} f \, dx \end{aligned}$$

after using the change of variables  $F \mapsto f = |J|^{1/q_*} F \circ \mathcal{S}^{-1}$ . The equality case is now achieved only by  $f = |J|^{1/q_*}$  for any  $q \in (2, q_*)$ , up to a multiplication by a constant, and the inequality is equivalent to (13).  $\square$

*Proof of Corollary 7.* Since  $\|g\|_{L_*^{2,2s}(\mathbb{R}^n)}^2 \leq \|g\|_{L^{2,2s}(\mathbb{R}^n)}^2$ , we deduce from (13) that

$$\|g\|_{L_*^{q,\beta}(\mathbb{R}^n)}^2 \leq \mathbf{a} \|g\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 + \mathbf{b} \|g\|_{L^{2,2s}(\mathbb{R}^n)}^2.$$

By applying this inequality to  $g = g_\lambda$  with  $g_\lambda(x) := \lambda^{(n-\beta)/q} g_1(\lambda x)$ , we observe that the r.h.s. can be written as

$$\mathbf{a} \|g_\lambda\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2 + \mathbf{b} \|g_\lambda\|_{L^{2,2s}(\mathbb{R}^n)}^2 = \mathbf{A} \lambda^{-a} + \mathbf{B} \lambda^b$$

where  $A = \mathbf{a} \|g_1\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^2$ ,  $B = \mathbf{b} \|g_1\|_{L^{2,2s}(\mathbb{R}^n)}^2$ ,  $b = n(1 - 2/q)$  and  $a = s - b$ . If we optimize the r.h.s. with respect to  $\lambda > 0$ , we obtain for  $\lambda = \lambda_*$  that it is given by  $\lambda_*^s = \frac{aA}{bB}$  and equal to  $A \lambda_*^{-a} + B \lambda_*^b = \frac{s}{a} \left(\frac{a}{b} A\right)^\vartheta B^{1-\vartheta}$  with  $\vartheta = b/(a+b)$ . Hence we have shown that

$$\left( \int_{\mathbb{R}^n} \frac{|g_1|^q}{(\lambda_*^2 + |x|^2)^{n(1-\frac{q}{q_*})}} dx \right)^{\frac{2}{q}} \leq K_{s,q} \|g_1\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^{2\vartheta} \|g_1\|_{L^{2,2s}(\mathbb{R}^n)}^{2(1-\vartheta)}$$

with  $K_{s,q} := \frac{s}{a} \left(\frac{a}{b} \mathbf{a}\right)^\vartheta \mathbf{b}^{1-\vartheta}$ . The r.h.s. is now invariant under the scaling corresponding to a second scaling given by  $g_1(x) := \mu^{(n-\beta)/q} f(\mu x)$ , and the inequality amounts, in terms of  $f$ , to

$$\left( \int_{\mathbb{R}^n} \frac{|f|^q}{(\mu^2 \lambda_*^2 + |x|^2)^{n(1-\frac{q}{q_*})}} dx \right)^{\frac{2}{q}} \leq K_{s,q} \|f\|_{\dot{H}^{s/2}(\mathbb{R}^n)}^{2\vartheta} \|f\|_{L^{2,2s}(\mathbb{R}^n)}^{2(1-\vartheta)}$$

for an arbitrary  $\mu > 0$ . We conclude by taking the limit as  $\mu \rightarrow 0_+$ .

In the limit case as  $q \rightarrow 2$ , the inequality becomes an equality, so that we can differentiate with respect to  $q$  and get a logarithmic Hardy inequality as in the  $s = 2$  case: see [20] for more details in a similar problem.  $\square$

#### 4. CONCLUDING REMARKS

A striking feature of Inequality (7) is that the optimal constant  $C_{q,s}$  is determined by a linear eigenvalue problem, although the problem is definitely nonlinear. This deserves some comments. Let  $q \in [1, 2) \cup (2, q_*)$  if  $s < n$  and  $q \in [1, 2) \cup (2, \infty)$  if  $s = n$ . With  $\mathcal{Q}$  defined by (9) on  $\mathcal{H}^{s/2}$ , the subset of the functions in  $H^{s/2}(\mathbb{S}^n)$  which are not almost everywhere constant, we investigate the relation

$$C_{q,s} \inf_{F \in \mathcal{H}^{s/2}} \mathcal{Q}[F] = 1.$$

Notice that both numerator and denominator of  $\mathcal{Q}[F]$  converge to 0 if  $F$  approaches a constant, so that  $\mathcal{Q}$  becomes undetermined in the limit. As we shall see next, this happens for a minimizing sequence and explains why a linearized problem appears in the limit.

By compactness of the Sobolev embedding  $H^{s/2}(\mathbb{S}^n) \hookrightarrow L^q(\mathbb{S}^n)$  (see [43, 1, 18] for fundamental properties of fractional Sobolev spaces, [22, sections 6 and 7] and [41] for application to variational problems), any minimizing sequence  $(F_n)_{n \in \mathbb{N}}$  for  $\mathcal{Q}$  is relatively compact if we assume that  $\|F_n\|_{L^q(\mathbb{S}^n)} = 1$  for any  $n \in \mathbb{N}$ . This normalization can be imposed without loss of generality because of the homogeneity of  $\mathcal{Q}$ . Hence  $(F_n)_{n \in \mathbb{N}}$  converges to a limit  $F \in H^{s/2}(\mathbb{S}^n)$ . Assume that  $F$  is not a constant. Then the denominator in  $\mathcal{Q}[F]$  is positive and by semicontinuity we know that

$$\int_{\mathbb{S}^n} F \mathcal{L}_s F d\mu \leq \lim_{n \rightarrow +\infty} \int_{\mathbb{S}^n} F_n \mathcal{L}_s F_n d\mu.$$

On the other hand, by compactness, up to the extraction of a subsequence, we have that

$$\|F\|_{L^2(\mathbb{S}^n)}^2 = \lim_{n \rightarrow +\infty} \|F_n\|_{L^2(\mathbb{S}^n)}^2 \quad \text{and} \quad \|F\|_{L^q(\mathbb{S}^n)}^2 = \lim_{n \rightarrow +\infty} \|F_n\|_{L^q(\mathbb{S}^n)}^2 = 1.$$

Hence  $F$  is optimal and solves the Euler-Lagrange equations

$$(q-2)C_{q,s}\mathcal{L}_s F + F = F^{q-1}.$$

Using Corollary 10, we also get that  $F$  lies in the kernel of  $\mathcal{R}_{q,s}$ , that is, the space generated by the spherical harmonics corresponding to  $k=0$  and  $k=1$ . From the Euler-Lagrange equations, we read that  $F$  has to be a constant. Because of the normalization  $\|F\|_{L^q(\mathbb{S}^n)} = 1$ , we obtain that  $F = 1$  a.e., a contradiction.

Hence  $(F_n)_{n \in \mathbb{N}}$  converges to 1 in  $H^{s/2}(\mathbb{S}^n)$ . With  $\varepsilon_n = \|1 - F_n\|_{H^{s/2}(\mathbb{S}^n)}$  and  $v_n := (F_n - 1)/\varepsilon_n$ , we can write that

$$F_n = 1 + \varepsilon_n v_n \quad \text{with} \quad \|v_n\|_{H^{s/2}(\mathbb{S}^n)} = 1 \quad \forall n \in \mathbb{N}$$

and

$$\lim_{n \rightarrow +\infty} \varepsilon_n = 0.$$

On the other hand, it turns out that  $(F_n)_{n \in \mathbb{N}}$  being a minimizing sequence,

$$C_{q,s}^{-1} = \lim_{n \rightarrow +\infty} \mathcal{Q}[F_n] = \lim_{n \rightarrow +\infty} \frac{\varepsilon_n^2 (q-2) \int_{\mathbb{S}^n} v_n \mathcal{L}_s v_n d\mu}{\|1 + \varepsilon_n v_n\|_{L^q(\mathbb{S}^n)}^2 - \|1 + \varepsilon_n v_n\|_{L^2(\mathbb{S}^n)}^2}.$$

If  $q > 2$ , an elementary computation shows that

$$(19) \quad \|1 + \varepsilon_n v_n\|_{L^q(\mathbb{S}^n)}^2 - \|1 + \varepsilon_n v_n\|_{L^2(\mathbb{S}^n)}^2 = (q-2) \varepsilon_n^2 \|v_n - \bar{v}_n\|_{L^2(\mathbb{S}^n)}^2 (1 + o(1))$$

as  $n \rightarrow +\infty$ , where  $\bar{v}_n := \int_{\mathbb{S}^n} v_n d\mu$ , so that

$$C_{q,s}^{-1} = \lim_{n \rightarrow +\infty} \mathcal{Q}[F_n] = \lim_{n \rightarrow +\infty} \frac{\int_{\mathbb{S}^n} v_n \mathcal{L}_s v_n d\mu}{\|v_n - \bar{v}_n\|_{L^2(\mathbb{S}^n)}^2}.$$

Details on the Taylor expansion used in (19) can be found in Appendix B. When  $q \in [1, 2)$ , we can estimate the denominator by restricting the integrals to  $\{x \in \mathbb{S}^n : \varepsilon_n |v_n| < 1/2\}$  and Taylor expand  $t \mapsto (1+t)^q$  on  $(1/2, 3/2)$ .

Notice that  $F_n$  being a function in  $\mathcal{H}^{s/2}$ , we know that  $\|v_n - \bar{v}_n\|_{L^2(\mathbb{S}^n)} > 0$  for any  $n \in \mathbb{N}$ , so that the above limit makes sense. With the notations of Section (2.1), we know that

$$C_{q,s}^{-1} \geq \inf_{v \in \mathcal{H}^{s/2}} \frac{\int_{\mathbb{S}^n} v \mathcal{L}_s v d\mu}{\|v - \bar{v}\|_{L^2(\mathbb{S}^n)}^2} \geq \lambda_1(\mathcal{L}_s) = \frac{2s\kappa_{n,s}}{n-s}$$

according to the Poincaré inequality of Lemma 8, which proves that we actually have equality in (16) and determines  $C_{q,s}$ .

Additionally, we may notice that  $(v_n)_{n \in \mathbb{N}}$  has to be a minimizing sequence for the Poincaré inequality, which means that up to a normalization and after the extraction of a subsequence,  $v_n - \bar{v}_n$  converges to a spherical harmonic function associated with the component corresponding to  $k=1$ . This explains why we obtain that  $C_{q,s} \lambda_1(\mathcal{L}_s) = 1$ .

The above considerations have been limited to the subcritical range  $q < q_*$  if  $s < n$  and  $q < +\infty$  if  $s = n$ . However, the critical case of the Sobolev inequality can be obtained by passing to the limit as  $q \rightarrow q_*$  (and even the Onofri type inequalities when  $s = n$ ) so that the optimal constants are also given by an eigenvalue in the critical case. However, due to the conformal invariance, the constant function  $F \equiv 1$



is not the only optimal function. At this point it should be noted that the above considerations heavily rely on Corollary 10 and, as a consequence, cannot be used to give a variational proof of Theorem 1.

Although the subcritical interpolation inequalities of this paper appear weaker than inequalities corresponding to a critical exponent, we are able to identify the equality cases and the optimal constants. We are also able to keep track of a remainder term which characterizes the functions realizing the optimality of the constant or, to be precise, the limit of any minimizing sequence and its first order correction. This first order correction, or equivalently the asymptotic value of the quotient  $\mathcal{Q}$ , determines the optimal constant and explains the role played by the eigenvalues in a problem which is definitely nonlinear.

#### APPENDIX A. THE SPECTRUM OF THE FRACTIONAL LAPLACIAN

The standard approach for computing  $\gamma_k$  in (6) relies on the Funk-Hecke formula as it is detailed in [33, Section 4]. In this appendix, for completeness, we provide a simple, direct proof of the expression of  $\gamma_k$ . For this purpose, we compute the eigenvalues  $\lambda_k = \lambda_k((-\Delta)^{s/2})$  of the fractional Laplacian on  $\mathbb{R}^n$ , that is,

$$(-\Delta)^{s/2} f_k = \frac{\lambda_k}{(1 + |x|^2)^s} f_k \quad \text{in } \mathbb{R}^n,$$

for any  $k \in \mathbb{N}$ . We shall then deduce the eigenvalues of  $\mathcal{L}_s$ . This determines the optimal constant in (5) and (7) without using Lieb's duality and without relying on the symmetry of the optimal case in (1) as in [38].

**Proposition 12.** *Given  $s \in (0, n)$ , the spectrum of the fractional Laplacian is*

$$\lambda_k((-\Delta)^{s/2}) = 2^s \frac{\Gamma(k + \frac{n}{q'})}{\Gamma(k + \frac{n}{q})} = 2^s \lambda_k(\mathcal{A}_s) = 2^s \frac{\Gamma(\frac{n}{q'})}{\Gamma(\frac{n}{q})} \lambda_k(\mathcal{K}_s).$$

*Proof.* Using the stereographic projection and a decomposition in spherical harmonics, we can reduce the problem of computing the spectrum to the computation of the spectrum associated with the eigenfunctions

$$f_k^\mu(x) = C_k^{(\alpha)}(z) (1 + |x|^2)^{-\mu} \quad \text{with } z = \frac{1 - |x|^2}{1 + |x|^2},$$

where  $\mu = \lambda/2 = (n - s)/2$ ,  $\alpha = (n - 1)/2$  and  $C_k^{(\alpha)}$  denotes the Gegenbauer polynomials. Let  $\hat{f}(\xi) = (\mathcal{F}f)(\xi) := \int_{\mathbb{R}^n} f(x) e^{-2\pi i \xi \cdot x} dx$  be the Fourier transform of a function  $f$ . The functions being radial, by the Hankel transform  $\mathcal{H}^{\frac{n}{2}-1}$  we get that

$$\widehat{f}_k^\mu(\xi) = \frac{2\pi}{|\xi|^{\frac{n}{2}-1}} \int_0^\infty f_k^\mu(r) J_{\frac{n}{2}-1}(2\pi r |\xi|) r^{\frac{n}{2}} dr$$

(cf. [35, Appendix B.5, p. 578]) where  $J_\nu$  is the Bessel function of the first kind.

The Fourier transform of  $f_0^\mu = (1 + |x|^2)^{-\mu}$  has been calculated, e.g., by E. Lieb in [38, p. 360, Eqs. (3.9)-(3.14)] in terms of the modified Bessel functions of the second kind  $K_\nu$  as

$$\widehat{f}_0^\mu(\xi) = \frac{\pi^{\frac{n}{2}} 2^{1+\frac{2}{n}-\mu}}{\Gamma(\mu)} (2\pi |\xi|)^{\mu-\frac{n}{2}} K_{\mu-\frac{n}{2}}(2\pi |\xi|).$$

This is a special case of the modified Weber-Schafheitlin integral formula in [44, Chapter XIII, 13.45]. Using the expansion of Gegenbauer polynomials, we get

$$\begin{aligned}
 \widehat{f}_k^\mu(\xi) &= \frac{2\pi}{\Gamma(\frac{n-1}{2})|\xi|^{\frac{n}{2}-1}} \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \sum_{l=0}^{k-2j} \left[ \frac{(-1)^{j+k-l} 2^{k+l-2j} \Gamma(\frac{n-1}{2} + k - j)}{j! k! (k-2j-l)!} \right. \\
 &\quad \left. \times \int_0^\infty (1+r^2)^{-(\mu+l)} J_{\frac{n}{2}-1}(2\pi r |\xi|) r^{\frac{n}{2}} dr \right] \\
 &= \frac{1}{\Gamma(\frac{n-1}{2})} \sum_{j=0}^{\lfloor \frac{k}{2} \rfloor} \sum_{l=0}^{k-2j} \frac{(-1)^{j+k-l} 2^{k+l-2j} \Gamma(\frac{n-1}{2} + k - j)}{j! l! (k-2j-l)!} \widehat{f_0^{\mu+l}}(\xi) \\
 &= \frac{2^{1+\frac{2}{n}-\mu} \pi^{\frac{n}{2}} (2\pi |\xi|)^{\mu-\frac{n}{2}}}{\Gamma(\frac{n-1}{2}) \Gamma(\mu+k)} I_{n,k}^\mu(|\xi|),
 \end{aligned}$$

where

$$\begin{aligned}
 I_{n,k}^\mu(|\xi|) &:= \sum_{l=0}^k c_{n,k,l} \frac{\Gamma(\mu+k)}{\Gamma(\mu+l)} (2\pi |\xi|)^l K_{\mu-\frac{n}{2}+l}(2\pi |\xi|), \\
 \text{and } c_{n,k,l} &:= \frac{1}{l!} \sum_{j=0}^{\lfloor \frac{k-l}{2} \rfloor} \frac{(-1)^{j+k-l} 2^{k-2j} \Gamma(\frac{n-1}{2} + k - j)}{j! (k-2j-l)!}.
 \end{aligned}$$

From the recurrence relation

$$x(K_{\nu-1} - K_{\nu+1}) = -2\nu K_\nu,$$

we deduce the identity

$$\sum_{l=0}^k c_{n,k,l} x^l \left( \frac{\Gamma(\nu + \frac{n}{2} + k)}{\Gamma(\nu + \frac{n}{2} + l)} K_{\nu+l}(x) - \frac{\Gamma(-\nu + \frac{n}{2} + k)}{\Gamma(-\nu + \frac{n}{2} + l)} K_{\nu-l}(x) \right) = 0 \quad \forall k \geq 0$$

and observe that

$$I_{n,k}^{\mu_1} = I_{n,k}^{\mu_2} \quad \forall k \in \mathbb{N}$$

if  $\mu_1 = \lambda/2$  and  $\mu_2 = \lambda/2 + s$ , so that  $\mu_1 + \mu_2 = n$  and  $\mu_1 - \mu_2 = -s$ . It remains to observe that

$$(2\pi |\xi|)^s \widehat{f_k^{\lambda/2}} = \lambda_k \mathcal{F} \left( f_k^{\lambda/2} (1+|x|^2)^{-s} \right) \quad \text{with } \lambda_k = 2^s \frac{\Gamma(k + \frac{n}{2})}{\Gamma(k + \frac{n}{2} + s)}.$$

□

## APPENDIX B. A TAYLOR FORMULA WITH INTEGRAL REMAINDER TERM

Let us define the function  $r : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$|1+t|^q = 1 + qt + \frac{1}{2} q(q-1)t^2 + r(t) \quad \forall t \in \mathbb{R}.$$

**Lemma 13.** *Let  $q \in (2, \infty)$ . With the above notations, there exists a constant  $C > 0$  such that*

$$|r(t)| \leq C |t|^3 \quad \text{if } |t| \leq 1 \quad \text{and} \quad |r(t)| \leq C |t|^q \quad \text{if } |t| \geq 1.$$

This result is elementary but crucial for the expansion of  $\|F\|_{L^q(\mathbb{S}^n)}^2 - \|F\|_{L^2(\mathbb{S}^n)}^2$  around  $F = 1$ . This is why we give a proof with some details, although we claim absolutely no originality for that. Similar computations have been repeatedly used in a related context, *e.g.*, in [15, 17, 40].

*Proof.* Using the Taylor formula with integral remainder term

$$f(t) = f(0) + f'(0)t + \frac{1}{2}f''(0)t^2 + \frac{1}{2}\int_0^t (t-s)^2 f'''(s) ds$$

applied to  $f(t) = (1+t)^q$  with  $q > 2$ , we obtain that

$$|1+t|^q = 1 + qt + \frac{1}{2}q(q-1)t^2 + r(t)$$

where the remainder term is given by

$$r(t) = \frac{1}{2}q(q-1)(q-2)t^q \int_0^1 (1-\sigma)^2 \left| \frac{1}{t} + \sigma \right|^{q-4} \left( \frac{1}{t} + \sigma \right) d\sigma.$$

Hence the remainder term can be bounded as follows:

(i) if  $t \geq 1$ , using  $\sigma < \frac{1}{t} + \sigma < 1 + \sigma$ , we get that

$$0 < r(t) < c_q t^q$$

with  $c_q = \frac{1}{2}q(q-1)(q-2)\int_0^1 (1-\sigma)^2 \max\{\sigma^{q-3}, (1+\sigma)^{q-3}\} d\sigma$ ,

(ii) if  $0 < t < 1$ , we get that

$$0 < r(t) < \frac{1}{6}q(q-1)(q-2) \max\{1, 2^{q-3}\} t^3$$

using  $\frac{1}{t} < \frac{1}{t} + \sigma < \frac{2}{t}$ ,

(iii) if  $-1 < t < 0$ , we get that

$$-\frac{1}{6}q(q-1)(q-2)|t|^3 < r(t) < 0$$

using  $\frac{1}{t} < \frac{1}{t} + \sigma < \frac{1}{t} + 1 < 0$ ,

(iv) if  $t \leq -1$ , we get that

$$-\frac{1}{2}(q-1)(q-2)t^q < r(t) < t^q$$

using  $\sigma - 1 < \frac{1}{t} + \sigma < \sigma$ .

This completes the proof.  $\square$

## APPENDIX C. NOTATIONS AND RANGES

For the convenience of the reader, this appendix collects various notations which are used throughout this paper and summarizes the ranges covered by the parameters.

The identity

$$\lambda = \frac{2n}{p'} \quad \text{where} \quad \frac{1}{p} + \frac{1}{p'} = 1$$

means that

$$p = \frac{2n}{2n - \lambda}.$$

With

$$\lambda = n - s,$$

we have

$$p = \frac{2n}{n + s} \quad \text{and} \quad p' = q_* = \frac{2n}{n - s}.$$

The limiting values of the parameters are summarized in Table 1.

$s$	0	2	$n$
$\lambda$	$n$	$n - 2$	0
$p$	2	$\frac{2n}{n+2}$	1
$p' = q_*$	2	$\frac{2n}{n-2}$	$+\infty$

TABLE 1. Correspondance of the limiting values of the parameters.

The coefficients  $\gamma_k$  and  $\delta_k$  defined by

$$\gamma_k(x) = \frac{\Gamma(x) \Gamma(n - x + k)}{\Gamma(n - x) \Gamma(x + k)}$$

$$\text{and} \quad \delta_k(x) = \frac{1}{\kappa_{n,s}} (\gamma_k(x) - 1) = \frac{\Gamma(n - x + k)}{\Gamma(x + k)} - \frac{\Gamma(n - x)}{\Gamma(x)}$$

are such that

$$\delta_k\left(\frac{n}{q_*}\right) = \frac{1}{\kappa_{n,s}} (\gamma_k\left(\frac{n}{q_*}\right) - 1) \quad \text{where} \quad \kappa_{n,s} = \frac{\Gamma\left(\frac{n}{q_*}\right)}{\Gamma\left(n - \frac{n}{q_*}\right)} = \frac{\Gamma\left(\frac{n-s}{2}\right)}{\Gamma\left(\frac{n+s}{2}\right)}.$$

We recall that  $\gamma_0(n/q) - 1 = 0$ ,  $\gamma_1(n/q) - 1 = q - 2$ ,  $\delta_k\left(\frac{n}{q_*}\right) = k(k + n - 1)$  and  $1/\kappa_{n,2} = \frac{1}{4}n(n - 2)$ . According to (17), we have that

$$\alpha_k(x) = -\frac{\gamma'_k(x)}{\gamma_k(x)} = \sum_{j=0}^{k-1} \beta_j(x) \quad \text{with} \quad \beta_j(x) = \frac{1}{n + j - x} + \frac{1}{j + x}$$

for any  $k \geq 1$ . With these notations, the eigenvalues of  $\mathcal{K}_s$ ,  $\mathcal{L}_s$  and  $\mathcal{K}'_0 = \mathcal{L}'_0$  are respectively given by  $\gamma_k(n/q_*(s)) = \gamma_k((n - s)/2)$ ,  $\kappa_{n,s}^{-1} (\gamma_k((n - s)/2) - 1)$  and  $\frac{1}{2} \alpha_k(n/2)$  with  $\alpha_k(n/2) = -\gamma'_k(n/2) = 4 \sum_{j=0}^{k-1} (n + 2j)^{-1}$ .




FIGURE 1. The functions  $x \mapsto \gamma_k(x)$  and  $q \mapsto \gamma_k(n/q)$  are both convex, and such that  $\gamma_k(n/2) = 1$ .

Finally, we recall that  $\mathcal{K}_s$ , the fractional Laplacian  $\mathcal{L}_s$  and the conformal fractional Laplacian  $\mathcal{A}_s$  satisfy the relations

$$\kappa_{n,s} \mathcal{A}_s = \mathcal{K}_s = \kappa_{n,s} \mathcal{L}_s + \text{Id}.$$

The results of Theorems 1 and 5 are summarized in Fig. 2.




FIGURE 2. The optimal constant  $C_{q,s}$  in (7) is independent of  $q$  and determined for any given  $s$  by the critical case  $q = q_*(s)$ : the Hardy-Littlewood-Sobolev inequality (1) if  $s \in (-n, 0)$  and the Sobolev inequality (5) if  $s \in (0, n)$ . The case  $s = 0$  is covered by Corollary 11, while  $q = 2$  corresponds to the fractional logarithmic Sobolev inequality (18) if  $s = 0$  and the subcritical fractional logarithmic Sobolev inequality by Corollary 4 if  $s \in (0, n]$ .

**Acknowledgements.** This work is supported by a public grant overseen by the French National Research Agency (ANR) as part of the “Investissements d’Avenir” program (A.Z., reference: ANR-10-LABX-0098, LabEx SMP) and by the projects *STAB* (J.D., A.Z.) and *Kibord* (J.D.) of the French National Research Agency (ANR). A.Z. thanks the ERC Advanced Grant BLOWDISOL (Blow-up, dispersion and solitons; PI: Frank Merle) # 291214 for support. The authors thank M.J. Esteban for fruitful discussions and suggestions.

© 2016 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

## REFERENCES

- [1] R. A. ADAMS, *Sobolev spaces*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. Pure and Applied Mathematics, Vol. 65.
- [2] A. ARNOLD, P. MARKOWICH, G. TOSCANI, AND A. UNTERREITER, *On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations*, Comm. Partial Differential Equations, 26 (2001), pp. 43–100.
- [3] D. BAKRY AND M. ÉMERY, *Diffusions hypercontractives*, in Séminaire de probabilités, XIX, 1983/84, vol. 1123 of Lecture Notes in Math., Springer, Berlin, 1985, pp. 177–206.
- [4] W. BECKNER, *Sobolev inequalities, the Poisson semigroup, and analysis on the sphere  $\mathbb{S}^n$* , Proc. Nat. Acad. Sci. U.S.A., 89 (1992), pp. 4816–4819.
- [5] ———, *Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality*, Ann. of Math. (2), 138 (1993), pp. 213–242.
- [6] ———, *Logarithmic Sobolev inequalities and the existence of singular integrals*, Forum Math., 9 (1997), pp. 303–323.
- [7] M. BERGER, P. GAUDUCHON, AND E. MAZET, *Le spectre d'une variété riemannienne*, Lecture Notes in Mathematics, Vol. 194, Springer-Verlag, Berlin, 1971.
- [8] M.-F. BIDAUT-VÉRON AND L. VÉRON, *Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations*, Invent. Math., 106 (1991), pp. 489–539.
- [9] L. CAFFARELLI, R. KOHN, AND L. NIRENBERG, *First order interpolation inequalities with weights*, Compositio Math., 53 (1984), pp. 259–275.
- [10] E. CARLEN AND M. LOSS, *Competing symmetries, the logarithmic HLS inequality and Onofri's inequality on  $\mathbb{S}^n$* , Geom. Funct. Anal., 2 (1992), pp. 90–104.
- [11] E. A. CARLEN AND M. LOSS, *Extremals of functionals with competing symmetries*, J. Funct. Anal., 88 (1990), pp. 437–456.
- [12] J. A. CARRILLO, Y. HUANG, M. C. SANTOS, AND J. L. VÁZQUEZ, *Exponential convergence towards stationary states for the 1D porous medium equation with fractional pressure*, J. Differential Equations, 258 (2015), pp. 736–763.
- [13] D. CHAFAÏ, *Entropies, convexity, and functional inequalities: on  $\Phi$ -entropies and  $\Phi$ -Sobolev inequalities*, J. Math. Kyoto Univ., 44 (2004), pp. 325–363.
- [14] L. CHEN, Z. LIU, AND G. LU, *Symmetry and regularity of solutions to the weighted Hardy-Sobolev type system*, Advanced Nonlinear Studies, 16 (2016), pp. 1–13.
- [15] S. CHEN, R. L. FRANK, AND T. WETH, *Remainder terms in the fractional Sobolev inequality*, Indiana Univ. Math. J., 62 (2013), pp. 1381–1397.
- [16] X. CHEN AND J. YANG, *Weighted fractional Sobolev inequality in  $\mathbb{R}^N$* , Advanced Nonlinear Studies, (2016).
- [17] M. CHRIST, *A sharpened Hausdorff-Young inequality*, [arXiv: 1406.1210](https://arxiv.org/abs/1406.1210), (2014).
- [18] P. D'ANCONA AND R. LUCÀ, *Stein-Weiss and Caffarelli-Kohn-Nirenberg inequalities with angular integrability*, J. Math. Anal. Appl., 388 (2012), pp. 1061–1079.
- [19] M. DEL PINO AND J. DOLBEAULT, *The Euclidean Onofri inequality in higher dimensions*, International Mathematics Research Notices, 2013 (2012), pp. 3600–3611.
- [20] M. DEL PINO, J. DOLBEAULT, S. FILIPPAS, AND A. TERTIKAS, *A logarithmic Hardy inequality*, Journal of Functional Analysis, 259 (2010), pp. 2045 – 2072.
- [21] J. DEMANGE, *Improved Gagliardo-Nirenberg-Sobolev inequalities on manifolds with positive curvature*, J. Funct. Anal., 254 (2008), pp. 593–611.
- [22] E. DI NEZZA, G. PALATUCCI, AND E. VALDINOCI, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math., 136 (2012), pp. 521–573.
- [23] J. DOLBEAULT, *Sobolev and Hardy-Littlewood-Sobolev inequalities: duality and fast diffusion*, Math. Res. Lett., 18 (2011), pp. 1037–1050.
- [24] J. DOLBEAULT, M. J. ESTEBAN, AND M. LOSS, *Nonlinear flows and rigidity results on compact manifolds*, J. Funct. Anal., 267 (2014), pp. 1338–1363.
- [25] ———, *Interpolation inequalities on the sphere: linear vs. nonlinear flows*, Annales de la Faculté des Sciences de Toulouse. Mathématiques. Série 6, [arXiv: 1509.09099](https://arxiv.org/abs/1509.09099), (2016).

- [26] J. DOLBEAULT, M. J. ESTEBAN, AND G. TARANTELO, *Optimal Sobolev inequalities on  $\mathbb{R}^N$  and  $S^n$ : a direct approach using the stereographic projection*. Unpublished.
- [27] J. DOLBEAULT AND G. JANKOWIAK, *Sobolev and Hardy–Littlewood–Sobolev inequalities*, J. Differential Equations, 257 (2014), pp. 1689–1720.
- [28] J. DOLBEAULT, B. NAZARET, AND G. SAVARÉ, *A new class of transport distances between measures*, Calc. Var. Partial Differential Equations, 34 (2009), pp. 193–231.
- [29] J. DOLBEAULT AND G. TOSCANI, *Improved interpolation inequalities, relative entropy and fast diffusion equations*, Annales de l’Institut Henri Poincaré (C) Non Linear Analysis, 30 (2013), pp. 917 – 934.
- [30] J. DOLBEAULT AND G. TOSCANI, *Stability results for logarithmic Sobolev and Gagliardo–Nirenberg inequalities*, International Mathematics Research Notices, (2015).
- [31] R. L. FRANK AND E. H. LIEB, *Inversion positivity and the sharp Hardy–Littlewood–Sobolev inequality*, Calc. Var. Partial Differential Equations, 39 (2010), pp. 85–99.
- [32] ———, *Spherical reflection positivity and the Hardy–Littlewood–Sobolev inequality*, in Concentration, functional inequalities and isoperimetry, vol. 545 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2011, pp. 89–102.
- [33] ———, *A new, rearrangement-free proof of the sharp Hardy–Littlewood–Sobolev inequality*, in Spectral theory, function spaces and inequalities, vol. 219 of Oper. Theory Adv. Appl., Birkhäuser/Springer Basel AG, Basel, 2012, pp. 55–67.
- [34] N. GHOUSSOUB AND S. SHAKERIAN, *Borderline variational problems involving fractional Laplacians and critical singularities*, Adv. Nonlinear Stud., 15 (2015), pp. 527–555.
- [35] L. GRAFAKOS, *Classical Fourier analysis*, vol. 249 of Graduate Texts in Mathematics, Springer, New York, third ed., 2014.
- [36] G. JANKOWIAK AND V. HOANG NGUYEN, *Fractional Sobolev and Hardy–Littlewood–Sobolev inequalities*, arXiv: 1404.1028, (2014).
- [37] T. JIN AND J. XIONG, *A fractional Yamabe flow and some applications*, J. Reine Angew. Math., 696 (2014), pp. 187–223.
- [38] E. H. LIEB, *Sharp constants in the Hardy–Littlewood–Sobolev and related inequalities*, Ann. of Math. (2), 118 (1983), pp. 349–374.
- [39] E. H. LIEB AND M. LOSS, *Analysis*, vol. 14 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2001.
- [40] H. LIU AND A. ZHANG, *Remainder terms for several inequalities on some groups of Heisenberg-type*, Sci. China Math., 58 (2015), pp. 2565–2580.
- [41] G. MOLICA BISCI, V. D. RADULESCU, AND R. SERVADEI, *Variational methods for nonlocal fractional problems*, vol. 162 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, 2016. With a foreword by Jean Mawhin.
- [42] C. MÜLLER, *Spherical harmonics*, vol. 17 of Lecture Notes in Mathematics, Springer-Verlag, Berlin-New York, 1966.
- [43] A. PERSSON, *Compact linear mappings between interpolation spaces*, Ark. Mat., 5 (1964), pp. 215–219 (1964).
- [44] G. N. WATSON, *A treatise on the theory of Bessel functions*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1995. Reprint of the second (1944) edition.
- [45] J. YANG, *Fractional Sobolev–Hardy inequality in  $\mathbb{R}^N$* , Nonlinear Anal., 119 (2015), pp. 179–185.

[J. DOLBEAULT] CEREMADE, UMR CNRS n° 7534, UNIVERSITÉ PARIS-DAUPHINE, PSL RESEARCH UNIVERSITY, PLACE DE LATTRE DE TASSIGNY, 75775 PARIS 16, FRANCE.

*E-mail address:* [dolbeaul@ceremade.dauphine.fr](mailto:dolbeaul@ceremade.dauphine.fr)

[A. ZHANG] CEREMADE, UMR CNRS n° 7534, UNIVERSITÉ PARIS-DAUPHINE, PSL RESEARCH UNIVERSITY, PLACE DE LATTRE DE TASSIGNY, 75775 PARIS 16, FRANCE.

*E-mail address:* [zhang@ceremade.dauphine.fr](mailto:zhang@ceremade.dauphine.fr)