



**HAL**  
open science

# Clustering rates and Chung type laws of the iterated logarithm for empirical and quantile processes.

Davit Varron

► **To cite this version:**

Davit Varron. Clustering rates and Chung type laws of the iterated logarithm for empirical and quantile processes.. *Annales de l'ISUP*, 2011, 55 (2-3), pp.3-26. hal-01358247

**HAL Id: hal-01358247**

**<https://hal.science/hal-01358247>**

Submitted on 21 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Pub. Inst. Stat. Univ. Paris*  
55, fasc. 2-3, 2011, 3 à 26

# Clustering rates and Chung type functional laws of the iterated logarithm for empirical and quantile processes

Davit Varron

*Laboratoire de Mathématiques de Besançon, UMR CNRS 6623,*

*Université de Franche-Comté*

*e-mail: davit.varron@univ-fcomte.fr*

**Abstract:** Following the works of Berthet [2, 3], we first obtain exact clustering rates in the functional law of the iterated logarithm for the uniform empirical and quantile processes and for their increments. In a second time, we obtain functional Chung-type limit laws for the local empirical process for a class of target functions on the border of the Strassen set.

**AMS 2000 subject classifications:** Primary 62G20, 62G30.

**Keywords and phrases:** Empirical processes, Strassen laws of the iterated logarithm, Clustering rates, Chung-Mogulskii limit laws.

## 1. Introduction

Define the uniform empirical process by  $\alpha_n(t) := n^{1/2}(F_n(t) - t)$ , where  $F_n(t) := n^{-1}\#\{i \in \{1, \dots, n\}, U_i \leq t\}$ ,  $t \in [0, 1]$ , and  $(U_i)_{n \geq 1}$  are independent, identically distributed (i.i.d) random variables uniformly distributed on  $[0, 1]$ . Define

the quantile process by

$$\beta_n(t) = n^{1/2} \left( F_n^{-1}(t) - t \right), \quad t \in [0, 1],$$

where  $F_n^{-1}(t) := \inf\{u : F_n(u) \geq t\}$ . In a metric space  $(\mathcal{E}, d)$  we write  $u_n \rightsquigarrow \mathcal{H}$  whenever  $u_n$  is relatively compact with limit set  $\mathcal{H}$  (see, e.g., [17]). The two above mentioned processes have been extensively investigated in the literature (see, e.g., [20] and [24] and the references therein). In a pioneering work, Finkelstein [10] has established the functional law of the iterated logarithm (FLIL) for  $\alpha_n$ . Namely, the author showed that, writing  $\log_2 u = \log(\log(u \vee e))$  and  $b_n = \sqrt{2 \log_2 n}$ , we have :

$$\frac{\alpha_n}{b_n} \rightsquigarrow_{a.s.} \mathcal{S}_2, \quad (1.1)$$

in the metric space  $(B[0, 1], \|\cdot\|)$ , where  $B[0, 1]$  stands for the set of bounded functions on  $[0, 1]$  and  $\|\cdot\|$  is the sup-norm over  $[0, 1]$ . The set  $\mathcal{S}_2$  in (1.1) is given by

$$\mathcal{S}_2 := \left\{ f(t) \in \mathcal{S}_1, f(1) = 0 \right\}, \quad (1.2)$$

where

$$\mathcal{S}_1 := \left\{ f \in B[0, 1], \exists f' \text{ Borel}, f(\cdot) := \int_0^\cdot f'(t) dt, \int_0^1 f'^2(t) dt \leq 1 \right\}. \quad (1.3)$$

Note that  $\mathcal{S}_2$  (resp.  $\mathcal{S}_1$ ) is the unit ball of the reproducing kernel Hilbert space of the Brownian bridge (resp. of the Wiener process) on  $[0, 1]$ . In the spirit of [10], Mason [17] has obtained the following FLIL for the local empirical process :

$$\frac{\alpha_n(a_n \cdot)}{\sqrt{a_n} b_n} \rightsquigarrow_{a.s.} \mathcal{S}_1. \quad (1.4)$$

$\infty$ . Deheuvels and Mason [8] have established a related uniform functional limit law for the following collections of random trajectories.

$$\Theta_n := \left\{ \frac{\alpha_n(t + a_n \cdot) - \alpha_n(t)}{\sqrt{2a_n \log(1/a_n)}}, t \in [0, 1 - a_n] \right\}.$$

They showed that, with probability one :

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{g_n \in \Theta_n} \inf_{f \in \mathcal{S}_1} \|g_n - f\| &= 0, \\ \lim_{n \rightarrow \infty} \sup_{f \in \mathcal{S}_1} \inf_{g_n \in \Theta_n} \|g_n - f\| &= 0, \end{aligned} \tag{1.5}$$

where  $a_n$  is a sequence of constants fulfilling  $a_n \downarrow 0$ ,  $na_n \uparrow \infty$ ,  $na_n/\log n \rightarrow \infty$ ,  $\log(1/a_n)/\log_2 n \rightarrow \infty$ . Berthet [2] refined (1.5) under slightly stronger conditions imposed upon  $a_n$ . Making use of sharp upper bounds for Gaussian measures due to Talagrand [22], he proved that for any  $\epsilon > \epsilon_0$  (where  $\epsilon_0$  is a universal constant), we have almost surely for all  $n$  large enough :

$$\Theta_n \subset \mathcal{S}_1 + \epsilon \log(1/a_n)^{-2/3} \mathcal{B}_0. \tag{1.6}$$

Here  $\mathcal{B}_0 := \{f \in B[0, 1] : \|f\| \leq 1\}$ . The first aim of the present article is to show that the techniques employed in the just-mentioned result can be adapted to some other random objects than that used for that given in (1.6) (see Theorems 1 and 2 in the sequel). Results of this kind are usually called *clustering rates*. Another related problem is finding rates of convergence of such random sequences to a specified function belonging to the cluster set. Such results are known under the name of functional *Chung-type limit laws*. We now focus on the local empirical process  $\alpha_n(a_n \cdot)$ , where  $a_n \downarrow 0$  as  $n \rightarrow \infty$ . The works of Csáki

[5], de Acosta [1], Grill [12], Gorn and Lifshits [11], and Berthet and Lifshits [4] on small ball probabilities for Wiener processes provide some crucial tools to establish such limit laws for  $(\alpha_n(a_n \cdot))_{n \geq 1}$ , as these are expected to asymptotically mimic their gaussian analogues (see Mason [17]). Along this line, Deheuvels [6] established Chung-type limit laws for  $(\alpha_n(a_n \cdot))_{n \geq 1}$ , by showing that, if  $a_n$  is a sequence of constants satisfying  $na_n \uparrow \infty$ ,  $a_n \downarrow 0$  and  $na_n/(\log_2 n)^3 \rightarrow \infty$ , we have, almost surely, for each  $f \in \mathcal{S}_1$  satisfying  $\|f\|_H^2 := \int_0^1 f'^2(t)dt < 1$ :

$$\liminf_{n \rightarrow \infty} (\log_2 n) \left\| \frac{\alpha_n(a_n \cdot)}{\sqrt{a_n b_n}} - f \right\| = \frac{\pi}{4\sqrt{1 - \|f\|_H^2}}.$$

The proof of this theorem relies on strong approximation methods in combination with the results of de Acosta [1]. The latter provides useful exponential bounds for

$$\mathbb{P}\left(\left\| \frac{W}{T} - f \right\| \leq \epsilon\right),$$

with a small  $\epsilon > 0$  and a large  $T$ . Here,  $W$  is a Wiener process on  $[0, 1]$  and  $f$  satisfies  $\|f\|_H^2 < 1$ . The study of related probabilities when  $\|f\|_H = 1$  has required different arguments. In [12], rough estimates are given. In [11] and [4], some exact rates are given, but only for functions with first derivatives having a variation either bounded or locally infinite. The sets of all functions of this type are called  $\mathcal{S}_1^{bv}$  and  $\mathcal{S}_1^{liv}$  respectively. In the present paper, we shall make use of the latter results to extend the work of Deheuvels [6] to the case where  $f \in \mathcal{S}_1^{bv} \cup \mathcal{S}_1^{liv}$ . The remainder of our paper is organized as follows. Our main results are stated in §2, Theorems 1, 2 and 3. In §3, the proofs of these theorems are provided.



## 2. Main Results

Our first result gives clustering rates in Finkelstein's FLIL [10].

**Theorem 1.** *There exists a universal constant  $\epsilon_0 > 0$  such that, for any choice of  $\epsilon > \epsilon_0$  we have almost surely, for all large  $n$*

$$\frac{\alpha_n}{(2 \log_2 n)^{1/2}} \in \mathcal{S}_2 + \epsilon (\log_2 n)^{-2/3} \mathcal{B}_0, \quad (2.1)$$

$$\frac{\beta_n}{(2 \log_2 n)^{1/2}} \in \mathcal{S}_2 + \epsilon (\log_2 n)^{-2/3} \mathcal{B}_0. \quad (2.2)$$

**Remark 2.0.1.** *The uniform Bahadur-Kiefer representation (see [13]) asserts that, almost surely :*

$$\limsup_{n \rightarrow \infty} n^{1/4} (\log n)^{-1/2} (\log_2 n)^{-1/4} \|\alpha_n + \beta_n\| = 2^{-1/4},$$

from where (2.2) is readily implied by (2.1).

Our second theorem concerns the FLIL for local increments of the empirical process.

**Theorem 2.** *Let  $a_n$  be positive real numbers satisfying, as  $n \rightarrow \infty$ ,*

$$na_n \uparrow \infty, \quad \frac{na_n}{(\log_2 n)^{7/3}} \rightarrow \infty, \quad a_n \downarrow 0. \quad (2.3)$$

*Then there exists a universal constant  $\epsilon_1 > 0$  such that, for any choice of  $\epsilon > \epsilon_1$  we have almost surely, for all large  $n$ ,*

$$\frac{\alpha_n(a_n \cdot)}{\sqrt{2a_n \log_2 n}} \in \mathcal{S}_1 + \epsilon (\log_2 n)^{-2/3} \mathcal{B}_0. \quad (2.4)$$

*If moreover  $na_n/(\log_2 n)^{11/3} \rightarrow \infty$  then we have, almost surely, ultimately as  $n \rightarrow \infty$ ,*

$$\frac{\beta_n(a_n \cdot)}{\sqrt{2a_n \log_2 n}} \in \mathcal{S}_1 + \epsilon (\log_2 n)^{-2/3} \mathcal{B}_0. \quad (2.5)$$

**Remark 2.0.2.** We shall use the fact (see e.g. [9], Theorem 5) that, under (2.3), we have almost surely

$$\limsup_{n \rightarrow \infty} (n/a_n)^{1/4} (\log_2 n)^{-1/4} (2 \log_2 n + \log(na_n))^{-1/2} \|\alpha_n(a_n \cdot) + \beta_n(a_n \cdot)\| \leq 2^{-1/4},$$

from where (2.5) is implied by (2.4) after straightforward computations.

In order to state our last result, we need to give some definitions. Recall that  $f \in \mathcal{S}_1^{bv}$  whenever  $f'$  has a derivative with bounded variation and  $\int_0^1 f'^2(t) dt = 1$ . Results on small ball probabilities for a Wiener process when  $f \in \mathcal{S}_1^{bv}$  have been established by Gorn and Lifshits [11]. For such a function  $f$ , we shall write  $\nabla_f(L) := L^{2/3}$ ,  $L > 0$  and we denote by  $\chi_f$  the constant which is the unique solution of equation (3.1) in [11] (we refer to the just mentioned paper for more details). The case where  $f \in \mathcal{S}_1^{liv}$  (i.e. where  $\int_0^1 f'^2(t) dt = 1$  and the derivative of  $f'$  admits a version with locally infinite variation) has been treated by Berthet and Lifshits [4]. For such a function  $f$ , we set  $\chi_f := 1$  and we denote by  $\nabla_f(L)$  the unique solution of equation (2.1) in [3]. Our third result is stated as follows.

**Theorem 3.** Let  $f \in \mathcal{S}_1^{bv} \cup \mathcal{S}_1^{liv}$  be arbitrary and let  $a_n$  be a sequence of real numbers satisfying, as  $n \rightarrow \infty$ ,

$$na_n \uparrow \infty, a_n \downarrow 0, a_n \log_2 n \rightarrow 0, \quad (2.6)$$

$$\lim_{n \rightarrow \infty} \frac{na_n}{\log_2 n \nabla_f^2(\log_2 n)} = \infty. \quad (2.7)$$

Then we have, almost surely :

$$\liminf_{n \rightarrow \infty} \nabla_f(\log_2 n) \left\| \frac{\alpha_n(a_n \cdot)}{\sqrt{2a_n \log_2 n}} - f \right\| = \chi_f.$$

**Remark 2.0.3.** *The conditions (2.6) and (2.7) imposed upon  $a_n$  turn out to be the best possible with respect to the methods used in the proof of Theorem 3. The latter combines poissonization techniques with strong approximation arguments. Deheuvels and Lifshits [7] and Shmileva [19] have provided new tools to estimate probabilities of shifted small balls for a Poisson process without making use of strong approximation techniques. These results show up to be powerful enough to investigate Chung-Mogulskii limit laws for  $\alpha_n(a_n \cdot)$  without making use of strong approximation techniques, and thus relaxing condition (2.6). However, the just-mentioned results do not cover the case where  $f \in \mathcal{S}_1^{liv}$ .*

### 3. Proofs

#### 3.1. Proof of Theorem 1

Select an  $\epsilon > 0$  and consider the sequence  $\epsilon_n := \epsilon(\log_2 n)^{-2/3}$ . The main tool to achieve our goal is the exponential inequality stated in the following fact, which follows directly from Talagrand [21]. Recall that  $\mathcal{B}_0$  is the unit ball for  $\|\cdot\|$ .

**Fact 3.1.** *Let  $B$  be a Brownian bridge. There exists three constants  $K$ ,  $L_0$  and  $u_0 > 0$  such that, for any  $0 < u < u_0$  and  $c > 0$ , we have :*

$$P(B \notin c\mathcal{S}_2 + u\mathcal{B}_0) \leq K \exp\left(\frac{L_0}{u^2} - \frac{cu}{2} - \frac{c^2}{2}\right). \quad (3.1)$$

Let  $W$  be a Wiener process on  $[0, 1]$ . There exist two constants  $u_1$  and  $L_1$  such that, for any  $0 < u < u_1$  and  $c > 0$ , we have

$$P(W \notin c\mathcal{S}_1 + u\mathcal{B}_0) \leq \exp\left(\frac{L_1}{u^2} - \frac{cu}{2} - \frac{c^2}{2}\right). \quad (3.2)$$



In the proof of Theorem 1, we will make use of blocking techniques (see, e.g., [8] and [2]). For any real number  $a$ , set  $[a]$  as the unique integer  $q$  fulfilling  $q \leq a < q + 1$ , and set

$$n_k := \left[ \exp \left( k \exp \left( -(\log k)^{1/6} \right) \right) \right], \quad k \geq 1.$$

Set  $N_k := \{n_k, \dots, n_{k+1} - 1\}$  for  $k \geq 5$ . Given an integer  $n \geq 1$ , we set  $k(n)$  as the unique integer  $k$  such that  $n \in N_k$ . We shall first study the following sequence of functions

$$g_n := (n_{k+1})^{-1/2} b_{n_{k+1}}^{-1} H_n, \quad k = k(n),$$

with  $H_n(t) := n(F_n(t) - t)$  and  $b_n := (2 \log_2 n)^{1/2}$ . Let  $p_1$  and  $q_1$  be two conjugates numbers (such that  $1/p_1 + 1/q_1 = 1$ ) with  $1 < p_1 < \infty$ . Set, for  $k \geq 1$ ,

$$m_{p_1, k} := \min_{n \in N_k} \mathbb{P} \left( \frac{1}{(n_{k+1})^{1/2} b_{n_{k+1}}} \| H_{n_{k+1}} - H_n \| \leq \frac{1}{p_1} \epsilon_{n_{k+1}} \right).$$

A standard blocking argument based upon Ottaviani's inequality (see, e.g., [8], Lemma 3.4) yields

$$\begin{aligned} & \mathbb{P} \left( \bigcup_{n \in N_k} \left\{ \frac{1}{(n_{k+1})^{1/2} b_{n_{k+1}}} H_n \notin \mathcal{S}_2 + \epsilon_{n_{k+1}} \mathcal{B}_0 \right\} \right) \\ & \leq \frac{1}{m_{p_1, k}} \mathbb{P} \left( \frac{1}{(n_{k+1})^{1/2} b_{n_{k+1}}} H_{n_{k+1}} \notin \mathcal{S}_2 + \frac{1}{q_1} \epsilon_{n_{k+1}} \mathcal{B}_0 \right). \end{aligned}$$

Let  $k$  be integer and select  $n \in N_k$ . By the Dvoretzky-Kiefer-Wolfowitz inequality

ity (see, e.g., [23]) we have :

$$\begin{aligned} & \mathbb{P}\left(\frac{1}{(n_{k+1})^{1/2}b_{n_{k+1}}}\|H_{n_{k+1}} - H_n\| \geq \frac{1}{p_1}\epsilon_{n_{k+1}}\right) \\ & \leq \mathbb{P}\left(\|\alpha_{n_{k+1}-n}\| \geq \frac{1}{p_1}\epsilon_{n_{k+1}}\left(\frac{1}{1-\frac{n_k}{n_{k+1}}}\right)^{1/2}b_{n_{k+1}}\right) \\ & \leq 3 \exp\left(-\frac{4\epsilon^2 \log_2(n_k)^{-1/3}}{p_1^2\left(1-\frac{n_k}{n_{k+1}}\right)}\right) \text{ for large enough } k, \end{aligned}$$

whence  $m_{p_1,k} \geq 1/2$  for all large  $k$  by routine analysis. Now let  $p_2, q_2 > 1$  be two conjugate numbers. For  $k \geq 1$  we have,  $B_{n_{k+1}}$  denoting a Brownian bridge,

$$\begin{aligned} & \mathbb{P}\left(\frac{\alpha_{n_{k+1}}}{b_{n_{k+1}}} \notin \mathcal{S}_2 + \frac{1}{q_1}\epsilon_{n_{k+1}}\mathcal{B}_0\right) \leq \mathbb{P}\left(\|\alpha_{n_{k+1}} - B_{n_{k+1}}\| \geq \frac{1}{p_2q_1}\epsilon_{n_{k+1}}b_{n_{k+1}}\right) \\ & \quad + \mathbb{P}\left(B_{n_{k+1}} \notin b_{n_{k+1}}\mathcal{S}_2 + \frac{1}{q_2q_1}\epsilon_{n_{k+1}}b_{n_{k+1}}\mathcal{B}_0\right) \\ & := \mathbb{P}_k^{KMT} + \mathbb{P}_k^{Tal}. \end{aligned}$$

Making use of the Komlós-Major-Tusnàdy approximation (see, e.g., [14]), we can choose a sequence  $(B_{n_k})_{k \geq 1}$  satisfying, for some universal constants  $C_2, C_3$  and for all  $k$  large enough,

$$\mathbb{P}_k^{KMT} \leq C_2 \exp\left(-C_3(n_{k+1})^{1/2}\frac{1}{2p_2q_1}\epsilon_{n_{k+1}}b_{n_{k+1}}\right).$$

On the other hand, by applying assertion (3.1) of Fact 3.1 we have, for all large  $k$ ,

$$\mathbb{P}_k^{Tal} \leq K \exp\left[-\left(\frac{\epsilon}{q_1q_2} - \frac{L_0(q_1q_2)^2}{2\epsilon^2}\right)(\log_2 n_{k+1})^{1/3} - \log_2 n_{k+1}\right].$$

Routine analysis shows that both  $\mathbb{P}_k^{KMT}$  and  $\mathbb{P}_k^{Tal}$  are sumable in  $k$  for any choice of  $\epsilon > (L_0/2)^{1/3} =: \epsilon_0$ , provided that  $q_1, q_2$  are chosen close enough to 1.

Now an application of (1.1) in combination with elementary properties of the

sequence  $(n_k)_{k \geq 1}$  shows that, almost surely, as  $n \rightarrow \infty$ ,

$$\|g_n - b_n^{-1} \alpha_n\| = o((\log_2 n)^{-2/3}).$$

### 3.2. Proof of Theorem 2

Recall that  $b_n := (2 \log_2 n)^{1/2}$ ,  $n \geq 1$ . Let  $p_1, q_1 > 1$  be two conjugate numbers.

Set, for  $k \geq 1$  :

$$m_{p_1, k} := \min_{n \in N_k} \mathbb{P} \left( \frac{1}{(n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}}} \|H_n(a_{n_{k+1}} \cdot) - H_{n_{k+1}}(a_{n_{k+1}} \cdot)\| \leq \frac{1}{p_1} \epsilon_{n_{k+1}} \right).$$

The same blocking argument as in §3.1 yields

$$\begin{aligned} & \mathbb{P} \left( \bigcup_{n \in N_k} \left\{ \frac{H_n(a_{n_{k+1}} \cdot)}{(n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}}} \notin \mathcal{S}_1 + \epsilon_{n_{k+1}} \mathcal{B}_0 \right\} \right) \\ & \leq \frac{1}{m_{p_1, k}} \mathbb{P} \left( \frac{H_{n_{k+1}}(a_{n_{k+1}} \cdot)}{(n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}}} \notin \mathcal{S}_1 + \frac{1}{q_1} \epsilon_{n_{k+1}} \mathcal{B}_0 \right). \end{aligned}$$

Now, for any integer  $k \geq 5$  and  $n \in N_k$ , we have

$$\begin{aligned} & \mathbb{P} \left( \frac{1}{\sqrt{n_{k+1}}} \|H_{n_{k+1}}(a_{n_{k+1}} \cdot) - H_n(a_{n_{k+1}} \cdot)\| \geq \frac{1}{p_1} \epsilon_{n_{k+1}} b_{n_{k+1}} \right) \\ & \leq \mathbb{P} \left( \sup_{t \leq a_{n_{k+1}}} \frac{|\alpha_{n_{k+1}-n}(t)|}{1-t} \geq \frac{1}{p_1} \epsilon_{n_{k+1}} b_{n_{k+1}} \left( \frac{n_{k+1} a_{n_{k+1}}}{n_{k+1} - n_k} \right)^{1/2} \right). \end{aligned}$$

It is well known (see, e.g., [20], Proposition 1, p. 133) that for each  $n$ , the process

$(1-t)^{-1} \alpha_n(t)$  is a martingale in  $t$ . The Doob-Kolmogorov inequality yields :

$$1 - m_{p_1, k} \leq \frac{p_1^2 (1 - a_{n_{k+1}}) (1 - \frac{n_k}{n_{k+1}}) (\log n_{k+1})^{1/3}}{2\epsilon^2}.$$

Hence for all  $k$  large enough we have  $m_{p_1, k} \geq 1/2$ . Now set for each integer

$n \geq 1$ ,

$$\tilde{\Pi}_n(t) := n^{-1/2} \left( \sum_{i=1}^{\eta_n} \mathbf{1}_{\{U_i \leq t\}} - t \right), \quad t \in [0, 1],$$

where  $\eta_n$  is a Poisson variable with expectation  $n$ , which is independent of  $(U_i)_{i \geq 1}$ . Let  $\tilde{\Pi}$  denote a standard centered Poisson process on  $\mathbb{R}^+$  and let  $W$  be a Wiener process that we assume to be constructed on the same underlying probability space as  $\tilde{\Pi}$ . Notice that  $\tilde{\Pi}_n(\cdot)$  and  $n^{-1/2}\tilde{\Pi}(n\cdot)$  are equal in distribution as processes on  $[0, 1]$ . Now let  $p_2, q_2 > 1$  be two conjugate numbers. By making use of Poissonization techniques (see, e.g., [8], Lemma 2.1 or [25], Proposition 2.1 for a more general form) we see that, for all sufficiently large  $k$  :

$$\begin{aligned} & \mathbb{P}\left(\frac{\alpha_{n_{k+1}}(a_{n_{k+1}} \cdot)}{a_{n_{k+1}}^{1/2} b_{n_{k+1}}} \notin \mathcal{S}_1 + \frac{1}{q_1} \epsilon_{n_{k+1}} \mathcal{B}_0\right) \\ & \leq 2\mathbb{P}\left(\frac{\tilde{\Pi}_{n_{k+1}}(a_{n_{k+1}} \cdot)}{a_{n_{k+1}}^{1/2} b_{n_{k+1}}} \notin \mathcal{S}_1 + \frac{1}{q_1} \epsilon_{n_{k+1}} \mathcal{B}_0\right) \\ & = 2\mathbb{P}\left(\frac{\tilde{\Pi}(n_{k+1} a_{n_{k+1}} \cdot)}{(2n_{k+1} a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \notin \mathcal{S}_1 + \frac{1}{q_1} \epsilon_{n_{k+1}} \mathcal{B}_0\right) \\ & \leq 2\mathbb{P}\left(\|W(n_{k+1} a_{n_{k+1}} \cdot) - \tilde{\Pi}(n_{k+1} a_{n_{k+1}} \cdot)\| \geq \frac{1}{q_1 p_2} (n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}} \epsilon_{n_{k+1}}\right) \\ & \quad + 2\mathbb{P}\left(\frac{W(n_{k+1} a_{n_{k+1}} \cdot)}{(n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}}} \notin \mathcal{S}_1 + \frac{1}{q_1 q_2} \epsilon_{n_{k+1}}\right) \\ & := \mathbb{P}_k^{KMT} + \mathbb{P}_k^{Tal}. \end{aligned}$$

Now, making use of the strong approximation theorem of Komlós-Major-Tusnády [15], we can assume that the process  $W$  involved in the former expression satisfies, for some universal constants  $C_1, C_2, C_3 > 0$ , and for all  $T > 0$ ,  $z > 0$ ,

$$\mathbb{P}(\| \tilde{\Pi}(T \cdot) - W(\cdot) \| \geq z + C_1 \log T) \leq C_2 \exp(-C_2 z). \quad (3.3)$$

Notice that, as  $k \rightarrow \infty$  :

$$\frac{(n_{k+1} a_{n_{k+1}})^{1/2} b_{n_{k+1}} \epsilon_{n_{k+1}}}{\log(n_{k+1} a_{n_{k+1}})} \rightarrow \infty.$$

Thus, we have, ultimately as  $k \rightarrow \infty$ ,

$$\mathbb{P}_k^{KMT} \leq C_2 \exp\left(-\frac{\epsilon C_3}{\sqrt{2}q_1 p_2} (n_{k+1} a_{n_{k+1}})^{1/2} (\log_2 n_{k+1})^{-1/6}\right). \quad (3.4)$$

Recalling the assumption  $na_n/(\log_2 n)^{7/3} \rightarrow \infty$  we see that  $\mathbb{P}_k^{KMT}$  is summable in  $k$ . Now, making use of assertion (3.2) of Fact 3.1 we have, for all large  $k$ ,

$$\begin{aligned} \mathbb{P}_k^{Tal} &= \mathbb{P}\left(W \notin b_{n_{k+1}} \mathcal{S}_1 + \frac{1}{q_1 q_2} \epsilon_{n_{k+1}} b_{n_{k+1}} \mathcal{B}_0\right) \\ &\leq \exp\left(-\left(\frac{\epsilon}{q_1 q_2} - \frac{L_1(q_1 q_2)^2}{2\epsilon^2}\right) (\log_2 n_{k+1})^{1/3} - \log_2 n_{k+1}\right). \end{aligned}$$

Now if  $\epsilon > (L_1/2)^{1/3} =: \epsilon_1$  and if  $q_1, q_2 > 1$  are chosen sufficiently small, then  $\mathbb{P}_k^{Tal}$  is summable in  $k$ . By the Borel-Cantelli lemma, we see that for any  $\epsilon > \epsilon_1$  we have almost surely, for all large  $n$ ,

$$g_n \in \mathcal{S}_1 + \epsilon_{n_{k+1}} \mathcal{B}_0,$$

where  $g_n := (n_{k+1} a_{n_{k+1}})^{-1/2} b_{n_{k+1}}^{-1} H_n(a_{n_{k+1}} \cdot)$ ,  $n \in N_k$ . To conclude the proof of Theorem 2, it remains to control the distance between  $a_n^{-1/2} b_n^{-1} \alpha_n(a_n \cdot)$  and  $g_n$ , which is the purpose of the following lemma.

**Lemma 3.1.** *We have almost surely :*

$$\limsup_{n \rightarrow \infty} (\log_2 n)^{2/3} \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} - g_n \right\| = 0.$$

**Proof :** Set  $\Gamma_n := 1 - (n/n_{k+1})^{1/2} (a_n/a_{n_{k+1}})^{1/2} (\log_2 n / \log_2 n_{k+1})^{1/2}$ . The



triangle inequality yields

$$\begin{aligned} \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} - g_n \right\| &\leq \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} \Gamma_n \right\| + \left\| \frac{H_n(a_n \cdot) - H_n(a_{n_{k+1}} \cdot)}{(2n_{k+1} a_{n_{k+1}} \log_2 a_{n_{k+1}})^{1/2}} \right\| \\ &:= A_n + B_n. \end{aligned} \tag{3.5}$$

Clearly we have, as  $k \rightarrow \infty$ ,

$$\max_{n \in N_k} (\log_2 n_{k+1})^{2/3} \Gamma_n \leq \left( 1 - \sqrt{\frac{n_k \log_2 n_k}{n_{k+1} \log_2 n_{k+1}}} \right) (\log_2 n_{k+1})^{2/3} \rightarrow 0.$$

Now, by applying (1.4) we have almost surely

$$\limsup_{n \rightarrow \infty} \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} \right\| = 1. \tag{3.6}$$

Obviously (3.6) implies that, almost surely :

$$\lim_{n \rightarrow \infty} (\log_2 n_{k+1})^{2/3} \max_{n \in N_k} A_n = 0.$$

We now focus on controlling  $B_n$ . Set  $\rho_k := a_{n_k}/a_{n_{k+1}}$  and notice that

$$\begin{aligned} &\mathbb{P} \left( \max_{n \in N_k} (\log_2 n)^{2/3} \left\| \frac{H_n(a_n \cdot) - H_n(a_{n_{k+1}} \cdot)}{(2n_{k+1} a_{n_{k+1}} \log_2 a_{n_{k+1}})^{1/2}} \right\| \geq \epsilon \right) \\ &\leq \mathbb{P} \left( \max_{n \in N_k} \sup_{1 \leq \rho \leq \rho_k, 0 \leq t \leq 1} (\log_2 n_{k+1})^{2/3} \left| \frac{\alpha_n(a_{n_{k+1}} \rho t) - \alpha_n(a_{n_{k+1}} t)}{(2a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \right| \geq \epsilon \right). \end{aligned}$$

Now consider the Banach space  $B([0, 1] \times [0, 2])$  of all real bounded functions on  $[0, 1] \times [0, 2]$ , endowed with the usual sup norm  $\| \cdot \|_{[0,1] \times [0,2]}$ . We shall now make use of the powerful maximal inequality of Montgomery-Smith. For fixed  $k \geq 1$ , we apply the just mentioned inequality to the finite sequence  $(X_i)_{i \in N_k}$ , with  $X_i(t, \rho) := \mathbb{1}_{[t, \rho t]}(U_i) - \rho t$ ,  $t \in [0, 1]$ ,  $\rho \in [1, \rho_k]$ ,  $\rho t \leq 1$  and  $X_i(t, \rho) = 0$  elsewhere. Hence, by a combination of Theorem 1 and Corollary 3 in [18], we

have :

$$\begin{aligned}
& \mathbb{P} \left( \max_{n \in N_k} \sup_{1 \leq \rho \leq \rho_k, 0 \leq t \leq 1} (\log_2 n_{k+1})^{2/3} \left| \frac{\alpha_n(a_{n_{k+1}} \rho t) - \alpha_n(a_{n_{k+1}} t)}{(2a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \right| \geq \epsilon \right) \\
& \leq 9 \mathbb{P} \left( \sup_{1 \leq \rho \leq \rho_k, 0 \leq t \leq 1} (\log_2 n_{k+1})^{2/3} \left| \frac{\alpha_{n_{k+1}}(a_{n_{k+1}} \rho t) - \alpha_{n_{k+1}}(a_{n_{k+1}} t)}{(2a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \right| \geq \epsilon/30 \right) \\
& \leq 18 \mathbb{P} \left( \left\| \tilde{\Pi}(n_{k+1} a_{n_k} \cdot) - W(n_{k+1} a_{n_k} \cdot) \right\| \geq \frac{\epsilon}{240} \frac{(2n_{k+1} a_{n_{k+1}} \log_2 n_{k+1})^{1/2}}{(\log_2(n_{k+1}))^{2/3}} \right) \\
& \quad + 18 \mathbb{P} \left( \left\| \frac{W(\rho_k \cdot) - W(\cdot)}{(2 \log_2 n_{k+1})^{1/2}} \right\| \geq \frac{\epsilon}{120 (\log_2 n_{k+1})^{2/3}} \right). \tag{3.7}
\end{aligned}$$

In the last expression (which is the combination of usual poissonization techniques with the triangular inequality),  $\tilde{\Pi}$  and  $W$  denote respectively a centered Poisson process and a Wiener process based on the same underlying probability space. By the Komlós-Major-Tusnàdy construction (see [15]),  $W$  can be constructed to satisfy (3.3). By making use of the same arguments as those invoked to obtain (3.4), we conclude that the first term in (3.7) is summable in  $k$ . To control the second term in (3.7), we shall make use of a well known inequality (see, e.g., [20], p. 536), with  $a := \rho_k - 1$ ,  $\lambda := (\rho_k - 1)^{-1/2} (\log_2 n_{k+1})^{-1/6} (\sqrt{2}\epsilon/120)$  and  $\delta := 1/2$ , to get

$$\begin{aligned}
& \mathbb{P} \left( \left\| \frac{W(\rho_k \cdot) - W(\cdot)}{\sqrt{2 \log_2 n_{k+1}}} \right\| \geq \frac{\epsilon}{120 (\log_2 n_{k+1})^{2/3}} \right) \\
& \leq \frac{30720}{\sqrt{2}\epsilon} (\rho_k - 1)^{-1/2} (\log_2 n_{k+1})^{1/6} \exp \left( - \frac{\epsilon^2}{19200} (\rho_k - 1)^{-1} (\log_2 n_{k+1})^{-1/3} \right).
\end{aligned}$$

This expression is summable in  $k$ , and hence  $\max_{n \in N_k} B_n \rightarrow 0$  almost surely as  $k \rightarrow \infty$ .  $\square$

### 3.3. Proof of Theorem 3

Recall that  $\chi_f$ ,  $\nabla_f$ ,  $\mathcal{S}_1^{bv}$  and  $\mathcal{S}_1^{liv}$  are defined in §2. The main tool to achieve the proof of Theorem 3 is the following inequality (see Berthet [3]), which sums up different results from Gorn and Lifshits [11], Berthet and Lifshits [4] and Grill [12] (see also de Acosta [1]).

**Inequality 3.1.** *For any  $f \in \mathcal{S}_1^{BV} \cup \mathcal{S}_1^{LIV}$  and  $\delta > 0$ , there exist  $\gamma^+ = \gamma^+(\delta, f) > 0$  and  $\gamma^- = \gamma^-(\delta, f) > 0$  such that for  $T$  sufficiently large :*

$$\begin{aligned} \mathbb{P}\left(\nabla_f\left(\frac{T^2}{2}\right)\left\|\frac{W}{T} - f\right\| \leq (1 + \delta)\chi_f\right) &\geq \exp\left(-\frac{T^2}{2} + \gamma^+ \frac{\nabla_f^2(T^2/2)}{T^2}\right), \\ \mathbb{P}\left(\nabla_f\left(\frac{T^2}{2}\right)\left\|\frac{W}{T} - f\right\| \leq (1 - \delta)\chi_f\right) &\leq \exp\left(-\frac{T^2}{2} - \gamma^- \frac{\nabla_f^2(T^2/2)}{T^2}\right). \end{aligned}$$

Select  $f \in \mathcal{S}_1^{BV} \cup \mathcal{S}_1^{LIV}$ . We remind the two following properties of  $\nabla_f$  (see [16]), namely  $\limsup_{L \rightarrow \infty} L^{-1}\nabla_f(L) < \infty$  and  $\liminf_{L \rightarrow \infty} L^{-2/3}\nabla_f(L) > 0$ .

We shall first show that, almost surely :

$$\liminf_{n \rightarrow \infty} \nabla_f(\log_2 n) \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} - f \right\| \geq \chi_f.$$

Let us fix  $\epsilon > 0$ . We start by applying poissonization techniques in combination with the Komlós-Major-Tusnàdy approximation.

$$\begin{aligned} &\mathbb{P}\left(\nabla_f(\log_2 n) \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} - f \right\| \leq \chi_f(1 - 2\epsilon)\right) \\ &\leq 2\mathbb{P}\left(\nabla_f(\log_2 n) \left\| \frac{W(na_n \cdot)}{(2na_n \log_2 n)^{1/2}} - f \right\| \leq \chi_f(1 - \epsilon)\right) \\ &\quad + 2\mathbb{P}\left(\left\| W(na_n \cdot) - \tilde{\Pi}(na_n \cdot) \right\| \geq \frac{\chi_f \epsilon (2na_n \log_2 n)^{1/2}}{\nabla_f(\log_2 n)}\right). \end{aligned}$$

These two terms are sumable along the subsequence  $n_k$ , the second term being controlled by the Komlós-Major-Tusnàdy approximation while the first one is

controlled by Inequality 3.1. Now the control between  $n_k$  and  $n_{k+1}$  follows the same line as in Lemma 3.1. We omit details for sake of brevity.

We now focus on showing that, almost surely,

$$\liminf_{n \rightarrow \infty} \nabla_f(\log_2 n) \left\| \frac{\alpha_n(a_n \cdot)}{(2a_n \log_2 n)^{1/2}} - f \right\| \leq \chi_f.$$

Set  $n_k := k^{2^k}$ ,  $v_k := n_{k+1} - n_k$  and

$$h_k := \frac{\sqrt{n_{k+1}} \alpha_{n_{k+1}}(a_{n_{k+1}} \cdot) - \sqrt{n_k} \alpha_{n_k}(a_{n_{k+1}} \cdot)}{\sqrt{2v_k a_{n_{k+1}} \log_2(v_k)}}. \quad (3.8)$$

Notice that the  $h_k$  are mutually independent processes, and that each  $h_k$  is distributed like  $(2a_{n_{k+1}} \log_2 v_k)^{-1/2} \alpha_{v_k}(a_{n_{k+1}} \cdot)$ . We now make use of the following "depoissonization" lemma. Recall that  $\tilde{\Pi}(\cdot)$  denotes a centered standard Poisson process on  $[0, \infty)$ .

**Lemma 3.2.** *Under assumptions (2.6) and (2.7), there exist two sumable positive sequences  $(c_k)_{k \geq 1}$ ,  $(b_k)_{k \geq 1}$  and an integer  $k_0 \geq 1$  such that, for any set  $A \subset B([0, 1])$  that is measurable for both empirical and Poisson processes and for all  $k \geq k_0$ ,*

$$\mathbb{P}\left(\tilde{\Pi}(v_k a_{n_{k+1}} \cdot) \in A\right) - c_k - b_k \leq 2\mathbb{P}\left(v_k^{1/2} \alpha_{v_k}(a_{n_{k+1}} \cdot) \in A\right).$$

**Proof :** Set  $u_k := (4 \log_2(v_k)/n_{k+1} a_{n_{k+1}})^{1/2}$ . By assumption (2.7) we have  $u_k \rightarrow 0$  as  $k \rightarrow \infty$ . Now set  $\Pi(t) := \tilde{\Pi}(t) + t$ ,  $\in [0, 1]$  and  $R_{1,k} := \Pi(v_k a_{n_{k+1}})$ ,  $R_{2,k} := \Pi(v_k) - \Pi(v_k a_{n_{k+1}})$ . For fixed  $k$ ,  $R_{1,k}$  and  $R_{2,k}$  are independent random variables and are distributed as Poisson variables with respective expectations  $v_k a_{n_{k+1}}$  and  $v_k(1 - a_{n_{k+1}})$ . Let  $A \subset B([0, 1])$  be an arbitrary set that we assume

to be measurable for  $\tilde{\Pi}$  and  $\alpha_n$ . Define the following events :

$$E_k := \left\{ \tilde{\Pi}(v_k a_{n_{k+1}} \cdot) \in A \right\}, \quad k \geq 1.$$

We have, for any integer  $k \geq 1$ ,

$$\begin{aligned} \mathbb{P}(E_k) &\leq \mathbb{P}\left(E_k \cap R_{1,k} \in [(1-u_k)v_k a_{n_{k+1}}, (1+u_k)v_k a_{n_{k+1}}]\right) \\ &\quad + \mathbb{P}\left(R_{1,k} < (1-u_k)v_k a_{n_{k+1}}\right) + \mathbb{P}\left(R_{1,k} > (1+u_k)v_k a_{n_{k+1}}\right). \end{aligned}$$

Denote by  $c_k$  and  $b_k$  the two last terms of the RHS of the preceding inequality. We shall show that these two sequences have finite sums. Making use of Chernoff's inequality, we have :

$$c_k \leq \exp\left(-v_k a_{n_{k+1}} \left((1+u_k) \log(1+u_k) - u_k\right)\right).$$

Since  $(1+u) \log(1+u) - u \sim \frac{u^2}{2}$  as  $u \rightarrow 0$ , it follows that for all large  $k$ ,

$$\begin{aligned} c_k &\leq \exp\left(-v_k a_{n_{k+1}} \frac{u_k^2}{2}\right) \\ &= \exp\left(-2 \log_2 n_k\right). \end{aligned}$$

We make use of a similar method to show that  $(b_k)_{k \geq 1}$  is sumable. It remains to show that, for all  $k \geq k_0$  (with  $k_0$  independent of  $A$ ), we have

$$\mathbb{P}\left(E_k \cap R_{1,k} \in [(1-u_k)v_k a_{n_{k+1}}, (1+u_k)v_k a_{n_{k+1}}]\right) \leq 2\mathbb{P}\left(E_k \mid \Pi(v_k) = v_k\right).$$

Now set

$$K_k := \inf \left\{ \frac{\mathbb{P}(R_{2,k} = v_k - j)}{\mathbb{P}(\Pi(v_k) = v_k)} j \in [(1-u_k)v_k a_{n_{k+1}}, (1+u_k)v_k a_{n_{k+1}}] \right\}. \quad (3.9)$$



We have

$$\begin{aligned}
& \mathbb{P}\left(E_k \cap R_{1,k} \in [(1-u_k)v_k a_{n_{k+1}}, (1+u_k)v_k a_{n_{k+1}}]\right) \\
& \leq \sum_{j=[(1-u_k)v_k a_{n_{k+1}}]^{+1}}^{[(1+u_k)v_k a_{n_{k+1}}]+1} \mathbb{P}(E_k \cap R_{1,k} = j) \\
& \leq K_k^{-1} \sum_{j=[(1-u_k)v_k a_{n_{k+1}}]^{+1}}^{[(1+u_k)v_k a_{n_{k+1}}]+1} \mathbb{P}(E_k \cap R_{1,k} = j) \frac{\mathbb{P}(R_{2,k} = v_k - j)}{\mathbb{P}(\Pi(v_k) = v_k)} \\
& = K_k^{-1} \sum_{j=[(1-u_k)v_k a_{n_{k+1}}]^{+1}}^{[(1+u_k)v_k a_{n_{k+1}}]+1} \frac{\mathbb{P}(E_k \cap R_{1,k} = j \cap R_{2,k} = v_k - j)}{\mathbb{P}(\Pi(v_k) = v_k)} \\
& \leq K_k^{-1} \mathbb{P}(E_k \mid \Pi(v_k) = v_k).
\end{aligned}$$

Hence, it suffices to show that  $K_k \rightarrow 1$ . For clarity of notations, set  $v'_k := v_k(1-a_{n_{k+1}})$ , recalling that  $R_{2,k}$  is a Poisson variable with parameter  $v'_k$ . Setting  $l = v_k - j$  in (3.9) we have, as  $k \rightarrow \infty$ ,

$$K_k = \inf \left\{ \frac{\mathbb{P}(R_{2,k} = l)}{\mathbb{P}(\Pi(v_k) = v_k)}, l \in [v'_k - v_k u_k a_{n_{k+1}}, v'_k + v_k u_k a_{n_{k+1}}] \right\}.$$

Now, by Stirling's formula, we have  $\mathbb{P}(\Pi(v_k) = v_k) \sim (2\pi v_k)^{-1/2}$  as  $k \rightarrow \infty$ . A routine study of the finite sequence

$$\left( \mathbb{P}(R_{2,k} = l), l \in [v'_k - v_k u_k a_{n_{k+1}}, v'_k + v_k u_k a_{n_{k+1}}] \right)$$

shows that

$$\mathbb{P}(\Pi(v_k) = v_k) K_k = \min(\mathbb{P}_{1,k}, \mathbb{P}_{2,k}), \text{ where}$$

$$\mathbb{P}_{1,k} := \mathbb{P}(R_{2,k} = [v'_k - v_k a_{n_{k+1}} u_k]), \text{ and}$$

$$\mathbb{P}_{2,k} := \mathbb{P}(R_{2,k} = [v'_k + v_k a_{n_{k+1}} u_k] + 1).$$

We set  $u'_k = a_{n_{k+1}} u_k v_k / v'_k \sim u_k a_{n_{k+1}}$ . Stirling's formula yields, ultimately as  $k \rightarrow \infty$ ,

$$\begin{aligned} \mathbb{P}_{1,k} &= \frac{v_k^{[v'_k - v_k a_{n_{k+1}} u_k]}}{[v'_k - v_k a_{n_{k+1}} u_k]!} \exp(-v'_k) \\ &\sim \left( \frac{v'_k}{v'_k (1 - u'_k)} \right)^{v'_k (1 - u'_k)} \frac{\exp(-v'_k)}{\exp(-v'_k + v'_k u'_k)} \sqrt{2\pi v'_k} \\ &\sim \sqrt{2\pi v_k} (1 - u'_k)^{-v'_k (1 - u'_k)} \exp(-v'_k u'_k) \\ &= \sqrt{2\pi v_k} \exp\left(-v'_k ((1 - u'_k) \log(1 - u_k) + u'_k)\right). \end{aligned}$$

Moreover, since  $(1 - \epsilon) \log(1 - \epsilon) + \epsilon \sim \frac{\epsilon^2}{2}$  as  $\epsilon \rightarrow 0$ , we have, for all large  $k$ ,

$$\exp\left(-2a_{n_{k+1}} \log_2(n_{k+1})\right) \leq \exp\left(-v'_k ((1 - u'_k) \log(1 - u_k) + u'_k)\right).$$

By assumption (2.6) we have  $a_n \log_2 n \rightarrow 0$ , which ensures that  $\mathbb{P}_{1,k} \sim \sqrt{2\pi v_k}$ .

The control of  $\mathbb{P}_{2,k}$  is very similar. This achieves the proof of Lemma 3.2.  $\square$

We now apply the preceding lemma in conjunction with the Komlós-Major-Tusnàdy approximation. Let  $W$  be a Wiener process constructed on the same underlying probability space as  $\tilde{\Pi}$ . For an arbitrary  $\delta > 0$ , we have (recall that  $h_k$  has been defined in (3.8))

$$\begin{aligned} &\mathbb{P}\left(\nabla_f(\log_2(v_k)) \parallel h_k - f \parallel \leq (1 + 2\delta)\chi_f\right) \\ &\geq \frac{1}{2}\mathbb{P}\left(\nabla_f(\log_2(v_k)) \parallel \frac{\tilde{\Pi}(v_k a_{n_{k+1}} \cdot)}{(2v_k a_{n_{k+1}} \log_2 v_k)^{1/2}} - f \parallel \leq (1 + 2\delta)\chi_f\right) - \frac{1}{2}c_k - \frac{1}{2}b_k \\ &\geq -\frac{1}{2}\mathbb{P}\left(\left\|\tilde{\Pi}(v_k a_{n_{k+1}} \cdot) - W(v_k a_{n_{k+1}} \cdot)\right\| \geq \frac{\delta\chi_f(2v_k a_{n_{k+1}} \log_2 v_k)^{1/2}}{\nabla_f(\log_2 v_k)}\right) \\ &\quad + \frac{1}{2}\mathbb{P}\left(\nabla_f(\log_2 v_k) \parallel \frac{W}{(2\log_2 v_k)^{1/2}} - f \parallel \leq \chi_f(1 + \delta)\right) - \frac{1}{2}c_k - \frac{1}{2}b_k \\ &=: -d_k + \frac{1}{2}\mathbb{P}_k - \frac{1}{2}c_k - \frac{1}{2}b_k. \end{aligned}$$

Since  $v_k \sim n_{k+1}$  it is easy to conclude that  $d_k$  is summable in  $k$ , by making use of the strong approximation (see [15]). Hence, making use of Inequality 3.1, we have asymptotically

$$\mathbb{P}_k \geq \exp\left(-\log_2 v_k\right).$$

But  $\log_2 v_k = \log(k+1) + \log_2 k + o(k^{-2}(\log k)^{-1})$  and hence

$$\sum_{k \geq 1} \mathbb{P}\left(\nabla_f(\log_2(v_k)) \parallel h_k - f \parallel \leq (1 + 2\delta)\chi_f\right) = \infty.$$

Applying the second half of the Borel Cantelli lemma, we deduce that, almost surely :

$$\liminf_{k \rightarrow \infty} \nabla_f(\log_2 v_k) \parallel h_k - f \parallel \leq \chi_f.$$

To conclude the proof, it is enough to show that, almost surely (recall that  $\lim_{L \rightarrow \infty} L^{-1} \nabla_f(L) > 0$ ),

$$\lim_{k \rightarrow \infty} (\log_2 n_{k+1}) \left\| h_k - \frac{\alpha_{n_{k+1}}(a_{n_{k+1}} \cdot)}{(2n_{k+1} a_{n_{k+1}} \log_2(n_{k+1}))^{1/2}} \right\| = 0.$$

Routine algebra shows that

$$\begin{aligned} & (\log_2 n_{k+1}) \left\| h_k - \frac{\alpha_{n_{k+1}}(a_{n_{k+1}} \cdot)}{(2a_{n_{k+1}} \log_2(n_{k+1}))^{1/2}} \right\| \\ & \leq (\log_2 n_{k+1}) \left( \left( \frac{n_{k+1} \log_2 n_{k+1}}{v_k \log_2(v_k)} \right)^{1/2} - 1 \right) \left\| \frac{\alpha_{n_{k+1}}(a_{n_{k+1}} \cdot)}{(2a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \right\| \\ & \quad + (\log_2 n_{k+1}) \left\| \frac{n_k^{1/2} \alpha_{n_k}(a_{n_{k+1}} \cdot)}{(2v_k a_{n_{k+1}} \log_2 n_{k+1})^{1/2}} \right\| \\ & := A_k + B_k. \end{aligned}$$

Applying theorem of Mason (1.4) we get  $A_k \rightarrow 0$  almost surely as  $k \rightarrow \infty$ . We

now apply Doob's inequality for positive submartingales to obtain

$$\begin{aligned} & \mathbb{P}\left(\left|(\log_2 n_{k+1})\right| \left\| n_k \frac{\alpha_{n_k}(a_{n_{k+1}} \cdot)}{((2v_k a_{n_{k+1}} \log_2 n_{k+1}))^{1/2}} \right\| \geq \epsilon\right) \\ &= \mathbb{P}\left(\sup_{0 \leq t \leq a_{n_{k+1}}} \left| \frac{\alpha_n(t)}{1-t} \right| \geq \frac{\epsilon(2v_k a_{n_{k+1}} \log_2 n_{k+1})^{1/2}}{n_k^{1/2} \log_2 n_{k+1}}\right) \\ &\leq \frac{1}{2\epsilon^2} (1 - a_{n_{k+1}}) \log_2 n_{k+1} \frac{n_k}{v_k}. \end{aligned}$$

Since  $n_k/v_k \sim 1/e^2 k^2$  as  $k \rightarrow \infty$ , we conclude the proof of the lower bound in

Theorem 3 with the Borel-Cantelli lemma.  $\square$

**Acknowledgements:** The author would like to thank P. Berthet and E. Shmil'eva for many fruitful conversations and advices on the subject.

## References

- [1] ACOSTA (DE), A. (1983). Small deviations in the functional central limit theorem with applications to the functional laws of the iterated logarithm. *Ann. Probab.* **11**, 2, 78–101.
- [2] BERTHET, P. (1997). On the rate of clustering to the Strassen set for increments of the empirical process. *J. Theoret. Probab.* **10**, 3, 557–579.
- [3] BERTHET, P. (2005). Inner rates of coverage of Strassen type sets by increments of the uniform empirical and quantile processes. *Stochastic Process. Appl.* **115**, 3, 493–537.
- [4] BERTHET, P. AND LIFSHITS, M. (2002). Some exact rates in the functional law of the iterated logarithm. *Ann. Inst. H. Poincaré Probab. Statist.* **38**, 811–824.

- [5] CSÁKI, E. (1980). A relation between Chung's and Strassen's laws of the iterated logarithm. *Z. Wahrsch. Verw. Gebiete* **54**, 3, 287–301.
- [6] DEHEUVELS, P. (2000). Chung type functional laws of the iterated logarithm for tail empirical processes. *Ann. Inst. H. Poincaré Probab. Statist.* **36**, 583–616.
- [7] DEHEUVELS, P. AND LIFSHITS, M. (2003). Probabilities of hitting of shifted small balls by centered Poisson processes. *J. Math. Sci. (N. Y.)* **118**, 6, 5541–5554.
- [8] DEHEUVELS, P. AND MASON, D. (1992). Functional laws of the iterated logarithm for the increments of empirical and quantile processes. *Ann. Probab.* **20**, 1248–1287.
- [9] EINMAHL, J. AND MASON, D. (1988). Strong limit theorems for weighted quantile processes. *Ann. Probab.* **16**, 4, 1626.
- [10] FINKELSTEIN, H. (1971). The law of iterated logarithm for empirical distributions. *Ann. Math. Statist.* **42**, 607–625.
- [11] GORN, N. AND LIFSHITS, M. (1999). Chung's law and the Csàki function. *J. Theoret. Probab.* **12**, 399–420.
- [12] GRILL, K. (1991). A lim inf result in Strassen's law of the iterated logarithm. *Probab. Theory Related Fields* **89**, 149–157.
- [13] KIEFER, J. (1967). On Bahadur's representation of sample quantiles. *Ann. Math. Statist.* **38**, 5, 1323–1342.
- [14] KOMLÓS, J., MAJOR, P., AND TUSNÁDY, G. (1975). An approximation of partial sums of independent r.v.'s and the sample d.f.I. *Z. Wahrsch. Verw.*



- Gebiete* 32, 111–131.
- [15] KOMLÓS, J., MAJOR, P., AND TUSNÁDY, G. (1977). An approximation of partial sums of independent r.v.'s and the sample d.f.II. *Z. Wahrsch. Verw. Gebiete* 34, 33–58.
- [16] KUELBS, J., LI, W., AND TALAGRAND, M. (1994). Liminf results for Gaussian samples and Chung's functional LIL. *Ann. Probab.* 22, 1879–1903.
- [17] MASON, D. (1988). A strong invariance principle for the tail empirical process. *Ann. Inst. H. Poincaré Probab. Statist.* 24, 491–506.
- [18] MONTGOMMERY-SMITH, J. (1993). Comparison of sums of identically distributed random vectors. *Probab. Math. Statist.* 14, 281–285.
- [19] SHMILEVA, E. Y. (2003). Small ball probabilities for a centered poisson process of high intensity. (russian). *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 298, 280–303.
- [20] SHORACK, G. AND WELLNER, J. (1986). *Empirical Processes and applications to statistics*. Springer.
- [21] TALAGRAND, M. (1992). On the rate of clustering in the law of the iterated logarithm for Brownian motion. *Probab. Banach Spaces*.
- [22] TALAGRAND, M. (1993). New Gaussian estimates for enlarged balls. *Geom. Funct. Anal.* 3, 5, 503–525.
- [23] TALAGRAND, M. (1994). Sharper bounds for Gaussian and empirical processes. *Ann. Probab.* 22, 28–76.
- [24] VAN DER VAART, A. AND WELLNER, J. (1996). *Weak convergence and empirical processes*. Springer.

- [25] VARRON, D. (2011). Some new almost sure results on the functional increments of the uniform empirical process. *Stochastic Processes and their Applications* **121**, 2, 337 – 356. <http://www.sciencedirect.com/science/article/B6V1B-51GHX0D-3/2/f3edcacf7b9948cc2a1690ad768fbf22>.