



**HAL**  
open science

## Equitability, edge-injectivity, and the 1-2-3 Conjecture

Julien Bensmail, Mohammed Senhaji, Kasper Szabo Lyngsie

► **To cite this version:**

Julien Bensmail, Mohammed Senhaji, Kasper Szabo Lyngsie. Equitability, edge-injectivity, and the 1-2-3 Conjecture. 2016. hal-01361482v1

**HAL Id: hal-01361482**

**<https://hal.science/hal-01361482v1>**

Preprint submitted on 7 Sep 2016 (v1), last revised 7 Aug 2017 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Equitability, edge-injectivity, and the 1-2-3 Conjecture

Julien Bensmail<sup>a</sup>, Mohammed Senhaji<sup>b</sup>, Kasper Szabo Lyngsie<sup>c</sup>

<sup>a</sup>*INRIA and Université Nice-Sophia-Antipolis  
I3S, UMR 7271  
06900 Sophia-Antipolis, France*

<sup>b</sup>*Université de Bordeaux  
LaBRI, UMR5800  
F-33400 Talence, France*

<sup>c</sup>*Department of Applied Mathematics and Computer Science  
Technical University of Denmark  
DK-2800 Lyngby, Denmark*

---

## Abstract

This paper is dedicated to studying the following question: Is it always possible to injectively assign the weights  $1, \dots, |E(G)|$  to the edges of any given graph  $G$  (with no component isomorphic to  $K_2$ ), so that every two adjacent vertices of  $G$  get distinguished by their sums of incident weights? We answer positively to this question for some classes of graphs, such as trees and regular graphs, while, for some other classes of graphs, such as 2-degenerate graphs and graphs with maximum average degree at most 3, we prove that, provided we use a constant number of additional edge weights, the claimed assignment always exists. Our investigations here, are strongly related to several aspects of the well-known 1-2-3 Conjecture and the Antimagic Labelling Conjecture.

*Keywords:* 1-2-3 Conjecture, Antimagic Labelling Conjecture, equitable edge-weightings, neighbour-sum-distinguishing edge-weightings.

---

## 1. Introduction

In order to present our investigations in this paper, as well as our motivations, we first need to introduce a few particular graph concepts and notions. We refer the reader to textbooks on graph theory for more details on any standard notion or terminology not introduced herein.

Given a (undirected, simple, loopless) graph  $G$  and an edge-weighting  $w$  of  $G$ , one can compute, for every vertex  $v$  of  $G$ , the sum  $\sigma(v)$  (or  $\sigma_w(v)$ , when more precision is needed) of weights assigned, by  $w$ , to the edges incident to  $v$ . That is,

$$\sigma_w(v) := \sum_{u \in N(v)} w(vu)$$

for every vertex  $v$  of  $G$ . In case we have  $\sigma_w(u) \neq \sigma_w(v)$  for every edge  $uv$  of  $G$ , we call  $w$  *neighbour-sum-distinguishing*. It can be observed that every graph with no connected component isomorphic to  $K_2$  admits neighbour-sum-distinguishing edge-weightings using sufficiently large weights. In the context of the current investigations, when speaking of a *nice graph*, we mean a graph with no connected component isomorphic to  $K_2$ . For a nice graph  $G$ , it hence makes sense to study the smallest number of consecutive weights  $1, \dots, k$

used in a neighbour-sum-distinguishing edge-weighting of  $G$ . We denote this chromatic parameter by  $\chi_{\Sigma}^e(G)$ .

Throughout this paper, we deal with edge-weightings that are not only neighbour-sum-distinguishing, but also do not assign any edge weight more than once. We say that such an edge-weighting is *edge-injective*. Still under the assumption that  $G$  is a nice graph, we denote by  $\chi_{\Sigma}^{e,1}(G)$  the smallest  $k$  such that  $G$  admits an edge-injective neighbour-sum-distinguishing  $k$ -edge-weighting.

In this paper, we consider the following conjecture. Our motivations for studying this conjecture, as well as our evidences to suspect that it might be true, are described below.

**Conjecture 1.1.** *For every nice graph  $G$ , we have  $\chi_{\Sigma}^{e,1}(G) = |E(G)|$ .*

By the edge-injectivity property, we note that  $|E(G)|$  is a lower bound on  $\chi_{\Sigma}^{e,1}(G)$  for every nice graph  $G$ . Conjecture 1.1, in brief words, hence asks whether, for every nice graph  $G$ , we can bijectively assign weights  $1, \dots, |E(G)|$  to the edges of  $G$ , so that no two adjacent vertices of  $G$  get the same value of  $\sigma$ .

Conjecture 1.1 is related to the well-known **1-2-3 Conjecture**, raised in 2004 by Karoński, Łuczak and Thomason [6], which states the following.

**1-2-3 Conjecture.** *For every nice graph  $G$ , we have  $\chi_{\Sigma}^e(G) \leq 3$ .*

Many aspects of the 1-2-3 Conjecture have been studied in literature. For an overview of those considered aspects, we refer the interested reader to the wide survey [7] by Seamone, which is dedicated to this topic. Our investigations in this paper, are mostly related to a recent equitable variant of the 1-2-3 Conjecture that was considered by Baudon, Piłśniak, Przybyło, Senhaji, Sopena and Woźniak in [1]. In this variant, the authors studied, for some families of nice graphs, the existence of neighbour-sum-distinguishing edge-weightings being *equitable*, i.e. in which any two distinct edge weights are assigned about the same number of times (being equal, or differing by 1). In particular, they introduced and studied, for any given graph  $G$ , the chromatic parameter denoted by  $\overline{\chi}_{\Sigma}^e(G)$ , being the smallest maximal weight in an equitable neighbour-sum-distinguishing edge-weighting of  $G$ . In brief words, they proved that, at least for particular common classes of nice graphs (such as complete graphs and some bipartite graphs), the two parameters  $\chi_{\Sigma}^e$  and  $\overline{\chi}_{\Sigma}^e$  are generally equal, except for a few exceptions.

Despite their results, the authors of [1] did not dare addressing a general conjecture on how should  $\overline{\chi}_{\Sigma}^e$  behave in general, or compared to  $\chi_{\Sigma}^e$  for a given nice graph. In particular, it does not seem obvious how big  $\overline{\chi}_{\Sigma}^e$  can be, neither whether this parameter can be arbitrarily large. This is one of our motivations for studying edge-injective neighbour-sum-distinguishing edge-weightings, as an edge-injective edge-weighting is always equitable. Thus,  $\overline{\chi}_{\Sigma}^e(G) \leq \chi_{\Sigma}^{e,1}(G)$  holds for every nice graph  $G$ . Hence, attacking Conjecture 1.1 can be regarded as a way to get progress towards all those questions.

Our second motivation for considering Conjecture 1.1, is that edge-injective neighbour-sum-distinguishing edge-weightings can be regarded as a weaker notion of well-known **antimagic labellings**. Formally, using our own terminology, an *antimagic labelling*  $w$  of a graph  $G$ , is an edge-injective  $|E(G)|$ -edge-weighting of  $G$  for which  $\sigma_w$  is injective, i.e. all vertices of  $G$  get a distinct sum of incident weights by  $w$ . We say that  $G$  is *antimagic*, if it admits an antimagic labelling. Many lines of research concerning antimagic labellings can be found in literature, most of which are related to the following conjecture, addressed by Hartsfield and Ringel in [4].

**Antimagic Labelling Conjecture.** *Every nice connected graph is antimagic.*

Despite lots of efforts (refer to the dynamic survey [3] by Gallian for an in-depth summary of the vast and rich literature on this topic), the Antimagic Labelling Conjecture is still open in general, even for common classes of graphs, such as nice trees. Conjecture 1.1, which is clearly much weaker than the Antimagic Labelling Conjecture, as the distinction condition here only concerns the adjacent vertices, hence sounds as a much easier challenge to us, in particular concerning classes of nice graphs that are not known to be antimagic.

Hence, every antimagic graph  $G$  agrees with Conjecture 1.1, implying, as described earlier, that

$$\overline{\chi}_{\Sigma}^e(G) \leq \chi_{\Sigma}^{e,1}(G) = |E(G)|$$

holds, thus providing an upper bound on  $\overline{\chi}_{\Sigma}^e(G)$  for  $G$ . This is of interest, as several classes of graphs, such as nice regular graphs and nice complete partite graphs, are known to be antimagic, see [3]. Conversely, proving that a graph  $G$  verifies  $\chi_{\Sigma}^{e,1}(G) = |E(G)|$ , and agrees with Conjecture 1.1, is similar to proving that, in some sense,  $G$  is “locally antimagic”.

Conjecture 1.1 can essentially be considered as a combination of the 1-2-3 Conjecture and the Antimagic Labelling Conjecture, as the notions behind it have flavours of both conjectures. As described earlier, proving Conjecture 1.1 for some classes of graphs has, to some extent, consequences on the 1-2-3 Conjecture and the Antimagic Labelling Conjecture, or at least on variants of these conjectures.

Our work in this paper, is focused on both proving Conjecture 1.1 for particular classes of nice graphs, and providing upper bounds on  $\chi_{\Sigma}^{e,1}$  for some classes of nice graphs. This paper is organized as follows. Tools and preliminary results we use throughout, are introduced in Section 2. After that, we start off by providing support to Conjecture 1.1 in Section 3, essentially by showing and pointing out that the conjecture holds for some classes of graphs, such as nice trees and regular graphs. Towards Conjecture 1.1, we then provide, in Section 4, general weaker upper bounds on  $\chi_{\Sigma}^{e,1}$ . These bounds are then improved for some classes of nice sparse graphs in Section 5. These classes include nice graphs with maximum average degree at most 3, and nice 2-degenerate graphs. Concluding comments are gathered in Section 6.

## 2. Preliminary remarks and results

In this section, we introduce several observations that will be of some use in the next sections. The reader should keep in mind that, because we focus on  $k$ -edge-weightings throughout this paper, hence on edge-weightings assigning positive weights only, every edge weight considered here is implicitly assumed to be positive.

We start off by pointing out a few situations, in which, for a given edge  $uv$  of any graph  $G$ , we necessarily get  $\sigma(u) \neq \sigma(v)$  by an edge-injective edge-weighting of  $G$ . We omit a formal proof, as it is easily seen that these claims are true. We note that the third item is more general, as it implies the other two.

**Observation 2.1.** *Let  $G$  be a graph, and  $w$  be an edge-injective edge-weighting of  $G$ . Then, for every edge  $uv$  of  $G$ , we have  $\sigma(u) \neq \sigma(v)$  in any of the following situations:*

1.  $d(u) = 1$  and  $d(v) \geq 2$ ;
2.  $d(u) = d(v) = 2$ ;

3.  $d(u) \geq d(v)$  and

$$\min \{w(uv') : v' \in N(u) \setminus \{v\}\} \geq \max \{w(vu') : u' \in N(v) \setminus \{u\}\}.$$

We now observe that, to be able to successfully extend a partial neighbour-sum-distinguishing edge-weighting to an edge, we need to have sufficiently distinct weights in hand for that purpose.

**Observation 2.2.** *Let  $G$  be a graph, and  $w$  be a partial neighbour-sum-distinguishing edge-weighting of  $G$ . Let  $uv$  be a non-weighted edge of  $G$  such that  $uv$  is the only non-weighted edge incident to  $u$  or  $v$ . If we currently have  $\sigma(u) \neq \sigma(v)$  by  $w$ , then  $w$  can be successfully extended to  $uv$ , provided we have a set  $W$  of at least  $d(u) + d(v) - 1$  distinct weights that can be assigned to  $uv$ .*

*Proof.* We note that  $w$  currently must satisfy  $\sigma(u) \neq \sigma(v)$ , as, otherwise, no matter what weight we assign to  $uv$ , we would eventually get  $\sigma_w(u) = \sigma_w(v)$ . Under that assumption, we note that weighting  $uv$  with any weight completely determines the value of both  $\sigma_w(u)$  and  $\sigma_w(v)$ . The value of  $\sigma_w(u)$  eventually has to be different from the sums of weights incident to the  $d(u) - 1$  neighbours of  $u$  different from  $v$ . Similarly, the value of  $\sigma_w(v)$  eventually has to be different from the sums of weights incident to the  $d(v) - 1$  neighbours of  $v$  different from  $u$ . The neighbours of  $u$  and  $v$  hence forbid us from assigning at most  $d(u) + d(v) - 2$  possible distinct weights to  $uv$ . Now, since weighting  $uv$  with distinct weights results in distinct values of  $\sigma_w(u)$  and  $\sigma_w(v)$ , it should be clear that we can find a correct weight for  $uv$  in  $W$ , provided  $W$  includes at least  $d(u) + d(v) - 1$  distinct weights.  $\square$

Throughout this paper, several of the proofs consist in deleting two adjacent edges  $vu_1$  and  $vu_2$  from  $G$ , edge-weighting the remaining graph, and correctly extending the weighting to  $vu_1$  and  $vu_2$ . In this regard, we will often refer to the following result, which is about the number of weights that are sufficient to weight  $vu_1$  and  $vu_2$ .

**Observation 2.3.** *Let  $G$  be a graph having two adjacent edges  $vu_1$  and  $vu_2$  such that  $G' := G - \{vu_1, vu_2\}$  admits a neighbour-sum-distinguishing edge-weighting  $w_{G'}$ . Assume further that  $d_G(u_1) \geq d_G(u_2)$ , and set*

$$\mu := (d_G(u_1) + 1) + \max \{0, d_G(v) + d_G(u_2) - d_G(u_1) - 1\}.$$

*Then, assuming we have a set  $W$  of at least  $\mu$  distinct weights, we can extend  $w_{G'}$  to a neighbour-sum-distinguishing edge-weighting of  $G$ , by assigning two distinct weights of  $W$  to  $vu_1$  and  $vu_2$ .*

*Proof.* We extend  $w_{G'}$  to a neighbour-sum-distinguishing edge-weighting  $w_G$  of  $G$ , by first assigning a weight of  $W$  to  $vu_1$ , and then assigning a distinct weight to  $vu_2$ . We determine, in this proof, the smallest number  $\mu$  of weights that  $W$  should contain, so that this strategy has sufficiently many weights to be successfully applied.

We note that extending  $w_{G'}$  to  $vu_1$ , completely determines the value of  $\sigma_{w_G}(u_1)$ , while the value of  $\sigma_{w_G}(v)$  is not determined until  $vu_2$  is also weighted. Hence, when first weighting  $vu_1$ , we mainly have to make sure that  $\sigma_{w_G}(u_1)$  does not get equal to the sum of weights incident to a neighbour of  $u_1$  different from  $v$ . Also, we should make sure that  $\sigma_{w_{G'}}(v) + w_G(vu_1)$  does not get equal to  $\sigma_{w_{G'}}(u_2)$ , as otherwise, we would necessarily get  $\sigma_{w_G}(v) = \sigma_{w_G}(u_2)$ , no matter how we weight  $vu_2$ . There are hence  $d_G(u_1)$  conflicts to take into account when weighting  $vu_1$ . Provided  $W$  includes at least  $d_G(u_1) + 1$  distinct weights, we

can hence weight  $vu_1$  correctly, i.e. so that we avoid all conflicts mentioned above, with one weight from  $W$ , since assigning different weights to  $vu_1$  alters  $\sigma_{w_G}(u_1)$  in distinct ways.

Now assume  $vu_1$  has been weighted, with the additional property that  $\sigma_{w_{G'}}(v) + w_G(vu_1) \neq \sigma_{w_{G'}}(u_2)$ . Since that property holds, Observation 2.2 tells us that we can correctly extend  $w_{G'}$  to  $vu_2$ , provided  $W \setminus \{w_G(vu_1)\}$  includes at least  $d_G(v) + d_G(u_2) - 1$  distinct weights. We hence need  $W \setminus \{w_G(vu_1)\}$  to include that many distinct weights.

As explained above,  $W$  necessarily includes at least  $d_G(u_1)$  weights that were not assigned to  $vu_1$ . Hence, to make sure, after weighting  $vu_1$ , that  $W$  still includes at least  $d_G(v) + d_G(u_2) - 1$  distinct weights, we need  $W$  to include at least

$$(d_G(v) + d_G(u_2) - 1) - d_G(u_1)$$

other weights. This quantity can be negative, as, notably,  $vu_1$  may need a lot of weights to be weighted. Hence

$$\mu = (d_G(u_1) + 1) + \max\{0, d_G(v) + d_G(u_2) - d_G(u_1) - 1\},$$

as claimed, and, under the assumption that  $W$  has size  $\mu$ , we can achieve the extension of  $w_{G'}$  to  $G$ , as described earlier.  $\square$

In our proofs, we will also use the fact that, in some situations, pendant edges can easily be weighted, assuming we are provided enough distinct weights.

**Observation 2.4.** *Let  $G$  be a graph having a pendant edge  $vu$ , where  $u$  is the degree-1 vertex, such that  $G' := G - uv$  admits a neighbour-sum-distinguishing edge-weighting  $w_{G'}$ . Then, assuming we have a set  $W$  of at least  $d_G(v)$  distinct weights, we can extend  $w_{G'}$  to a neighbour-sum-distinguishing edge-weighting of  $G$ , by assigning a weight of  $W$  to  $vu$ .*

*Proof.* Following Observation 2.1, when extending  $w_{G'}$  to  $vu$ , we do not have to care whether  $\sigma(u)$  gets equal to  $\sigma(v)$ . We thus just have to make sure that  $\sigma(v)$  does not get equal to the sum of weights incident to one of its neighbours in  $G'$ . Recall that assigning distinct weights to  $vu$  results in different sums as  $\sigma(v)$ . Therefore, since  $v$  has  $d_G(v) - 1$  neighbours in  $G'$ , while  $W$  has size at least  $d_G(v)$ , there is necessarily a weight in  $W$  that can be assigned to  $vu$ , such that no conflict is created. An extension of  $w_{G'}$  to  $G$  hence exists.  $\square$

### 3. Classes of graphs agreeing with Conjecture 1.1

As mentioned in Section 1, we directly benefit, in the context of Conjecture 1.1, from the investigations on antimagic labellings, as antimagic graphs verify Conjecture 1.1. Following the survey [3] by Gallian, the following classes of nice graphs hence agree with Conjecture 1.1.

**Theorem 3.1.** *The classes of known antimagic graphs notably include:*

- nice paths,
- wheels,
- nice regular graphs,
- nice complete partite graphs.

Consequently, every of these graphs  $G$  verifies  $\chi_{\Sigma}^{e,1}(G) = |E(G)|$ .

When it comes to nice graphs with maximum degree 2, it is easily seen, as we are assigning strictly positive weights only, that any edge-injective edge-weighting is neighbour-sum-distinguishing. Disjoint unions of nice paths and cycles hence agree with Conjecture 1.1.

**Observation 3.2.** *Let  $G$  be a nice graph with  $\Delta(G) = 2$ . Then any edge-injective edge-weighting of  $G$  is neighbour-sum-distinguishing.*

One of the main lines of research concerning antimagic labellings, is to determine whether nice trees are all antimagic. In the following result, we prove that this question can be answered positively when relaxed to edge-injective neighbour-sum-distinguishing edge-weightings.

**Theorem 3.3.** *For every nice forest  $F$ , we have  $\chi_{\Sigma}^{e,1}(F) = |E(F)|$ .*

*Proof.* If  $\Delta(F) = 2$ , then the result follows from Observation 3.2. So the claim holds whenever  $F$  has size 2. Assume now that the claim is false, and let  $F$  be a counterexample that is minimum in terms of  $n_F + m_F$ , where  $n_F := |V(F)|$  and  $m_F := |E(F)|$ . By the remark above, we have  $m_F \geq 3$ . Furthermore, due to the minimality of  $F$ , all connected components of  $F$  include a vertex with degree at least 3, as otherwise we could remove a component  $C$  with maximum degree 2 from  $F$ , find an edge-injective neighbour-sum-distinguishing ( $|E(F - C)|$ )-edge-weighting of  $F - C$ , and easily extend the weighting to  $C$ , according to Observation 3.2.

We now successively show that  $F$ , because it is a counterexample to the claim, cannot contain certain structures, until we reach the point where  $F$  is shown to not exist at all, a contradiction. In particular, we focus on the length of the pendant paths of  $F$ , where a *pendant path* of  $F$  is a maximal path  $v_k \dots v_1$ , where  $k \geq 2$ , such that  $d(v_k) \geq 3$ ,  $d(v_{k-1}) = \dots = d(v_2) = 2$ , and  $d(v_1) = 1$ . In the case where  $k = 2$ , we note that the pendant path is a pendant edge, in which case  $v_k = v_2$  and we have  $d(v_2) \geq 3$ . Since every connected component  $C$  of  $F$  has a vertex with degree at least 3, every  $C$  has at least three pendant paths.

We start off by showing that the pendant paths of  $F$  all have length at most 2.

**Claim 3.4.** *Every pendant path of  $F$  has length at most 2.*

*Proof.* Assume  $F$  has a pendant path  $P := v_k \dots v_1$  with  $k \geq 4$ , where  $d(v_k) \geq 3$ . In this case, let  $F' := F - \{v_{k-1}v_{k-2}, \dots, v_2v_1\}$  be the forest obtained by removing all edges of  $P$ , but the one incident to  $v_k$ , from  $F$ . Clearly,  $F'$  is nice, and, due to the minimality of  $F$ , there exists an edge-injective neighbour-sum-distinguishing  $m_{F'}$ -edge-weighting  $w_{F'}$  of  $F'$ , where  $m_{F'} := |E(F')|$ . To prove that the claim holds, we just have to prove that we can extend  $w_{F'}$  to the edges  $v_{k-1}v_{k-2}, \dots, v_2v_1$ , hence to  $F$ , using weights  $m_{F'} + 1, \dots, m_F$ , so that we get an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting of  $F$ , a contradiction.

Due to the length of  $P$ , we have  $|\{m_{F'} + 1, \dots, m_F\}| \geq 2$ . When weighting the edges  $v_{k-1}v_{k-2}, \dots, v_2v_1$ , we note that we cannot create any sum conflicts involving any two consecutive vertices in  $\{v_1, \dots, v_{k-1}\}$ . This is according to Observation 2.1, because we are assigning weights injectively. Hence, when extending  $w_{F'}$ , we just have to make sure that  $\sigma(v_{k-1})$  gets different from  $\sigma(v_k)$ , which is possible as we have at least two distinct edge weights to work with. So we can assign a weight to  $v_{k-1}v_{k-2}$  which avoids that conflict, and then arbitrarily extend the weighting to the edges  $v_{k-2}v_{k-3}, \dots, v_2v_1$ . This yields an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting of  $F$ .  $\square$

Now consider any connected component  $C$  of  $F$  (in case  $F$  is a tree, we have  $C = F$ ), and designate a vertex  $r$  with degree at least 3 of  $C$  as being the *root* of  $C$ . This naturally

defines, in the usual way, an orientation of  $C$  from its root to its leaves. For every vertex  $v$  of  $C$ , the *father*  $f(v)$  of  $v$  is the neighbour of  $v$  which is the closest from  $r$  (if any). Conversely, the *descendants* of  $v$  are all vertices, different from  $v$ , in the subtree of  $C$  rooted at  $v$  (if any). We note that  $r$  has no father, while the leaves of  $C$  have no descendants. The descendants of  $v$  adjacent to  $v$  (if any), are called its *children*.

A *multifather*  $v$  of  $C$ , is a vertex with degree at least 3, i.e. having at least two children. In case all descendants of  $v$  have degree at most 2, we call  $v$  a *last multifather* of  $C$ . In other words, a last multifather is a vertex with at least two pendant paths attached. Since  $\Delta(C) \geq 3$ , there are last multifathers in  $C$ .

To further study the structure of  $F$ , we now focus on  $C$ , and prove properties of its last multifathers, still under the assumption that  $C$  is rooted at a vertex  $r$  with degree at least 3.

**Claim 3.5.** *Vertex  $r$  is not a last multifather.*

*Proof.* Assume the contrary. Then  $r$  is the only vertex with degree at least 3 of  $C$ . In other words,  $C$  is a subdivided star. Let  $F' := F - C$ , and set  $m_{F'} := |E(F')|$ . In case  $F'$  is not empty, due to the minimality of  $F$ , and because  $F'$  remains nice, there exists an edge-injective neighbour-sum-distinguishing  $m_{F'}$ -edge-weighting of  $F'$ . We show below that  $w_{F'}$  can be extended to an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting  $w_F$  of  $F$ . This is a contradiction.

So that  $w_F$  is edge-injective, we must extend  $w_{F'}$  to the edges of  $C$  using the weights among  $\{m_{F'} + 1, \dots, m_F\}$  injectively. Since  $r$  is the only vertex with degree at least 3 in  $C$ , it should be clear that assigning the weights  $m_F, m_F - 1, \dots, m_{F'} + 1$ , following this ordering, to the edges of  $C$  as they are encountered during a breadth-first search algorithm performed from  $r$ , results in a neighbour-sum-distinguishing edge-weighting of  $C$ . To be convinced of this statement, one can e.g. just refer to Observation 2.1.

As a concluding remark, let us just note that, in case  $F'$  is empty, i.e.  $F = C$ , the weighting scheme above can be applied on  $F$  from scratch, resulting in an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting of  $F$ .  $\square$

Due to Claim 3.5, we may assume that the root  $r$  of  $C$  is not a last multifather. Then, all last multifathers of  $C$  (there are some), are different from  $r$ , and hence have a father. We now refine Claim 3.4 to the following.

**Claim 3.6.** *Every pendant path attached to a last multifather of  $C$  has length 1.*

*Proof.* Let  $v \neq r$  be a last multifather of  $C$ , and assume  $v$  is incident to pendant paths with length 2. We recall that all pendant paths attached to  $v$  have length at most 2 (Claim 3.4), and, since  $v$  is a last multifather, it is incident to at least two pendant paths. Let  $F'$  be the forest obtained from  $F$  by removing all pendant paths attached to  $v$ . Because  $m_{F'} := |E(F')|$  is smaller than  $m_F$ , there exists an edge-injective neighbour-sum-distinguishing  $m_{F'}$ -edge-weighting  $w_{F'}$  of  $F'$ . For contradiction, we prove below that  $w_{F'}$  can be extended correctly to the pendant paths attached to  $v$ , using the weights among  $\{m_{F'} + 1, \dots, m_F\}$  injectively.

Let  $b \geq 1$  be the number of pendant paths of length 2 attached to  $v$  in  $C$ , and let  $vx_1y_1, \dots, vx_by_b$  denote those paths (so that the  $x_i$ 's have degree 2 in  $C$ , while the  $y_i$ 's have degree 1). Vertex  $v$  is also adjacent to  $c \geq 0$  leaves  $x_{b+1}, \dots, x_{b+c}$ , which are, in some sense, pendant paths of length 1. Since  $v$  is a multifather, we recall that  $b + c = d_F(v) - 1 \geq 2$ .

We extend  $w_{F'}$  to the edges of the pendant paths attached to  $v$ , in the following way. First, we injectively arbitrarily assign the  $d_F(v) - 1$  weights in  $\{m_F - d_F(v) + 2, \dots, m_F\}$  to the edges  $vx_2, \dots, vx_{b+c}$ . After that, we assign to the edge  $vx_1$  one of the weights



$m_F - d_F(v) + 1$  or  $m_F - d_F(v)$ , chosen so that  $\sigma_{w_F}(v)$  is different from the sum of weights, by  $w_{F'}$ , incident to  $f(v)$ , the father of  $v$ . We then assign to  $x_1y_1$  the one weight of  $m_F - d_F(v) + 1$  or  $m_F - d_F(v)$  not assigned to  $vx_1$ . We note that no matter how we complete the extension of  $w_{F'}$ , eventually  $\sigma_{w_F}(v)$  will be strictly bigger than  $\sigma_{w_F}(x_1)$ .

We finish the extension of  $w_{F'}$  to  $C$ , by arbitrarily injectively assigning the remaining non-used smaller weights to the edges  $x_1y_1, \dots, x_by_b$ . Because all the  $x_i$ 's have degree 2 and the  $y_i$ 's have degree 1, no conflict may arise between those vertices (Observation 2.1). Furthermore, since the degree of  $v$  is larger than the degree of the  $x_i$ 's, and the weights assigned to the  $vx_i$ 's are bigger than the weights assigned to the  $x_iy_i$ 's (with possibly the exception of  $vx_1$  and  $x_1y_1$ , which we have discussed above), it should be clear that no conflict may arise between  $v$  and the  $x_i$ 's (again according to Observation 2.1). So we eventually get an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting of  $F$ , a contradiction.  $\square$

We finally study last multifathers of  $C$  being at maximum distance from  $r$ . We call these vertices the *deepest last multifathers* of  $C$ . From now on, we focus on a fixed deepest last multifather  $v^*$  of  $C$ , which we choose arbitrarily. In the upcoming proof, for any vertex  $v$  of  $C$ , we denote by  $C_v$  the subtree of  $C$  rooted at  $v$ . Recall that all children of a last multifather are leaves (Claim 3.6).

**Claim 3.7.** *Every last multifather  $v$  of  $C_{f(v^*)}$  is a child of  $f(v^*)$ . In other words,  $v$  is a deepest last multifather of  $C$ .*

*Proof.* The claim follows from the fact that if there exists a descendant  $v \neq v^*$  of  $f(v^*)$  being at distance at least 2 from  $f(v^*)$ , then  $v$  would, in  $C$ , be at greater distance from  $r$  than  $v^*$  is. This would contradict the fact that  $v^*$  is a deepest last multifather.  $\square$

Recall that  $f(v^*)$  cannot be incident, in  $C$ , to a pendant path with length at least 3 (Claim 3.4). Hence, every child of  $f(v^*)$  is either a leaf (type-1), a degree-2 vertex adjacent to a leaf (type-2, i.e. the inner vertex of a pendant path with length 2), or a deepest last multifather (type-3). Furthermore, we know that  $f(v^*)$  is adjacent to at least one type-3 vertex, which is  $v^*$ . In the following proof, we show that  $v^*$  is actually the only child of  $f(v^*)$  in  $C$ .

**Claim 3.8.** *Vertex  $v^*$  is the only child of  $f(v^*)$  in  $C$ .*

*Proof.* Suppose the claim is false, and let  $v \neq v^*$  be another child of  $f(v^*)$ . Let  $x_1$  and  $x_2$  be two leaves adjacent to  $v^*$ , which exist since  $v^*$  is a last multifather, and all pendant paths attached to  $v^*$  have length 1 (Claim 3.6).

Assume first that  $v$  is type-2 or type-3, or, in other words, that  $d_F(v) \geq 2$ . In that case,  $v$  is adjacent to at least one leaf, say  $y$ . We here consider  $F' := F - \{vy, v^*x_1, v^*x_2\}$ . Note that  $F'$  remains nice, and has fewer edges than  $F$ . Due to the minimality of  $F$ , there hence exists an edge-injective neighbour-sum-distinguishing  $m_{F'}$ -edge-weighting  $w_{F'}$  of  $F'$ , where  $m_{F'} := |E(F')|$ . We show below that  $w_{F'}$  can be extended to the three removed edges, with injectively using the three edge weights  $m_F - 2, m_F - 1, m_F$ , yielding an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting  $w_F$  of  $F$ , a contradiction.

We first assign a weight to  $v^*x_1$ , based on the conflicts that may happen when weighting  $vy$ . When assigning any of the three weights to  $vy$ , the only problem which may occur, recall Observation 2.1, is that  $\sigma_{w_F}(v)$  gets equal to  $\sigma_{w_{F'}}(f(v^*))$ . If assigning one of the three weights  $m_F - 2, m_F - 1, m_F$  to  $vy$  indeed results in that conflict, we assign that weight to  $v^*x_1$ . Otherwise, we assign any of the three weights to  $v^*x_1$ . In any case, no conflict may arise, as  $\sigma_{w_F}(v^*)$  is still not determined.

We are now left with two weights, which we must assign to  $v^*x_2$  and  $vy$ . Due to the choice of the weight assigned to  $v^*x_1$ , we note that no problem may occur when weighting  $vy$ . Hence, we just have to weight  $v^*x_2$  correctly, and assign the remaining weight to  $vy$ . When weighting  $v^*x_2$ , the only problem which may occur, according to Observation 2.1, is that  $\sigma_{w_F}(v^*)$  gets equal to  $\sigma_{w_{F'}}(f(v^*))$ . But, since we have two distinct weights to work with, one of them can be assigned to  $v^*x_2$  so that this conflict is avoided. Thus we can weight  $v^*x_2$  correctly, and eventually weight  $vy$  with the remaining weight, resulting in the claimed  $w_F$ .

We may now assume that all children, including  $v$ , of  $f(v^*)$  different from  $v^*$  are type-1, i.e. leaves. The contradiction can then be obtained quite similarly as in the previous case, but with setting  $F' := F - \{f(v^*)v, v^*x_1, v^*x_2\}$ . When weighting  $f(v^*)v$ , we have to make sure, if  $f(v^*) \neq r$ , that  $\sigma_{w_F}(f(v^*))$  does not get equal to  $\sigma_{w_{F'}}(f(f(v^*)))$ . Note that if  $f(v^*) = r$ , then the situation is actually easier, as there is one less conflict to consider. If one of the three available weights  $m_F - 2, m_F - 1, m_F$ , when assigned to  $f(v^*)v$ , yields a conflict involving  $f(v^*)$  and  $f(f(v^*))$ , then we assign that weight to  $v^*x_1$ . Otherwise, we assign any weight to  $v^*x_1$ . This ensures that, when assigning any of the two remaining weights to  $f(v^*)v$ , no conflict may involve  $f(v^*)$  and  $f(f(v^*))$ . We finally arbitrarily assign the two remaining weights to  $v^*x_2$  and  $f(v^*)v$ . If this results in a neighbour-sum-distinguishing edge-weighting  $w_F$  of  $F$ , then we are done. Otherwise, it means that  $\sigma_{w_F}(v^*) = \sigma_{w_F}(f(v^*))$ . In that case, note that, because all assigned edge weights are distinct, when swapping the values assigned to  $v^*x_2$  and  $f(v^*)v$  by  $w_F$ , that conflict cannot remain. Furthermore, according to the remarks above, we still do not create any sum conflict involving  $f(v^*)$  and  $f(f(v^*))$ . After the swapping operation,  $w_F$  hence gets neighbour-sum-distinguishing.  $\square$

We are now ready to finish off the proof, by showing that, under all information we have obtained,  $F$  actually admits an edge-injective neighbour-sum-distinguishing  $w_F$ -edge-weighting, a contradiction.

From Claim 3.8, we get that  $d_F(f(v^*)) = 2$ , as  $v^*$  is not the root of  $C$ , so  $f(f(v^*))$  exists. Let  $x_1, \dots, x_k$  be the  $k \geq 2$  leaves attached to  $v^*$  in  $C$ , which exist since  $v^*$  is a type-3 vertex. Now consider the forest  $F' := F - \{v^*x_1, \dots, v^*x_k\}$  with size  $m_{F'} := |E(F')|$ . Due to the minimality of  $F$ , there exists an edge-injective neighbour-sum-distinguishing  $m_{F'}$ -edge-weighting  $w_{F'}$  of  $F'$ . We extend  $w_{F'}$  to the  $k$  removed edges, so that an edge-injective neighbour-sum-distinguishing  $m_F$ -edge-weighting  $w_F$  of  $F$  is obtained, a contradiction. To that aim, we arbitrarily injectively assign the weights  $m_F - k + 1, \dots, m_F$  to the pendant edges  $v^*v_1, \dots, v^*v_k$  attached to  $v^*$ . Recall that we cannot get sum conflicts involving  $v^*$  and the  $v_i$ 's, according to Observation 2.1. Furthermore, we have  $d_F(v^*) \geq 3$ , while  $d_F(f(v^*)) = 2$  (Claim 3.8), and we have used the  $k$  biggest weights among  $\{1, \dots, m_F\}$  to weight the edges incident to  $v^*$ . From this and Observation 2.1, we get that, necessarily,  $\sigma_{w_F}(v^*) > \sigma_{w_{F'}}(f(v^*))$ . So  $w_F$  is neighbour-sum-distinguishing.  $\square$

#### 4. General upper bounds

Towards Conjecture 1.1, we start off by exhibiting, for any nice graph  $G$ , an upper bound on  $\chi_{\Sigma}^{e,1}(G)$  of the form  $k \cdot |E(G)|$ , where  $k$  is a fixed constant.

It turns out, first, that some results towards the 1-2-3 Conjecture can be extended to the edge-injective context, hence yielding bounds to our context. This is in particular the case of the weighting algorithm by Kalkowski, Karoński and Pfender from [5], which was designed to prove that  $\chi_{\Sigma}^e(G) \leq 5$  holds for every nice graph  $G$ . In very brief words, this algorithm initially assigns the list of weights  $\{1, 2, 3, 4, 5\}$  to every edge of  $G$ , which

basically contains the possible weights that any edge can be assigned at any moment of the algorithm. The algorithm then linearly processes the vertices of  $G$ , with possibly adjusting some incident edges weights, and staying in the list  $\{1, 2, 3, 4, 5\}$ , so that sum conflicts are avoided around any vertex considered during the course.

It is an easy exercise to check that this algorithm also works, under the assumption that every edge of  $G$  is assigned a (possibly unique) list of five allowed consecutive weights  $\{\alpha - 2, \alpha - 1, \alpha, \alpha + 1, \alpha + 2\}$ . In particular, when applied with non-intersecting such lists assigned to the edges, the algorithm yields an edge-injective neighbour-sum-distinguishing edge-weighting, as every edge weight can be assigned to at most one edge. So, applying the algorithm on a nice graph  $G$  with edges  $e_0, \dots, e_{m-1}$  where each edge  $e_i$  is assigned the list  $\{5i + 1, 5i + 2, 5i + 3, 5i + 4, 5i + 5\}$ , results in an edge-injective neighbour-sum-distinguishing  $(5 \cdot |E(G)|)$ -edge-weighting of  $G$ . From this, we get that  $\chi_{\Sigma}^{e,1}(G) \leq 5 \cdot |E(G)|$  holds for every nice graph  $G$ .

The  $5 \cdot |E(G)|$  bound on  $\chi_{\Sigma}^{e,1}(G)$  above, can actually be improved down to  $2 \cdot |E(G)|$ , by means of a careful inductive proof scheme, which we describe in the following proof.

**Theorem 4.1.** *For every nice graph  $G$ , we have  $\chi_{\Sigma}^{e,1}(G) \leq 2 \cdot |E(G)|$ .*

*Proof.* The proof is by induction on  $n_G + m_G$ , where  $n_G := |V(G)|$  and  $m_G := |E(G)|$ . As it can easily be checked that the claim is true for small values of  $n_G + m_G$ , we proceed to the induction step. Consider hence a value of  $n_G + m_G$ , such that the claim is true for smaller values of this sum.

Set  $\Delta := \Delta(G)$ . Since we may assume that  $m_G \geq 4$ , and  $G$  is nice, we clearly have  $\Delta \geq 2$ . We may even assume that  $\Delta \geq 3$ , as otherwise  $G$  would admit an edge-injective neighbour-sum-distinguishing  $m_G$ -edge-weighting, according to Observation 3.2. Consider any vertex  $v^*$  of  $G$  verifying  $d_G(v^*) = \Delta$ , and denote by  $u_1, \dots, u_{\Delta}$  the neighbours of  $v^*$  in  $G$ . Let further  $C$  denote the connected component of  $G$  containing  $v^*$ , and set  $C' := C - v^*$ . Note that  $C'$  may include connected components isomorphic to  $K_2$ , and thus be not nice. In this context, we say that a component of  $C'$  is *empty* if it has no edge, *bad* if it has size 1, and *good* otherwise. Basically, a bad component of  $C'$  is an edge to which  $v^*$  is joined in  $C$ : either  $v^*$  is adjacent to the two ends of that edge, or  $v^*$  is adjacent to only one of the two ends.

If  $C'$  does not have good components, then  $C$  is basically a connected graph whose only vertex with degree at least 3 is  $v^*$ , such that  $C'$  consists of isolated vertices and isolated edges only. In particular, all vertices of  $C$  but  $v^*$  have degree at most 2, and every degree-2 vertex  $u_i$  adjacent to  $v^*$  is either adjacent to another degree-2 vertex  $u_j$  adjacent to  $v^*$ , or adjacent to a degree-1 vertex. In such a situation, it can easily be seen that assigning  $|E(C)|$  consecutive decreasing weights  $\alpha, \alpha - 1, \dots, \alpha - |E(C)| + 1$ , following this order, to the edges of  $C$  as they are encountered while performing a breadth-first search algorithm from  $v^*$ , results in an edge-injective neighbour-sum-distinguishing edge-weighting of  $C$ . This notably follows as a consequence of Observation 2.1. If  $G = C$ , then this procedure yields an edge-injective neighbour-sum-distinguishing  $m_G$ -edge-weighting of  $G$ . Now, if  $G \neq C$ , then we can just consider  $G' := G - C$ , apply induction on  $G'$ , and extend an edge-injective neighbour-sum-distinguishing  $2m_{G'}$ -edge-weighting of  $G'$  to  $G$ , where  $m_{G'} := |E(G')|$ , by just assigning  $|E(C)|$  new consecutive weights to the edges of  $C$ , as described above.

Hence we may assume that  $C'$  has good connected components  $C'_1, \dots, C'_k$ . Let  $G'$  denote the union of the  $C'_i$ 's and of the connected components of  $G$  different from  $C$ , and set  $m_{G'} := |E(G')|$ . Since the  $C'_i$ 's and all connected components of  $G$  are nice, so is  $G'$ . Furthermore, we have that  $m_{G'} < m_G$ . According to the induction hypothesis, there hence

exists an edge-injective neighbour-sum-distinguishing  $2m_{G'}$ -edge-weighting  $w_{G'}$  of  $G'$ . In order to get an edge-injective neighbour-sum-distinguishing  $2m_G$ -edge-weighting  $w_G$  of  $G$ , we eventually just need to extend  $w_{G'}$  to the remaining edges of  $G$ , i.e. to the  $v^*u_i$ 's and the edges of the bad components of  $C'$ .

To that aim, we restrict ourselves to injectively using weights among  $\{2m_{G'}+1, \dots, 2m_G\}$ , i.e. we do not use non-used weights among  $\{1, \dots, 2m_{G'}\}$ . The main reason for doing so, is that, according to Observation 2.1, and because  $d_G(v^*) = \Delta(G)$  and  $w_{G'}$  is a  $2m_{G'}$ -edge-weighting of  $G'$ , this will guarantee that  $\sigma_{w_G}(v^*) > \sigma_{w_G}(u_i)$  holds for every  $u_i$  in  $G'$ . In other words, when extending  $w_{G'}$  to  $w_G$  by using weights among  $\{2m_{G'}+1, \dots, 2m_G\}$ , no sum conflicts involving  $v^*$  and any  $u_i \in V(G')$  can arise.

We start by extending  $w_{G'}$  to the edges of  $G$  belonging to the bad components, or being incident to the bad components (i.e. being incident to  $v^*$ ). Assume there are  $m'$  of them. Then, as in a previous case, we assign weights  $2m_{G'}+m', \dots, 2m_{G'}+1$  to these edges, following this order, as they are encountered during a breadth-first search algorithm performed from  $v^*$ . It can easily be checked that, since some of the  $u_i$ 's belong to  $G'$ , no matter how we complete the extension of  $w_{G'}$  to  $w_G$ , no conflict involving  $\sigma_{w_G}(v^*)$  and  $\sigma_{w_G}(u_i)$  for any  $u_i \notin V(G')$  can occur. This is because  $\sigma_{w_G}(v^*)$  will eventually be big enough.

We are now left with weighting the edges of  $G$  of the form  $v^*u_i$  with  $u_i \in V(G')$ , and we use the weights among  $\{2m_{G'}+m'+1, \dots, 2m_G\}$  for that. According to previous arguments, when weighting such an edge  $v^*u_i$ , we just have to make sure that  $\sigma_{w_G}(u_i)$  does not hit the current value of  $\sigma_{w_G}(u')$  for every neighbour  $u' \neq v^*$  of  $u_i$ . This is clearly always possible, as

$$|\{2m_{G'}+m'+1, \dots, 2m_G\}| \geq 2 \cdot |\{u_i : u_i \in V(G')\}|,$$

while  $d_{G'}(u_i) \leq \Delta - 1$ . So, we have that

$$|\{2m_{G'}+m'+1, \dots, 2m_G\}| - (\Delta - 1) > \Delta,$$

which implies that, even for the last  $u_i$  in  $G'$  to be considered, there are still at least  $\Delta + 1$  available weights that can be assigned to the incident edge  $v^*u_i$ , while only at most  $\Delta - 1$  adjacent sums are forbidden around  $u_i$ . This is basically a straight application of Observation 2.2. For these reasons,  $w_{G'}$  can hence be extended to every edge  $v^*u_i$  of  $G$ , implying that the claimed edge-weighting  $w_G$  of  $G$  exists. This concludes the proof.  $\square$

We now provide a second upper bound on  $\chi_{\Sigma}^{e,1}(G)$  of the form  $|E(G)| + k$ , for every nice graph  $G$ . Here, our  $k$  is a small linear function of  $\Delta(G)$ , making the bound 1) mostly interesting in the context of nice graphs with bounded maximum degree, and 2) generally better than the bound in Theorem 4.1 (except in some cases to be discussed later). The proof scheme we employ here, is different from the one used to prove Theorem 4.1.

**Theorem 4.2.** *For every nice graph  $G$ , we have  $\chi_{\Sigma}^{e,1}(G) \leq |E(G)| + 2\Delta(G)$ .*

*Proof.* We here focus on the case where  $G$  is connected, as it can easily be observed that, when  $G$  is connected, the proof scheme below can be applied whenever the edges of  $G$  are injectively weighted with weights from a set of at least  $|E(G)| + 2\Delta(G)$  distinct weights. In other words, if  $G$  is not connected, then the weighting strategy below can be repeated for every connected component of  $G$ , and, to ensure edge-injectivity, we can just make sure to use any set of non-used weights for every component to be weighted.

Set  $\Delta := \Delta(G)$ , and let  $n := |V(G)|$  and  $m := |E(G)|$  denote the order and size, respectively, of  $G$ . First choose a vertex  $v^*$  with degree  $\Delta$  in  $G$ , and let  $T$  be a spanning tree of  $G$  including all edges incident to  $v^*$ . From  $T$ , we deduce a partition  $V_0 \cup \dots \cup V_k$  of

$V(G)$ , where each part  $V_i$  includes the vertices of  $G$  being at distance  $i$  from  $v^*$  in  $T$ . In particular,  $V_0 = \{v^*\}$ , and, for every vertex  $u$  in a part  $V_i$  with  $i \neq 0$ , there is exactly one edge from  $u$  to  $V_{i-1}$  in  $T$ . We call this edge the *private edge* of  $u$ .

We now describe how to obtain an edge-injective neighbour-sum-distinguishing  $(m + 2\Delta)$ -edge-weighting of  $G$ . We start by assigning the edge weights  $1, \dots, m - (n - 1)$  to the edges of  $E(G) \setminus E(T)$  in an arbitrary way. This leaves us with all edges of  $T$  to be weighted, which includes at least one incident edge for every vertex different from  $v^*$ , and all edges incident to  $v^*$ . To weight these edges without creating any conflict, we first consider all vertices of  $V_k$  and weight their private edges carefully, then do the same for all vertices of  $V_{k-1}$ , and so on layer by layer until all edges of  $T$  are weighted. Fixing any ordering over the vertices of  $V_k, \dots, V_1$ , this weighting scheme yields an ordering  $u_1, \dots, u_{n-1}$  in which the vertices are considered (i.e. the  $|V_k|$  first  $u_i$ 's belong to  $V_k$ , the  $|V_{k-1}|$  next  $u_i$ 's belong to  $V_{k-1}$ , and so on; the  $|V_1|$  last  $u_i$ 's belong to  $V_1$ ). We note that the private edges of the  $|V_1|$  last  $u_i$ 's, go to  $v^*$ .

Before giving technical details, we first explicit the conflicts that may arise when weighting the private edges. Assume we are currently dealing with a vertex  $u_i \in V_j$ , meaning that the private edge of each of  $u_1, \dots, u_{i-1}$  has been weighted successfully, hence that for any  $u_{i'}$  of these vertices, we have that  $\sigma(u_{i'})$  is determined, and  $\sigma(u_{i'})$  is different from any  $\sigma(u_{i''})$ , where  $i'' < i$ , such that  $u_{i'}$  and  $u_{i''}$  are neighbours. In case  $u_i \neq u_{n-1}$ , we note that the only constraint we need to take into account when weighting the private edge of  $u_i$ , is that the value of  $\sigma(u_i)$  must get different from  $\sigma(u_{i'})$  for every  $u_{i'}$  neighbouring  $u_i$ , where  $i' < i$ . Hence, to be successful in weighting the private edge of  $u_i$ , we just need to have at least  $d_G(u_i)$  non-used weights in hand, as the private edge of  $u_i$  is its only incident edge being not weighted yet, and, since  $u_i \neq u_{n-1}$ , there are at most  $\Delta - 1$  vertices  $u_{i'}$  neighbouring  $u_i$ , and verifying  $i' < i$ .

In the case where  $u_i = u_{n-1}$ , we note that weighting the private edge of  $u_i$  also determines  $\sigma(v^*)$ , which must be different from the sums of weights incident to the vertices in  $V_1$ . The trick here, is to weight the private edges going to  $v^*$ , i.e. all edges from  $V_1$  to  $v^*$ , with weights that are bigger than those used on the edges not incident to  $v^*$ . In doing so, and because  $d(v^*) = \Delta$ , we necessarily get that  $\sigma(v^*)$  is bigger than the sums of weights incident to the vertices in  $V_1$ , and, hence, when weighting the private edge of  $u_{n-1}$ , we only have to make sure that  $\sigma(u_{n-1})$  gets different from the incident sums of at most  $\Delta - 1$  of the  $u_i$ 's. As a consequence, we have, before the weighting procedure, to save at least  $2\Delta$  big weights so that the private edges from  $V_1$  to  $v^*$  can be weighted in the manner described above, according to Observation 2.2.

We are now ready to give more precise details. To extend the edge-injective neighbour-sum-distinguishing edge-weighting to the edges of  $T$  correctly, we apply the procedure above, with sticking to the following rules:

- 1) to every edge of  $T$  not incident to  $v^*$ , we assign a weight among  $\{m - (n - 1) + 1, \dots, m\}$ ;
- 2) to every other edge of  $T$ , we assign a weight among  $\{m + 1, \dots, m + 2\Delta\}$ .

We note that, when considering any vertex  $u_i$  not in  $V_1$ , the number of non-used weights in  $\{m - (n - 1) + 1, \dots, m\}$  is at least  $\Delta + 1$ . According to the arguments above, we can hence always correctly weight the private edge of  $u_i$ . Similarly, by the explanations above,  $2\Delta$  big weights are sufficient to weight the private edges of the  $u_i$ 's in  $V_1$ . The weighting process can hence always be achieved successfully, which concludes the proof.  $\square$

As a concluding remark, we would like to point out that the  $2 \cdot |E(G)|$  bound from Theorem 4.1, can, in several situations, be better than the  $|E(G)| + 2\Delta(G)$  bound from Theorem 4.2. To be convinced of that statement, just consider the class of graphs obtained by starting from any star with  $\Delta$  leaves  $u_1, \dots, u_\Delta$ , and adding no more than  $\Delta - 1$  edges joining pairs of vertices among  $\{u_1, \dots, u_\Delta\}$ .

## 5. Refined bounds for particular classes of sparse graphs

We now improve the bounds in Section 4 to bounds of the form  $|E(G)| + k$ , where  $k$  is a small constant, for several classes of nice graphs  $G$ . Our weighting strategy here, relies on removing some edges from  $G$ , then deducing a correct edge-weighting of the remaining graph, and extending that weighting to  $G$ . So that this weighting strategy applies, we focus on rather sparse graph classes with particular properties inherited by their subgraphs. In that respect, we give a special focus to nice 2-degenerate graphs, and nice graphs with maximum average degree at most 3. It is worth recalling that these graphs may have arbitrarily large maximum degree, so Theorem 4.2 does not provide the kind of bound we are here interested in.

Throughout this section, when speaking of a  $k$ -vertex, we mean a degree- $k$  vertex. By a  $k^-$ -vertex (resp.  $k^+$ -vertex), we refer to a vertex with degree at most (resp. at least)  $k$ .

### 5.1. 2-degenerate graphs

A graph  $G$  is said to be  $k$ -degenerate, if every subgraph of  $G$  has a  $k^-$ -vertex. In the next result, we focus on nice 2-degenerate graphs, and exhibit an upper bound on their value of  $\chi_\Sigma^{e,1}$ .

**Theorem 5.1.** *For every nice 2-degenerate graph  $G$ , we have  $\chi_\Sigma^{e,1}(G) \leq |E(G)| + 4$ .*

*Proof.* Assume the claim is false, and let  $G$  be a counterexample that is minimal in terms of  $n_G + m_G$ , where  $n_G := |V(G)|$  and  $m_G := |E(G)|$ . We show below that  $G$  cannot be a counterexample, and thereby get a contradiction. This is done by showing that we can always remove some edges from  $G$ , while keeping the graph nice, then deduce an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting  $w_{G'}$  of the remaining graph  $G'$ , where  $m_{G'} := |E(G')|$ , and finally extend  $w_{G'}$  to get an edge-injective neighbour-sum-distinguishing  $(m_G + 4)$ -edge-weighting  $w_G$  of  $G$ .

We start by pointing out properties of  $G$  we may assume. According to Observation 3.2, we may assume that  $\Delta(G) \geq 3$ , and, therefore, that  $m_G \geq 4$ , as otherwise  $G$  would be a tree, in which case  $\chi_\Sigma^{e,1}(G) = m_G$ , according to Theorem 3.3. We note as well that if  $G$  has 1-vertices, then those vertices must be adjacent to vertices with sufficiently large degree:

**Claim 5.2.** *Every 1-vertex of  $G$  is adjacent to a  $6^+$ -vertex.*

*Proof.* Assume for contradiction that  $G$  has a 1-vertex  $u$  adjacent to a  $5^-$ -vertex  $v$ , and consider  $G' := G - uv$ . Clearly  $G'$  is nice, and 2-degenerate. So  $G'$  admits an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting  $w_{G'}$ , where  $m_{G'} := m_G - 1$ . According to Observation 2.4, we can correctly extend  $w_{G'}$  to  $uv$ , hence to  $G$ , since we have at least five distinct weights available for that. This is a contradiction.  $\square$

From Claim 5.2, we also deduce the following claim concerning the 2-vertices of  $G$ .

**Claim 5.3.** *Graph  $G$  has no edge  $uv$  with  $d_G(u) = d_G(v) = 2$ .*

*Proof.* Assume  $G$  has such an edge, and let us denote by  $u'$  and  $v'$  the neighbour of  $u$  and  $v$ , respectively, different from  $v$  and  $u$ , respectively. Then  $d_G(u'), d_G(v') \geq 2$ , as otherwise  $G$  would have a 1-vertex contradicting Claim 5.2. Set  $G' := G - uv$  and  $m_{G'} := |E(G')|$ . Since  $d_G(u'), d_G(v') \geq 2$ , we note that  $G'$  is nice. Furthermore,  $G'$  is 2-degenerate, so there exists an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting of  $G'$ .

We observe that  $w_{G'}$  can be extended to an edge-injective neighbour-sum-distinguishing  $(m_G + 4)$ -edge-weighting  $w_G$  of  $G$ , a contradiction. Since  $w_{G'}(uu') \neq w_{G'}(vv')$  by edge-injectivity, we note that we cannot eventually get  $\sigma_{w_G}(u) = \sigma_{w_G}(v)$ . Hence, we just have to make sure that  $\sigma_{w_G}(u)$  does not get equal to  $\sigma_{w_{G'}}(u')$ , and that  $\sigma_{w_G}(v)$  does not get equal to  $\sigma_{w_{G'}}(v')$ . This is clearly possible, since we have five distinct ways to weight  $uv$ .  $\square$

We are now ready to start off the proof. Let  $S_1$  denote the set of  $2^-$ -vertices of  $G$ , and set  $G_1 := G - S_1$ . Since  $\Delta(G) \geq 3$ , graph  $G_1$  has vertices. In particular, since  $G_1$  is 2-degenerate, it has a  $2^-$ -vertex  $v$ . Let us denote as  $d^+(v)$  the number of neighbours, in  $G$ , of  $v$  in  $S_1$ . Then  $d_G(v) = d^+(v) + d_{G_1}(v)$ .

First assume that  $d^+(v) \geq 3$ , and let  $v_1, v_2, v_3$  be three neighbours of  $v$  in  $S_1$ . We here consider  $G' := G - \{vv_1, vv_2, vv_3\}$ . Note that  $G'$  has to be nice, as otherwise  $G$  would have a 1-vertex violating Claim 5.2, or an edge violating Claim 5.3. Due to the minimality of  $G$ , and because  $G'$  is a nice 2-degenerate graph, there exists an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting  $w_{G'}$  of  $G'$ . We extend  $w_{G'}$  to  $vv_1, vv_2, vv_3$ , thus to  $G$ , assigning weights among a set of seven weights including those among  $\{m_G + 2, m_G + 3, m_G + 4\}$ , in the following way.

We first assign a weight  $\alpha_1$  from  $\{m_G + 3, m_G + 4\}$  to the edge  $vv_1$ , so that we do not create a sum conflict involving  $v_1$  and its neighbour different from  $v$  (if any), which is clearly possible with two distinct weights. Similarly, we then assign a weight  $\alpha_2$  from  $\{m_G + 2, m_G + 3, m_G + 4\} \setminus \{\alpha_1\}$  to  $vv_2$ , so that we do not create a sum conflict involving  $v_2$  and its neighbour different from  $v$  (if any). Note that, due to the choice of  $\alpha_1$  and  $\alpha_2$ , which are strictly bigger than the weights among  $\{1, \dots, m_{G'} + 4\}$ , no matter how we extend the weighting to  $vv_3$ , it cannot occur that  $\sigma_{w_G}(v)$  gets equal to the sum of weights incident to a  $2^-$ -vertex neighbouring  $v$ . Hence, when extending  $w_{G'}$  to  $vv_3$ , we just have to make sure that  $\sigma_{w_G}(v_3)$  does not get equal to the sum of weights incident to the neighbour of  $v_3$  different from  $v$  (if any), and that  $\sigma_{w_G}(v)$  does not get equal to the sums of weights incident to its at most two neighbours in  $G_1$ . So there are at most three conflicts to take into account, while we have five weights in hand to weight  $vv_3$ . Clearly, this is sufficient to extend the weighting.

Assume now that  $d^+(v) = 2$  and let  $v_1, v_2 \in S_1$  denote the two neighbours of  $v$  with degree at most 2 in  $S_1$ . We here consider  $G' := G - \{vv_1, vv_2\}$ , which is again nice (due to Claims 5.2 and 5.3) and 2-degenerate, and hence admits an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting  $w_{G'}$ , where  $m_{G'} := m_G - 2$ . Recall that  $d_G(v_1), d_G(v_2) \leq 2$ , and that  $d_G(v) \leq 4$ . According to Observation 2.3, we can correctly extend  $w_{G'}$  to  $vv_1$  and  $vv_2$  provided we have at least six distinct weights in hand. Since this is precisely the case here, an extension of  $w_{G'}$  to  $G$  exists.

The last case to consider, is when  $d^+(v) = 1$ , which we cannot directly treat using similar arguments as above. We may however assume that all  $2^-$ -vertices  $v$  of  $G_1$  verify  $d^+(v) = 1$ , as otherwise one of the previous situations would apply. Furthermore, these vertices have degree exactly 3, i.e. they each have exactly two neighbours in  $G_1$ , as otherwise they would belong to  $S_1$ . Now let  $S_2$  denote the set of all  $2^-$ -vertices of  $G_1$ , and set  $G_2 := G - \{S_1, S_2\}$ . We fix a vertex  $v^*$  for the rest of the proof, chosen as follows. If  $G_2$  has vertices, then we just choose, as  $v^*$ , a vertex of  $G_2$  verifying  $d_{G_2}(v^*) \leq 2$  (which exists,

as  $G_2$  is 2-degenerated). Otherwise, we choose as  $v^*$  one vertex verifying  $0 < d_{G_1}(v^*) \leq 2$ . In the latter case, note that  $v^*$  belongs to  $S_2$ .

Now, consider the following sets:

$$V_1 := \{v \in S_1 \cap N_G(v^*)\} \quad \text{and} \quad V_2 := \{v \in S_2 \cap N_G(v^*)\},$$

and set  $d_1^+ := |V_1|$  and  $d_2^+ := |V_2|$ . Due to our choice of  $v^*$ , we have  $d_2^+ \geq 1$ . Furthermore, all vertices in  $V_2$  are 3-vertices adjacent to  $v^*$  and to a  $2^-$ -vertex in  $S_1$ , while all vertices in  $V_1$  are  $2^-$ -vertices adjacent to  $v^*$ . Also, we have  $d_1^+ + d_2^+ \leq d_G(v^*) \leq d_1^+ + d_2^+ + 2$ .

First assume that  $d_1^+ + d_2^+ \geq 4$ , and let  $v_1, v_2, v_3, v_4$  be any four distinct neighbours of  $v^*$  in  $V_1 \cup V_2$ . We here set  $G' = G - \{v^*v_1, v^*v_2, v^*v_3, v^*v_4\}$ . Since the  $v_i$ 's are  $3^-$ -vertices in  $G$ , it should be clear, according to Claims 5.2 and 5.3, that  $G'$  is nice. As it is also 2-degenerated, by minimality of  $G$ , there exists an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting  $w_{G'}$ , where  $m_{G'} := m_G - 4$ .

We now have to prove that we can extend  $w_{G'}$  to  $w_G$ , using at most eight distinct weights, including those among  $\{m_G + 1, m_G + 2, m_G + 3, m_G + 4\}$ . Since  $d_2^+ \geq 1$ , some of the  $v_i$ 's belong to  $V_2$ ; assume  $v_1$  is one such vertex. We first assign a weight  $\alpha_1$  to  $v^*v_1$ , from the set  $\{m_G + 2, m_G + 3, m_G + 4\}$ , so that no conflict involving  $v_1$  and one of its two neighbours different from  $v^*$  arises. This is clearly possible with at least three distinct weights. Similarly, we assign two weights  $\alpha_2$  and  $\alpha_3$ , from the set  $\{m_G + 1, m_G + 2, m_G + 3, m_G + 4\} \setminus \{\alpha_1\}$ , to  $v^*v_2$  and  $v^*v_3$ , respectively, so that no conflict involving  $v_2$  or  $v_3$  and one of their at most two neighbours different from  $v^*$  arises. We note that this is possible since, though  $v_2$  and  $v_3$  might be 3-vertices, they are adjacent to a 2-vertex in that case. Under the assumption that we assign a weight among  $\{m_G + 1, m_G + 2, m_G + 3, m_G + 4\}$  to  $v^*v_2$  and  $v^*v_3$ , we cannot create any sum conflict involving  $v_2$  or  $v_3$  and a neighbouring 2-vertex. In other words, only one conflict involving  $v_2$  or  $v_3$  may arise here.

We finally have to extend  $w_{G'}$  to  $v^*v_4$ . Note that, due to the choice of  $\alpha_1, \alpha_2, \alpha_3$ , and because  $v_4$  is a  $3^-$ -vertex in  $G$ , it cannot be that, currently, the sum of weights incident to  $v^*$  is exactly the sum of weights incident to  $v_4$ . Furthermore, for the same reasons, no matter how we weight  $v^*v_4$ , it cannot happen that, eventually,  $\sigma_{w_G}(v^*)$  gets equal to the sum of weights incident to any vertex in  $V_1 \cup V_2$ . Hence, when weighting  $v^*v_4$ , we just have to make sure that  $\sigma_{w_G}(v^*)$  does not get equal to the sums of weights incident to the at most two other neighbours of  $v^*$  (i.e. those not in  $V_1 \cup V_2$ , unless  $G_2$  is empty in which case all neighbours of  $v^*$  belong to  $V_1 \cup V_2$ ), and that  $\sigma_{w_G}(v_4)$  does not get equal to the sums of weights incident to the at most two neighbours of  $v_4$  different from  $v^*$ . Since we have five distinct weights left to weight  $v^*v_4$ , necessarily one of these weights respect these conditions. The claimed extension of  $w_{G'}$  hence exists.

To complete the proof, we have to consider the cases where  $d_1^+ + d_2^+ \leq 3$ . Denote by  $v_1$  one neighbour of  $v^*$  in  $V_2$ , which exists since  $d_2^+ \geq 1$ . Since  $v_1$  belongs to  $V_2$ , we know that  $v_1$  is a 3-vertex adjacent to a 2-vertex, say  $u_1$ , in  $S_1$ . Set  $G' := G - \{v^*v_1, v_1u_1\}$ . Again,  $G'$  is nice according to Claims 5.2 and 5.3, and 2-degenerate. So let  $w_{G'}$  be an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 4)$ -edge-weighting of  $G'$ , which exists due to the minimality of  $G$ , where  $m_{G'} := m_G - 2$ . For contradiction, we show that  $w_{G'}$  can be extended to  $G$ , and that we can do it with six distinct weights.

The degree properties here, are that  $d_G(v^*) \leq 5$ ,  $d_G(v_1) = 3$  and  $d_G(u_1) = 2$ . It can be observed, under those assumptions, that the quantity

$$\mu := (d_G(v^*) + 1) + \max\{0, d_G(v_1) + d_G(u_1) - d_G(v^*) - 1\}$$

is bounded above by 6. From Observation 2.3, we hence know that  $w_{G'}$  can be extended to  $v^*v_1$  and  $v_1u_1$ , as claimed. This completes the proof.  $\square$



## 5.2. Graphs with maximum average degree at most 3

We recall that, for any given graph  $G$ , the *maximum average degree* of  $G$ , denoted  $\text{mad}(G)$ , is defined as the maximum average degree of a subgraph of  $G$ , that is

$$\text{mad}(G) := \max \left\{ \frac{2|E(H)|}{|V(H)|} : H \text{ is a non-empty subgraph of } G \right\}.$$

In the next result, we prove an upper bound on  $\chi_{\Sigma}^{e,1}$  for every nice graph with maximum average degree at most 3.

**Theorem 5.4.** *For every nice graph  $G$  with  $\text{mad}(G) \leq 3$ , we have  $\chi_{\Sigma}^{e,1}(G) \leq |E(G)| + 6$ .*

*Proof.* Assume there exists a counterexample to the claim, that is, there exists a nice graph  $G$  for which we have  $\text{mad}(G) \leq 3$ , but  $\chi_{\Sigma}^{e,1}(G) > |E(G)| + 6$ . We consider  $G$  minimum in terms of  $n_G + m_G$ , where  $n_G := |V(G)|$  and  $m_G := |E(G)|$ . Our ultimate goal in this proof, is to show that  $G$  cannot exist. The strategy we employ to this end, is essentially to show that  $G$  has a nice subgraph  $H$ , with order  $n_H$  and size  $m_H$ , such that  $H$  has an edge-injective neighbour-sum-distinguishing  $(m_H + 6)$ -edge-weighting  $w_H$  that can be extended to an edge-injective neighbour-sum-distinguishing  $(m_G + 6)$ -edge-weighting  $w_G$  of  $G$ , contradicting the fact that  $G$  is a counterexample. The main tool we want to use, in order to show that  $H$  has such an edge-weighting, is Theorem 4.2. Since  $G$  is a counterexample to the claim, note that Theorem 4.2 already implies that  $\Delta(G) \geq 4$ .

The subgraph  $H$  we consider, is obtained by removing all 1-vertices from  $G$ . Of course, we have  $\text{mad}(H) \leq 3$ , and it may happen that  $G = H$ . We may as well assume that  $H$  remains nice, as, if it is not the case, then  $G$  would be a tree (a bistar) and hence  $\chi_{\Sigma}^{e,1}(G) \leq m_G$ , according to Theorem 3.3. We may hence further assume that  $G$  is not a tree.

In the following result, we observe that, by showing that  $H$  verifies  $\Delta(H) \leq 3$ , then we will get our conclusion.

**Proposition 5.5.** *If  $\Delta(H) \leq 3$ , then  $\chi_{\Sigma}^{e,1}(G) \leq m_G + 6$ , and  $G$  is not a counterexample.*

*Proof.* If  $G = H$ , then  $G$  admits an edge-injective neighbour-sum-distinguishing  $(m_G + 6)$ -edge-weighting, according to Theorem 4.2, since we would have  $\Delta(G) \leq 3$ . So assume that  $G$  has 1-vertices. Since we assume that  $\Delta(H) \leq 3$ , there exists an edge-injective neighbour-sum-distinguishing  $(m_H + 6)$ -edge-weighting  $w_H$  of  $H$ , still according to Theorem 4.2.

We now extend  $w_H$  to the pendant edges of  $G$ . We successively consider every vertex  $v$  of  $H$  incident to a pendant edge. We start by assigning an arbitrary non-used weight to every pendant edge incident to  $v$ , but one, say  $vu$ .

We claim that we can find a correct weight for  $vu$ . First, we note, according to Observation 2.1, that only the neighbours of  $v$  in  $H$  can eventually cause sum conflicts. Hence, when extending  $w_H$  to  $vu$ , we just have to make sure, since  $vu$  is the last non-weighted pendant edge incident to  $v$ , that  $\sigma(v)$  does not meet any of the determined sums of the vertices adjacent to  $v$  in  $H$ . By our assumption on  $\Delta(H)$ , there are at most three such vertices, while we have at least seven ways to weight  $vu$  (among  $\{1, \dots, m_G + 6\}$ ), each determining a distinct value for  $\sigma(v)$ . We can hence find a correct non-used weight for  $vu$ .

Since the process above can be applied for all vertices of  $H$  incident to a pendant edge in  $G$ , weighting  $w_H$  can hence be extended to all pendant edges of  $G$ . Thus  $w_H$  can be extended to an edge-injective neighbour-sum-distinguishing  $(m_G + 6)$ -edge-weighting of  $G$ , as claimed.  $\square$

It remains to show that  $\Delta(H) \leq 3$ . This is proved by getting successive information concerning the structure of  $H$ , so that classical discharging arguments can eventually be employed.

**Claim 5.6.** *If  $v \in V(H)$  is adjacent to 1-vertices in  $G$ , then  $d_H(v) \geq 7$ .*

*Proof.* This follows from Observation 2.4, as, when removing a pendant edge from  $G$ , applying induction, and putting the edge back, we then have seven distinct weights to achieve the extension to  $G$ .  $\square$

**Claim 5.7.** *We have  $\delta(H) \geq 2$ .*

*Proof.* If  $\delta(H) = 0$ , then  $G$  is a star, contradicting one of our initial assumptions. Now, if  $\delta(H) = 1$ , then  $G$  includes a vertex  $v$  such that  $d_H(v) = 1$  and  $v$  is incident to pendant edges in  $G$ . But this is impossible, as such a  $v$  would not meet the condition in Claim 5.6. So  $\delta(H) \geq 2$ .  $\square$

**Claim 5.8.** *Graph  $H$  has no two adjacent 2-vertices.*

*Proof.* Suppose that  $H$  has an edge  $uv$  such that  $d_H(u) = d_H(v) = 2$ . Recall that, according to Claim 5.6, we have  $d_G(u) = d_G(v) = 2$ . In this case, we consider the graph  $G' := G - uv$  with size  $m_{G'} := |E(G')|$ . Clearly  $G'$  remains nice (otherwise Claim 5.6 would be violated), has  $\text{mad}(G') \leq 3$ , and, due to the minimality of  $G$ , graph  $G'$  admits an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 6)$ -edge-weighting  $w_{G'}$ .

In  $G'$ , we have  $d_{G'}(u) = d_{G'}(v) = 1$ . Let  $u'$  and  $v'$  be the neighbours of  $u$  and  $v$ , respectively, in  $G'$ . Since  $w_{G'}$  is edge-injective, we have  $w_{G'}(uu') \neq w_{G'}(vv')$ . We now note that, under all those assumptions, weighting  $w_{G'}$  can easily be extended to an edge-injective neighbour-sum-distinguishing  $(m_G + 6)$ -edge-weighting  $w_G$  of  $G$ , i.e. to the edge  $uv$ , a contradiction. We note that, because  $w_{G'}(uu') \neq w_{G'}(vv')$  and  $d_G(u) = d_G(v) = 2$ , we cannot get  $\sigma_{w_G}(u) = \sigma_{w_G}(v)$  when assigning any weight to  $uv$ , recall Observation 2.1. So the only constraints we have, are that  $\sigma_{w_G}(u)$  has to be different from  $\sigma_{w_G}(u')$  (which is exactly  $\sigma_{w_{G'}}(u')$ ) and  $\sigma_{w_G}(v)$  must be different from  $\sigma_{w_G}(v')$  (which is exactly  $\sigma_{w_{G'}}(v')$ ). These constraints forbid us from assigning, to  $uv$ , at most two of the seven weights that have not been used yet. So we can extend  $w_{G'}$  to  $w_G$ .  $\square$

**Claim 5.9.** *Graph  $H$  has no 2-vertex adjacent to two 3-vertices.*

*Proof.* Assume  $H$  has such a vertex  $v$  with  $d_H(v) = 2$ , and  $v$  has two neighbours  $u_1$  and  $u_2$  verifying  $d_H(u_1) = d_H(u_2) = 3$ . According to Claim 5.6, we have  $d_G(v) = 2$ ,  $d_G(u_1) = 3$  and  $d_G(u_2) = 3$ . Let  $G' := G - \{vu_1, vu_2\}$  and  $m_{G'} := |E(G')|$ . Clearly,  $G'$  remains nice, with  $\text{mad}(G') \leq 3$ , and, by the minimality of  $G$ , there exists an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 6)$ -edge-weighting  $w_{G'}$ . According to Observation 2.3, weighting  $w_{G'}$  can be extended to an edge-injective neighbour-sum-distinguishing edge-weighting of  $G$ , provided we have at least five distinct edge weights in hand. Since we here have eight non-used edge weights dedicated to weighting  $vu_1$  and  $vu_2$ , the extension of  $w_{G'}$  to  $G$  hence exists.  $\square$

**Claim 5.10.** *Graph  $H$  has no 3-vertex adjacent to two  $3^-$ -vertices.*

*Proof.* The proof is similar to that of the previous claim. Assume  $H$  has such a 3-vertex  $v$ , being adjacent to at least two  $3^-$ -vertices  $u_1$  and  $u_2$ . Again, we set  $G' := G - \{vu_1, vu_2\}$ , and let  $w_{G'}$  be an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 6)$ -edge-weighting of  $G'$ , where  $m_{G'} := |E(G')|$ . Still according to Observation 2.3, we know that an extension

exists, provided we have at least six weights available. So  $w_{G'}$  can correctly be extended to  $vu_1$  and  $vu_2$ , as eight edge weights can be used in the present context.  $\square$

Before getting our conclusion, we prove two last claims, which are a bit more general than what we actually need.

**Claim 5.11.** *Graph  $H$  has no 6-vertex adjacent to two 2-vertices.*

*Proof.* Assume  $H$  has such a 6-vertex  $v$ , and let  $u_1$  and  $u_2$  denote any two of its neighbouring 2-vertices. Recall that  $d_H(v) = d_G(v)$ ,  $d_H(u_1) = d_G(u_1)$  and  $d_H(u_2) = d_G(u_2)$ , according to Claim 5.6. Let  $G' := G - \{vu_1, vu_2\}$ , and set  $n_{G'} := |V(G')|$  and  $m_{G'} := |E(G')|$ . Clearly  $G'$  is nice (Claims 5.6 and 5.8), with  $\text{mad}(G') \leq 3$ , and, since  $n_{G'} + m_{G'} < n_G + m_G$ , there exists an edge-injective neighbour-sum-distinguishing  $(m_{G'} + 6)$ -edge-weighting  $w_{G'}$  of  $G'$ . Again according to Observation 2.3, under these conditions, we know that  $w_{G'}$  can be extended to  $vu_1$  and  $vu_2$  provided we have at least eight weights available. Since this is precisely the case, we are done.  $\square$

**Claim 5.12.** *Graph  $H$  has no 4- or 5-vertex adjacent to at least two  $3^-$ -vertices.*

*Proof.* The proof is very similar to that of Claim 5.11, and can be mimicked by just letting  $u_1$  and  $u_2$  be two  $3^-$ -vertices adjacent to  $v$ . We then get the same conclusion, from Observation 2.3.  $\square$

We are now ready to prove that  $H$  has maximum degree 3.

**Claim 5.13.** *We have  $\Delta(H) \leq 3$ .*

*Proof.* Assume the contrary, namely that  $\Delta(H) \geq 4$ . We prove the claim by means of the so-called discharging method, through a discharging procedure, based on the following rules.

To every vertex  $v$  of  $H$ , we assign an initial charge  $\omega(v)$  being  $d_H(v) - 3$ . Since  $\text{mad}(H) \leq 3$ , we have

$$\sum_{v \in V(H)} d_H(v) \leq 3 \cdot n_H,$$

which implies that

$$\sum_{v \in V(H)} \omega(v) \leq 0.$$

Without creating or deleting any amount of charge assigned to the vertices, we now transfer a part of the assigned charges from neighbours to neighbours, through three discharging rules applied in two successive steps.

In the sequel, by a *weak 3-vertex* of  $H$ , we refer to a 3-vertex neighbouring a 2-vertex (recall that a 3-vertex of  $H$  is adjacent to at most one 2-vertex, according to Claim 5.10). The first discharging step, consists in applying the following rule:

- (R1) Every  $4^+$ -vertex transfers  $\frac{1}{4}$  to every adjacent weak 3-vertex.

Once the first discharging step has been performed, we then apply the second step, which consists in applying the following two discharging rules:

- (R2) Every weak 3-vertex transfers  $\frac{1}{2}$  to every adjacent 2-vertex.  
(R3) Every  $4^+$ -vertex transfers  $\frac{1}{2}$  to every adjacent 2-vertex.

We now compute the final charge  $\omega^*(v)$  that every vertex  $v$  of  $H$  gets, once the two steps above have been performed. Recall that  $\delta(H) \geq 2$ , according to Claim 5.7.

1. If  $v$  is a 2-vertex, then  $v$  is adjacent to a  $4^+$ -vertex, and either a weak 3-vertex or a  $4^+$ -vertex, according to Claims 5.8 and 5.9. Through Rules (R2) and (R3), the two neighbours of  $v$  both transfer  $\frac{1}{2}$  to  $v$ . Hence,  $\omega^*(v) = \omega(v) + 2 \times \frac{1}{2} = 0$ .
2. If  $v$  is a 3-vertex, then  $v$  is either weak, or not. If  $v$  is not weak, it is not concerned by any of Rules (R1), (R2) and (R3), so  $\omega^*(v) = \omega(v) = 0$ . Now assume  $v$  is a weak 3-vertex. According to Claim 5.10, vertex  $v$  is adjacent to a 2-vertex  $u$ , and two  $4^+$ -vertices  $z_1$  and  $z_2$ . Through Rule (R1), vertex  $v$  receives  $\frac{1}{4}$  from each of  $z_1$  and  $z_2$ , while, through Rule (R2), vertex  $v$  then transfers  $\frac{1}{2}$  to  $u$ . Therefore,  $\omega^*(v) = \omega(v) + 2 \times \frac{1}{4} - \frac{1}{2} = 0$ .
3. If  $v$  is a 4- or 5-vertex, then  $v$  is adjacent to at most one vertex being either a 2-vertex or weak 3-vertex  $u$ , according to Claim 5.12. The case where  $\omega^*(v)$  is minimum, is basically when  $v$  is a 4-vertex, and  $u$  is a 2-vertex, in which case  $v$  transfers  $\frac{1}{2}$  to  $u$ . In that case, through Rule (R3), we get  $\omega^*(v) = \omega(v) - \frac{1}{2} = \frac{1}{2}$ . So, whenever  $v$  is a 4- or 5-vertex, we get  $\omega^*(v) > 0$ .
4. If  $v$  is a 6-vertex, then  $v$  is adjacent to at most one 2-vertex, according to Claim 5.11. The case where  $\omega^*(v)$  gets minimum, is essentially when  $v$  neighbours one 2-vertex and five weak 3-vertices. In that case, following Rules (R1) and (R3), we get  $\omega^*(v) = \omega(v) - 5 \times \frac{1}{4} - \frac{1}{2} = \frac{5}{4}$ . Hence, we always get  $\omega^*(v) > 0$  in that case.
5. If  $v$  is a  $7^+$ -vertex, then  $v$  transfers most charge when  $v$  is adjacent to  $d_H(v)$  2-vertices. In that case, following Rule (R3), we deduce that  $\omega^*(v) = \omega(v) - d_H(v) \times \frac{1}{2}$ . Under the assumption that  $d_H(v) \geq 7$ , observe that  $\omega(v) > d_H(v) \times \frac{1}{2}$ . So, again, we always have  $\omega^*(v) > 0$  in this case.

From the analysis above, we get, because  $\Delta(H) \geq 4$ , that

$$\sum_{v \in V(H)} \omega(v) \leq 0 < \sum_{v \in V(H)} \omega^*(v),$$

which is impossible, as we did not create any new amount of charge when applying the discharging procedure. Hence, we have  $\Delta(H) \leq 3$ . □

The result now follows from Proposition 5.5. □

Theorem 5.4 applies to all nice graphs with maximum average degree at most 3. Among the classes of such graphs, we would like to highlight the class of nice planar graphs with girth at least 6, where the *girth*  $g(G)$  of a graph  $G$ , is the length of its smallest cycles. We refer the reader to e.g. [2], wherein the authors noticed that, for every planar graph  $G$ , we have

$$\text{mad}(G) < \frac{2g(G)}{g(G) - 2}.$$

This gives that every planar graph  $G$  with  $\text{mad}(G) \leq 3$ , indeed has  $g(G) \geq 6$ .

**Corollary 5.14.** *For every nice planar graph  $G$  with  $g(G) \geq 6$ , we have  $\chi_{\Sigma}^{e,1}(G) \leq |E(G)| + 6$ .*

## 6. Discussion

In this work, we have introduced and studied Conjecture 1.1, which stands, in some sense, as a combination of the 1-2-3 Conjecture and the Antimagic Labelling Conjecture. In particular, as a support to Conjecture 1.1, we have pointed out that some families of nice graphs agree with it, or sometimes almost agree with it, i.e. up to an additive constant term. Although these results can be regarded as a first step towards Conjecture 1.1, it is worth emphasizing that our work does not bring anything new towards attacking the 1-2-3 Conjecture and the Antimagic Labelling Conjecture, but rather concerns some side aspects of these two conjectures.

As further work towards Conjecture 1.1, it would be interesting exhibiting, for all nice graphs  $G$ , bounds on  $\chi_{\Sigma}^{e,1}(G)$  of the form  $|E(G)| + k$  for a fixed constant  $k$ . One could as well try to get a better bound of the form  $k \cdot |E(G)|$ , for some  $k$  in between 1 and 2. Obtaining one such of these two bounds, would already improve the ones we have exhibited in Section 4. It is worth mentioning that our bounds in that section, can slightly be improved by making some choices in a more clever way. But these improvements would allow us to save a small constant number of weights only, which is far from the desired improvement we have mentioned earlier.

As another direction, we would also be interested in knowing other classes of nice graphs agreeing with Conjecture 1.1, and being not known to be antimagic yet. Among such classes, let us mention the case of nice bipartite graphs  $G$ , for which we did not manage to come up with an  $|E(G)| + k$  bound on  $\chi_{\Sigma}^{e,1}(G)$ , for any constant  $k$ . Another such class that would be interesting investigating, is the one of nice subcubic graphs. We already know that cubic graphs agree with Conjecture 1.1, recall Theorem 3.1. Furthermore, we also know that nice subcubic graphs  $G$ , in general, verify  $\chi_{\Sigma}^{e,1}(G) \leq |E(G)| + 6$ , recall Theorems 4.2 and 5.4. It nevertheless does not seem obvious how these results can be used, in order to show that nice subcubic graphs agree with Conjecture 1.1. Such a result, though, would be one natural step following Observation 3.2. Nice planar graphs would also be interesting candidates to investigate, as we have been mostly successful with sparse classes of nice graphs. Our result in Corollary 5.14, may be regarded as a first step towards that direction.

Our results in this paper may also be subject to further investigations. In particular, there is still a gap, for nice 2-degenerate graphs and graphs with maximum average degree at most 3, between our bounds in Section 5 and the bound in Conjecture 1.1. One could as well wonder how to generalize our results to nice  $k$ -degenerate graphs and graphs with maximum average degree at most  $k$ , for larger fixed values of  $k$ . In particular, it could be interesting to exhibit, for these graphs  $G$ , a general upper bound on  $\chi_{\Sigma}^{e,1}(G)$  of the form  $|E(G)| + \mathcal{O}(k)$ , involving a small function of  $k$ .

## Acknowledgements

This work was initiated when the second author visited the Technical University of Denmark. Support from the ERC Advanced Grant GRACOL, project no. 320812, is gratefully acknowledged.

## References

- [1] O. Baudon, M. Piłśniak, J. Przybyło, M. Senhaji, É. Sopena, M. Woźniak. Equitable neighbour-sum-distinguishing edge and total colourings. hal-01330531, 2016.

- [2] O.V. Borodin, A.V. Kostochka, J. Nešetřil, A. Raspaud, É. Sopena. On the maximum average degree and the oriented chromatic number of a graph. *Discrete Mathematics*, 206:77-89, 1999.
- [3] J.A. Gallian. A Dynamic Survey of Graph Labeling. *Electronic Journal of Combinatorics*, DS6, 1997.
- [4] N. Hartsfield, G. Ringel. *Pearls in Graph Theory*. Academic Press, San Diego, 1990.
- [5] M. Kalkowski, M. Karoński, F. Pfender. Vertex-coloring edge-weightings: towards the 1-2-3 Conjecture. *Journal of Combinatorial Theory, Series B*, 100:347-349, 2010.
- [6] M. Karoński, T. Łuczak, A. Thomason. Edge weights and vertex colours. *Journal of Combinatorial Theory, Series B*, 91:151–157, 2004.
- [7] B. Seamone. The 1-2-3 Conjecture and related problems: a survey. arXiv:1211.5122, 2012.