



HAL
open science

A polynomial-time recursive algorithm for some unconstrained quadratic optimization problems

Walid Ben-Ameur, José Neto

► **To cite this version:**

Walid Ben-Ameur, José Neto. A polynomial-time recursive algorithm for some unconstrained quadratic optimization problems. CTW 2009 : 8th Cologne-Twente Workshop on Graphs and Combinatorial Optimization, Jun 2009, Paris, France. pp.105 - 108. hal-01368385

HAL Id: hal-01368385

<https://hal.science/hal-01368385v1>

Submitted on 19 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A polynomial-time recursive algorithm for some unconstrained quadratic optimization problems

Walid Ben-Ameur,^a José Neto^a

^a*Institut TELECOM, TELECOM & Management SudParis, CNRS Samovar,
9, rue Charles Fourier, 91011 Evry, France
{Walid.Benameur, Jose.Neto}@it-sudparis.eu*

Key words: arrangements, unconstrained quadratic programming, complexity

1. Introduction

Consider a quadratic function $q : \mathbb{R}^n \rightarrow \mathbb{R}$ given by: $q(x) = x^t Q x$, with $Q \in \mathbb{R}^{n \times n}$. An unconstrained $(-1, 1)$ -quadratic optimization problem can be expressed as follows:

$$(QP) Z^* = \min\{q(x) \mid x \in \{-1, 1\}^n\},$$

where $\{-1, 1\}^n$ denotes the set of n -dimensional vectors with entries either equal to 1 or -1 . We consider here that the matrix Q is symmetric and given by its spectrum, i.e. the set of its eigenvalues and associated unit pairwise orthogonal eigenvectors.

Problem (QP) is a classical combinatorial optimization problem with many applications, e.g. in statistical physics and circuit design [2; 8; 10]. It is well-known that any $(0,1)$ -quadratic problem expressed as: $\min\{x^t A x + c^t x \mid x \in \{0, 1\}^n\}$, $A \in \mathbb{R}^{n \times n}$, $c \in \mathbb{R}^n$, can be formulated in the form of problem (QP) and conversely [9; 4].

The contribution of this work is 3-fold:

- (i) We slightly extend the known polynomially solvable cases of (QP) to when the matrix Q has fixed rank and the number of positive diagonal entries is $\mathcal{O}(\log(n))$.
- (ii) We introduce a new (to our knowledge) polynomial-time algorithm for solving problem (QP) when it corresponds to such a polynomially solvable case.
- (iii) Preliminary experiments indicate that the proposed method may be computationally efficient. [7].

2. Properties of optimal solutions for peculiar instances of (QP)

Let us firstly introduce some notation to be used hereafter. The eigenvalues of the matrix Q will be noted $\lambda_1(Q) \leq \lambda_2(Q) \leq \dots \leq \lambda_n(Q)$ (or more simply $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ when clear from the context) and the corresponding unit (in Euclidean norm) and pairwise orthogonal eigenvectors: v_1, \dots, v_n . The j -th entry of the vector v_i is noted v_{ij} . Given some set of vectors $a_1, \dots, a_q \in \mathbb{R}^n$, $q \in \mathbb{N}$, we note $Lin(a_1, \dots, a_q)$ the subspace spanned by these vectors.

In this section we shall make the following assumptions on the matrix Q :

- (i) Q has rank $p \leq n$,
- (ii) Q has nonpositive diagonal entries only, and
- (iii) Q is given by its set of rational eigenvalues and eigenvectors: $Q = \sum_{i=1}^p \lambda_i v_i v_i^t$.

Any optimal solution y^* to the problem $\min_{y \in \{-1, 1\}^n} y^t Q y$ can be shown to satisfy the following implication:

$$\sum_{i=1}^p \lambda_i \alpha_i v_{ij} > 0 \Rightarrow y_j^* = -1 \quad (2.1)$$

And analogously:

$$\sum_{i=1}^p \lambda_i \alpha_i v_{ij} < 0 \Rightarrow y_j^* = 1. \quad (2.2)$$

From this simple property we can namely show that in order to find an optimal solution of problem (QP) , it is sufficient to enumerate over all vectors $y \in \{-1, 1\}^n$ for which there exists a vector $\alpha \in \mathbb{R}^p$ such that $y_j = -\text{sign}(\sum_{i=1}^p \lambda_i \alpha_i v_{ij})$ (or equivalently $y_j = \text{sign}(\sum_{i=1}^p \lambda_i \alpha_i v_{ij})$, see hereafter), $\sum_{i=1}^p \lambda_i \alpha_i v_{ij} \neq 0, \forall j \in \{1, \dots, n\}$, with $\text{sign}(x) = 1$ if $x > 0$ and -1 if $x < 0$. In the next section we focus on finding such a set of vectors.

3. Determining cells in an arrangement of n hyperplanes

Let $v_1, \dots, v_p \in \mathbb{R}^n$ denote p independent vectors. Let $V \in \mathbb{R}^{n \times p}$ denote the matrix whose columns correspond to the vectors v_1, \dots, v_p and V_i the i -th row of V . From this set of vectors we define n hyperplanes in \mathbb{R}^p : $H_j = \{\alpha \in \mathbb{R}^p \mid V_j \cdot \alpha = 0\}$ with $j \in \{1, \dots, n\}$. Then we can notice that there is a one-to-one correspondence between the set of vectors in $\{-1, 1\}^n$ for which there exists a vector $\alpha \in \mathbb{R}^p$ such that $y_j = \text{sign}(\sum_{i=1}^p \alpha_i v_{ij})$, with $\sum_{i=1}^p \alpha_i v_{ij} \neq 0, \forall j \in \{1, \dots, n\}$ and the cells (i.e. the full dimensional regions) in \mathbb{R}^p of the hyperplane arrangement $\mathcal{A}(H)$ that is defined by the family of hyperplanes $(H_j)_{j=1}^n$. To see this just interpret the sign vector y as the position vector of the corresponding cell c w.r.t. an orientation of

the space by the vector V_j : cell c is *above* hyperplane H_j iff $y_j > 0$ and *under* otherwise.

For a general arrangement in \mathbb{R}^p that is defined by n hyperplanes (see e.g. [6; 11] for further elements on arrangements), the number of cells is upper bounded by $\sum_{i=0}^p \binom{n}{i}$ (which is in $\mathcal{O}(n^p)$). (For a proof we refer the reader e.g., to Lemma 1.2 in [6]). In our case, since all the hyperplanes considered contain the origin (i.e. the arrangement is *central*), this number reduces to $\mathcal{O}(n^{p-1})$ (see Section 1.7 in [6]).

We have introduced [3] a simple procedure with time complexity lying between the time complexity of the incremental algorithm [6; 5] (in $\mathcal{O}(n^{p-1})$) and that of the reverse search algorithm [1; 7] (in $\mathcal{O}(n LP(n, p) C)$ where C denotes the number of cells in $\mathcal{A}(H)$ and $LP(n, p)$ is the time needed to solve a linear program with n inequalities and p variables) in order to compute a set of vectors in $\{-1, 1\}^n$ corresponding to a description of a set containing the cells of the arrangement $\mathcal{A}(H)$. Space complexity can be shown to be polynomially bounded by the output size. To our view the interest of the proposed method by comparison with the former ones is 2-fold:

- (i) it is very easy to understand and implement,
- (ii) computationally, by using proper data structures (to be specified latter) we could solve instances of the same magnitude as the ones reported in [7], without parallelization and substantially improved computation times.

The basic principle of the proposed method may be expressed as follows. Given some integer $q \in \{1, \dots, n\}$, let $\mathcal{B}^q(H)$ denote the arrangement in the subspace $\{\alpha \in \mathbb{R}^p \mid V_q \alpha = 0\}$ that is defined by the hyperplanes (in \mathbb{R}^{p-1}) $\{H_j \cap H_q \mid j \in \{1, \dots, n\} \text{ and } j \neq q\}$. Any cell of $\mathcal{B}^q(H)$ (which is a region of dimension $p - 1$) corresponds to a facet of exactly two cells of the arrangement $\mathcal{A}(H)$ i.e. one on each side of the hyperplane H_q . The other cells of $\mathcal{A}(H)$ i.e. those not intersecting H_q , are cells of the arrangement $\mathcal{C}^q(H)$ in \mathbb{R}^p which is defined by the $n - 1$ hyperplanes $\{H_j \mid j \in \{1, \dots, n\} \text{ and } j \neq q\}$. Since each cell of $\mathcal{A}(H)$ intersects at least one of the hyperplanes $(H_j)_{j=1}^n$, it follows that all the cells of $\mathcal{A}(H)$ can be derived from the ones of all the arrangements $\mathcal{B}^q(H)$, $q = 1, \dots, n$. A recursive use of this argument leads to the generation of all the cells of $\mathcal{A}(H)$.

From a complexity study of the proposed method we can show the following result.

Theorem 3.1. For a fixed integer $p \geq 2$, if the matrix Q (given by its nonzero eigenvalues and associated eigenvectors) has rank at most p and $\mathcal{O}(\log(n))$ positive diagonal entries, then problem (QP) can be solved in strongly polynomial time.

4. Conclusion

We propose a new (up to our knowledge) approach for solving in polynomial time some unconstrained quadratic optimization problems. Preliminary computational results illustrate that the recursive procedure briefly presented here can be a valuable approach on some instances by comparison with a reverse search w.r.t. computation times.

Further computational studies are under work and could involve a parallelization of the code in order to deal with larger instances.

References

- [1] D. Avis and K. Fukuda: Reverse search for enumeration. *Discrete Applied Mathematics* **65**, 21-46 (1996)
- [2] F. Barahona and M. Grötschel and M. Jünger and G. Reinelt: An application of Combinatorial Optimization to Statistical Physics and Circuit Layout Design. *Operations Research* **36**, 493–513 (1988)
- [3] W. Ben-Ameur and J. Neto: Spectral bounds for the maximum cut problem. Institut TELECOM, TELECOM & Management SudParis, Evry. Rapport de recherche 08019-RS2M (2008)
- [4] C. De Simone: The cut polytope and the Boolean quadric polytope. *Discrete Mathematics* **79**, 71-75 (1990)
- [5] H. Edelsbrunner and J. O'Rourke and R. Seidel: Constructing arrangements of lines and hyperplanes with applications. *SIAM Journal on Computing* **15**, 341-363 (1986)
- [6] H. Edelsbrunner: *Algorithms in Combinatorial Geometry*. Springer (1987)
- [7] J.A. Ferrez and K. Fukuda and T.M. Liebling: Solving the fixed rank convex quadratic maximization in binary variables by a parallel zonotope construction algorithm. *European Journal of Operational Research* **166**(1), 35-50 (2005)
- [8] M. Grötschel and M. Jünger and G. Reinelt: Via minimization with pin preassignment and layer preference. *Zeitschrift für Angewandte Mathematik und Mechanik*, **69**, 393–399 (1989)
- [9] P. Hammer: Some network flow problems solved with pseudo-Boolean programming. *Operations Research* **32**, 388-399 (1965)
- [10] R.Y. Pinter: Optimal Layer Assignment for Interconnect. *J. VLSI Comput. Syst.* **1**, 123–137 (1984)
- [11] G. M. Ziegler: *Lectures on Polytopes*, Graduate Texts in Mathematics **152**. Springer-Verlag New-York (1995)