

Geomorphic impacts of the April 25, 2015 earthquake in Nepal: Preliminary assessments

Monique Fort

▶ To cite this version:

Monique Fort. Geomorphic impacts of the April 25, 2015 earthquake in Nepal: Preliminary assessments. 5th National Meeting of the Italian Association of Physical Geography and Geomorphology: "Geomorphology for Society from risk knowledge to landscape heritage", Sep 2015, Cagliari, Italy. hal-01373154

HAL Id: hal-01373154 https://hal.science/hal-01373154

Submitted on 28 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



GEOMORPHIC IMPACTS OF THE APRIL 25, 2015 EARTHQUAKE IN NEPAL: PRELIMINARY ASSESSMENTS, A REVIEW by Monique FORT

CHINA

Dept of Geography, CNRS UMR 8586 PRODIG, CC. 7001, Univ. Paris-Diderot-SPC, F - 75 251 PARIS Cedex 05, France (fort@univ-paris-diderot.fr)

TIBET

DAKISTA

PLATEAU

1. THE EVENT

A magnitude Mw 7.8 earthquake struck Central Nepal on April 25, 2015 at 11:41 am (local time), followed by another Mw 7.4 one on May 10, 2014 at 12:30 pm (local time). The first rupture took place along the Main Frontal Thrust, a main structure interface between Indian and Eurasian plates; it has occurred at a depth varying from 10 km (USGS) to 29 km (Geoscope) corresponding to a relative slip of 4-5 m. It was a forseeable disaster, with continuous seismic monitoring (Seismology Laboratory of Kathmandu) and preparedness actions such as infrastructure strengthening, development of early warning systems, plans for evacuation and recovery (thanks to the NSET NGO). Still it was very damageous.

About ¼ Nepal population was affected (~8 M people), both in the mountains and Kathmandu vallev (~2,5 M people). ~9000 people lost their lives. The economic damage -about \$10 billion- is almost half of the country's GDP.





2. GEOMORPHIC IMPACTS Though many fatalities are due to buildings collapses, a large part of them are also related to slope processes: snow avalanches, rock falls (including very large blocks), debris avalanches and landslides.





AlGeo

ICIMOD

25 April 2015 Nepal Earthquake Landslide Intensity

GEOMORPHOLOGY GEOMORPHOLOGIE

IAG MEMBERS



Aggravating factors are many folds. (1) In the Kathmandu valley, the Quaternary lacustrine substrate amplified the vibrations, hence substantially damaged the buildings. (2) After a dry winter, snowfalls and rainfalls were abundant during March and April, hence increasing the pore pressure and the potential instability of slopes and/or snow cover. (3 During the last decades, the construction of earthy, fragile roads accentuated, slope steepness. (4) Very short time left between the earthquake and the onset of the next Indian monsoon rains.





how the ground shifted during the earthquake. In addition to vertical movements, the researchers also detected north-south horizontal shifts of nearly 2 meters (German Aerospace Center - DLR)

onsoon rains: another aggravating factor, leaving little chance e population to recover and to live under safer condition





avalanche that came from a glacier several hundred meters above,



was provided by Laurent Bollinger (CEA/DASE). My very deep appreciation goes to him and Dr. Som Sapkota (Dpt of Mines and Geology. KTM) for their help.

on the Langtang Lirung peak (7227 m); apparently the debris were taken away from the moraines and carried downward by the avalanche

WHAT'S NEXT?







The village of Langtang was totally buried under a rock-snow

