

Physical, social and institutional vulnerability assessment in small Alpine communities. Results of the SAMCO-ANR project in the Upper Guil Valley (French Southern Alps)

Benoit Carlier, Constance Dujarric, Nikita Frison-Bruno, Anne Puissant, Candide Lissak, Malika Madelin, Vincent Viel, François Bétard, Monique Fort, Gilles Arnaud-Fassetta

▶ To cite this version:

Benoit Carlier, Constance Dujarric, Nikita Frison-Bruno, Anne Puissant, Candide Lissak, et al.. Physical, social and institutional vulnerability assessment in small Alpine communities. Results of the SAMCO-ANR project in the Upper Guil Valley (French Southern Alps). EGU General Assembly 2016, Apr 2016, Vienne, Austria. hal-01373208

HAL Id: hal-01373208

https://hal.science/hal-01373208

Submitted on 28 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Physical, social and institutional vulnerability assessment in small Alpine communities. Results of the SAMCO-ANR project in the Upper Guil Valley (French Southern Alps)






Benoit Carlier ¹⁻⁴, Constance Dujarric ¹⁻⁴, Nikita Frison-Bruno ¹⁻⁴, Anne Puissant ²⁻⁴, Candide Lissak, ³⁻⁴, Malika Madelin ¹⁻⁴, Vincent Viel ¹⁻⁴, François Bétard ¹⁻⁴, Monique Fort ¹⁻⁴, Gilles Arnaud-Fassetta ¹⁻⁴


1 Université Paris-Diderot, UMR CNRS 8586 PRODIG ; 2 Université de Strasbourg, UMR CNRS 7362 LIVE ; 3 Université de Caen, Basse Normandie, UMR CNRS 6554 LETG-Caen Geophen ; 4 ANR 12 SENV-0004 SAMCO

1- Background


The Guil catchment is particularly prone to torrential and gravitational hazards such as floods, debris flows, landslides or avalanches due to several predisposing factors (bedrock supplying abundant debris, strong hillslope-channel connectivity) in a context of summer Mediterranean rainstorms as triggers. Since the second half of the 20th century, the progressive decline of agropastoralism and the development of tourism activities led to a concentration of human stakes on alluvial cones and valley bottom, therefore an increase of vulnerability for mountainous communities. Following the 1957 and 2000 catastrophic floods and the 1948 and 2008 avalanche episodes, some measures were taken to reduce exposure to risks (engineering works, standards of construction, rescue training...). Nevertheless, in front of urban expansion (land pressures and political pressures) and obsolescence of the existing protective measures, it is essential to reassess the vulnerability of the stakes exposed to hazards. In the frame of the SAMCO project designed for mountain risk assessment in a context of global change, we developed a systemic approach to assess three specific components of vulnerability physical, social and institutional – for the six municipalities of the Upper Guil catchment: Ristolas, Abriès, Aiguilles, Château-Ville-Vieille, Molines-en-Queyras and St-Véran (Fig. 1).





Physical vulnerability (i.e. total potential consequences of hazards on stakes) was estimated and mapped via GIS model from Potential Damage Index (PDI) (Fig. 2). This index allowed us to quantify and describe both direct physical injury, structural and functional impacts - and indirect consequences - socio-economic impacts - in duced by hazards; this by combining weighted parameters reflecting the exposure of elements at risk: buildings, network and land cover (Fig. 3). At least 1890 buildings, 367 km² of land cover and 902 km of network were considered. Vulnerability maps were then crossed to hazard map reflecting different scenarios of exposure. To take into account the temporal variability of vulnerability, we produced different maps for summer and winter periods. To assess social and institutional vulnerability we realized questionnaires (5% of the total population investigated), interviews and mind-maps (80 collected) dealing with risk perception, mitigation measures and confidence in the actors of risk management.




3- Results: physical vulnerability assessement









summer torrential risk for 1957 flood extension and winter availanche risk. The highest degree of potential lized and cultural space. Regarding building, we observe a high degree of vulnerability on recent housing and networks close and 9). Considering winter risk for availanches, we count about 665 ha of land and 127 physical injury (Fig. 4) for flooding is preferentially located in recent settlement on the outskirts of his- housing, store and public services. Concerning mostly in the upper part of Guil catchment in torical villages which are often close to torrential rivers. Potential structural and functional vulnera- vulnerable on many points and particularly near torrential confluence areas. Socio-economic 100 year 1957 flood extension, we observed that more than 411 Ristolas municipality and in the adjacent valley of Aigues in the municipalities of Mo**bility** map for flooding (Fig. 5) put forward **urbanized and cultural space**.


1957 flood extension


For the sake of clarity for readers we present here only few scenarios: summer torrential vulnerability for flooding (Fig. 7) is sexpected, total potential vulnerability for flooding (Fig. 5) put forward urbanlines-en-Queyras and St-Véran (Figs. 10 and 11).


4- Results: Social and institutional vulnerability assessement


maps (Fig. 6) bring to light touristic issues such as shops, camping grounds and lifts.

Questionnaires and interviews suggest that Queyras inhabitants are globally aware and well informed about risk in their municipality (Fig. 12 and 13). According to their response, floods and avalanches are the major risks to consider. $\begin{bmatrix} 0 \\ 50 \end{bmatrix}$ Many of them (Fig. 14) have experienced a natural disaster, which suggest a good memory of risk. Mind-maps (Figs. 16 and 17) lead in that direction. Nevertheless, just few of them considered that they could be directly and physically affected by risk (Fig. 13); and there is still **7%** of people which are not informed about risk. These are often seasonal workers or newcomers such as retired people. Regarding to confidence towards stakeholder we observe a clear next flood preference for local authorities and civil security which are the most visible actors in those isolated regions (Fig. 15).









5- Conclusion

This work remains part of a large study on risk in mountainous region that should lead to a **web demonstrator** intended for risk stakeholders. We expect that these first results on vulnerability will contribute to a better assessment of the global vulnerability of the upper Queyras region to hydrogeomorphic hazards. This work must help the development of **better land use** and could be used to help local authorities to improve and update their **Emergency Action** Number of inhabitants Plan or their Prevention Plan. The next step of this work will be to try

to elaborate a method combining all these maps to produce a **global risk** map for mountain risks.