



# Mapping agricultural phenology using repetitive optical remote sensing over a peri-urban region

Nicolas Delbart, Emmanuelle Vaudour, Mihaela Dragoi, O. Maignan,  
Catherine Ottle

## ► To cite this version:

Nicolas Delbart, Emmanuelle Vaudour, Mihaela Dragoi, O. Maignan, Catherine Ottle. Mapping agricultural phenology using repetitive optical remote sensing over a peri-urban region. EGU General Assembly 2016, Apr 2016, Vienne, Austria. , 2016. hal-01375859

HAL Id: hal-01375859

<https://hal.science/hal-01375859>

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

Nicolas DELBART<sup>1</sup>, Emmanuelle VAUDOUR<sup>2</sup>, Mihaela DRAGOI<sup>3</sup>, Fabienne MAIGNAN<sup>4</sup>, Catherine OTTLE<sup>4</sup>

<sup>1</sup> PRODIG (UMR 8586), Université Paris-Diderot, CUE Sorbonne-Paris-Cité.

<sup>2</sup> ECOSYS UMR AgroParisTech, INRA, Université Paris-Saclay, 78850, Thiverval-Grignon, France

<sup>3</sup> Master Course student (TGAE Remote Sensing and Geomatics applied to the Environment)

<sup>4</sup> LSCE (UMR8212), Orme des Merisiers, Gif-sur-Yvette, France

nicolas.delbart@univ-paris-diderot.fr

**Framework :** map soil organic carbon content, using remote sensing, ground and laboratory measurements, modelling. Project **SURFAC\_EGC\_SENTINEL\_PLEIADES\_CO**, financed by CNES (French National Space Agency).

**Objectives of this work :** determine typology of crop rotations as a further input to modelling.

**Remote sensing data used :** Landsat-8 surface reflectance, PROBA-V, SPOT4-TAKE5, Pleiades.



### Data presentation and preparation




Figure 1 : NDVI from Landsat-8 surface reflectance (L8SR), subsetted and masked for clouds and strong aerosols loads.




Figure 2 : temporal variability based colour composite image, based on NDVI min, max and amplitude within 3 years. Indicative range : Yellow : [0.2-0.8]. Red : [-0.6-0.1]. Dark blue : [0.2-0.3]. Cyan : [0.6-0.8].




Figure 3 : crop mask, selected as  $NDVI_{min} > 0.3$  and  $NDVI_{max} > 0.6$ . Morphological reconstruction dilatation filter was applied to remove isolated pixels.

Vector layer : stands from national land parcel registry subdivided by individual crops based on high resolution imagery (Vaudour et al. 2015).



Figure 4 : Comparative dataset, Vaudour et al. (2015). See area in Fig.3. Crop map obtained from the SVM classifier implemented on the bi-temporal 3 and 24 April 2013 Pléiades images.

### Typology of NDVI temporal variations in NDVI for agricultural areas.

Constraint : incomplete time series (clouds, aerosols).

Methods for 2013 :

1/ Missing data reduction → Maximum value composite in July 2013 (DOY 191-200), December 2013 (344-360), September 2014 (251-258).

2/ Kmeans classification on pixels with only valid data (2013<sub>DOY104</sub>, 2013<sub>July</sub>, 2013<sub>DOY248</sub>, 2013<sub>DOY280</sub>, 2013<sub>Dec</sub>, 2014<sub>DOY75</sub>, 2014<sub>DOY107</sub>, 2014<sub>DOY139</sub>, 2014<sub>Sept</sub>, 2014<sub>DOY315</sub>, 2015<sub>DOY110</sub>, 2014<sub>DOY270</sub>). 60 classes → 60 temporal profiles.

2/ Assignment of all pixels (including pixels with missing values) to a temporal profile.

3/ Clustering temporal profiles according to their mutual correlations, and grouping the classified pixels on this basis.

Results :




Figure 5: temporal variations in cultures.

Top-left: map of temporal types, classes correspond to the clustered profiles shown on the right panel.

Bottom-left : Dark green show pixels for which NDVI increases in spring 2013, plateaus in summer 2013, and then decreases during autumn 2013. This groups the above classes 3, 12, 18 and 19.






Figure 6 : comparison of the temporal classes (Fig. 5) with classes obtained from multi-spectral classification in April 2013 (Fig. 4)

### Conclusion

Pixel aggregation based on NDVI temporal variations across 3 years (cultural cycle of 4 years in the study area).

**Some agreement with multipectral classification in spring 2013 :** e.g. NDVI variation type 3 matches well to spring cultures.

**Some disagreement :** e.g. NDVI variation type 4 is not specific to any crop mapped.

### Perspective 1 : Densifying time series




Figure 7: NDVI temporal variations, stand-wise. Landat, PROBA-V (100m central camera), SPOT4-TAKE5. 9 stands from Fig. 3.

### Perspective 2 : input to crop models for soil organic carbon modelling

