

# Comparative study of trophic transfer of the essential metals Co and Zn in two tropical fish: a radiotracer approach

Simon Pouil, Jean-Louis Teyssié, Claude Rouleau, Scott W. Fowler, Marc Metian, Paco Bustamante, Michel W Warnau

# ▶ To cite this version:

Simon Pouil, Jean-Louis Teyssié, Claude Rouleau, Scott W. Fowler, Marc Metian, et al.. Comparative study of trophic transfer of the essential metals Co and Zn in two tropical fish: a radiotracer approach. Journal of Experimental Marine Biology and Ecology, 2017, 486, pp.42-51. 10.1016/j.jembe.2016.09.005. hal-01377863

# HAL Id: hal-01377863 https://hal.science/hal-01377863

Submitted on 7 Oct 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Comparative study of trophic transfer of the essential metals Co and Znin two tropical fish: a radiotracerapproach

Simon Pouil<sup>a,b</sup>, Jean-Louis Teyssié<sup>a</sup>, Claude Rouleau<sup>c</sup>, Scott W. Fowler<sup>d</sup>, Marc Metian<sup>a</sup>, PacoBustamante<sup>b</sup>, Michel Warnau<sup>a</sup>,

- <sup>a</sup> International Atomic Energy Agency EnvironmentLaboratories (IAEA-EL), 4a, Quai
  Antoine Ier, MC-98000, Principality of Monaco, Monaco
- <sup>b</sup> Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, F-17000 La Rochelle, France
- <sup>c</sup> Institut Maurice-Lamontagne, Pêches et Océans Canada, 850, Route de la Mer, C.P.
  1000, Mont-Joli G5H 3Z4, Québec, Canada
- <sup>d</sup> Institute Bobby, 8 Allée des Orangers, Cap d'Ail, 06320 France
- \* Corresponding author: Dr Marc Metian

Radioecology Laboratory

IAEA EnvironmentLaboratories

4a Quai Antoine 1er

MC-98000 Principality of Monaco

Telephone: +377 97 97 72 17

E-mail: m.metian@iaea.org

#### Abstract

The trophic transfer of two essential metals (Co and Zn)was investigated in twotropical euryhalinefish species (*Monodactylus argenteus* and *Scatophagusargus*), at both the adult and juvenilestages, using the pulse-chase feeding method and radiotracers ( $^{57}$ Co and  $^{65}$ Zn).The food selected wasbrine shrimp (*Artemiasalina*) previously exposed to dissolved radiotracers. Depuration kinetics of both elements were followed for 45d. During this period,  $^{57}$ Co and  $^{65}$ Zn distribution was also determined at different time intervals among 15 different body compartments. Results showed that, in juveniles, the ingested Co was poorly assimilated by both species (AE <6%), whereas *S. argus* assimilated  $^{65}$ Zn more efficiently than *M. argenteus*(AE= 24% *vs.* 15%). In terms of bodydistribution of these essential elements, the trends were similar between adults and juveniles of both species:Co was concentrated and mainly distributed in the liver in *S.argus*,highlighting the role of this organ in the storage of Co. Conversely, in both species, ingested Zn was more diffusely distributed throughout the body.The differences observed between species could be related to a difference in the digestive enzymatic system ineach studied species,which in turn is related to their slightly different feeding habits in the natural environment.

Key words: Cobalt, Zinc, Assimilation, Kinetics, Tissue distribution, Interspeciesdifferences

#### **1. Introduction**

Essential metals are part of the functional groups of various enzymes, play a structural role in respiratory pigments and metalloenzymes, and can act as activating co-factors for various enzymes (see e.g. Simkiss 1979; Williams 1981). For example, Co is a key component of vitamin B12 (cyanocobalamin), which is a coenzyme in a number of cellular processes including the oxidation of fatty acids and the synthesis of DNA (e.g. Blust 2011). Zn has structural and catalytic roles in many proteins and almost 10% of all genes in sequenced fish genomes carry the annotation of Zn binding (e.g. Hogstrand 2011). Healthy conditions of fish are optimal if these elements are present in sufficient amounts in their tissues: depletion in these elements can provoke pathological damages and/or physiological alterations and their excess can also lead totoxic effects (e.g.Rainbow 2002; Förstner&Wittmann 2012).

In fish, foodappears increasingly as an important pathway for acquiring essential metals (e.g.Xu & Wang 2002; Mathews & Fisher 2009). For example, the metal intake from the diet in the teleost *Scophthalmus maximus* was estimated to be responsible for more than 70% of the body burden of Co, Mn and Zn (Mathews &Fisher 2009).Despite a growing dataset showing the major role that foodplays in essential and non-essential metal accumulation in fish (e.g. Spry et al. 1988; Zhang &Wang 2005), there is still limited understanding of the factors that influence metal assimilation at the interspecific level. Among these factors, feeding strategies have been reported to be crucial in the differential metal assimilation incoexisting species (e.g. Ni et al. 2000).

A key parameter for understanding and modeling metal trophic transfer in fish is the assimilation efficiency (AE). This variablecan be easily determined under laboratory conditions, using radiotracer techniques (e.g. Mathews et al. 2008; Pouil et al. 2015). A complementary approach is the determination of the dynamics of inter-organ metal transfer, based on the measurements of metal in the different organs and tissues at different time

intervals during the depuration phase. Although of interest, this approach is not very commonly reported in the literature. It has however been applied fortoxicokineticstudies in fish (e.g. Hogstrand et al. 2003) or with other aquatic aquatic active predatorslike cephalopods(Bustamante et al. 2002; 2004).

In this context, the present work investigated the trophic transfer of two essential elements (Co and Zn) in adults and juveniles of two tropicaleuryhalinefish species, the silvermoony*Monodactylus argenteus* and the spotted scat*Scatophagusargus*,using gamma-emitting radiotracers(<sup>57</sup>Co and <sup>65</sup>Zn).Thisbiokinetic study has focused on obtaining transfer rate data for tropical fish species given the fact that the majority of availableinformation has to date come from studies carried out with temperate species. Recent reviews on the topic of trace element and radionuclide transfer in marine organisms have stressed the need for more data on tropical species in order to make valid comparisons on a global scale (see e.g. Fowler &Fisher 2005).

Two levels of biological organization were considered in this study, i.e. the whole organism and the different organs and tissues, in order to evaluate the biokinetic parameters of the metal depuration and to identify metal transfer dynamics among the body compartments during the depuration phase.

# 2. Materials and Methods

# 2.1. Acclimation of organisms

One hundred juveniles and 20 adults of each species (silvermoony *M. argenteus* and spotted scat *S. argus*)were shipped to the International Atomic Energy Agency-Environment Laboratories' premises in the Principality of Monaco. Fish were acclimated for 3 months to laboratory conditions, adjusted to replicate as closely as possible their natural tropical

environment (2000-L tank for adults and 700-L aquarium for juveniles;open circuit: 200 L h<sup>-1</sup> in each tank;0.45-µm filtered seawater; salinity: 35p.s.u.; temperature:  $25 \pm 0.3$ °C; pH: 8.1 ± 0.1; light/dark: 12h/12h). During the acclimation period for both species, the juvenile fish were fed 3 to 4 times per day with adult brine shrimp (*Artemiasalina*) at a daily ration of 130-290 mg wet weight (wwt)per individual. Adults were fed4 times per day with compounded pellets (JBL Mariperls<sup>®</sup>) at a daily ration of 1.6-2 g dry weight (dwt)per individual. Virtually no mortality was observed during the acclimationperiod. Since body size (age) is known to affect metal bioaccumulation in marine organisms (Boyden 1974; Warnau et al. 1996a; Hédouin et al. 2006), only individuals with homogeneous sizes were used in the experiments.

#### 2.2. Experimental Procedure

# 2.2.1. Radiolabelling of brine shrimp

In order to investigate the trophic transfer of the two essential elements (Co and Zn) by *M*. *argenteus* and *S. argus*, adult brine shrimp (*Artemiasalina*) were used as prey. Preparation of the radiolabelled brine shrimp was carried out by exposing them for 18 h in aerated 20-L aquaria(approx. 300gwwt of adult brine shrimp per aquarium). At the end of the exposure period, brine shrimp were brieflyrinsed and stored at -20°C. Exposure was repeated using 6 batches of brine shrimp in order to obtain enough radiolabelled food for trophic transfer experiments (with both juvenile and adult fish). Radiotracers of high specific activity were purchased from CERCA, France (<sup>57</sup>Co, [T<sub>1/2</sub>] = 271.8 days) and Amersham, UK (<sup>65</sup>Zn, [T<sub>1/2</sub>] = 243.9 days). Stock solutions of radiotracers were prepared in 0.1 M HCl.

Seawater was spiked with the radiotracers (nominal activity of  $3kBq L^{-1}$  for Co and  $2 kBq L^{-1}$  for Zn). Small volumes (1 mL) of the diluted radiotracer solution were added to the aquaria (close circuit), and no change in pH and salinity was detectable after the tracer addition. At

the end of the exposure period, average activities in the brine shrimp were 239 Bq  $g^{-1}$  wwt for <sup>57</sup>Co and 40 Bq  $g^{-1}$  wwt for <sup>65</sup>Zn.

#### 2.2.2. Exposure of juvenile fish

One batch of 86 juvenile silvermoonies and one of 96 juvenile spotted scats ( $8.4 \pm 1.4$  g and  $4.8 \pm 1.4$  g wwt, respectively) were each transferred into a 70-L aquarium (open circuit: 100 L h<sup>-1</sup>; aerated, 0.45-µm filtered seawater; salinity: 35 p.s.u.; temperature:  $25 \pm 0.3$ °C; pH:  $8.1 \pm 0.1$ ; light/dark: 12h/12h)prior to the exposure to radiolabelled brine shrimp.

The experiment consisted of a single exposure to radiolabelled brine shrimp via ingestion (thawed prey, single feeding method also called pulse-chase feeding; see e.g. Metian et al. 2010; Pouil et al. 2015). Juvenile fishwere fed *ad libitum* for 1.25 h in a closed circuit system (247gwwt and 255g wwtof radiolabelled brine shrimp distributed to silver moony and spotted scat, respectively). Two hours after the beginning of ingestion, all fish were whole-body  $\gamma$ -counted alive and each batch of fish was removed and divided into two groups. Twenty-one individuals were transferred into a 20-L aquarium containing clean, flowing seawater (parameters as previously described). The remainder of the fish were returned into the initial 70-L aquarium. No regurgitation of the ingested radiolabelled brine shrimp was observed.

The 21 individuals from the 20-L aquarium were weighed separately (for individual recognition) and  $\gamma$ -counted alive (see section 2.3)at different time intervals over 45d to follow the whole-body depuration kinetics of <sup>57</sup>Co and <sup>65</sup>Zn. The aquarium was cleaned during each counting period to avoid contamination from radiotracerscontained in the faeces.Fish from the 70-L aquarium were regularly sampled (n=3) using the same protocol, anesthetized and dissected (see section 2.2.4) after  $\gamma$ -counting.

#### 2.2.3. Exposure of adult fish

Eighteenadult silvermooniesand 13 adult spotted scats ( $81 \pm 12g$  and  $193 \pm 35g$  wwt, respectively) were fed *ad libitum* in a 70-L aquariumduring 6 h with thawed radiolabelled brine shrimp (i.e. 11 feedings for 30min each time) in order to maximise the ingested activity in the fish (total distribution of 795g of radiolabelled brine shrimp for both species). Due to the size of adult individuals,  $\gamma$ -counting was not performed with livingwhole organisms;thus, 3 fish were randomly collected at different time intervals and dissected during the 45-d depuration period. For this purpose, fish were anesthetized, sacrificed and dissected into 15 body compartments (see details in section 2.2.4).

#### 2.2.4. Body distribution of metals in fish

Juvenile fish maintained in the 70-L aquarium and all adults were regularly sampled, anesthetized usingeugenol, and dissected in order to compare the metal organotropism in the tissues of both species. At each sampling time, 3 individuals (for both juvenilesand adults) were dissected into the followingbody compartments: (1) muscles with the skin, (2) stomach and pyloric caeca, (3) intestine, (4) liver, (5) gall bladder, (6) pancreas and spleen, (7) kidneys, (8) heart, (9) gills, (10) eyes, (11) brain, (12) skeleton (13) head, and in addition for adult fish (14) gonads and (15) red gland. All compartments were separated, weighed (wwt) and radio-analysed (see section 2.3) to determine the radiotracer distribution among the various tissues. During all the experiments, no mortality occurred.

# 2.3. Radiotracers and Counting

The radioactivity of the tracers was measured using a high-resolution  $\gamma$ -spectrometer system composed of 3 Germanium N type detectors (EGNC 33-195-R, Intertechnique<sup>®</sup>) connected to

a multi-channel analyser and a computer equipped with spectra analysis software (Interwinner 4, Intertechnique<sup>®</sup>). The radioactivity in live organisms and tissue samples was determined by comparison with known standards of appropriate geometry (calibration and counting). Measurements were corrected for background and physical radioactive decay. Liveorganisms were placed in polystyrene counting tubes (diameter: 80 mm, height: 50 mm), filled with clean seawaterduring the counting period. The volume of seawater was kept as low as possible to ensure atthe same time the welfare of the fish and a constant geometry during the counting period(i.e. reduced fish movement). The counting time was kept as short as possible and adjusted to obtain a propagated counting error less than 5% (e.g. Rodriguez y Baena et al. 2006; Metian et al. 2009). In the case of radioanalysis oflive fish, counting timesgenerally ranged between 15 and 55 min.Tests were performed prior to the experiments, where fish were placed in similar counting conditions in order to observe their behaviour, i.e. in a counting box for 1 hour in the dark. Dissolved O<sub>2</sub> concentration was monitored throughout these tests and was always > 3mg L<sup>-1</sup>. No alteration in organism health or behaviourwasobserved.

#### 2.4. Data treatment and statistical analyses

#### 2.4.1. Whole-body depuration kinetics in juvenile fish

Depuration of radiotracers was expressed as the percentage of remaining radioactivity (radioactivity at time t divided by the initial radioactivity measured in the organism at the beginning of the depuration period \* 100; Warnau et al. 1996b). The depuration kinetics of the radiotracers were best fitted using a two-component exponential model(Eq. 1), with a decision based on F test and ANOVA tables (Hédouin et al. 2010):

(1) 
$$A_t = A_{0s} \cdot e^{-k_{es}t} + A_{0l} \cdot e^{-k_{el}t}$$

where  $A_t$  and  $A_0$  are the remaining activities (%) at time t (d) and 0, respectively;  $k_e$  is the depuration rate constant (d<sup>-1</sup>); 's' and 'l' are the subscripts for the 'short-lived' and 'long-lived' components, respectively. The short-lived component represents the depuration kinetics of the radiotracer fraction that is weakly associated with the organisms and rapidly eliminated (e.g. fraction in faeces), whereas the long-lived component describes the depuration kinetics of the radiotracer fraction that is assimilated by and tightly bound to the organism (Warnau et al. 1996b). The long-lived component allows assessing the assimilation efficiency (AE) of the radiotracer ingested with food (AE =  $A_{01}$ ). For each exponential component (s and l), a biological half-life can be calculated ( $T_{bt/2s}$  and  $T_{bt/2i}$ ) from the corresponding depuration rate constants ( $k_{es}$  and  $k_{el}$ , respectively) according to the relation  $T_{bt/2} = \ln 2/k_e$ . The model was fitted using non-linear routines and model constants were estimated by iterative adjustment of the model, using the Quasi-Newton method in the Statistica<sup>®</sup> software 7.0.

# 2.4.2. Distribution among body compartments (juveniles and adults)

For juveniles and adults of the two fish species, distribution of Co and Zn among the body compartments was determined over time and compared using the non-parametric Mann-Whitney U test (Zar 1996). Furthermore, in order to quantify the transfer of each element over time among the internal body compartments, the ratio of the concentration of Co and Zn in each organ and/or tissue (Bq g<sup>-1</sup>wwt) over the whole-body (Bq g<sup>-1</sup>wwt) was calculated (sicconcentration index, I<sub>c</sub>, as defined by Rouleau et al. 2000). Values of I<sub>c</sub>> 1 indicate that a tissue is enriched in Co or Zn compared to the average whole-body metal concentration. This indexcomplements the information provided by metal body distribution and is useful forquantifying and comparing the storage capacities of the different body compartments in fish whose weight and ingested activity were variable. Comparisons of I<sub>c</sub> values amongbody compartments werecarried out using theKruskall-Wallis non-parametric test,followed by a multiple-comparison test of Siegel and Castellan (Zar 1996).

The level of significance for all statistical analyses was always set at  $\alpha = 0.05$ . All the statistical analyses were performed using either the Statistica<sup>®</sup> software 7.0 or R freeware 3.0.1 (R Development Core Team 2014).

# 3. Results

# 3.1. Depuration kinetics and assimilation efficiencies of <sup>57</sup>Co and <sup>65</sup>Zn in juveniles

The whole-body depuration kinetics of <sup>57</sup>Co and <sup>65</sup>Zn following the ingestion of radiolabeled *A. salinas*by the juveniles of both fish species were best described by a twocomponent exponential model (Table 1 and Fig. 1). Assimilation efficiency (AE) and retention capacities are metal- and species- dependent. Both species assimilated poorly<sup>57</sup>Co (AE <6%), whereas spotted scat assimilated <sup>65</sup>Zn more efficiently than silver moony (AE = 24.2 ±  $1.0\% vs.14.7 \pm 0.9\%$ , respectively; p<0.05). Assimilated <sup>57</sup>Co and <sup>65</sup>Zn were slowly eliminated, with biological half-life (T<sub>b/2l</sub>) values ranging from 33 to 55 d for <sup>57</sup>Co and from 67 to 83 d for <sup>65</sup>Zn (Table 1).

# 3.2. Body distribution and body dynamic of metals

During the 45-d depuration phase, dissections were carried outregularly in order to determine the metal distribution among selected body compartments (see section 2.2.4). Figures 2, 3 and 4 describe the distribution of Co and Zn,respectively, in the 4 main body compartments (i.e. stomach, intestine, liver, and muscles with skin representing 50-98% and 40-99% of total activity for Co and Zn, respectively) and in the remaining parts (i.e. the sum of the remaining body compartments) during the depuration phase in juveniles and adults of both species. The detailed distributions in each body compartment are provided in the supplementary material for this article. Considering elementdistribution, the depuration may be divided into three phases. The first phase (days 0-1), very rapid, corresponds to the passage of the food into the stomach which containsup to 80% of the radioactivity body burden at the end of the feeding period (85 min). The second phase (days1-4) is characterized by the occurrence of the digestive processes. During phases 1 and 2 (i.e. the first 4d of depuration), almost all the activity was distributed in the stomach and the intestine (up to 95% of the total body burden). Then, after4 days a third phase can be depicted and corresponds to elementtranslocations among the main compartments when whole-body activities are stable (i.e. when the absorbed fractionhas stabilized; Fig. 1). Gradually, the liver and the muscles with skin becomethen the predominant compartments in terms of trace element burden.Distributions arehoweverrelatively stable after day 10 except for Co inS. argus. Indeed, the percentage of <sup>57</sup>Co body burden in the liver tended to decrease between days 10 and 45 in both juvenile and adult fish, whereas a concomitant increase was observed in the muscles with skin(Fig. 2 and Fig. 4).Distribution patterns over time were similar between the species, juveniles or adults, for <sup>65</sup>Zn (Fig. 3 and Fig. 4), but not for<sup>57</sup>Coin whichcase the distribution in the liver was significantly higher in S. argus than in M. argenteus(Fig.2 and Fig. 4; p<0.05).

In terms of concentration index,  $I_c$ , Tables 2 and 3 provide a ranking of the  $I_c$  for Co and Zn, respectively, in each body compartment during the depuration phase in juveniles and adults of both species. The results indicate that after  $2 days^{57}$ Co was mainly concentrated in the liverfor *S. argus* juveniles and adults(Table 2). In the case of *M. argenteus*, there was no significant difference between the calculated  $I_c$  values of juveniles and adults for all the compartments, although the liver and the pancreas seem to play an important role in the storage of  ${}^{57}$ Co during the depuration phase. For  ${}^{65}$ Zn, regardless of the species and the life stage, the highest

values of  $I_c$  were always measured in the stomach and the intestine during the first 24 https://doi.org/10.001/10.0000/10000/1000/1000/1000/1000/1000/1000/10000/1000/1000/100

# 4. Discussion

The tropical silver moony,*Monodactylus argenteus*, and the spotted scat,*Scatophagusargus*, were selected to investigate theassimilation,retention and transfer among organs and tissues of Co and Zn ingested from radiolabeled food under controlled laboratory conditions. Experimental approaches commonly used, based on whole-body kinetics (e.g. Zhao et al. 2001; Mathews &Fisher 2009; Pouil et al. 2015), do not allow easily discerningthe dynamics of metal transfer among organs and tissues. This fact underscores the originality of the present study since it combines different approaches to provide new information regarding the mechanisms of assimilation and the subsequent dynamics of translocation of these trace metals among the tissues and organs of these euryhaline fish.

At the whole-body level, results show that AEs of Co are quite similarfor the two fish species whereas theydiffer for Zn. Indeed, ranges of AE for <sup>57</sup>Co and <sup>65</sup>Zn in silver moony and spotted scat were 5-6% and 15-24%, respectively. These AEs are in accordance with those reported in the literature for other species of marine fish, for which AEs ranged from 2 to 44% for Co and 1 to 52% for Zn (Table 4).

Many factors could explain the variations of AEs. Wang (2001) proposed in his review a ranking of these factors classified in 3 different categories: (1) environmental quality (such as food quality and quantity), (2) metal geochemistry and (3) feeding physiology and biology (such asingestion rate and gut passage time). Although only bivalves were considered in that review, these factors are generally relevant for fishas well. In the present study, interspecific difference in Zn AE was observed. Indeed, results indicated a higher AE for *S. argus*.Exposureconditions (i.e. the use of radiolabeled brine shrimp as unique food for trophic transfer in both species maintained under the same experimental conditions) did not allow

addressing categories (1) and (2) mentioned by Wang (2001). Although both species are omnivorous, as indicated by their identical Fishbasetrophic levels (3.0 for the both species; Froese and Pauly 2015a, b), differences exist in their trophic ecology.Indeed, analyses of the stomach contents from field surveys indicated that detritus is the main diet of *M. argenteus*(Blaber 1980; Rainboth 1996),whereas *S. argus* feedsmainly onzoobenthos such as worms and nekton such as finfish (Mills &Vevers 1989; Monkolprasit 1994; Jeyasselan 1998). These observations suggestthat spotted scat has a stronger predatory behavior than silver moony.Dietary habits are related to specific mixes of digestive enzymes or at least to variability of some enzymes'activity (e.g. Furne et al. 2005). For example, among 6 fish species, Hidalgo et al. (1999) have demonstrated that amylase displayedhigher enzymatic activity in omnivorous species than in carnivorous predators.Thus, in the presentstudy, the higher Zn AE measured in *S. argus* could be due to a more efficient enzymatic system for the digestion of brine shrimp than in silver moony.In short, spotted scat might be better "equipped" to deal with digesting complex multicellular prey.

This suggested interspecific variation in the fish digestive enzyme array cannot explain the trends observed for Co. For this element, AEs were very low and very similar between the two fish species. Differences,however, occurredin terms of body distribution and I<sub>C</sub>.The results highlightthe major role played by the liver in Co storage in spotted scat.This organ is involved in the mechanisms of detoxification and storage of Co,as previously shown in the European plaice*Pleuronectesplatessa* fedradiolabeled annelid worms *Arenicola marina*, where Amiard-Triquet&Amiard (1974) reported a low Co AE with almost all the activity found inliver and kidney.Thus, trophic ecology could lead to the occurrence of specific mechanisms of storage and detoxification. Further investigation is howeverneeded to obtain a betterunderstanding of the mechanisms of detoxification and storage of Co in the twospeciesexamined in this study.

Regarding the potential effect of thelife-stage, juveniles and adults of both specieshad comparable patterns of distribution and concentrations (i.e. "concentrationindex") for Co and Zn,i.e.the major fraction of both elements was found in the same organs and the metal concentrations in the organs were similar between the two life-stages. These comparable patterns can result from asimilar metabolism in juveniles and adults(viz. similar digestive activity and AE) although there were differences in size (adults of silver moony and spotted scat were respectively 10 and 40 times larger than juveniles). In addition, it is important to note that adults of both species were notsexually matureat the moment of the experiment. This would potentially affect the results due to the non-negligible allocation of some essential elements to reproductive organs, especially in mature females with oocyte production(e.g. Protasowicki 1986; Rajkowska&Protasowicki 2013).

#### Conclusion

The biokinetic approach at two levels of biological organization, (1) the whole organism and (2) the different organs and tissues, has provided animproved understanding of the mechanisms responsible forthe difference in Co and Zn assimilation efficiency observed between*M. argenteus* and *S. argus*.Dissections carried out at different times during the depuration provided new information about the dynamics of metal translocation in the different organs and tissues. Thus, changes observed in the Co distribution between the liver and the muscles in *S. argus* are a good example of the contribution of this radiotracerbasedmethodology. These also demonstrated interspecific variations in the assimilation and tissue distribution of Co and Zn in tropicalmarine fish.

# References

- Amiard-Triquet, C., Amiard, J.C., 1974. Contamination de chaînes trophiques marines par le cobalt 60 (in french).Rev. Intern. Ocean. Med.33,49–59.
- Blaber, S.J.M., 1980. Fish of the Trinity Inlet System of North Queensland with notes on the ecology of fish faunas of tropical Indo-Pacific estuaries. Aust. J. Mar.Freshwat. Res. 31,137-46.
- Blust, R., 2011.Cobalt,in: Wood, C.M., Farrell, A.P., Brauner, C.J. (Eds.), Fish physiology: homeostasis and toxicology of essential metals, 1<sup>st</sup> edition. Elsevier, Amsterdam, pp.291–326.
- Boyden, C.R., 1974. Trace element content and body size in molluscs.Nature251,311–314.
- Bustamante, P., Teyssié, J.-L., Fowler, S.W., Cotret, O., Danis, B., Miramand, P., Warnau, M., 2002.Biokinetics of zinc and cadmium accumulation and depuration at different stages in the life cycle of the cuttlefish *Sepia officinalis*.Mar. Ecol.Prog. Ser.231,167–177.
- Bustamante, P., Teyssié, J.-L., Danis, B., Fowler, S.W., Miramand, P., Cotret, O., Warnau, M., 2004. Uptake, transfer and distribution of silver and cobalt in tissues of the common cuttlefish *Sepia officinalis* at different stages of its life cycle.Mar. Ecol.Prog. Ser.269,185–195.
- Förstner, U., Wittmann, G.T., 2012. Metal pollution in the aquatic environment. Springer Science & Business Media, Berlin, Germany.
- Fowler, S.W., Fisher, N.S. 2005. Radionuclides in the biosphere, in: Livingston, H.D. (Eds.), Radioactivity in the environment. Elsevier, Amsterdam, pp. 167–203.
- Froese, R., Pauly, D., 2015a.FishBaseMonodactylus argenteus, Silver moony: fisheries, aquarium.http://wwwfishbaseorg/summary/Monodactylus-argenteushtml#comment (accessed 16.04.11).
- Froese, R., Pauly, D., 2015b.FishBaseScatophagusargus, Spotted cat: fisheries, aquaculture, aquarium. http://wwwfishbaseorg/summary/Scatophagus-argushtml#comment(accessed 16.04.11).
- Froese, R., Pauly, D., 2015c.FishBase http://wwwfishbase.org (accessed 16.04.11).
- Furne, M., Hidalgo, M.C., Lopez, A., Garcia-Gallego, M., Morales, A.E., Domezain, A, Domezaine, J.,Sanz, A., 2005.Digestive enzyme activities in Adriatic sturgeon *Acipensernaccarii* and rainbow trout *Oncorhynchus mykiss*: A comparative study.Aquaculture250,391–398.
- Hédouin, L., Metian, M., Teyssié, J.-L., Fowler, S.W., Fichez, R., Warnau, M., 2006. Allometric relationships in the bioconcentration of heavy metals by the edible tropical clam *Gafrariumtumidum*.Sci. Total Environ.366,154–163.
- Hédouin, L., Metian, M., Lacoue-Labarthe, T., Fichez, R., Teyssié, J.-L., Bustamante, P.,Warnau, M., 2010.Influence of food on the assimilation of selected metals in tropical bivalves from the New Caledonia lagoon: Qualitative and quantitative aspects.Mar.Pollut. Bull.61,568–575.
- Hidalgo, M.C., Urea, E.,Sanz, A. 1999. Comparative study of digestive enzymes in fish with different nutritional habits: Proteolytic and amylase activities.Aquaculture170,267– 283.
- Hogstrand, C., 2011.Zinc,in: Wood, C.M., Farrell, A.P., Brauner, C.J. (Eds.), Fish physiology: homeostasis and toxicology of essential metals, 1<sup>st</sup> edition. Elsevier, Amsterdam, pp. 135–200.
- Hogstrand, C.,Grosell, M., Wood, C.M., Hansen, H., 2003. Internal redistribution of radiolabelled silver among tissues of rainbow trout (*Oncorhynchus mykiss*) and European eel (*Anguilla anguilla*): The influence of silver speciation.Aquat.Toxicol.63,139–157.

- Jeffree, R.A., Warnau, M., Teyssié, J.-L., Markich, S.J., 2006. Comparison of the bioaccumulation from seawater and depuration of heavy metals and radionuclides in the spotted dogfish *Scyliorhinuscanicula* (Chondrichthys) and the turbot *Psetta maxima* (Actinopterygii: Teleostei).Sci. Total Environ.368,839–852.
- Jeyaseelan, M.J.P., 1998. Manual of fish eggs and larvae from Asian mangrove waters. United Nations Educational, Scientific and Cultural Organization. Paris. 193 p.
- Mathews, T., Fisher, N.S., 2008. Trophic transfer of seven trace metals in a four-step marine food chain.Mar. Ecol.Prog. Ser.367,23–33.
- Mathews, T., Fisher, N.S., 2009. Dominance of dietary intake of metals in marine elasmobranch and teleost fish.Sci. Total Environ.407,5156–5161.
- Mathews, T., Fisher, N.S., Jeffree, R.A., Teyssié, J.-L., 2008. Assimilation and retention of metals in teleost and elasmobranch fishes following dietary exposure.Mar. Ecol.Prog. Ser.360,1–12.
- Metian, M., Warnau, M., Oberhänsli, F., Bustamante, P., 2009.Delineation of Pb contamination pathways in two Pectinidae: The variegated scallop *Chlamysvaria* and the king scallop *Pecten maximus*.Sci. Total Environ.407,3503–3509.
- Metian, M., Hédouin, L., Eltayeb, M.M., Lacoue-Labarthe, T., Teyssié, J-L., Mugnier, C., Bustamante, P., Warnau, M., 2010. Metal and metalloid bioaccumulation in the Pacific blue shrimp *Litopenaeusstylirostris* (Stimpson) from New Caledonia: Laboratory and field studies.Mar.Pollut. Bull. 61,576–584.
- Mills, D., Vevers, G., 1989. The Tetra encyclopedia of freshwater tropical aquarium fishes, Tetra Press, New Jersey.
- Monkolprasit, S., 1994. Fish composition and food habits in mangrove forests at Phang-Nga Bay and Ban Don Bay, Thailand.Kasetsart. Univ. Fish Res. Bull. 20,1–21.
- Ni, I.-H., Wang, W.-X., Yin, K.T., 2000.Transfer of Cd, Cr and Zn from zooplankton prey to mudskipper *Periophthalmuscantonensis* and glassy *Ambassisurotaenia* fishes.Mar. Ecol.Prog. Ser.194,203–210.
- Pouil, S., Warnau, M., Oberhänsli, F., Teyssié J.-L., Bustamante P., Metian M., (*in press*). Influence of food on the assimilation of essential elements (Co, Mn& Zn) by turbot *Scophthalmus maximus*. Mar. Ecol.Prog. Ser.550, 207–218.
- Pouil, S., Warnau M., Oberhänsli, F., Teyssié, J.-L., Metian, M., 2015. Trophic transfer of <sup>110m</sup>Ag in the turbot *Scophthalmus maximus* through natural prey and compounded feed. J. Environ. Radioact. 150, 189–194.
- Protasowicki, M.,1986. Sex effects on Cd, Pb, Cu and Zn contents in selected fish organs.Balt. Sea Environ. Proc.19,433–441.
- R Development Core Team, 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rainboth, W.J., 1996. Fishes of the Cambodian Mekong FAO species identification field guide for fishery purposes.Food and Agriculture Organization of the United Nations, Rome, Italy.
- Rainbow, P.S., 2002. Trace metal concentrations in aquatic invertebrates: Why and so what? Environ.Pollut.120,497–507.
- Rajkowska, M., Protasowicki, M., 2013. Distribution of metals (Fe, Mn, Zn, Cu) in fish tissues in two lakes of different trophy in Northwestern Poland.Environ.Monit. Assess.185,3493–3502.
- Reinfelder, J.R., Fisher, N.S., 1994. Retention of elements absorbed by juvenile fish (*Menidiamenidia, Menidiaberyllina*) from zooplankton prey.Limnol.Oceanogr.39,1783–1789.

- Rodriguez y Baena, A.M., Miquel, J.C., Masqué, P., Povinec, P.P., La Rosa, J., 2006. A single vs double spike approach to improve the accuracy of <sup>234</sup>Th measurements in small-volume seawater samples. Mar. Chem. 100, 269–281.
- Rouleau, C., Gobeil, C., Tjälve, H., 2000. Accumulation of silver from the diet in two marine benthic predators: The snow crab (*Chionoecetesopilio*) and American plaice (*Hippoglossoidesplatessoides*). Environ. Toxicol. Chem. 19,631–637.
- Simkiss, K., 1979. Metal ions in cells.Endeavour3,2-6.
- Spry, D.J., Hodson, P.V., Wood, C.M., 1988. Relative contributions of dietary and waterborne zinc in the rainbow trout, *Salmo gairdneri*.Can. J. Fish Aquat. Sci.45,32– 41.
- Wang, W.-X., 2001. Comparison of metal uptake rate and absorption efficiency in marine bivalves.Environ. Toxicol.Chem.20,1367–1373.
- Wang, W.-X., Onsanit, S., Dang, F., 2012.Dietary bioavailability of cadmium, inorganic mercury, and zinc to a marine fish: Effects of food composition and type.Aquaculture356–357,98–104.
- Warnau, M., Teyssié, J.-L., Fowler, S.W., 1996a.Biokinetics of selected heavy metals and radionuclides in the common Mediterranean echinoid *Paracentrotuslividus*: Sea water and food exposures.Mar. Ecol.Prog. Ser.141,83–94.
- Warnau, M., Fowler, S.W., Teyssié, J.-L., 1996b. Biokinetics of selected heavy metals and radionuclides in two marine macrophytes: The seagrass *Posidoniaoceanica* and the alga *Caulerpataxifolia*. Mar. Environ. Res.41,343–362.
- Williams, R.J.P., 1981. The Bakerian lecture, 1981: Natural selection of the chemical elements. Proc. R. Soc.Lond. B Biol. Sci.213, 361–397.
- Xu, Y., Wang, W.-X., 2002. Exposure and potential food chain transfer factor of Cd, Se and Zn in marine fish *Lutjanusargentimaculatus*.Mar. Ecol.Prog. Ser.238,173–186.
- Zar, H.J., 1996.Biostatistical analysis: Third edition, Prentice-Hall, New Jersey.
- Zhang, L., Wang, W.-X., 2005. Effects of Zn pre-exposure on Cd and Zn bioaccumulation and metallothionein levels in two species of marine fish.Aquat.Toxicol.73,353–369.
- Zhang, L., Wang, W.-X., 2006. Significance of subcellular metal distribution in prey in influencing the trophic transfer of metals in a marine fish.Limnol.Oceanogr.51,2008–2017.
- Zhang, L., Wang, W.-X., 2007. Gastrointestinal uptake of cadmium and zinc by a marine teleost *Acanthopagrusschlegeli*.Aquat.Toxicol.85,143–153.
- Zhao, X., Wang, W.-X., Yu, K.N., Lam, P.K.S., 2001. Biomagnification of radiocesium in a marine piscivorous fish.Mar. Ecol.Prog. Ser.222,227–237.



Figure 1. Whole-body depuration of <sup>57</sup>Co and <sup>65</sup>Zn after a single-feeding with radiolabelled brine shrimp in juvenile silver moony (*Monodactylus argenteus*, n=21) and spotted scat (*Scatophagusargus*, n=21) expressed as percentage of remaining activities (means  $\pm$  SD). Parameters of depuration kinetics and their statistics are given in Table 1.



Figure 2.<sup>57</sup>Co distribution among 4 body compartments (intestine, stomach, liver, and muscles with skin) of juvenile silver moony (*M. argenteus*) and spotted scat (*S.argus*) over the depuration phase after feeding with radiolabelled brine shrimp. At each time three fish were dissected. All the values are expressed as percentage of the whole-body activity (means + SD).



Figure 3.<sup>65</sup>Zn distribution among 4 body compartments (intestine, stomach, liver, and muscles with skin) of juvenile silver moony (*M. argenteus*) and spotted scat (*S.argus*) over the depuration phase after feeding with radiolabelled brine shrimp. At each time three fish were dissected. All the values are expressed as percentage of the whole-body activity (means + SD).



Figure 4.<sup>57</sup>Co and <sup>65</sup>Zn distribution among 4 body compartments (intestine, stomach, liver and muscles with skin) of adult silver moony (*M. argenteus*) and spotted scat (*S.argus*) over the depuration phase after feeding with radiolabelled brine shrimp. At each time three fish were dissected. All the values are expressed as percentage of the whole-body activity (means + SD).

Table 1. Estimated depuration kinetic parameters of <sup>57</sup>Co and <sup>65</sup>Zn insilver moony (*M. argenteus*) and spotted scat (*S. argus*) after a single feeding with radiolabelled brine shrimp and then maintained for 45d in a clean flowing seawater. Depuration parameters:  $A_{0s}$  and  $A_{0l}$ : activity (%) lost according to the short-and the long-lived exponential component,respectively;the  $A_{0l}$  value is also the assimilation efficiency (AE; %), $T_{b'_2}$ : biological half-life (d).ASE: asymptotic standard error;  $R^2$ : determination coefficient of kinetics.

		Short-term		Long-term					
Tracer	Species	$A_{0s} \pm ASE$ (%)	$T_{b\frac{1}{2s}} \pm ASE$	$A_{0l}$ (=AE) ± ASE (%)	$T_{b\frac{1}{2}l} \pm ASE$	$\mathbf{R}^2$			
<sup>57</sup> Co	M.argenteus	95.4 ± 0.7***	0.42 ± 0.01***	4.58 ± 0.44***	32.7 ± 9.0***	0.99			
	S. argus	94.2 ± 0.6***	$0.28 \pm 0.01^{***}$	$5.77 \pm 0.30 ***$	54.5 ± 12.5***	0.99			
<sup>65</sup> Zn	M.argenteus	85.3 ± 1.9***	0.12 <sup>NS</sup>	14.7 ± 0.9**	66.8 ± 21.2***	0.95			
	S. argus	75.8 ± 1.9***	$0.27 \pm 0.04^{***}$	$24.2 \pm 1.0$ ***	83.1 ± 19.3***	0.94			

Significance of estimated parameters:

<sup>NS</sup> p >005 (not significant)

\*\* p < 001

\*\*\* p< 0001

Table 2.Concentration index (I<sub>c</sub>) of <sup>57</sup>Co, throughoutthe 45-d depuration of juvenile silver moony (*M. argenteus*) and spotted scat (*S. argus*) after a single feeding with radiolabelled brine shrimp.I<sub>c</sub> is the ratio between the activity concentration (Bq g<sup>-1</sup>wwt) in the considered compartment and the activity in the whole fish (Bq g<sup>-1</sup>wwt) multiplied by 100.A selection of body compartments having the highest and lowest I<sub>c</sub> values (p<005) has been made. EYE: eyes, GAL: gall bladder, GIL: gills, GON: gonads, HRT: heart, INT: intestine, KID: kidney, LIV: liver, MUS: muscle, PAN: pancreas + spleen, RED: red gland, SKE: skeleton, STO: stomach. Values are means or means  $\pm$  SD; n=3. Details of the data are available in supplementary materials.

	_	Time (d)																
Species	-	0	1	2	3		4	7	10		14	17	18	23		30	38	45
		Juvenile	Juvenile	Juvenile	Juvenile	Adult	Juvenile	Juvenile	Juvenile	Adult	Juvenile	Adult	Juvenile	Juvenile	Adult	Juvenile	Juvenile	Juvenile
W. argenteus M. argenteus Highest Lovalues	[ <sub>c</sub> values	Max: STO (12±2)	Max:	Max: INT (19±11)	<b>Max:</b> KID (13±1)	Max:	<b>Max:</b> HRT (14±6)	<b>Max:</b> HRT (28±19)	<b>Max:</b> GAL (21±13)	<b>Max:</b> LIV (21±13)	<b>Max:</b> PAN (30±28)	_	<b>Max:</b> PAN (47±36)	<b>Max:</b> LIV (18±2)	<b>Max:</b> LIV (16±11)	<b>Max:</b> PAN (60±9)	<b>Max:</b> PAN (48±2)	<b>Max:</b> PAN (51±34)
	Highest ]	Min: (39 INT (8±2)	(39±3)	Min: PAN (7±2)	<b>Min:</b> STO (4±1)	$\begin{array}{c} \text{Min:} \\ (63\pm4) \\ \text{LIV} \\ (10\pm2) \end{array}$	Min: LIV (15±1)	Min: STO (4±1)	<b>Min:</b> HRT (7±2)	Min: LIV (16±5)		<b>Min:</b> LIV (12±3)	Min: HRT (10±2)	Min: STO (6±3)	Min: LIV (8±2)	Min: LIV (8±3)	<b>Min:</b> LIV (7±2)	
	values	<b>Max:</b> PAN (<1)	<b>Max:</b> KID (2)	<b>Max:</b> STO (3)	<b>Max:</b> GIL (2)	<b>Max:</b> GAL (6)	<b>Max:</b> INT (10)	<b>Max:</b> KID (7)	<b>Max:</b> GIL (2)	Max: GON (5)	<b>Max:</b> GIL (2)	<b>Max:</b> GON (6)	<b>Max:</b> STO (4)	<b>Max:</b> GAL (7)	<b>Max:</b> GON (6)	<b>Max:</b> STO (4)	<b>Max:</b> STO (4)	<b>Max:</b> STO (4)
	Lowest I <sub>(</sub>	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)
S. argus Hickney Liveration	Highest I <sub>c</sub> values	Max: STO (9±2) Min: INT (3±2)	Max: INT (13±1) Min: LIV (8±3)	Max: LIV (13±6) Min: INT (10±7)	<b>Max:</b> LIV (14±2)	-	<b>Max:</b> LIV (18±1)	<b>Max:</b> LIV (26±3)	<b>Max:</b> LIV (18±4)	<b>Max:</b> LIV (30±2)	<b>Max:</b> LIV (33±6)	<b>Max:</b> LIV (21±1)	<b>Max:</b> LIV (21±1)	<b>Max:</b> LIV (25±4)	<b>Max:</b> LIV (29±6)	<b>Max:</b> LIV (17±2)	<b>Max:</b> LIV (17±3)	<b>Max:</b> LIV (15±2)
	values	<b>Max:</b> LIV (<1)	<b>Max:</b> KID (<1)	<b>Max:</b> KID (4)	<b>Max:</b> INT (7)	<b>Max:</b> LIV (10)	<b>Max:</b> HRT (4)	<b>Max:</b> INT (3)	<b>Max:</b> KID (2)	<b>Max:</b> INT (9)	<b>Max:</b> HRT (3)	<b>Max:</b> BRA (5)	<b>Max:</b> HRT (3)	<b>Max:</b> PAN (4)	<b>Max:</b> RED (4)	<b>Max:</b> HRT (4)	<b>Max:</b> HRT (5)	<b>Max:</b> HRT (6)
	Lowest I <sub>c</sub>	<b>Min:</b> EYE (0)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)

Table 3.Concentrationindex (I<sub>c</sub>) of <sup>65</sup>Zn, throughout the 45-d depuration of juvenile silver moony (*M argenteus*) and spotted scat (*S argus*) after a single feeding with radiolabelled brine shrimp.I<sub>c</sub> is the ratio between the activity (Bq g<sup>-1</sup>wwt) in the considered compartment and the total activity in the fish (Bq g<sup>-1</sup>wwt) A selection of the compartments having the highest and lowest of I<sub>c</sub> values (p<005) has been made.BRA: brain, EYE: eyes, GAL: gall bladder, GIL: gills, HRT: heart, HED: head, INT: intestine,KID: kidney, LIV: liver, MUS: muscle, PAN: pancreas + spleen, RED: red gland, SKE: skeleton, STO: stomach. Values are means or means  $\pm$  SD; n=3. Details of the data are available in supplementary materials.

		Time (d)																
Species		0	1	2	3		4	7	10		14	17	18	23		30	38	45
		Juvenile	Juvenile	Juvenile	Juvenile	Adult	Juvenile	Juvenile	Juvenile	Adult	Juvenile	Adult	Juvenile	Juvenile	Adult	Juvenile	Juvenile	Juvenile
<i>W. argenteus</i>	t I <sub>c</sub> vaues	Max: STO (13±1) Min:	Max: KID (40±27) Min:	<b>Max:</b> GAL (47±19) <b>Min:</b>	<b>Max:</b> PAN (25±13)	-	<b>Max:</b> GAL (34±23) Min:	<b>Max:</b> GAL (61±41) Min:	<b>Max:</b> GAL (79±72) Min:	-	<b>Max:</b> PAN (50±22)	<b>Max:</b> PAN (198±93) Min:	<b>Max:</b> PAN (88±61) <b>Min</b> :	Max: PAN (53±44) Min:	-	<b>Max:</b> PAN (102±78) Min:	<b>Max:</b> PAN (100±71) <b>Min:</b>	<b>Max:</b> PAN (88±20) <b>Min:</b>
	Highes	INT (7±3)	GAL (22±13)	PAN (26±16)	GAL (22±6)		HRT (13±8)	HRT (26±15)	KID (20±3)		GAL (29±7)	HRT (22±3)	KID (25±8)	GAL (31±28)		HRT (14±10)	HRT (31±14)	KID (17±7)
	values	<b>Max:</b> PAN (3)	<b>Max:</b> HRT (21)	<b>Max:</b> INT (4)	<b>Max:</b> HRT (10)	<b>Max:</b> PAN (50)	<b>Max:</b> BRA (4)	<b>Max:</b> KID (7)	<b>Max:</b> LIV (7)	<b>Max:</b> GAL (24)	<b>Max:</b> STO (4)	<b>Max:</b> BRA (4)	<b>Max:</b> BRA (7)	<b>Max:</b> LIV (3)	<b>Max:</b> RED (66)	<b>Max:</b> BRA (6)	<b>Max:</b> BRA (4)	<b>Max:</b> BRA (6)
	Lowest I <sub>c</sub>	<b>Min:</b> SKE (0)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> EYE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (1)	<b>Min:</b> MUS (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> HED (<1)
S. argus	Highest I <sub>c</sub> values	Max: STO (7±0) Min: INT (5±3)	Max: INT (6±2) Min: LIV (2±1)	-	-	-	-	-	-		-	-	-	<b>Max:</b> HRT (6±2)		Max: PAN (11±7) Min: MUS (3±0)	-	-
	values	<b>Max:</b> HRT (<1)	<b>Max:</b> HED (1)	<b>Max:</b> KID (9)	<b>Max:</b> PAN (6)	<b>Max:</b> KID (7)	<b>Max:</b> PAN (5)	<b>Max:</b> PAN (5)	<b>Max:</b> HRT (3)	<b>Max:</b> LIV (4)	<b>Max:</b> PAN (4)	<b>Max:</b> HRT (8)	<b>Max:</b> GAL (7)	<b>Max:</b> MUS (4)	<b>Max:</b> HRT (5)	<b>Max:</b> EYE (2)	<b>Max:</b> HRT (5)	<b>Max:</b> PAN (13)
	Lowest I,	<b>Min:</b> SKE (<1)	<b>Min:</b> BRA (<1)	<b>Min:</b> BRA (<1)	<b>Min:</b> BRA (<1)	<b>Min:</b> SKE (<1)	<b>Min:</b> BRA (<1)	<b>Min:</b> BRA (1)	<b>Min:</b> GAL (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> BRA (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> INT (<1)	<b>Min:</b> INT (<1)	<b>Min:</b> MUS (<1)	<b>Min:</b> INT (<1)	<b>Min:</b> INT (<1)	<b>Min:</b> INT (<1)

Table 4.Comparison of Co and Zn dietary assimilation efficiencies (AEs; means in %) in marineand brackish water fish species.

Species	Trophic level*	Food	Metal	(AE)	References
			Co	Zn	
Marine					
Acanthopagrusschlegeli	3.2±0.45	Artificialdiets		15-26	Wang et al. (2012)
		Mullet muscle		42	Wang et al. (2012)
		Mussel tissue		25	Wang et al. (2012)
		Squidviscera		14	Wang et al. (2012)
		Oyster tissue		4	Zhang & Wang (2007)
Scophthalmus maximus	$4.4 \pm 0.0$	Seabream juveniles	27	22	Mathews et al (2008)
		Ragworms	5	17	Pouil et al (2016)
		Seabream juveniles	44	23	Pouil et al (2016)
		Shrimp	16	32	Pouil et al (2016)
		Commercial pellets	1	13	unpublished data
Dicentrarchuslabrax	3.5±0.5	Seabreamjuveniles	21	38	Mathews& Fisher (2008)
Lutjanusargentimaculatus	3.6±0.5	Brineshrimp		15	Xu & Wang (2002)
		Clam tissue		30	Xu & Wang (2002)
		Copepods		20	Xu & Wang (2002)
		Manilla clam		19	Xu & Wang (2002)
Sparusaurata	3.7±0.0	Brine shrimp nauplii	21	18	Mathews et al. (2008)
Brackish					
Ambassisurotaenia	3.4±0.4	Brine shrimp nauplii		15-17	Ni et al. (2000)
		Copepods		5-9	Ni et al. (2000)
Menidiasp	2.6-3.9	Copepods	2	6	Reinfelder& Fisher (1994)
Monodactylus argenteus	$3.0 \pm 0.33$	Brine shrimp	5	15	this study
Periophthalmusmodestus	3.2±0.4	Brine shrimp larvae		11-21	Ni et al. (2000)
		Copepods		21-31	Ni et al. (2000)
Scatophagusargus	3.0±0.35	Brine shrimp	6	24	this study
Teraponjarbua	3.9±0.5	Barnacles		2	Zhang & Wang (2006)
		Copepods		23	Zhang & Wang (2006)
		Clams		36	Zhang & Wang (2006)
		Fish viscera		52	Zhang & Wang (2006)
		Mussels		22	Zhang & Wang (2006)

\* Data obtained from FishBase (Froese &Pauly, 2015c)