



HAL
open science

ON EXPONENTIAL FUNCTIONALS OF PROCESSES WITH INDEPENDENT INCREMENTS

Paavo P. Salminen, L Vostrikova

► **To cite this version:**

Paavo P. Salminen, L Vostrikova. ON EXPONENTIAL FUNCTIONALS OF PROCESSES WITH INDEPENDENT INCREMENTS. 2016. hal-01388080v1

HAL Id: hal-01388080

<https://hal.science/hal-01388080v1>

Preprint submitted on 26 Oct 2016 (v1), last revised 14 Jul 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON EXPONENTIAL FUNCTIONALS OF PROCESSES WITH INDEPENDENT INCREMENTS

P. Salminen, Department of Natural Sciences Åbo
Akademi University FIN-20500 Åbo Finland
and
L. Vostrikova ¹, LAREMA, Département de
Mathématiques, Université d'Angers, 2, Bd Lavoisier
49045, ANGERS CEDEX 01

ABSTRACT. In this paper we study the exponential functionals of the processes X with independent increments, namely

$$I_t = \int_0^t \exp(-X_s) ds, \quad t \geq 0,$$

and also

$$I_\infty = \int_0^\infty \exp(-X_s) ds.$$

When X is a semi-martingale with absolutely continuous characteristics, we derive necessary and sufficient conditions for the existence of the Laplace exponent of I_t , and also the sufficient conditions of finiteness of the Mellin transform $\mathbf{E}(I_t^\alpha)$ with $\alpha \in \mathbb{R}$. We give a recurrent integral equations for this Mellin transform. Then we apply these recurrent formulas to calculate the moments. We present also the corresponding results for the exponentials of Levy processes, which hold under less restrictive conditions than in [4]. In particular, we obtain an explicit formula for the moments of I_t and I_∞ , and we precise the exact number of finite moments of I_∞ .

MSC 2010 subject classifications: 60G51, 91G80

1. INTRODUCTION

The exponential functionals arise in many areas, in particular in the theory of self-similar Markov processes, mathematical finance, random processes in random environment and so on, as mentioned in [4]. More

¹¹This work is supported in part by DEFIMATHS project of the Research Federation of "Mathématiques de Pays de la Loire" and PANORisk project of Pays de la Loire region.

precisely, see [13], self-similar Markov processes are related with exponential functionals by Lamperti transform, namely self-similar Markov process can be written as an exponential of Levy process time changed by the inverse of exponential functional of the same Levy process. In mathematical finance the question is related to the perpetuities containing the liabilities, the perpetuities subjected to the influence of economical factors (see, for example, [11]), and also with the price of Asian options and related questions (see, for instance, [8], [10] and references therein).

In the case of Levy processes, asymptotic behaviour of exponential functionals was studied in [6], in particular for α -stable Levy processes. The authors also give an integro-differential equation for the density of the law of exponential functionals, when this density w.r.t the Lebesgue measure exists. For Levy subordinators they give the formulas for the positive and negative moments, the Mellin transform and the Laplace transform. The questions related with the characterisation of the law of exponential functionals by the moments was studied in [4]. In [3] the authors study continuity properties of exponential functionals, in particular they give sufficient conditions for the existence of the density of the law of I_∞ w.r.t. the Lebesgue measure. For Levy processes see also [17], [2], [12].

Exponential functionals for diffusions was studied in [14]. The authors considered exponential functionals stopped at first hitting time and they derive the Laplace transform of these functionals. To find laws of such exponential functionals, the authors perform numerical inversion of the Laplace transform. The relations between hitting times and occupation times for exponential functionals was considered in [16], where the versions of identities in law such as Dufresne's identity, Ciesielski-Taylor's identity, Biane's identity, LeGall's identity was considered.

The case when X is a non-homogeneous process with independent increments (PII in short) has not been considered up to now. At the same time PII models for logarithme of the prices are quite natural in mathematical finance, it is the case of non-homogeneous Poisson process, Levy process with deterministic time change, the integrals of Lévy processes with deterministic integrands, hitting times for diffusions and so on (see for instance [18], [7], [1]).

The aim of this paper is to study the exponential functionals of the processes X with independent increments, namely

$$(1) \quad I_t = \int_0^t \exp(-X_s) ds, \quad t \geq 0,$$

and also

$$I_\infty = \int_0^\infty \exp(-X_s) ds,$$

and give such important characteristics of these exponential functionals as the moments and the Laplace transforms and the Mellin transforms of mentioned functionals.

We consider a real valued process $X = (X_t)_{t \geq 0}$ with independent increments and $X_0 = 0$, which is a semi-martingale with respect to its natural filtration. We denote by (B, C, ν) a semi-martingale triplet of this process, which can be chosen deterministic (see [9], Ch. II, p.106). We suppose that $B = (B_t)_{t \geq 0}$, $C = (C_t)_{t \geq 0}$ and ν are absolutely continuous with respect to the Lebesgue measure in t , i.e. that X is an Ito process such that

$$B_t = \int_0^t b_s ds, \quad C_t = \int_0^t c_s ds, \quad \nu(dt, dx) = dt K_t(dx)$$

with measurable functions $b = (b_s)_{s \geq 0}$, $c = (c_s)_{s \geq 0}$, and the kernel $K = (K_t(A))_{t \geq 0, A \in \mathcal{B}(\mathbb{R} \setminus \{0\})}$. For more information about the semi-martingales and the Ito processes see [9].

We assume that the compensator of the measure of the jumps ν verify the usual condition: for each $t \in \mathbb{R}^+$

$$(2) \quad \int_0^t \int_{\mathbb{R} \setminus \{0\}} (x^2 \wedge 1) K_s(dx) ds < \infty.$$

We recall that the characteristic function of X_t

$$\phi_t(\lambda) = \mathbf{E} \exp(i\lambda X_t)$$

is defined by the following expression: for $\lambda \in \mathbb{R}$

$$\phi_t(\lambda) = \exp\left\{i\lambda B_t - \frac{1}{2}\lambda^2 C_t + \int_0^t \int_{\mathbb{R} \setminus \{0\}} (e^{i\lambda x} - 1 - i\lambda x \mathbf{1}_{\{|x| < 1\}}) K_s(dx) ds\right\}$$

which can be easily obtained by the Ito formula for semimartingales. We recall also that X is a semi-martingale if and only if for all $\lambda \in \mathbb{R}$ the characteristic function of X_t is of finite variation in t on finite intervals (cf. [9], Ch.2, Th. 4.14, p.106). Moreover, the process X always can be written as a sum of a semi-martingale and a deterministic function which is not necessarily of finite variation on finite intervals.

From the formula for the characteristic function we can easily find the Laplace transform of X_t , if it exists:

$$\mathbf{E}(e^{-\alpha X_t}) = e^{-\Phi(t, \alpha)}$$

Putting $\lambda = i\alpha$ in the previous formula, we get that

$$\Phi(t, \alpha) = \alpha B_t - \frac{1}{2} \alpha^2 C_t - \int_0^t \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x \mathbf{1}_{\{|x| < 1\}}) K_s(dx) ds.$$

As known, in the case when X is a Levy process with parameters (b_0, c_0, K_0) , it holds

$$\mathbf{E}(e^{-\alpha X_t}) = e^{-t\Phi(\alpha)}$$

with

$$\Phi(\alpha) = \alpha b_0 - \frac{1}{2} \alpha^2 c_0 - \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x \mathbf{1}_{\{|x| < 1\}}) K_0(dx).$$

In this article we establish necessary and sufficient conditions of the existence of the Laplace exponent, and also the sufficient conditions for finiteness of $\mathbf{E}(I_t^\alpha)$ with fixed $t > 0$ (see Proposition 1). Next, we use time reversal of the process X at fixed time $t > 0$ to introduce a new process $Y = (Y_s)_{0 \leq s \leq t}$ with $Y_s = X_t - X_{(t-s)-}$. In Lemma 1 we show that

$$I_t = e^{-Y_t} \int_0^t e^{Y_s} ds.$$

Then, we prove that Y is a process with independent increments and we identify its semi-martingale characteristics. This time reversal plays an important role and permits to replace the process $(I_t)_{t \geq 0}$ which is not Markov, by a family of Markov processes $V = (V_s)_{0 \leq s \leq t}$ with $t > 0$, where $V_s = e^{-Y_s} \int_0^s e^{Y_u} du$.

In Theorem 1 we consider the case of $\alpha \geq 0$ and we give a recurrent integral equation for the Mellin transform of I_t . In the Corollaries 2 and 3 we consider the case when X is a Levy process. We give the formulas for positive moments of I_t and I_∞ and also the formula for the Laplace transform. The results for I_∞ coincide, of course, with the ones given in [4] for Levy subordinators. But it holds under less restrictive integrability conditions on I_∞ and less restrictive condition on Levy measure at zero. We can also precise the number of finite moments of I_∞ (cf. Corollary 3 and Corollary 5). In Theorem 2 and Corollaries 4 and 5 we present analogous result for the case $\alpha < 0$.

2. FINITENESS OF $\mathbf{E}(I_t^\alpha)$ FOR FIXED $t > 0$.

Now, we fix $t > 0$. In the following proposition we give necessary and sufficient conditions for the existence of the Laplace transform of X_t , and also sufficient conditions for the existence of the Mellin transform of I_t . In what follows we assume that K verify the following stronger condition then (2): for $t > 0$

$$(3) \quad \int_0^t \int_{\mathbb{R} \setminus \{0\}} (x^2 \wedge |x|) K_s(dx) ds < \infty.$$

This condition says, roughly speaking, that the "big" jumps of the process X are integrable, and it ensure that finite variation part of semi-martingale decomposition of X remains deterministic. Moreover, the truncation of the jumps is no more necessary.

Proposition 1. *Let $\alpha \in \mathbb{R}$ and $t > 0$. If X is integrable, i.e. $\mathbf{E}(|X_t|) < \infty$ for $\forall t > 0$, then under (3), the condition*

$$(4) \quad \int_0^t \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds < \infty,$$

is equivalent to

$$(5) \quad \mathbf{E}(e^{-\alpha X_t}) < \infty.$$

Otherwise, (3) and (4) implies (5). Moreover, the condition (4) implies for $\alpha \geq 1$ and $\alpha \leq 0$ that

$$(6) \quad \mathbf{E}(I_t^\alpha) < \infty.$$

In addition, if (4) is valid for $\alpha = 1$, then for $0 \leq \alpha \leq 1$ we have (6).

Remark 1. *Assume that X has only positive jumps, then for $\alpha > 0$ the condition (4) is always satisfied. In the same way, if X has only negative jumps, then, of course, for $\alpha < 0$ the condition (4) is satisfied. If X has bounded jumps, the condition (4) is also satisfied. In general case, the condition (4) is equivalent to the one of the following conditions: if $\alpha > 0$, then*

$$(7) \quad \int_0^t \int_{x < -1} e^{-\alpha x} K_s(dx) ds < \infty,$$

and if $\alpha < 0$, then

$$(8) \quad \int_0^t \int_{x > 1} e^{-\alpha x} K_s(dx) ds < \infty.$$

In the case of Levy processes these conditions coincide with the ones given in [12], p. 79.

Remark 2. It should be noticed that if the condition (4) is verified for $\alpha' = \alpha + \delta$ with $\delta > 0$ and $\alpha > 0$, then it is verified also for α . To see this, apply Hölder inequality to the integrals in (7) and (8) with the parameters $p = \frac{\alpha+\delta}{\alpha}$ and $q = \frac{\alpha+\delta}{\delta}$. In fact, since $\frac{1}{p} + \frac{1}{q} = 1$ and $\nu([0, t] \times] - \infty, -1]) < \infty$ we have for (7):

$$\int_0^t \int_{x < -1} e^{-\alpha x} K_s(dx) ds \leq \left(\int_0^t \int_{x < -1} e^{-\alpha p x} K_s(dx) ds \right)^{\frac{1}{p}} \left(\int_0^t \int_{x < -1} K_s(dx) ds \right)^{\frac{1}{q}} < \infty$$

The same can be made for $\alpha < 0$. In this case we take $\delta < 0$.

Proof. Since X is assumed to be a process with independent increment which is a semi-martingale, it can be written as the sum of deterministic process B of finite variation on finite intervals and local martingale M :

$$X_t = B_t + M_t$$

with $B_0 = 0$ and $M_0 = 0$. Notice that, the local martingale property of M follows from the independence of the increments and the fact that PII is a strong Markov process. Moreover, local martingale M will be a martingale when X is integrable. Notice also that to prove (5) is equivalent proving

$$(9) \quad \mathbf{E}(e^{-\alpha M_t}) < \infty$$

since B_t is deterministic.

For local martingale M we take a localizing sequence $(s_n)_{n \geq 1}$, and we introduce for $n \geq 1$ the stopping times

$$\tau_n = \inf\{s \geq 0 : |M_s| \geq n\} \wedge s_n$$

with $\inf\{\emptyset\} = +\infty$. By the Ito formula we have:

$$e^{-\alpha M_{t \wedge \tau_n}} = e^{-\alpha M_0} - \alpha \int_0^{t \wedge \tau_n} e^{-\alpha M_{s-}} dM_s + \frac{1}{2} \alpha^2 \int_0^{t \wedge \tau_n} e^{-\alpha M_{s-}} d \langle M^c \rangle_s + \int_0^{t \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} e^{-\alpha M_{s-}} (e^{-\alpha x} - 1 + \alpha x) \mu_M(ds, dx),$$

where μ_M is the jump measure of M and $M_0 = 0$. We remark that on stochastic interval $\llbracket 0, t \wedge \tau_n \rrbracket$, $0 \leq e^{-\alpha M_{s-}} \leq e^{|\alpha|n}$ and then,

$$\mathbf{E} \int_0^{t \wedge \tau_n} e^{-\alpha M_{s-}} dM_s = 0.$$

We take the mathematical expectation in the previous decomposition and we use the projection theorem (see [9], Ch. 2, Th. 1.8, p.66) together with the facts that $d < M^c >_s = c_s$ and $\Delta M_s = \Delta X_s$, to obtain

$$(10) \quad \mathbf{E}(e^{-\alpha M_{t \wedge \tau_n}}) = 1 + \frac{1}{2} \alpha^2 \mathbf{E} \int_0^{t \wedge \tau_n} e^{-\alpha M_s} c_s ds \\ + \mathbf{E} \left(\int_0^{t \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} e^{-\alpha M_{s-}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds \right)$$

We also remark that the both terms on the r.h.s. of the last equality are positive since $e^{-\alpha x} - 1 + \alpha x \geq 0$ for all $x \in \mathbb{R}$. Take $n \rightarrow +\infty$, then $\tau_n \rightarrow +\infty$, and by the Fatou lemma and the Fubini theorem we get that

$$(11) \quad \mathbf{E}(e^{-\alpha M_t}) \leq 1 + \frac{1}{2} \alpha^2 \int_0^t \mathbf{E}(e^{-\alpha M_s}) c_s ds \\ + \int_0^t \mathbf{E}(e^{-\alpha M_s}) \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds$$

Then, by the Gronwall lemma

$$\mathbf{E}(e^{-\alpha M_t}) \leq \exp \left(\int_0^t \left[\frac{1}{2} \alpha^2 c_s + \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) \right] ds \right) \\ = \exp \left(\frac{1}{2} \alpha^2 C_t + \int_0^t \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds \right)$$

and using (4), we get (5) via (9).

Assume next that (9) holds and X is integrable. From (10) we get

$$\mathbf{E}(e^{-\alpha M_{t \wedge \tau_n}}) \geq 1 + \mathbf{E} \left(\int_0^{t \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} e^{-\alpha M_{s-}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds \right)$$

Again, the Fatou lemma applied to the r.h.s. of the previous inequality and the Jensen inequality applied to the l.h.s. with $M_{t \wedge \tau_n} = \mathbf{E}(M_t | \mathcal{F}_{t \wedge \tau_n})$ gives us:

$$\mathbf{E}(e^{-\alpha M_t}) \geq 1 + \int_0^t \mathbf{E}(e^{-\alpha M_s}) \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds$$

Then, using the Gronwall lemma again

$$\mathbf{E}(e^{-\alpha M_t}) \geq \exp\left(\int_0^t \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx) ds\right)$$

Then, we have the condition (4).

Let τ be uniformly distributed on $[0, t]$ random variable independent of X and $\alpha \geq 1$ or $\alpha < 0$. Then,

$$\mathbf{E}(e^{-X\tau}) = \mathbf{E}(I_t/t)$$

and by the Jensen inequality

$$(I_t/t)^\alpha \leq \frac{1}{t} \int_0^t e^{-\alpha X_s} ds$$

and

$$\mathbf{E}(I_t^\alpha) \leq t^{\alpha-1} \int_0^t \mathbf{E}(e^{-\alpha X_s}) ds$$

Then, since $(e^{-\alpha M_s})_{0 \leq s \leq t}$ is a sub-martingale and (9) holds, we have

$$\begin{aligned} \int_0^t \mathbf{E}(e^{-\alpha X_s}) ds &= \int_0^t e^{-\alpha B_s} \mathbf{E}(e^{-\alpha M_s}) ds \\ &\leq \mathbf{E}(e^{-\alpha M_t}) \int_0^t e^{-\alpha B_s} ds < \infty \end{aligned}$$

and this yields (6).

If $0 \leq \alpha \leq 1$, then by the Jensen inequality again $\mathbf{E}(I_t^\alpha) \leq (\mathbf{E}I_t)^\alpha < \infty$. \square

Let us give some examples useful in mathematical finance.

Example 1. Let X be non-homogeneous Poisson process with intensity $(\lambda_s)_{s \geq 0}$. Then, $K_s(A) = \lambda_s \cdot \delta_{\{1\}}(A)$ for all $A \in \mathcal{B}(\mathbb{R} \setminus \{0\})$, where $\delta_{\{1\}}$ is delta-function at 1, and the condition (4) is satisfied as soon as $\int_0^t \lambda_s ds < \infty$. Hence, under the last condition, $\mathbf{E}(I_t^\alpha) < \infty$ for all $\alpha \geq 1$ and $\alpha < 0$.

Example 2. Let L be Levy process with generating triplet (b_0, c_0, K_0) and X be L time changed by deterministic continuously differentiable process $(\tau(t))_{t \geq 0}$, i.e. $X_t = L_{\tau(t)}$. Then, for $t \geq 0$

$$\mathbf{E}e^{-\alpha L_{\tau(t)}} = e^{-\Phi(\alpha)\tau(t)}$$

and $b_t = b_0\tau(t)$, $c_t = c_0\tau(t)$, and for all $A \in \mathcal{B}(\mathbb{R} \setminus \{0\})$, $K_t(A) = K_0(A)\tau'(t)$. The condition (4) is satisfied whenever

$$(12) \quad \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 - \alpha x) K_0(dx) < \infty.$$

Under this condition $\mathbf{E}(I_t^\alpha) < \infty$ for $\alpha \geq 1$ and $\alpha < 0$.

Example 3. Let L be Levy process with generating triplet (b_0, c_0, K_0) and

$$X_t = \int_0^t g_s dL_s$$

where $g = (g_s)_{s \geq 0}$ is a measurable positive function. Then, $b_t = b_0 \int_0^t g_s ds$, $c_t = c_0 \int_0^t g_s^2 ds$ and since $\Delta X_s = g_s \Delta L_s$ for $s > 0$, we get that $K_s(dx) = g_s K_0(dx)$. The condition (4) is satisfied whenever we have (12) and $\int_0^t g_s^2 ds < \infty$ in case $c_0 \neq 0$ and $\int_0^t |g_s| ds < \infty$ in case $c_0 = 0$. Under these conditions, $\mathbf{E}(I_t^\alpha) < \infty$ for all $\alpha \geq 1$ and $\alpha < 0$.

3. TIME REVERSAL PROCEDURE

We introduce, for fixed $t > 0$, a new process $Y^{(t)} = (Y_s^{(t)})_{0 \leq s \leq t}$ with $Y_s^{(t)} = X_t - X_{(t-s)-}$. To simplify the notations we anyway omit the index (t) and write Y_s instead of $Y_s^{(t)}$.

First of all we establish the relation between I_t and the process Y .

Lemma 1. *For $t > 0$ the following equality holds:*

$$I_t = e^{-Y_t} \int_0^t e^{Y_s} ds$$

Proof. Using the definition of the process Y and the assumption that $X_0 = 0$ we have

$$\begin{aligned} e^{-Y_t} \int_0^t e^{Y_s} ds &= \int_0^t e^{-Y_t + Y_s} ds = \int_0^t e^{-X_{(t-s)-} + X_0} ds \\ &= \int_0^t e^{-X_{t-s}} ds = \int_0^t e^{-X_s} ds = I_t, \end{aligned}$$

since we are integrating with respect to the Lebesgue measure. \square

We will show that the process Y is PII and we will give its semi-martingale triplet with respect to its natural filtration. For that we put

$$\bar{b}_u = \begin{cases} b_{t-u} & \text{for } 0 \leq u < t, \\ b_t & \text{for } u = t. \end{cases}$$

which can be written also as

$$\bar{b}_u = \mathbf{1}_{\{t\}}(u)(b_t - b_0) + b_{t-u},$$

where $\mathbf{1}_{\{t\}}$ is indicator function of the set $\{t\}$. We do the similar definitions for \bar{c} and \bar{K} :

$$\begin{aligned} \bar{c}_u &= \mathbf{1}_{\{t\}}(u)(c_t - c_0) + c_{t-u}, \\ \bar{K}_u(A) &= \mathbf{1}_{\{t\}}(u)(K_t(A) - K_0(A)) + K_{t-u}(A) \end{aligned}$$

for all $A \in \mathcal{B}(\mathbb{R} \setminus \{0\})$.

Lemma 2. *The process Y is a process with independent increments, it is a semi-martingale with respect to its natural filtration, and its semi-martingale triplet $(\bar{B}, \bar{C}, \bar{\nu})$ is given by :*

$$(13) \quad \bar{B}_s = \int_0^s \bar{b}_u du, \quad \bar{C}_s = \int_0^s \bar{c}_u du, \quad \bar{\nu}(ds, dx) = \bar{K}_s(dx) ds,$$

where $0 \leq s \leq t$.

Proof Let us take $0 = s_0 < s_1 < s_2 < \dots < s_n = t$ with $n \geq 2$. Then the increments $(Y_{s_k} - Y_{s_{k-1}})_{1 \leq k \leq n}$ of the process Y are equal to $(X_{(t-s_{k-1})-} - X_{(t-s_k)-})_{1 \leq k \leq n}$ and $0 = t - s_n < t - s_{n-1} < \dots < t - s_0 = t$. From the fact that X is the process with independent increments, the characteristic function of the vector $(X_{t-s_{k-1}-h} - X_{t-s_k-h})_{1 \leq i \leq n}$ with small $h > 0$, can be written as a product of the corresponding characteristic functions. Namely, for any real constants $(\lambda_k)_{0 \leq k \leq n}$ we get:

$$\mathbf{E} \exp\left(i \sum_{k=1}^n \lambda_k (X_{t-s_{k-1}-h} - X_{t-s_k-h})\right) = \prod_{k=1}^n \mathbf{E} \exp(i \lambda_k (X_{t-s_{k-1}-h} - X_{t-s_k-h}))$$

Then, passing to the limit as $h \rightarrow 0+$,

$$\mathbf{E} \exp\left(i \sum_{k=1}^n \lambda_k (X_{(t-s_{k-1})-} - X_{(t-s_k)-})\right) = \prod_{k=1}^n \mathbf{E} \exp(i \lambda_k (X_{(t-s_{k-1})-} - X_{(t-s_k)-}))$$

Hence, Y is a process with independent increments.

We know that we can identify the semi-martingale characteristics w.r.t. the natural filtration of the process with independent increments from characteristic function of this process. We notice that $Y_s = X_t - X_{(t-s)-}$ and by the independence of Y_s and $X_{(t-s)-}$

$$\mathbf{E}(e^{i\lambda X_t}) = \mathbf{E}(e^{i\lambda Y_s}) \mathbf{E}(e^{i\lambda X_{(t-s)-}})$$

Then,

$$\mathbf{E} \exp(i\lambda Y_s) = \mathbf{E} \exp(i\lambda X_t) / \mathbf{E} \exp(i\lambda X_{(t-s)-}) =$$

$$\exp\left\{i\lambda \int_{t-s}^t b_u du - \frac{1}{2}\lambda^2 \int_{t-s}^t c_u du + \int_{t-s}^t \int_{\mathbb{R}\setminus\{0\}} (e^{i\lambda x} - 1 - i\lambda x) K_u(dx) du\right\}$$

We substitute u by $u' = t - u$ in the integrals to obtain

$$\begin{aligned} & \mathbf{E} \exp(i\lambda Y_s) = \\ & \exp\left\{i\lambda \int_0^s \bar{b}_u du - \frac{1}{2}\lambda^2 \int_0^s \bar{c}_u du + \int_0^s \int_{\mathbb{R}\setminus\{0\}} (e^{i\lambda x} - 1 - i\lambda x) \bar{K}_u(dx) du\right\} \end{aligned}$$

Therefore, the characteristics of Y are in (13) and the proof is complete. \square

4. RECURRENT FORMULAS FOR THE MELLIN TRANSFORM OF I_t WITH $\alpha \geq 0$.

Let us consider two important processes related with the process Y , namely the process $V = (V_s)_{0 \leq s \leq t}$ and $J = (J_s)_{0 \leq s \leq t}$ defined via

$$V_s = e^{-Y_s} J_s, \quad J_s = \int_0^s e^{Y_u} du.$$

We underline that the both processes depend of the parameter t .

We remark that according to Lemma 1, $I_t = V_t$ for each $t \geq 0$. For $\alpha \geq 0$ and $t \geq 0$ we introduce the Mellin transform of I_t of the parameter α :

$$m_t^{(\alpha)} = \mathbf{E}(I_t^\alpha) = \mathbf{E}(e^{-\alpha Y_t} J_t^\alpha)$$

and the Mellin transform for shifted process:

$$m_{s,t}^{(\alpha)} = \mathbf{E} \left[\left(\int_s^t e^{-(X_u - X_{s-})} du \right)^\alpha \right]$$

Notice that $m_{0,t}^{(\alpha)} = m_t^{(\alpha)}$. Notice also that

$$m_{s,t}^{(\alpha)} = \mathbf{E} \left[\left(\int_s^t e^{-(X_u - X_s)} du \right)^\alpha \right]$$

In fact, $X_u - X_s = X_u - X_{s-} - \Delta X_s$ and

$$\int_s^t e^{-(X_u - X_{s-})} du = e^{-\Delta X_s} \int_s^t e^{-(X_u - X_s)} du$$

Since ΔX_s and $(X_u - X_s)_{u \geq s}$ are independent, and $\mathbf{E}(e^{-\alpha \Delta X_s}) = 1$, we get the equality of two expressions for $m_{s,t}^{(\alpha)}$.

We introduce also two functions: for $0 \leq s \leq t$

$$(14) \quad H_s^{(\alpha)} = \alpha b_s - \frac{1}{2}\alpha^2 c_s - \int_{\mathbb{R}\setminus\{0\}} (e^{-\alpha x} - 1 + \alpha x) K_s(dx)$$

and

$$(15) \quad \bar{H}_s^{(\alpha)} = \mathbf{1}_{\{t\}}(s)(H_t^{(\alpha)} - H_0^{(\alpha)}) + H_{t-s}^{(\alpha)}$$

These functions represent the derivatives w.r.t. s , of the Laplace exponents $\Phi(s, \alpha)$ and $\bar{\Phi}(s, \alpha)$. We recall that

$$\Phi(s, \alpha) = \alpha B_s - \frac{\alpha^2}{2} C_s - \int_0^s \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) \nu(du, dx)$$

and

$$\bar{\Phi}(s, \alpha) = \alpha \bar{B}_s - \frac{\alpha^2}{2} \bar{C}_s - \int_0^s \int_{\mathbb{R} \setminus \{0\}} (e^{-\alpha x} - 1 + \alpha x) \bar{\nu}(du, dx)$$

where ν and $\bar{\nu}$ are the compensators of the jump measure of X and Y respectively. We notice, that these functions are well-defined under condition (4). We also notice that

$$\bar{\Phi}(s, \alpha) = \Phi(t, \alpha) - \Phi(t - s, \alpha)$$

Our aim now is to obtain a recurrent integral equation for the Mellin transform of I_t . For condition (16) below see Remark 1.

Theorem 1. *Let $\alpha \geq 1$ be fixed and assume that for $t > 0$ there exists $\delta > 0$ such that*

$$(16) \quad \int_0^t \int_{x < -1} e^{-(\alpha + \delta)x} K_s(dx) ds < \infty.$$

Then, $m_t^{(\alpha)}$ is well-defined and the following recurrent integral equation holds

$$(17) \quad m_t^{(\alpha)} = \alpha \int_0^t m_{u,t}^{(\alpha-1)} e^{-\Phi(u, \alpha)} du$$

If X is Levy process, then for all $t > 0$

$$(18) \quad m_t^{(\alpha)} = \alpha e^{-\Phi(\alpha)t} \int_0^t m_s^{(\alpha-1)} e^{\Phi(\alpha)s} ds$$

Moreover,

$$(19) \quad \frac{d}{dt} [m_t^{(\alpha)}] = -m_t^{(\alpha)} \Phi(\alpha) + \alpha m_t^{(\alpha-1)}$$

Proof. From Lemma 2 we know that $Y = (Y_s)_{0 \leq s \leq t}$ is a process with independent increments which is a semi-martingale, i.e. $Y_s = \bar{B}_s + \bar{M}_s$, where \bar{B} is a deterministic process of finite variation on finite intervals

and \bar{M} is a local martingale. Let $(\bar{s}_n)_{n \geq 1}$ be localizing sequence for \bar{M} . For $n \geq 1$ we introduce the stopping times

$$\tau_n = \inf\{0 \leq s \leq t : V_s \geq n \text{ or } \exp(-Y_s) \geq n\} \wedge \bar{s}_n$$

with $\inf\{\emptyset\} = +\infty$. For fixed s , $0 < s < t$, we write the Ito formula for $V_{s \wedge \tau_n}^\alpha$:

$$(20) \quad \begin{aligned} V_{s \wedge \tau_n}^\alpha &= \alpha \int_0^{s \wedge \tau_n} V_{u-}^{\alpha-1} dV_u + \frac{1}{2} \alpha (\alpha - 1) \int_0^{s \wedge \tau_n} V_{u-}^{\alpha-2} d \langle V^c \rangle_u \\ &+ \int_0^{s \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} ((V_{u-} + x)^\alpha - V_{u-}^\alpha - \alpha V_{u-}^{\alpha-1} x) \mu_V(du, dx) \end{aligned}$$

where μ_V is the measure of the jumps of V . Using integration by part formula, we have:

$$(21) \quad dV_u = du + J_u d(e^{-Y_u})$$

Now, again by the Ito formula, we get

$$(22) \quad \begin{aligned} e^{-Y_u} &= e^{-Y_0} - \int_0^u e^{-Y_{v-}} dY_v + \frac{1}{2} \int_0^u e^{-Y_{v-}} d \langle Y^c \rangle_v \\ &+ \int_0^u \int_{\mathbb{R} \setminus \{0\}} e^{-Y_{v-}} (e^{-x} - 1 + x) \mu_Y(dv, dx) \end{aligned}$$

Then, putting (22) into (21), we obtain

$$(23) \quad \begin{aligned} dV_u^c &= -e^{-Y_{u-}} J_u dY_u^c = -V_{u-} dY_u^c, \\ d \langle V^c \rangle_u &= V_{u-}^2 d \langle Y^c \rangle_u \end{aligned}$$

and

$$(24) \quad \Delta V_u = e^{-Y_{u-}} J_u (e^{-\Delta Y_u} - 1) = V_{u-} (e^{-\Delta Y_u} - 1),$$

where $\Delta V_u = V_u - V_{u-}$ and $\Delta Y_u = Y_u - Y_{u-}$. The previous relations imply that

$$(25) \quad \begin{aligned} V_{s \wedge \tau_n}^\alpha &= \alpha \int_0^{s \wedge \tau_n} V_{u-}^{\alpha-1} du \\ &+ \alpha \int_0^{s \wedge \tau_n} J_u V_{u-}^{\alpha-1} d(e^{-Y_u}) + \frac{1}{2} \alpha (\alpha - 1) \int_0^{s \wedge \tau_n} V_{u-}^\alpha d \langle Y^c \rangle_u \\ &+ \int_0^{s \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} V_{u-}^\alpha (e^{-\alpha x} - 1 - \alpha(e^{-x} - 1)) \mu_Y(du, dx) \end{aligned}$$

To use in efficient way the Ito formula for e^{-Y_u} given before, we introduce the processes $A = (A_u)_{0 \leq u \leq t}$ and $N = (N_u)_{0 \leq u \leq t}$ via

$$A_u = \int_0^u e^{-Y_{v-}} [-d\bar{B}_v + \frac{1}{2} d\bar{C}_v] + \int_0^u \int_{\mathbb{R} \setminus \{0\}} e^{-Y_{v-}} (e^{-x} - 1 + x) \bar{\nu}(dv, dx)$$

$$N_u = - \int_0^u e^{-Y_v} d\bar{M}_v + \int_0^u \int_{\mathbb{R} \setminus \{0\}} e^{-Y_v} (e^{-x} - 1 + x) [\mu_Y(dv, dx) - \bar{\nu}(dv, dx)]$$

We notice that A is a process of locally bounded variation and N is a local martingale with localizing sequence $(\tau_n)_{n \geq 1}$, since \bar{B} , \bar{C} are of bounded variation and

$$\int_0^u \int_{\mathbb{R} \setminus \{0\}} (e^{-x} - 1 + x) \bar{K}_s(dx) ds < \infty.$$

From (22) we get that

$$(26) \quad e^{-Y_u} = e^{-Y_0} + A_u + N_u.$$

We incorporate this semi-martingale decomposition into (25) and we take mathematical expectation. Using the fact that the expectations of martingales starting from zero are equal to zero and also applying the projection theorem, we obtain:

$$(27) \quad \mathbf{E}(V_{s \wedge \tau_n}^\alpha) = \alpha \mathbf{E} \left(\int_0^{s \wedge \tau_n} V_u^{\alpha-1} du \right) + \alpha \mathbf{E} \left(\int_0^{s \wedge \tau_n} V_{u-}^{\alpha-1} J_u dA_u \right) + \frac{1}{2} \alpha (\alpha - 1) \mathbf{E} \left(\int_0^{s \wedge \tau_n} V_u^\alpha d\bar{C}_u \right) + \mathbf{E} \left(\int_0^{s \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} V_{u-}^\alpha [e^{-\alpha x} - 1 - \alpha(e^{-x} - 1)] \bar{\nu}(du, dx) \right)$$

and, hence,

$$(28) \quad \mathbf{E}(V_{s \wedge \tau_n}^\alpha) = \alpha \mathbf{E} \left(\int_0^{s \wedge \tau_n} V_u^{\alpha-1} du \right) - \mathbf{E} \left(\int_0^{s \wedge \tau_n} V_{u-}^\alpha d\bar{\Phi}(u, \alpha) \right)$$

We remark that $\tau_n \rightarrow +\infty$ (\mathbf{P} - a.s.) as $n \rightarrow +\infty$. To pass to the limit as $n \rightarrow \infty$ in r.h.s. of the above equality, we use the Lebesgue monotone convergence theorem for the first term and the Lebesgue dominated convergence theorem for the second term. In fact, for the second term we have using (14) and (15) :

$$\left| \int_0^{s \wedge \tau_n} V_{u-}^\alpha d\bar{\Phi}(u, \alpha) \right| \leq \int_0^t V_{u-}^\alpha |\bar{H}_u^{(\alpha)}| du$$

In addition,

$$\mathbf{E} \left(\int_0^t V_{u-}^\alpha |\bar{H}_u^{(\alpha)}| du \right) \leq \sup_{0 \leq u \leq t} \mathbf{E}(V_u^\alpha) \int_0^t |\bar{H}_u^{(\alpha)}| du$$

The function $(\bar{H}_u^{(\alpha)})_{0 \leq u \leq t}$ is deterministic function, integrable on finite intervals. Hence, it remains to show that

$$(29) \quad \sup_{0 \leq s \leq t} \mathbf{E}(V_s^\alpha) < \infty.$$

By the Jensen inequality

$$V_s^\alpha \leq s^{\alpha-1} \int_0^s e^{\alpha(Y_u - Y_s)} du$$

We remark that $Y_u - Y_s = X_{(t-s)-} - X_{(t-u)-}$, and then

$$(30) \quad \mathbf{E}(V_s^\alpha) \leq s^{\alpha-1} \int_0^s \mathbf{E}(e^{\alpha(X_{t-s}-X_{t-u})}) du$$

Since the process X is a process with independent increments, we have for $0 \leq u \leq s < t$

$$\begin{aligned} \mathbf{E}(e^{\alpha(X_{t-s}-X_{t-u})}) &= \mathbf{E}(e^{-\alpha X_{t-u}}) / \mathbf{E}(e^{-\alpha X_{t-s}}) = \\ \exp\left\{-\int_0^{t-u} H_r^{(\alpha)} dr + \int_0^{t-s} H_r^{(\alpha)} dr\right\} &\leq \exp\left\{\int_0^t |H_r^{(\alpha)}| dr\right\} \end{aligned}$$

Due to the Remark 1 and the condition (16), $H^{(\alpha)}$ is integrable function on finite intervals, and hence,

$$\sup_{0 \leq s \leq t} \mathbf{E}(V_s^\alpha) \leq t^\alpha \exp\left\{\int_0^t |H_r^{(\alpha)}| dr\right\} < \infty$$

To pass to the limit in l.h.s. of (28), we show that the family of $(V_{s \wedge \tau_n})_{n \geq 1}$ is uniformly integrable, uniformly in $0 \leq s \leq t$. For that we can prove in the same way as before for (43) that

$$\sup_{0 \leq s \leq t} \mathbf{E}(V_s^{\alpha+\delta}) < \infty.$$

As is well known, this condition implies uniform integrability of the family. In fact, take $g(x) = x^{\frac{\alpha+\delta}{\alpha}}$ and verify that $\sup_{0 \leq s \leq t} \mathbf{E}(g(V_s^\alpha)) < \infty$.

After limit passage, we get that

$$(31) \quad \mathbf{E}(V_s^\alpha) = - \int_0^s \mathbf{E}(V_u^\alpha) d\bar{\Phi}(u, \alpha) + \alpha \int_0^s \mathbf{E}(V_u^{\alpha-1}) du$$

We see that each term of this equation is differentiable w.r.t. s for $s < t$. In fact, the family $(V_s^\alpha)_{0 \leq s \leq t}$ is uniformly integrable, the function $(\mathbf{E}(V_s^\alpha))_{0 \leq s \leq t}$ is continuous in s as well as $(\bar{\Phi}(s, \alpha))_{0 \leq s < t}$. We calculate the derivatives in s of both sides of the above equation and we solve the corresponding linear equation to obtain:

$$\mathbf{E}(V_s^\alpha) = \alpha e^{-\bar{\Phi}(s, \alpha)} \int_0^s \mathbf{E}(V_u^{\alpha-1}) e^{\bar{\Phi}(u, \alpha)} du$$

Now, we write that $\bar{\Phi}(u, \alpha) - \bar{\Phi}(s, \alpha) = -\Phi(t - u, \alpha) + \Phi(t - s, \alpha)$ and we let $s \rightarrow t-$ to get

$$\mathbf{E}(V_t^\alpha) = \alpha \int_0^t \mathbf{E}(V_u^{\alpha-1}) e^{-\Phi(t-u, \alpha)} du$$

Notice that $I_t = V_t$ and, hence, $\mathbf{E}(V_t^\alpha) = m_t^{(\alpha)}$. Since

$$V_u = \int_0^u e^{Y_v - Y_u} dv = \int_0^u e^{-(X_{(t-v)-} - X_{(t-u)-})} dv = \int_{t-u}^t e^{-(X_v - X_{(t-u)-})} dv$$

we also have $\mathbf{E}(V_u^{\alpha-1}) = m_{t-u, t}^{(\alpha-1)}$. Then, we obtain (17) after the change of variables in integrals replacing $t - u$ by u .

In particular case of Levy process, we use (17) and take into account that $m_{u, t}^{(\alpha-1)} = m_{t-u}^{(\alpha-1)}$ and that $\Phi(u, \alpha) = \Phi(\alpha) u$. Then, after the change of variables, the formula (18) follows as well as (19). \square

Remark 3. *It should be noticed that the relation (17) can be obtained with another technique (see [15]), based on the approach of [4].*

Now we will apply our results to calculate the moments.

Example 4. We consider Levy process $L = (L_t)_{t \geq 0}$ with generating triplet (b_0, c_0, K_0) starting from 0, and time changed by deterministic process $(\tau_t)_{t \geq 0}$ with $\tau_t = r \ln(1 + t)$ with $r > 0$, i.e. $X_t = L_{\tau_t}$ for $t \geq 0$. Then, by change of variables $u = r \ln(1 + s)$ we get

$$I_t = \int_0^t e^{-L_{r \ln(1+s)}} ds = \frac{1}{r} \int_0^{r \ln(1+t)} e^{-(L_u - u/r)} du = \frac{1}{r} \int_0^{r \ln(1+t)} e^{-\tilde{L}_u} du$$

where \tilde{L} is Levy process with generating triplet $(b_0 - \frac{1}{r}, c_0, K_0)$. We denote by $\tilde{\Phi}$ the Laplace exponent of \tilde{L} . Then for $k \geq 0$, $\tilde{\Phi}(k) = \Phi(k) - \frac{k}{r}$ and

$$\mathbf{E}(I_t) = \frac{1}{r} \int_0^{r \ln(1+t)} e^{-\tilde{\Phi}(1)u} du = \frac{1 - (1+t)^{-\Phi(1) + \frac{1}{r}}}{r\Phi(1) - 1}$$

For shifted process $X_u^{(s)} = L_{\tau_u} - L_{\tau_s}$, $u \geq s$ the corresponding moments

$$m_{s, t}^{(n)} = \mathbf{E} \left[\left(\int_s^t e^{-(L_{\tau_u} - L_{\tau_s})} du \right)^n \right] = \mathbf{E} \left[\left(\int_s^t e^{-L_{(\tau_u - \tau_s)}} du \right)^n \right]$$

since L is homogeneous process and $(L_{\tau_u} - L_{\tau_s})_{u \geq s} \stackrel{\mathcal{L}}{=} (L_{\tau_u - \tau_s})_{u \geq s}$.

We change the variable putting $\tau_u - \tau_s = r \ln(1+u) - r \ln(1+s) = v - s$ where v is new variable. We denote $v(s, t) = r \ln\left(\frac{1+t}{1+s}\right)$ and we remark that $1+u = (1+s) \exp((v-s)/r)$. Then,

$$\begin{aligned} \mathbf{E} \left[\left(\int_s^t e^{-(L\tau_u - \tau_s)} du \right)^n \right] &= \mathbf{E} \left[\left(\frac{1+s}{r} \int_s^{s+v(s,t)} e^{-\tilde{L}_{v-s}} dv \right)^n \right] \\ &= \frac{(1+s)^n}{r^n} \mathbf{E} \left[\left(\int_0^{v(s,t)} e^{-\tilde{L}_u} du \right)^n \right] \end{aligned}$$

Finally, we get for $n \geq 0$:

$$m_{s,t}^{(n)} = \frac{(1+s)^n}{r^n} \tilde{m}_{v(s,t)}^{(n)}$$

where $\tilde{m}_{v(s,t)}^{(n)}$ is n -th moment of exponential functional of \tilde{L} on $[0, v(s, t)]$.

We suppose for simplicity that $\tilde{\Phi}$ is strictly monotone on the interval $[0, n+1]$. Then, using the integral equation of Theorem 1 and the expression for the moments of Levy processes given in Corollary 1 below, we get for $n \geq 1$:

$$m_t^{(n+1)} = \frac{(n+1)!}{r^n} \sum_{k=0}^n \int_0^t (1+s)^{q_n} \frac{e^{-\tilde{\Phi}(k)v(s,t)} - e^{-\tilde{\Phi}(n)v(s,t)}}{\prod_{0 \leq i \leq n, i \neq k} (\tilde{\Phi}(i) - \tilde{\Phi}(k))} ds$$

where $q_n = n - r\Phi(n+1)$ and $\tilde{\Phi}(k)$ is the Laplace exponent of Levy process \tilde{L} , $\tilde{\Phi}(k) = \Phi(k) - \frac{k}{r}$, $1 \leq k \leq n$. To express final result we put

$$\rho(k) = r\Phi(k) - k,$$

$$\gamma(n, k) = n - k - r(\Phi(n+1) - \Phi(k)), \quad \beta(n, k) = r\Phi(k) - k$$

and

$$Q_t(n, k) = \frac{(1+t)^{\gamma(n,k)+1} - 1}{(\gamma(n, k) + 1)(1+t)^{\beta(n,k)}}$$

Then, after integration we find that for $n \geq 1$

$$\mathbf{E}(I_t^{n+1}) = m_t^{(n+1)} = (n+1)! \sum_{k=0}^n \frac{Q_t(n, k) - Q_t(n, n)}{\prod_{0 \leq i \leq n, i \neq k} (\rho(i) - \rho(k))}$$

Remark 4. *It is clear that the explicit formulas for the moments in non-homogeneous case will be rather exceptional. For numerical calculus the following formula could be useful: for all $0 \leq s \leq t$*

$$m_{s,t}^{(\alpha)} = \alpha \int_s^t m_{u,t}^{(\alpha-1)} e^{\Phi(u,\alpha) - \Phi(s,\alpha)} du$$

To obtain this formula it is sufficient to find the Laplace exponent of shifted process $(X_u^{(s)})_{s \leq u \leq t}$ with $X_u^{(s)} = X_u - X_s$. Since X is a PII, the variables X_s and $X_u - X_s$ are independent, and

$$\mathbf{E}(e^{-\alpha X_u}) = \mathbf{E}(e^{-\alpha X_s}) \mathbf{E}(e^{-\alpha(X_u - X_s)})$$

and then, the Laplace exponent of shifted process $X^{(s)}$ is given by:

$$\Phi^{(s)}(u, \alpha) = \Phi(u, \alpha) - \Phi(s, \alpha).$$

5. POSITIVE MOMENTS OF I_t AND I_∞ FOR LEVY PROCESSES

Now we suppose that X is Levy process with the parameters (b_0, c_0, K_0) . In this case the condition (3) become:

$$(32) \quad \int_{\mathbb{R} \setminus \{0\}} (x^2 \wedge |x|) K_0(dx) < \infty.$$

It should be noticed that in [4] the condition on Levy measure was

$$\int_{\mathbb{R} \setminus \{0\}} (|x| \wedge 1) K_0(dx) < \infty$$

and this condition is stronger at zero and weaker at infinity than (32). When X is Levy process, the condition (16) become: for $\delta > 0$

$$(33) \quad \int_{x < -1} e^{-(\alpha + \delta)x} K_0(dx) < \infty.$$

Corollary 1. *Let $n \geq 1$ and suppose that Φ is bijective on $[0, n] \cap \mathbb{N}$. Then,*

$$\mathbf{E}(I_t^n) = n! \sum_{k=0}^n \frac{e^{-\Phi(k)t} - e^{-\Phi(n)t}}{\prod_{0 \leq i \leq n, i \neq k} (\Phi(i) - \Phi(k))}$$

Proof. The formula follows from (18) by induction. □

We present here two examples, one of them is related with Brownian motion, and second one with compound Poisson process.

Example 5. Let us consider as X Brownian motion with drift $\mu > 0$ and diffusion coefficient $\sigma > 0$, i.e. $X_t = \mu t + \sigma W_t$ where $W = (W_t)_{t \geq 0}$ is standard Brownian motion. Then, $\Phi(\alpha) = \alpha\mu - \frac{\alpha^2 \sigma^2}{2}$ and if $\frac{2\mu}{\sigma^2}$ is not an integer, we get:

$$\mathbf{E}(I_t^n) = n! \sum_{k=0}^n \frac{e^{-(k\mu - k^2 \sigma^2 / 2)t} - e^{-(n\mu - n^2 \sigma^2 / 2)t}}{\prod_{0 \leq i \leq n, i \neq k} (i - k)(\mu - (i + k)\sigma^2 / 2)}$$

Example 6. Let X be compound Poisson process such that $X_t = \sum_{k=1}^{N_t} U_k$ where $(U_k)_{k \geq 0}$ is a sequence of independent random variables with distribution function F and N is homogeneous Poisson process with intensity $\lambda > 0$. Then, $\Phi(\alpha) = \lambda \int_{\mathbb{R} \setminus \{0\}} (1 - e^{-\alpha x}) F(dx)$ and the moments can be calculated by the formula of Corollary 1. In particular, if the U_k 's are standard normal variables, we get that $\Phi(\alpha) = \lambda(1 - e^{\alpha^2/2})$ and

$$\mathbf{E}(I_t^n) = n! \sum_{k=0}^n \frac{\exp(\lambda t(1 - e^{k^2/2})) - \exp(\lambda t(1 - e^{n^2/2}))}{\lambda^n \prod_{0 \leq i \leq n, i \neq k} (e^{k^2/2} - e^{i^2/2})}$$

For $m_t^{(\alpha)}$ we introduce the Laplace-Carson transform of the parameter $q > 0$:

$$\hat{m}_q^{(\alpha)} = \int_0^\infty q e^{-qt} m_t^{(\alpha)} dt$$

This integral is always well-defined in general sense, since the integrand is positive.

Corollary 2. (cf. [4]) Let X be a Levy process which verifies (32) and (33), and such that for fixed $\alpha \geq 1$, $m_\infty^{(\alpha)} < \infty$. Then the Laplace-Carson transforms $\hat{m}_q^{(\alpha)}$ and $\hat{m}_\infty^{(\alpha-1)}$ of $m_\infty^{(\alpha)}$ and $m_\infty^{(\alpha-1)}$ respectively, are well-defined and we have a recurrent formula:

$$(34) \quad \hat{m}_q^{(\alpha)} (q + \Phi(\alpha)) = \alpha \hat{m}_q^{(\alpha-1)}$$

In particular, for integer $n \geq 1$ such that $m_\infty^{(n)} < \infty$ we get:

$$(35) \quad \hat{m}_q^{(n)} = \frac{n!}{\prod_{k=1}^n (q + \Phi(k))}$$

As a consequence,

$$\mathbf{E}(I_\infty^n) = \frac{n!}{\prod_{k=1}^n \Phi(k)}$$

Moreover, if all positive moments of I_∞ exist, then the Laplace transform of I_∞ of parameter $\beta > 0$ is given by:

$$(36) \quad \mathbf{E}(e^{-\beta I_\infty}) = \sum_{n=0}^{\infty} \frac{(-1)^n \beta^n}{\prod_{k=1}^n \Phi(k)}$$

Proof. The first equality for the Laplace-Carson transforms follows directly from (19). The second equality can be obtained as particular case from the first one, by recurrence.

For the third one we prove that $\hat{m}_q^{(n)} \rightarrow m_\infty^{(n)}$ as $q \rightarrow 0$. In fact, $m_t^{(n)} \rightarrow m_\infty^{(n)}$ as $t \rightarrow +\infty$ and

$$\lim_{t \rightarrow +\infty} \frac{1}{t} \int_0^t m_s^{(n)} ds = m_\infty^{(n)}.$$

Let us denote $M_t^{(n)} = \int_0^t m_s^{(n)} ds$. We have for each $t > 0$ by integration by part formula:

$$\int_0^t q e^{-qs} m_s^{(n)} ds = q e^{-qt} M_t^{(n)} + \int_0^t q^2 e^{-qs} M_s^{(n)} ds$$

Then, since $M_t^{(n)}/t \rightarrow m_\infty^{(n)}$ as $t \rightarrow +\infty$,

$$\hat{m}_q^{(n)} = \int_0^\infty q e^{-qs} m_s^{(n)} ds = q^2 \int_0^\infty e^{-qs} M_s^{(n)} ds$$

Since $q^2 \int_0^\infty s e^{-qs} ds = 1$, we get

$$\hat{m}_q^{(n)} - m_\infty^{(n)} = q^2 \int_0^\infty e^{-qs} (M_s^{(n)} - s m_\infty^{(n)}) ds$$

For each $\epsilon > 0$ there exists t_ϵ such that for $s \geq t_\epsilon$, $|\frac{M_s^{(n)}}{s} - m_\infty^{(n)}| \leq \epsilon$. Then,

$$\begin{aligned} |\hat{m}_q^{(n)} - m_\infty^{(n)}| &\leq q^2 \int_0^{t_\epsilon} e^{-qs} |M_s^{(n)} - s m_\infty^{(n)}| ds + q^2 \int_{t_\epsilon}^\infty s e^{-qs} \left| \frac{M_s^{(n)}}{s} - m_\infty^{(n)} \right| ds \leq \\ & q^2 \int_0^{t_\epsilon} e^{-qs} |M_s^{(n)} - s m_\infty^{(n)}| ds + \epsilon \end{aligned}$$

We notice that

$$\lim_{q \rightarrow 0} q^2 \int_0^{t_\epsilon} e^{-qs} |M_s^{(n)} - s m_\infty^{(n)}| ds = 0$$

Then, taking $\lim_{\epsilon \rightarrow 0} \lim_{q \rightarrow 0}$ in the previous inequality we get that $\hat{m}_q^{(n)} \rightarrow m_\infty^{(n)}$ as $q \rightarrow 0$. Finally, we take the limit as $q \rightarrow 0$ in second equality, to obtain the third one. The formula for the Laplace transform of I_∞ can be proved by using Taylor expansion with remainder in Lagrange form. \square

Example 7. Let X be homogeneous Poisson process of intensity λ . Then all positive moments of I_∞ exist, and we have:

$$\mathbf{E}(e^{-\beta I_\infty}) = \sum_{n=0}^{\infty} \frac{(-1)^n \beta^n}{\lambda^n \prod_{k=1}^n (1 - e^{-k})}$$

The Laplace transform can be extended for $\beta = -i\gamma$, $\gamma \in \mathbb{R}$, since the series in the right-hand side is absolutely converging. As a conclusion, the law of I_∞ is entirely characterised by its moments.

Corollary 3. *Let $\alpha_0 = \inf\{\alpha > 0 \mid \Phi(\alpha) \leq 0\}$ with $\inf\{\emptyset\} = +\infty$. Then, $\mathbf{E}(I_\infty^n) < \infty$ if and only if $1 \leq n < \alpha_0$. In particular, for Brownian motion with drift coefficient b_0 and diffusion coefficient $c_0 \neq 0$,*

$$\Phi(n) = nb_0 - \frac{1}{2}n^2c_0$$

and the moment of I_∞ of order $n \geq 1$ will exist if $n < \frac{2b_0}{c_0}$.

If X is a subordinator with Levy measure K_0 such that (32) holds, then

$$\Phi(n) = n[b_0 - \int_{\mathbb{R}^+ \setminus \{0\}} xK_0(dx)] - \int_{\mathbb{R}^+ \setminus \{0\}} (e^{-nx} - 1)K_0(dx),$$

and under the condition

$$b_0 - \int_{\mathbb{R}^+ \setminus \{0\}} xK_0(dx) \geq 0,$$

all moments of I_∞ exist.

Proof. Let $n = \sup\{k \geq 1 : \mathbf{E}(I^k) < \infty\}$. If $n = +\infty$, then $\Phi(k) > 0$ for all $k \geq 1$ and $\alpha_0 = +\infty$. If $n < +\infty$, from Corollary 2 we get that $\Phi(n) > 0$. Since $\Phi(\alpha)$ is concave function such that $\Phi(0) = 0$, $n < \alpha_0$. Conversely, substituting $t - s$ by s in (18) of Theorem 1 we get:

$$\mathbf{E}(I_t^n) \leq n \mathbf{E}(I_\infty^{n-1}) \int_0^t e^{-\Phi(n)s} ds$$

If $n < \alpha_0$, $\Phi(n) > 0$, and the integral on the r.h.s. of this inequality converge as $t \rightarrow \infty$. By induction, it gives that $\mathbf{E}(I_\infty^n) < \infty$. Moreover, the results for continuous case and the case when X is a subordinator, follow immediately from the expression of $\Phi(k)$. \square

Example 8. Let X be time changed Brownian motion, namely $X_t = \mu\tau_t + \sigma W_{\tau_t}$ where $W = (W_t)_{t \geq 0}$ is standard Brownian motion, $\mu \in \mathbb{R}$, $\sigma > 0$ and τ_t is first hitting time of the level t of the independent (from W) standard Brownian motion $B = (B_t)_{t \geq 0}$ with the drift coefficient $b > 0$. Then, as known, $\Phi(\alpha) = (b^2 + 2\alpha\mu - \alpha^2\sigma^2)^{1/2} - b$ with $b^2 + 2\alpha\mu - \alpha^2\sigma^2 > 0$ (see for instance [18]). Then, $\mathbf{E}(I_\infty^n) < \infty$ if and only if $2\mu - n\sigma > 0$.

Example 9. Let X be pure discontinuous Levy process with Levy measure

$$K_0(dx) = \frac{c \exp(-Mx)}{x^{1+\beta}} \mathbf{1}_{]0, +\infty[}(x) dx$$

where $c > 0, M > 0, 0 < \beta < 1$. Then, $\Phi(\alpha) = \frac{c\Gamma(1-\beta)}{\beta}((M+\alpha)^\beta - M^\beta)$ (see for instance [18]). Then, $\Phi(\alpha) > 0$ for $\alpha \geq 1$, and all moments of I_∞ exist.

6. RECURRENT FORMULAS FOR THE MELLIN TRANSFORM OF I_t WITH $\alpha < 0$.

In the following Theorem 2, we derive the integro-differential equation for the Mellin transform $m_t^{(\alpha)}$ of I_t with $\alpha < 0$.

Theorem 2. *Let $\alpha < 0$ be fixed and*

$$(37) \quad \int_0^t \int_{x>1} e^{(|\alpha|+1)x} K_s(dx) ds < \infty.$$

Then, for $s > 0$, $m_{s,t}^{(\alpha)}$ is well-defined as well as $m_{s,t}^{(\alpha-1)}$ and we get the following recurrent differential equation:

$$(38) \quad m_{s,t}^{(\alpha-1)} = \frac{1}{\alpha} \left(m_{s,t}^{(\alpha)} H_s^\alpha - \frac{d}{ds} m_{s,t}^{(\alpha)} \right)$$

In the case of Levy process X we have:

$$(39) \quad m_s^{(\alpha-1)} = \frac{1}{\alpha} \left(m_s^{(\alpha)} \Phi(\alpha) + \frac{d}{ds} m_s^{(\alpha)} \right)$$

Proof. The proof of this result is similar to the proof of Theorem 1. Let $(\bar{s}_n)_{n \geq 1}$ be localizing sequence for \bar{M} . For $n \geq 1$ we introduce the stopping times

$$\tau_n = \inf\{u \geq s : V_u \leq \frac{1}{n} \text{ or } \exp(-Y_u) \geq n\} \wedge \bar{s}_n$$

with $\inf\{\emptyset\} = +\infty$. Then from the Ito formula similarly to (20) we get: for $0 < s < t$

$$(40) \quad \begin{aligned} V_{t \wedge \tau_n}^\alpha &= V_s^\alpha + \alpha \int_s^{t \wedge \tau_n} V_{u-}^{\alpha-1} dV_u + \frac{1}{2} \alpha(\alpha-1) \int_s^{t \wedge \tau_n} V_{u-}^{\alpha-2} d \langle V^c \rangle_u \\ &+ \int_s^{t \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} ((V_{u-} + x)^\alpha - V_{u-}^\alpha - \alpha V_{u-}^{\alpha-1} x) \mu_V(du, dx) \end{aligned}$$

where μ_V is the measure of the jumps of V . Using (21), (23), (24) we have

$$(41) \quad V_{t \wedge \tau_n}^\alpha = V_s^\alpha + \alpha \int_s^{t \wedge \tau_n} V_{u-}^{\alpha-1} du$$

$$\begin{aligned}
& +\alpha \int_s^{t \wedge \tau_n} J_u V_{u-}^{\alpha-1} d(e^{-Y_u}) + \frac{1}{2} \alpha (\alpha - 1) \int_s^{t \wedge \tau_n} V_{u-}^\alpha d \langle Y^c \rangle_u \\
& + \int_s^{t \wedge \tau_n} \int_{\mathbb{R} \setminus \{0\}} V_{u-}^\alpha (e^{-\alpha x} - 1 - \alpha(e^{-x} - 1)) \mu_Y(du, dx)
\end{aligned}$$

where μ_Y the measure of the jumps of Y . Taking in account (26) we, finally, find that

$$(42) \quad \mathbf{E}(V_{t \wedge \tau_n}^\alpha) = \mathbf{E}(V_s^\alpha) + \alpha \mathbf{E} \left(\int_s^{t \wedge \tau_n} V_{u-}^{\alpha-1} du \right) - \mathbf{E} \left(\int_s^{t \wedge \tau_n} V_{u-}^\alpha d\bar{\Phi}(u, \alpha) \right)$$

We remark that $\tau_n \rightarrow +\infty$ ($\mathbf{P} - a.s.$) as $n \rightarrow +\infty$. To pass to the limit as $n \rightarrow \infty$ in r.h.s. of the above equality, we use the Lebesgue monotone convergence theorem for the first term and the Lebesgue dominated convergence theorem for the second term. In fact, for second term we have:

$$\left| \int_s^{t \wedge \tau_n} V_{u-}^\alpha d\bar{\Phi}(u, \alpha) \right| \leq \int_s^t V_{u-}^\alpha |\bar{H}_u^{(\alpha)}| du$$

In addition,

$$\mathbf{E} \left(\int_s^t V_{u-}^\alpha |\bar{H}_u^{(\alpha)}| du \right) \leq \sup_{s \leq u \leq t} \mathbf{E}(V_u^\alpha) \int_s^t |\bar{H}_u^{(\alpha)}| du$$

The function $(\bar{H}_u^{(\alpha)})_{0 \leq u \leq t}$ is deterministic function, integrable on finite intervals. Hence, it remains to show that

$$(43) \quad \sup_{s \leq u \leq t} \mathbf{E}(V_u^\alpha) < \infty.$$

By the Jensen inequality

$$V_u^\alpha \leq u^{\alpha-1} \int_0^u e^{\alpha(Y_v - Y_u)} dv$$

Then, in the same way as in Theorem 1 we get that

$$\sup_{s \leq u \leq t} \mathbf{E}(V_u^\alpha) \leq s^\alpha \exp \left\{ \int_s^t |H_r^{(\alpha)}| dr \right\}$$

To pass to the limit in l.h.s. of (42), we prove that the family of $(V_{u \wedge \tau_n}^\alpha)_{n \geq 1}$ is uniformly integrable, uniformly in $u \in [s, t]$. For that we can prove in the same way as for (43) that

$$\sup_{s \leq u \leq t} \mathbf{E}(V_u^{\alpha-1}) < \infty.$$

As well known, this condition implies uniform integrability of the family. In fact, take $g(x) = x^{\frac{\alpha-1}{\alpha}}$ and verify that $\sup_{s \leq u \leq t} \mathbf{E}(g(V_u^\alpha)) < \infty$.

After limit passage we get that

$$(44) \quad \mathbf{E}(V_t^\alpha) = \mathbf{E}(V_s^\alpha) - \int_s^t \mathbf{E}(V_u^\alpha) d\bar{\Phi}(u, \alpha) + \alpha \int_s^t \mathbf{E}(V_u^{\alpha-1}) du$$

We differentiate each term of this equality to obtain that

$$\frac{d}{ds} \mathbf{E}(V_s^\alpha) + \mathbf{E}(V_s^\alpha) \bar{H}_s^{(\alpha)} - \alpha \mathbf{E}(V_s^{\alpha-1}) = 0$$

We take in account that $\mathbf{E}(V_s^\alpha) = m_{t-s,t}^{(\alpha)}$, $\mathbf{E}(V_s^{\alpha-1}) = m_{t-s,t}^{(\alpha-1)}$ and that $\bar{H}_s^{(\alpha)} = H_{t-s}^{(\alpha)}$. This gives us that

$$\frac{d}{ds} m_{t-s,t}^{(\alpha)} + m_{t-s,t}^{(\alpha)} H_{t-s}^{(\alpha)} - \alpha m_{t-s,t}^{(\alpha-1)} = 0$$

Finally, replacing $t - s$ by s we get (38). In the case of Levy processes $m_{t-s,t}^{(\alpha)} = m_s^{(\alpha)}$ due to homogeneity, and $H_{t-s}^{(\alpha)} = \Phi(\alpha)$, and this gives (39). \square

To present the results about negative moments of Levy process X with the parameters (b_0, c_0, K_0) , we introduce the condition: for $\delta > 0$

$$(45) \quad \int_{x>1} e^{(|\alpha|+1)x} K_0(dx) < \infty.$$

Corollary 4. (cf. [4]) *Let X be Levy process verifying (32) and (45), and let $\alpha \leq -1$ be fixed. Suppose that $m_\infty^{(\alpha-1)} < \infty$. Then the Laplace-Carson transforms of $m_\infty^{(\alpha)}$ and $m_\infty^{(\alpha-1)}$ are well-defined and we have a recurrent formula:*

$$(46) \quad \hat{m}_q^{(\alpha-1)} = \frac{1}{\alpha} (\hat{m}_q^{(\alpha)} \Phi(\alpha) + q \hat{m}_q^{(\alpha)})$$

and, hence,

$$\hat{m}_q^{(\alpha-1)} = \frac{1}{\alpha} \hat{m}_q^{(\alpha)} (q + \Phi(\alpha))$$

In particular, under above conditions, for integer $n \geq 2$ and $\alpha = -n$ we get:

$$(47) \quad \hat{m}_q^{(-n)} = \hat{m}_q^{(-1)} \cdot \frac{(-1)^{n-1}}{(n-1)!} \prod_{k=1}^{n-1} (q + \Phi(-k))$$

As a consequence,

$$(48) \quad \mathbf{E}(I_\infty^{-n}) = \mathbf{E}(I_\infty^{-1}) \cdot \frac{(-1)^{n-1}}{(n-1)!} \prod_{k=1}^{n-1} \Phi(-k)$$

Proof. Two first equalities given above follow directly from Theorem 2. The third one can be proved in the same way as in Corollary 2 by letting $q \rightarrow 0$. \square

Example 10. For compound Poisson process presented in Example 5 with U_k 's which follows standard normal distribution, we get for $n \geq 1$

$$\mathbf{E}(I_\infty^{-n}) = \frac{\mathbf{E}(I_\infty^{-1})}{(n-1)!} \prod_{k=1}^{n-1} (e^{\frac{k^2}{2}} - 1)$$

Corollary 5. *Let us suppose that $\mathbf{E}(I_\infty^{-1}) < \infty$ and $n > 1$. Then, $\mathbf{E}(I_\infty^{-n}) < \infty$ if and only if $\Phi(-1) < 0$. In particular, for Brownian motion with drift coefficient b_0 and diffusion coefficient $c_0 \neq 0$, it is the case when $\frac{b_0}{c_0} > 1$. If X is a subordinator with $\int_{\mathbb{R}^+ \setminus \{0\}} x K_0(dx) < \infty$, then*

$$\Phi(-1) = -[b_0 - \int_{\mathbb{R}^+ \setminus \{0\}} x K_0(dx)] - \int_{\mathbb{R}^+ \setminus \{0\}} (e^x - 1) K_0(dx),$$

and under the condition $b_0 - \int_{\mathbb{R}^+ \setminus \{0\}} x K_0(dx) \geq 0$ all negative moments of I_∞ exist.

Proof. From $\mathbf{E}(I_\infty^{-n}) < \infty$ by Cauchy-Schwartz inequality we get that for all k , $1 \leq k \leq n$, $\mathbf{E}(I_\infty^{-k}) < \infty$. Then the formula (48) yields that $\Phi(-k) < 0$ for $1 \leq k \leq n$. In particular, $\Phi(-1) < 0$. If $\Phi(-1) < 0$, then from convexity of the function Φ and $\Phi(0) = 0$, for $1 \leq k \leq n$ $\Phi(-k) < 0$. Then by recurrence, the formula (46) gives the existence of Laplace-Carson transform of $\mathbf{E}(I_\infty^{-n})$. Then by (48) we get that $\mathbf{E}(I_\infty^{-n}) < \infty$. \square

Example 11. For time changed Brownian motion considered in Example 8 we get that $\Phi(\alpha) \leq 0$ whenever $-b^2 \leq 2\alpha\mu - \alpha^2\sigma^2 < 0$. Hence, the h -th negative moments of I_∞ exists if $\mathbf{E}(I_\infty^{-1}) < \infty$ and $2\mu + n\sigma^2 > 0$. For pure discontinuous Levy process considered in Example 9, if $\mathbf{E}(I_\infty^{-1}) < \infty$, then all negative moments exist, since $\Phi(\alpha) \leq 0$ for all $\alpha < 0$.

REFERENCES

- [1] O. Barndorff-Nielsen, A.N. Shiryaev. *Change of Time and Change of Measure*. World Scientific, 2010.
- [2] J. Bertoin. *Lévy processes*, Cambridge University Press, 1996, P.266.
- [3] J. Bertoin, A. Lindler, R. Maller (2006) *On continuity Properties of the Law of Integrals of Levy Processes*, Séminaire de probabilités XLI, 1934, (2008), 137–159. Math. Review 2483729.

- [4] J. Bertoin, M. Yor. (2005) *Exponential functionals of Lévy processes*, Probability Surveys, 191-212.
- [5] A. Borodin, P. Salminen. (2002) *Handbook of Brownian motion - Facts and Formulae*, Birkhäuser Verlag, Basel-Boston-Berlin, 672.
- [6] P. Carmona, F. Petit, M. Yor (2004) *On the distribution and asymptotic results for exponential functionals of Lévy processes*, in "Exponential functionals and principal values related to Brownian motion", Biblioteca de la Revista Matemática Iberoamericana (1997) ISSN 0207-2230.
- [7] P. Carr, L. Wu. (2004) *Time-changed Lévy processes and option pricing*, Journal of Financial Economics 71,113–141.
- [8] D. Dufresne (1990) *The distribution of a perpetuity, with applications to risk theory and pension funding*. Scand. Actuarial J., 1-2, 39-79.
- [9] J. Jacod, A. Schiryaev. (1987) *Limit theorems for Stochastic Processes*, Springer-Verlag, 606.
- [10] M. Jeanblanc, M. Yor, M. Chesnay. (2009) *Mathematical Methods for Financial Markets*, Springer Finance Textbook.
- [11] C. Kardaras, S. Robertson (2014) *Continuous time perpetuities and time reversal of diffusions* ARXIV: 1411.7551v1.
- [12] A. Kyprianou. (2014) *Fluctuations of Lévy processes with applications*, Springer-Verlag, Berlin, Heidelberg, second addition, 2014.
- [13] J. Lamperti. (1972) *Semi-stable Markov Processes*. Z. Wahrscheinlichkeitstheorie verw. Gebiete, 22, 205-225.
- [14] P. Salminen, O. Wallin. (2005) *Perpetual integral functional of diffusions and their numerical computations*, Dept. of Math, Univ. of Oslo, Pure Mathematics, 35.
- [15] P. Salminen, L. Vostrikova. (2016) *On moments of exponential functionals of additive processes*, Peprint.
- [16] P. Salminen, M. Yor. (2005) *Perpetual Integral Functionals as Hitting and Occupation Times*, Electronic Journal of Probability, Vol. 10, Issue 11, 371-419.
- [17] K. Sato. *Lévy Processes and Infinitely Divisible Distributions*, Cambridge University Press, second edition, 2013.
- [18] A.N. Shiryaev. (1999) *Essentials of Stochastic Finance: Facts, Models, Theory*, World Scientific, 834.