



**HAL**  
open science

## Residual stress assessment in human facial tissue due to isotropic gradual growth

Mohammad Amin Askari, Mohammad Ali Nazari, Pascal Perrier, Yohan Payan

► **To cite this version:**

Mohammad Amin Askari, Mohammad Ali Nazari, Pascal Perrier, Yohan Payan. Residual stress assessment in human facial tissue due to isotropic gradual growth. 22nd Congress of the European Society of Biomechanics, Jul 2016, Lyon, France. hal-01390181

**HAL Id: hal-01390181**

**<https://hal.science/hal-01390181v1>**

Submitted on 31 Oct 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# RESIDUAL STRESS ASSESSMENT IN HUMAN FACIAL TISSUE DUE TO ISOTROPIC GRADUAL GROWTH

M. A. Askari (1), M. A. Nazari (1), P. Perrier (2), Y. Payan (3)

1. Mechanical Engineering Department, Faculty of Engineering, University of Tehran, Tehran, Iran

2. Univ. Grenoble Alpes, UMR CNRS 5216, GIPSA-Lab, F-38400 St.Martin d'Hères, France

3. Université Joseph Fourier, UMR CNRS 5525, TIMC-IMAG, F-38700 La Tronche, France

## Introduction

Growth and remodeling are among the main features of living tissues. Within the growth mechanics, living systems can be formulated by mechanical phenomena and continuum mechanics. As shown by Fung's experiments, biological tissues are not stress-free even when entirely unloaded. Moreover ignoring the residual stress and its effects may lead to erroneous results [1]. The exact origin of residual stresses in living tissues is not clear, but one of their main causes is tissue growth. In this research, we have implemented the concept of fictitious configuration for growth mechanics in order to determine the residual stresses due to growth in the tissues. Tissue growth itself occurs in response of the changes in the stress state, but at the same time growth under loading changes the equilibrium state of stress. Hence determining the residual stresses due to growth requires an iterative method. We propose a gradual growth method together with a loading-growth-unloading procedure to estimate tissues residual stresses. The method is applied to a model of healthy human facial tissues.

## Method

Let  $F$  shows the deformation gradient tensor. Using multiplicative decomposition, we introduce the elastic ( $F_e$ ) and growth ( $F_g$ ) part of deformation as:

$$F = F_e F_g \quad (1)$$

To model the behaviour of soft tissues, we use a 5-parameter Mooney-Rivlin function (see Table 1) as the energy strain potential with an incompressible constraint, which represents a nonlinear isotropic hyperelastic material. Since only the elastic tensor  $F_e$  generates stress, we use this tensor instead of the total deformation gradient  $F$  in the expression of strain energy potential [2].

$C_1$	$C_2$	$C_3$	$C_4$	$C_5$
2.5E+3	0	1.175E+3	0	0

Table 1: Mooney-Rivlin Material Constants Used in This Paper

We use the isotropic growth law, which is one of the most widely used models in the literature. We thus parameterize the growth tensor,  $F_g$ , as a multiple of a growth multiplier  $\theta_g$  by second order identity tensor  $I$ ,

$$F_g = \theta_g I \quad (3)$$

To determine the residual stresses due to growth, we propose a gradual growth method together with a loading-growth-unloading protocol. Based on this, we first simulated the behaviour of solve the finite element model under external mechanical loading and boundary

conditions. Then, under loading condition, growth is applied by using the growth multiplier  $\theta_g$ . In each step, this growth multiplier is increased to a maximum value with a specified velocity. Finally, in the unloading step, the external loads on the grown model are removed. The remained stresses are thus supposed to represent the residual stresses due to growth.

## Results

The loading-growth-unloading sequence was evaluated on a finite element model of face [3] with a loading representing gravity. After this loading step, growth was applied gradually to the maximum value of  $\theta_g=2$  (Fig. 1) which seems reasonable according to the literature [4]. After removing the effect of gravity, the residual stresses due to growth in the facial tissue is then computed (Fig. 2 and Table2).




Figure 1: Von-Mises stress distribution in grown mechanically loaded configuration




Figure 2: Von-Mises stress distribution in grown mechanically unloaded configuration (residual stress)

From Fig.1 to Fig.2, the stress field has been made more uniform and homogenous as a consequence of growth.

	Comp. stress	Tens. stress
<b>Loading</b>	3.17	963.56
<b>growth</b>	4.97	1225.65
<b>Unloading(res. Stress)</b>	1.21	58.79

Table 2: max and min value of stress in each step (pa)

## References

1. F. Morin et al, Computer Methods in Biomechanics and Biomedical Engineering, 18:2006-2007,2015
2. M Genet et al, J Biomech, 48:2080-2089, 2015.
3. M.A. Nazari et al. , Computer Methods in Biomechanics and Biomedical Engineering, 13:469-482, 2010.
4. A.M. Zöllner, Biomechanics and Modeling in Mechanobiology, 11:855-867,2012

