



HAL
open science

Age-related differences in affective responses to and memory for emotions conveyed by music: a cross-sectional study

Sandrine Vieillard, Anne-Laure K Gilet

► To cite this version:

Sandrine Vieillard, Anne-Laure K Gilet. Age-related differences in affective responses to and memory for emotions conveyed by music: a cross-sectional study. *Frontiers in Psychology*, 2013, 4, pp.711-10.3389/fpsyg.2013.00711 . hal-01392594

HAL Id: hal-01392594

<https://hal.science/hal-01392594>

Submitted on 4 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Age-related differences in affective responses to and memory for emotions conveyed by music: a cross-sectional study

Sandrine Vieillard^{1*} and Anne-Laure Gilet²

¹ Laboratoire de Psychologie (EA 3188), Psychology, Université de Franche-Comté, Besançon, France

² Laboratoire de Psychologie des Pays de la Loire (EA 4638), Université de Nantes, Nantes, France

Edited by:

Petri Laukka, Stockholm University, Sweden

Reviewed by:

Jonna K. Vuoskoski, University of Oxford, UK

Michaela Riediger, Max Planck Institute for Human Development, Germany

*Correspondence:

Sandrine Vieillard, Laboratoire de Psychologie (EA 3188), Université de Franche-Comté, 30 Rue Mégevand, 25030 Besançon, France
e-mail: sandrine.vieillard@univ-fcomte.fr

There is mounting evidence that aging is associated with the maintenance of positive affect and the decrease of negative affect to ensure emotion regulation goals. Previous empirical studies have primarily focused on a visual or autobiographical form of emotion communication. To date, little investigation has been done on musical emotions. The few studies that have addressed aging and emotions in music were mainly interested in emotion recognition, thus leaving unexplored the question of how aging may influence emotional responses to and memory for emotions conveyed by music. In the present study, eighteen older (60–84 years) and eighteen younger (19–24 years) listeners were asked to evaluate the strength of their experienced emotion on happy, peaceful, sad, and scary musical excerpts (Vieillard et al., 2008) while facial muscle activity was recorded. Participants then performed an incidental recognition task followed by a task in which they judged to what extent they experienced happiness, peacefulness, sadness, and fear when listening to music. Compared to younger adults, older adults (a) reported a stronger emotional reactivity for happiness than other emotion categories, (b) showed an increased zygomatic activity for scary stimuli, (c) were more likely to falsely recognize happy music, and (d) showed a decrease in their responsiveness to sad and scary music. These results are in line with previous findings and extend them to emotion experience and memory recognition, corroborating the view of age-related changes in emotional responses to music in a positive direction away from negativity.

Keywords: aging, musical emotions, emotional responses, facial muscle activity, incidental recognition, positivity effect

INTRODUCTION

Research on age differences in emotion processing has been mostly restricted to visual stimuli (e.g., facial expression, video, words, and pictures) but a growing body of research converges in indicating that music also serves as a powerful emotional trigger. For example, Blood et al. (1999) have shown that classical musical stimuli which are selected to elicit intensely pleasant emotional responses engage neural networks that are implicated in reward. Neuropsychological studies have also demonstrated that the amygdala is recruited when processing scary music (e.g., Gosselin et al., 2005, 2007). Physiological data have also put in evidence that music is a strong emotion inducer (e.g., Khalfa et al., 2002). Furthermore, music has the clear advantage of maintaining attention toward the emotions conveyed because it does not allow perceptual attention to be redirected (except if the listener takes off his/her headphones). For these reasons, music appears as a viable method to test age-related changes in emotion processing.

Among the past studies that have addressed the question of age-related changes in musical emotion processing, most of them have focused on the people's ability to recognize musical emotions (Allen and Brosigole, 1993; Laukka and Juslin, 2007; Drapeau et al., 2009; van Tricht et al., 2010; Lima and Castro, 2011; Vieillard et al., 2012). For instance, Drapeau et al.

(2009) compared healthy elderly adults and elderly adults with Alzheimer's disease in their ability to rate the extent to which the selected musical stimuli communicating happiness, peacefulness, sadness and fear (Vieillard et al., 2008) expressed each of these four emotions. Their findings showed that recognition performances of healthy older adults were relatively preserved with the highest recognition accuracy for happy stimuli. Laukka and Juslin (2007) compared young and older adults' ability to recognize anger, fear, happiness, sadness, and neutrality in short melodies performed on an electric guitar with different degrees of expressivity. In their study, the participants judged the emotional expression of each stimulus in a forced choice task comprised of anger, fear, happiness, sadness, neutral, and "other emotion" alternatives. Compared to young adults, older adults were less accurate in recognizing negative emotions such as sadness and fear, but their ability to recognize other emotion categories was spared. More recently, Lima and Castro (2011) went one step further by examining age-related changes in emotion recognition among three age groups of participants (i.e., young, middle-aged, and old). Selecting the same musical excerpts as those used in Drapeau et al. (2009)'s study, the authors asked the participants to judge the emotional intensity perceived for each stimulus on four rating scales (i.e., happiness, sadness, peacefulness, and fear)

presented simultaneously. Consistent with previous findings, the authors found an emotion-specific age-related change characterized by a stable recognition of happiness and peacefulness categories but a gradual decline in responsiveness to sad and scary music from young adulthood to older age.

As an alternative to the hypotheses of age-related differences in cognitive functioning¹ or hearing loss², the above findings have been interpreted as being the result of a combination of age-related changes in brain structure and functioning and in motivational goals. On the one hand, empirical evidence suggest that the observed decline found in older adults in negative emotion recognition may be the result of a linear reduction in the volume of the amygdala (e.g., Zimmerman et al., 2006) and/or of a decrease of the reactivity to negative information in the amygdala (e.g., Mather et al., 2004). On the other hand, Socioemotional Selectivity Theory (Carstensen et al., 1999; Carstensen, 2006) suggests that the decline in the recognition of negative emotions would reflect a motivational shift toward emotionally meaningful goals due to an increased awareness of the limited perspective of time. This so-called “positivity effect” which refers to all combinations of *enhanced* processing of positive information and *reduced* processing of negative information, has been thought of as an emotion regulation strategy to preserve high levels of well-being in later life. In a recent literature review, Reed and Carstensen (2012) showed that this positivity effect (1) requires cognitive resources (e.g., Mather and Knight, 2005), (2) is sensitive to the experimental context (Kensinger et al., 2002; Grühn et al., 2005), and (3) is adaptive, i.e., it emerges when emotional well-being is prioritized. Consequently, the authors claimed that the positivity effect would represent a controlled shift in attentional resources rather than an *automatic* process associated with the neuronal degeneration in brain regions. Such view is compatible with the idea that the positivity effect may have a cognitive counterpart.

Recently, Vieillard et al. (2012) conducted a study in order to test for age-related differences in the psychological structure of musical emotions, and to assess whether these changes may be associated with a decrease in emotional complexity. In this research, younger and older participants were presented with musical excerpts conveying different emotions such as happiness, peacefulness, sadness, and fear (Vieillard et al., 2008). Participants were asked to perform an emotional judgment task using different rating scales (i.e., valence, hedonic value, arousal, and liking) as well as a free categorization task in which they freely created emotional categories based on the perceived acoustical cues. Findings showed age-related differences characterized by a reduced processing of arousal for scary music, an increased focus on happy music, and an emotional dedifferentiation corresponding to a decrease in differentiation between the arousal and valence dimensions. Such results have been explained within the framework of the Dynamic Integration Theory (Labouvie-Vief, 2009; Labouvie-Vief et al., 2010) postulating that the degradation of emotional complexity would be the cognitive counterpart of the

older adults’ attempt to maximize positive affects and minimize negative affect in order to preserve well-being.

In short, the studies reviewed above show converging evidence for a positivity effect in how emotions in music are perceived, categorized, and recognized with advancing age. However, several questions remain open. First, little is known about age-related changes regarding the emotions *experienced* while listening to music. Previous findings have suggested that participants were more accurate in their judgment of intended emotions in musical excerpts when focusing on their own emotional experience (Vieillard et al., 2008). One of the goals of the present study was thus to investigate the influence of aging on emotion processing while being personally engaged in musical listening. This is an important question because it has been suggested that emphasizing on emotion rather than on knowledge may be more meaningful to older adults (e.g., Mikels et al., 2010). Second, as far as we know, the possibility of age-related changes in memory recognition for positive musical stimuli has yet to be examined. Past research showed that age-related changes in emotional goals influence memory. A positivity effect in memory recognition tasks has already been shown in older adults for affective pictures (e.g., Charles et al., 2003; Mather and Carstensen, 2005) and for words (e.g., Kensinger, 2008). The main explanation was that memory can work as an elaborative process to regulate emotions such that the older adults’ goal to maintain well-being would influence mental constructions of the past and thus lead to a positivity effect in the way they remember events. In line with this hypothesis, it has been showed that memory for negative pictures decreased in older adults both in recall and recognition tasks (e.g., Charles et al., 2003). Kensinger (2008) found a positivity effect in older adults for non-arousing words, explaining this as an age-related difference in the way positive information was primarily processed as a function of differences in motivational goals at each age level. To date, the question remains open whether age-related differences in the elaborative processing of memory may be observed in a non-verbal channel of emotion communication such as music. Third, to our knowledge, no previous study has investigated age differences in facial muscle activity when listening to music. Past studies examining age-related differences in facial expressiveness have found that young and older adults express similar patterns of facial responding to visual stimuli such as emotional scenes, objects or faces (Reminger et al., 2000; Smith et al., 2005; Bailey et al., 2009). However, older adults compared to younger adults may exhibit diminished reactivity in facial expressiveness (Smith et al., 2005; Burriss et al., 2007). The reduction of facial expressiveness in the elderly has been thought to be a possible consequence of general physiological losses in the nervous system. However, another explanation suggests that the reduction of facial expressiveness may reflect an attempt to regulate emotion since facial expression may be motivationally driven (Smith et al., 2005). Given such emotion regulation hypothesis and in line with the embodiment theory of emotion (Niedenthal et al., 2005), one can imagine that facial expressions may not only help to down-regulate emotion (by displaying less facial expressions), but also allow to modify the emotional reaction (by displaying a facial expression contrary to what one feels). In this perspective, facial electromyogram (EMG) appears as an interesting indicator

¹ Emotion categories that were less accurately recognized by older adults were not those known to be the hardest to recognize.

² Age groups did not differ or only differed marginally on self-reported hearing loss.

of whether there is congruence between facial expressivity and musical emotion in both young and older adults, or do older adults express positive facial expression as a means to counteract negative emotions.

There is an agreement to consider that emotional responding is a multi-component process, giving rise to affective experiences, physiological adjustments and expressive behaviors (e.g., Scherer, 2005). These various aspects of emotion may be differentially influenced by age. Therefore, a unifying view of these changes is necessary to give more insight into the lifespan developmental course of emotion, particularly in the musical domain in which this topic has remained unexplored. To this end, we focused on different indexes of emotional response to music, namely subjective experience, emotion expression and memory recognition for musical excerpts that conveyed different emotions.

Finally, although the positivity effect has been observed across a number of experimental paradigms such as dot-probe tasks (Charles et al., 2003), eye-tracking paradigms (Isaacowitz et al., 2006a,b), working memory (Mikels et al., 2005), memory recognition and free recall tasks (Charles et al., 2003), and across a variety of stimuli (e.g., pictures, word lists, facial expressions), the robustness of this phenomenon has been mainly demonstrated through the visual channel of emotion communication. Consequently, research regarding the effect of age on emotion processing in music is needed to test for the generalizability of the positivity effect.

CURRENT RESEARCH

The present study was designed to further extend previous studies and expand experimental designs to the domain of music. Our aim was twofold: first, to investigate the effects of aging on the emotion felt when listening to music and second, to address age-related differences on memory recognition for musical excerpts as a function of their intended emotion. To this end, we used a set of musical stimuli expressing happiness, peacefulness, sadness, and fear which were all controlled for valence and arousal (Vieillard et al., 2008). We used a rating task focused on the emotion experienced by the participants rather than on the emotion recognized by the participants. In order to address more extensively the effect of aging on the experienced emotion, we designed the experiment so that the subjective report of the intensity of the emotion felt was coupled with a recording of participants' facial expressions. These particular indexes were chosen since past research has shown that facial expressions, measured by the corrugator (i.e., frowning) and zygomatic (i.e., smiling) muscle activity, were mostly related to the valence in music: positive emotions generally lead to increased zygomatic activity, while negative ones were associated more with increased corrugator activity (e.g., Witvliet and Vrana, 2007; Khalifa et al., 2008). Because it has been suggested that facial EMG may also be voluntarily modulated to serve emotion regulation goal (Smith et al., 2005), we used this index to examine to what extent and how older adults show positive or negative facial expression as a function of musical emotions.

Based on the hypothesis postulating a motivated attention toward positivity with advancing age, and given the view that situations relevant to a person's motivational goals may elicit more intense emotional experience (e.g., Charles and Piazza,

2007), we expected that older adults, compared to younger adults, would judge their feeling as more intense when listening to positive musical excerpts (especially happy music that is more arousing than peaceful one) than when listening to negative musical excerpts. Since no age-related changes in facial expressivity were found in previous research (e.g., Levenson et al., 1991; Tsai et al., 2000; Magai et al., 2006), we also expected older adults to be spared in their facial expressions (i.e., corrugators and zygomatic muscle activity). More specifically, if older adults have spontaneous facial activity, it is expected that they would display a greater zygomatic activity for positive music in comparison with their younger counterparts. At the same time, it is also expected that older adults would display voluntary facial expression as a tool to manage emotion. In this hypothesis, older adults would show reduced expressivity or incongruent expression, in particular in response to negativity. Moreover, in view of the scarce data available on the influence of aging on memory recognition for musical excerpts that convey different emotions, and because the memory elaborative processes for music are based on more abstract information than those involved in the memory for visual and autobiographical material, it is difficult to predict the nature of the effects likely to be observed. Consequently, we conducted an exploratory approach to test whether the positivity effect may be generalized to memory recognition for musical excerpts conveying different emotions. We expected that compared to younger adults, older adults should better recognize positive musical excerpts than negative ones.

METHOD

PARTICIPANTS

A total of 40 native French speaking volunteers (22% amateur musicians³) participated in the present study. Exclusion criteria included the presence of uncorrected hearing, medical or psychiatric antecedents, psychotic symptoms, and history of substance abuse. As a result, the data of 18 young adults (19–24 years, $M = 21$ years; 61 % females) and 18 older adults (60–84 years, $M = 66$ years; 83% females) was analyzed. Younger and older adults were recruited, respectively at the psychology department of the University of Franche-Comté and through senior social programs in Besançon. Participants did not receive financial compensation for their participation.

APPARATUS

Participants were tested individually in a quiet room at stable ambient temperature at the University. Facial muscle activity was monitored continuously during the listening and rating phases using an MP150 Biopac system (Biopac Systems, Inc., Goleta, CA) at a sampling rate of 500 Hz and processed using AcqKnowledge software. Eprime software (Schneider et al., 2002) was used for excerpts presentation and ratings recording. Musical excerpts were presented binaurally through Professional 240 Sennheiser headphones.

³The musical training was measured as the proportion of participants who received at least 3 years of formal training and who were still practicing a musical instrument without reaching professional levels.

MATERIALS

Forty short musical excerpts, computer-generated in a piano timbre and taken from Vieillard et al. (2008) set of unfamiliar musical stimuli were selected for their power to convey four distinct emotions (i.e., happiness, peacefulness, sadness, and fear). Musical excerpts were controlled for their valence (unpleasant vs. pleasant) and arousal (low vs. high). Each emotion category included ten musical excerpts that lasted an average of 10 s. The happy excerpts were written in a major mode at an average tempo of 137 Metronome Markings (MM range: 92–196), with the melodic line lying in the medium high pitch range (the pedal was not used). The peaceful excerpts were composed in a major mode, had an intermediate tempo (mean: 74 MM, range: 54–100), and were played with pedal and arpeggio accompaniment. The sad excerpts were written in a minor mode at an average slow tempo of 46 MM (range: 40–60), with the pedal. The scary excerpts were composed with minor chords on the third and sixth degree, hence implying the use of many out-of-key notes. Although most scary excerpts were regular and consonant, a few had irregular rhythms and were dissonant. Their tempo varied from 44 to 172 MM. Examples can be heard at www.brams.umontreal.ca/peretz. A previous study that was conducted to examine the effect of age on emotion perception in music demonstrated that older listeners successfully distinguished happiness, peacefulness, sadness, and fear conveyed by these musical excerpts (Vieillard et al., 2012). In a study phase described below, participants were presented with 20 musical excerpts (i.e., 5 happy, 5 peaceful, 5 sad, and 5 scary) and were instructed to indicate what they experienced in terms of Emotional Intensity. The 20 remaining musical excerpts were then used as lures in the incidental recognition task.

PROCEDURE

The experiment was divided into two sessions separated by an interval of ~1 week. During the first session, participants completed a consent form and were asked about their age, musical listening, education level, self-reported health, visual and auditory acuity, and medical history. Auditory perception was controlled using free AudioTest software (www.cotral.com). More specifically, it was assessed by presenting pure tones at intervals between 500 and 8000 Hz to both ears through Professional 240 Sennheiser headphones. For each participant, the lowest sound pressure level at which each frequency was detected was recorded. In addition, several tests assessing general cognitive function (MMSE, Petit et al., 1998), fluid intelligence (Raven's progressive matrices, set I, Raven et al., 1998, updated 2003), and working memory (letter-number sequencing from WAIS-III, Wechsler, 2000) were administered. The first session lasted about an hour.

During the second session, physiological sensors were attached while the participants sat comfortably in a quiet room in the presence of the experimenter. To prevent participants from focusing on their facial muscles, they were informed that the electrodes placed on their face were used to record their electrodermal activity during the experiment. At the beginning of the session, two musical excerpts different from those used in the experiment were used in order to adjust the volume of the headphones

for each participant. In the study phase, two practice excerpts (1 happy and 1 peaceful) following by 20 excerpts (5 of each emotion) were then presented binaurally. After each trial, participants were asked to rate the intensity of the emotion felt using a 10-point scale ranging from 0 "weak" to 9 "strong." Facial muscle activity was also recorded. The excerpts were presented in two pseudo-randomized orders that were created to ensure that no more than two excerpts of the same emotion category were presented consecutively. Each musical excerpt was preceded and followed by two baseline periods of at least 10 s of silence.

Before the incidental recognition task, participants completed two questionnaires to assess depression (Beck et al., 1998) and anxiety (Spielberger, 1993; Beck et al., 1998). In the recognition task participants were asked to indicate whether an excerpt had been heard before ("old") or not ("new") by pressing the appropriate key. In this phase, 40 musical excerpts (i.e., 20 old excerpts and 20 new excerpts) were randomly presented.

Finally, participants were instructed to listen to the same set of 40 musical excerpts presented binaurally in a randomized order. After each musical excerpt, participants were asked to judge to what extent they experienced "happiness," "peacefulness," "sadness," or "fear" using a 10-point scale ranging from 0 "not at all" to 9 "a lot." Accordingly, each excerpt was presented four times; each presentation was associated with one of the four emotion scales. The presentation order of each musical excerpt and each rating scale was fully randomized across participants. The second session lasted about 2 h. At the end of the session, participants were fully debriefed.

DATA ACQUISITION AND TRANSFORMATION

Facial EMG activity (μ Volts) was recorded over the left corrugator and zygomatic sites, using two pairs of 8 mm Ag/AgCl shielded electrodes filled with isotonic gel. The EMG data were band-pass filtered from 100 to 500 Hz and processed with a root mean square algorithm over 20 samples (with a 100-ms window). Recording artifacts were visually identified and discarded from the sample. These corresponded to less than 0.5% of all measurements.

RESULTS

SAMPLE CHARACTERISTICS

Younger adults reported more years of education than the older adults, $t_{(34)} = -2.33, p < 0.05$. A chi-square goodness-of-fit test (χ^2) indicated no significant differences between age groups in the proportion of formal musical training of at least 3 years $\chi^2(1, N = 36) = 4.5, p > 0.05$. Age groups did not differ regarding their depression, state anxiety, or trait anxiety scores⁴.

⁴BDI-II scores indicated that 75% ($n = 27$) of the participants scored below the cut-off (score of 11) for a minimum depression symptomatology and that 19.44% ($n = 7$) of the participants scored below the cut-off (score of 19) for at least mild depression. The remaining two participants (one young and one older adult) scored between 22 and 24, which indicates moderate depression symptomatology. STAI state anxiety scores indicated that 83.33% ($n = 30$) of the participants scored below the cut-off (score of 35) for very low state anxiety, 13.89% ($n = 5$) of them scored below the cut-off (score of 45) for low state anxiety, and that 2.78% ($n = 1$) of them scored below the cut-off (score of 55) for mild state anxiety. STAI trait anxiety scores indicated that 30.56% ($n = 11$) of the participants scored below the cut-off (score of 35) for very low

457 Non-parametric Mann-Whitney U -test⁵ performed on the audi- 514
 458 tory thresholds (dB) did not show statistically significant differ- 515
 459 ences between younger and older adults ($U = 10.50, z = 0.77,$ 516
 460 $p = 0.44$ for 500 Hz; $U = 0, z = 0, p = 1$ for 1000 Hz; $U =$ 517
 461 $10, z = 0.84, p = 0.40$ for 2000 Hz; $U = 8, z = 1.12, p = 0.26$ 518
 462 for 4000 Hz; $U = 33, z = 0.23, p = 0.81,$ for 8000 Hz). As was 519
 463 expected, younger adults scored better than older adults on fluid 520
 464 intelligence (Raven's progressive matrices, set I, Raven et al., 1998, 521
 465 updated 2003), $t_{(34)} = -2.84, p < 0.05$. Younger adults tended 522
 466 to perform better than older adults on a working memory test 523
 467 (Digit Span from WAIS-III, Wechsler, 2000), $t_{(34)} = -1.94, p =$ 524
 468 0.06 . There were no significant age differences on self-reported 525
 469 health, $t_{(34)} = -0.81, p = 0.42$. Finally, the Mini Mental State 526
 470 Examination (MMSE; Petit et al., 1998) scores for the older adults 527
 471 suggested no apparent signs of dementia ($M = 29.7, 28-30$). 528
 472 Sample characteristics are detailed in **Table 1**. 529

473 **EMOTION INTENSITY FELT**

474 A mixed model analysis of variance was conducted on the mean 530
 475 score of the Emotional Intensity Felt with Age Group (younger 531
 476 adults, older adults) as the between-subjects factor and Intended 532
 477 Emotion (happiness, peacefulness, sadness, fear) as the within- 533
 478 subjects factor. 534

479 **Table 1 | Sample characteristics.**

	Age group	
	Young (18)	Older (18)
DEMOGRAPHIC CHARACTERISTICS		
Age (year)	19–24	61–74
Sex	61% Women	83% Women
Education (year)*	15 (3.17)	13 (1.57)
Musical training (proportion)	0.22	0
Self reported health (Max. 5)	4.44 (0.86)	4.22 (0.79)
COGNITIVE SCORES		
Advanced progressive matrices (Set 1, Max. 12)*	10.00 (1.91)	7.78 (2.71)
Letter-number sequencing (Max. 21)	11.89 (3.05)	10.17 (2.23)
MMSE (Max. 30)	–	29.7
AFFECTIVE SCORES		
Depression (BDI-II, Max 63)	8.33 (6.79)	8.67 (5.30)
State Anxiety (STAI-Y, Max 80)	30.55 (6.35)	27.83 (6.72)
Trait Anxiety (STAI-Y, Max 80)	40.83 (10.26)	40.17 (9.57)

480 *Standard deviations are listed in parentheses. *Significant difference at $p < 0.05$.*

481 state anxiety, 41.67% ($n = 15$) scored below the cut-off (score of 45) for low 535
 482 trait anxiety, and 19.44% ($n = 7$) scored below the cut-off (score of 55) for 536
 483 moderate trait anxiety. The remaining two participants (one young and one 537
 484 older adult) scored between 57 and 58, which correspond for a high anxiety 538
 485 state. 539

486 ⁵Because the audio test software did not provide absolute thresholds, we used 540
 487 the lowest sound pressure level (dB) at which each frequency was detected 541
 488 as an ordinal data. For this reason, the non-parametric Mann-Whitney test 542
 489 (independent samples) was used. Although hearing thresholds varied as a 543
 490 function of age, the Mann-Whitney test did not reveal significant differences 544
 491 between age groups probably because of the variability observed within the 545
 492 groups, in particular in the older adults. 546

Emotions (happiness, peacefulness, sadness, fear) as the within- 514
 subjects factor. 515

As illustrated in **Figure 1**, we found a significant Age Group by 516
 Intended Emotion interaction, $F_{(3, 102)} = 3.75, p < 0.05, \eta_G^2 =$ 517
 0.06^6 . In order to test our hypothesis, we computed a planned 518
 comparison between the emotion intensity felt by young adults 519
 and that felt by older adults when listening to happy music. As 520
 expected, older adults compared to young adults reported experi- 521
 encing higher emotional intensity when listening to happy music, 522
 $F_{(1, 34)} = 5.57, p < 0.05, \eta_G^2 = 0.14$. This older adults' reactivity 523
 for positivity was also confirmed by another set of planned com- 524
 parisons indicating that older adults reported experiencing higher 525
 emotional activation when listening to happy music than when 526
 listening to sad music, $F_{(1, 34)} = 7.34, p < 0.05, \eta_G^2 = 0.18,$ or 527
 scary music, $F_{(1, 34)} = 8.76, p < 0.05, \eta_G^2 = 0.20,$ while younger 528
 adults did not. No other significant effect was found. A separ- 529
 ate analysis with years of education, fluid intelligence scores, and 530
 working memory performances (i.e., factors that were found to 531
 be different between the two age groups) as covariates indicated 532
 that the Age Group by Intended Emotion interaction remained 533
 significant, $F_{(3, 93)} = 3.14, p < 0.05, \eta_G^2 = 0.05$. 534

535 **FACIAL MUSCLE ACTIVITY**

536 Facial EMG responses were calculated as the difference between 537
 the signal (Area under the curve, $\mu V \cdot sec$) over the time course 538
 of the musical excerpt and a baseline EMG level measured from 539
 1s prior to the onset of the excerpt (time -1 to 0 s) to the begin- 540
 ning of the excerpt. The area under the curve was extracted 541
 within these two time windows and was averaged for each con- 542
 dition and for each participant. Two participants were excluded 543
 (1 younger and 1 older adults) from the initial sample due to 544
 technical problems. Analyses were then conducted on seventeen 545
 younger and seventeen older adults. The Shapiro-Wilk normality 546
 test reached significance for the sample set of EMG data mean- 547
 ing that the assumption of normality has to be rejected. As a 548
 result, statistical analyses were performed using non-parametric 549
 tests. First, the differences between age groups were tested sep- 550
 arately for zygomatic and corrugator using the Mann-Whitney 551
 U -test. Results showed a significant effect of Age Group both 552
 for zygomatic ($U = 42, z = 3.51, p < 0.001$) and for corrugator 553
 muscle ($U = 83, z = 2.10, p < 0.05$) indicating that facial activ- 554
 ity was more important in older adults than in their younger 555
 counterparts. Friedman repeated measures analyses of variance 556
 (RM-ANOVA) were conducted separately on zygomatic and on 557
 corrugator activity to test the effect of Intended Emotion factor 558
 for each Age Group. For zygomatic muscle, data revealed a signifi- 559
 cant effect of Intended Emotion in older adults, $\chi^2 = 12.46, df =$ 560
 $3, p < 0.05,$ but not in young adults, $\chi^2 = 2.01, df = 3, p = 0.57$. 561
 As shown in **Figure 1**, older adults showed an increased zyo- 562
 gomatic activity, in particular for scary music. Regarding corrugator 563
 activity, older adults showed an increased activity, in particular for 564
 scary music. 565

⁶We computed generalized eta squared statistics (η_G^2) with the aim to yield 566
 measures of effect size comparable across a wide variety of research designs 567
 (Bakeman, 2005), regardless of whether the factor is between or within sub- 568
 jects. These effect-size measures provide indices of effect that are consistent 569
 with Cohen (1988) guidelines indicating that $\eta_G^2 = 0.01$ corresponds to a 570
 small effect, $\eta_G^2 = 0.09$ to a medium effect, and $\eta_G^2 = 0.25$ to a large effect.




FIGURE 1 | Mean score and standard error of emotion intensity felt, zygomatic and corrugator muscle activity as a function of intended emotions (happiness, peacefulness, sadness, fear) and age group (younger adults, older adults).

activity, no significant effect of Intended Emotion was found either in older adults, $\chi^2 = 3.00$, $df = 3$, $p = 0.39$, or in younger adults, $\chi^2 = 5.68$, $df = 3$, $p = 0.13$.

INCIDENTAL RECOGNITION TASK

Proportions of hits, false alarms, and corrected recognition scores (hits minus false alarms) are reported in **Table 2**. An analysis of variance was conducted on corrected recognition scores with Age Group (younger adults, older adults) as the between-subjects factor and Intended Emotion (happiness, peacefulness, sadness, fear) as the within-subjects factor. Analysis showed a significant main effect of Age Group indicating that younger adults recognized more musical stimuli than older adults, $F_{(1, 34)} = 30.87$, $p < 0.001$, $\eta_G^2 = 0.50$. There was also a significant main effect of Intended Emotion, $F_{(3, 102)} = 4.86$, $p < 0.05$, $\eta_G^2 = 0.12$, indicating that scary music stimuli were better recognized than peaceful and sad music stimuli. This was confirmed by *post-hoc* Bonferroni comparisons ($ps < 0.05$). No significant Age Group by Intended Emotion interaction was observed, $F_{(3, 102)} = 0.27$, $p = 0.84$, $\eta_G^2 = 0.01$. Younger adults' recognition performances were at chance level or above only for the highly arousing musical stimuli such as happy (52%) and scary (61%) ones, while older adults' recognition performances varied between 10 and 39% through the four intended emotions.

Additional separate analyses conducted on hits and false alarms revealed that older adults generated more false alarms than younger adults, $F_{(1, 34)} = 18.47$, $p < 0.001$, $\eta_G^2 = 0.35$. There was also a significant main effect of Intended Emotion, $F_{(3, 102)} = 22.02$, $p < 0.001$, $\eta_G^2 = 0.36$, as well as a significant Intended Emotion by Age Group interaction for false alarm rates, $F_{(3, 102)} = 4.85$, $p < 0.05$, $\eta_G^2 = 0.08$. *Post-hoc* Bonferroni comparisons showed that the older listeners, compared with their younger counterparts, had more difficulty to correctly reject new happy musical stimuli ($p < 0.05$). Moreover, older adults were better to correctly reject new scary musical stimuli in comparison to all other intended emotions ($ps < 0.001$) while younger adults

were better to correctly reject new scary musical stimuli in comparison to only low arousing stimuli like peaceful ($p < 0.05$) and sad ($p < 0.05$) ones. No significant effect or interaction was found for hits rates. Again, the analysis of covariance with years of education, fluid intelligence scores, and working memory performances as covariates showed that the Intended Emotion by Age Group interaction remained significant, $F_{(3, 93)} = 3.70$, $p < 0.05$, $\eta_G^2 = 0.07$.

D-prime was calculated using tables for d-prime and beta available in Hochhaus (1972) and analyzed using another mixed ANOVA. This indicates the ability to discriminate between true targets and false targets (Green and Swets, 1966), with Age Group (younger adults, older adults) as the between-subjects factor and Intended Emotion (happiness, peacefulness, sadness, fear) as the within-subjects factors. We obtained a significant main effect of Age, $F_{(1, 34)} = 26.41$, $p < 0.001$, $\eta_G^2 = 0.44$ indicating that, overall, older adults showed lower d-prime score ($M = 0.43$, $SE = 0.15$) than younger adults ($M = 1.50$, $SE = 0.15$). This suggests a lower sensitivity in the discrimination of true musical excerpts from false musical excerpts in older adults. No other significant interaction was found with d-prime as the dependent variable. The beta value that indicates the minimum level of activation necessary for a participant to respond to a true target (Green and Swets, 1966) was also calculated (Hochhaus, 1972). No significant main effect or interaction was found with beta value as the dependent variable.

DIFFERENTIATION IN EMOTION FELT

As in previous studies (Vieillard et al., 2008), we derived the best label attributed to each musical excerpt by each participant. This was done selecting the label (i.e., happy, peaceful, sad, scary) that had received the maximal rating. When the maximal rating corresponded to the label that matched the intended emotion, a score of 1 was given. When the maximal rating did not correspond to the emotion, a score of 0 was given. When the highest rating was given for more than one label, the response was considered

Table 2 | Hit rates, False Alarms Rates (Fas) and Corrected Recognition (CR) by Emotions Category and Group Age.

Age group	Intended emotions											
	Happiness			Peacefulness			Sadness			Fear		
	Hits	FA	CR	Hits	FA	CR	Hits	FA	CR	Hits	FA	CR
YOUNG (n = 18)												
M	0.64	0.12	0.52	0.70	0.31	0.39	0.70	0.32	0.38	0.64	0.03	0.61
SE	0.05	0.06	0.05	0.06	0.07	0.07	0.07	0.07	0.09	0.07	0.03	0.07
OLDER (n = 18)												
M	0.74	0.56	0.18	0.66	0.56	0.10	0.63	0.50	0.13	0.46	0.07	0.39
SE	0.05	0.06	0.05	0.06	0.07	0.07	0.07	0.07	0.09	0.07	0.03	0.07

Table 3 | Mean percentage of the label that received the maximal rating of Emotion Felt by younger and older listeners as a function of the Intended Emotions.

Intended Emotion	Emotion felt				
	Happiness	Peacefulness	Sadness	Fear	Ambivalent
Young adults					
Happiness	81	5	2	1	11
Peacefulness	13	37	28	1	21
Sadness	1	20	56	10	13
Fear	10	2	9	72	7
Older adults					
Happiness	73	1	1	1	24
Peacefulness	5	37	22	7	29
Sadness	3	24	35	6	32
Fear	9	4	28	34	25

Bold indicates the match between Emotion Felt and Intended Emotions. Ambivalent responses correspond to highest ratings given to more than one label.

ambivalent and received a score of 0. For example, when an excerpt was perceived eliciting both peacefulness and sadness to the same degree (e.g., with a rating of 7), it was considered as ambivalent. Best labels scores are presented in **Table 3**.

A mixed model analysis of variance was conducted on the mean Best Label with Age Group (younger adults, older adults) as a between-subjects factor and Intended Emotions (happiness, peacefulness, sadness, fear) and Emotion Felt (happiness, peacefulness, sadness, fear) as within-subjects factors. Significant main effects of Age Group, $F_{(1, 34)} = 7.43, p < 0.05, \eta_G^2 = 0.18$, and Emotion Felt, $F_{(3, 102)} = 5.87, p < 0.001, \eta_G^2 = 0.14$ were found. As expected, results also indicated a significant Intended Emotion by Emotion Felt interaction, $F_{(9, 306)} = 71.45, p < 0.001, \eta_G^2 = 0.58$, as well as a significant Intended Emotion by Emotion Felt by Age Group, $F_{(9, 306)} = 5.10, p < 0.001; \eta_G^2 = 0.09$. No other significant main effect or interaction was observed. We first compared the experience emotion between younger and older adults for each intended emotion. The results indicated that older adults, compared to younger adults, reported experiencing lower levels of sadness when listening to sad music, $F_{(1, 34)} = 5.44, p < 0.05, \eta_G^2 = 0.14$, but reported higher levels of sadness when listening to scary music, $F_{(1, 34)} = 6.35, p < 0.05, \eta_G^2 = 0.16$. Moreover,

when listening to scary music, older adults reported experiencing lower levels of fear, $F_{(1, 34)} = 18.78, p < 0.001, \eta_G^2 = 0.36$, than their younger counterparts. The second set of comparisons was conducted to compare the emotion experienced for each intended emotion within each age group. The results showed that older adults reported similar levels of sadness and peacefulness when listening to peaceful music, $F_{(1, 34)} = 1.92, p = 0.18, \eta_G^2 = 0.05$, and sad music, $F_{(1, 34)} = 0.85, p = 0.36, \eta_G^2 = 0.03$, as well as similar levels of sadness and fear when listening to scary music, $F_{(1, 34)} = 0.44, p = 0.51, \eta_G^2 = 0.01$. Younger adults reported similar levels of sadness and peacefulness only when listening to peaceful music, $F_{(1, 34)} = 0.73, p = 0.40, \eta_G^2 = 0.02$. Happy music was the only music that primarily elicited happiness (when compared with the level of peacefulness felt) in both young and older adults, $F_{(1, 34)} = 117.63, p < 0.001, \eta_G^2 = 0.78$, and $F_{(1, 34)} = 105.92, p < 0.001, \eta_G^2 = 0.76$, respectively]. The Intended Emotion by Emotion Felt by Age Group interaction remained significant when years of education, fluid intelligence scores, and working memory performances were entered as covariates, $F_{(9, 279)} = 2.90, p < 0.05, \eta_G^2 = 0.06$.

THE RELATIONSHIP BETWEEN AGE AND DEPENDENT MEASURES

In order to check for any relationships between age and the different dependent measures (i.e., emotion intensity felt, physiological responses to music, recognition accuracy, and the type of emotion felt) for each of the four intended emotions, we computed a series of correlations. Because age was significantly correlated with years of education, $r_{(34)} = -0.41, p < 0.05$, fluid intelligence, $r_{(34)} = -0.48, p < 0.05$, working memory, $r_{(34)} = -0.35, p < 0.05$, and each of the five measures of auditory thresholds, $r_{(34)} = 0.56, p < 0.001$ for 500 Hz; $r_{(34)} = 0.54, p < 0.001$ for 1000 Hz; $r_{(34)} = 0.61, p < 0.001$ for 2000 Hz; $r_{(34)} = 0.42, p < 0.05$ for 4000 Hz; $r_{(34)} = 0.75, p < 0.001$ for 8000 Hz, these variables were controlled for in partial correlations. Results indicated that the mean scores of hits, corrected recognition, and d-prime for peaceful music were negatively and significantly correlated with age, $r_{(34)} = -0.34, p < 0.05$ for hits, $r_{(34)} = -0.51, p < 0.05$ for corrected recognition, and $r_{(34)} = -0.53, p < 0.05$ for d-prime. Similarly, age was negatively and significantly correlated with the mean score of corrected recognition and d-prime for happy music, $r_{(34)} = -0.69, p < 0.001$ for corrected recognition, and $r_{(34)} = -0.65, p < 0.001$ for d-prime, while it was positively and significantly correlated with the mean score of false alarms, $r_{(34)} = 0.63, p < 0.001$. Altogether, the data indicated that the older the people are, the lower their ability to

799 discriminate studied positive musical excerpts conveying peace- 856
800 fulness and happiness from unstudied ones. The mean score of 857
801 Beta index for sad music was also positively and significantly cor- 858
802 related with age, $r_{(34)} = 0.50$, $p < 0.05$, suggesting that the older 859
803 the people are, the more conservative they are to discriminate 860
804 studied stimuli from unstudied musical excerpts conveying sad- 861
805 ness. Moreover, data indicated that the older the people are, the 862
806 stronger their experience of sadness while listening to peaceful 863
807 music, $r_{(34)} = 0.46$, $p < 0.05$, and the weaker their experience of 864
808 fear while listening to scary music, $r_{(34)} = -0.49$, $p < 0.05$. No 865
809 other significant correlations were found. 866

810 **THE RELATIONSHIP BETWEEN EMOTION INTENSITY FELT, FACIAL** 811 **MUSCLE ACTIVITY, AND RECOGNITION PERFORMANCES**

812 For each age group, we investigated to what extent the emotion 868
813 intensity felt during the first presentation of musical stimuli was 869
814 linked to the physiological responses as well as to the subsequent 870
815 cognitive performances on the incidental recognition task. The 871
816 relationship between physiological reactions and recognition per- 872
817 formances was also examined. In younger adults, results indicated 873
818 that the stronger the emotion intensity felt in response to sad 874
819 music, the higher the hits, $r_{(14)} = 0.54$, $p < 0.05$. In older adults, 875
820 results showed that the stronger the emotion intensity felt for 876
821 happy music, the higher the false alarms, $r_{(14)} = 0.54$, $p < 0.05$. 877
822 No other significant correlations were found. 878
823 879
824 880

825 **DISCUSSION**

826 In this study, we investigated how the emotional experience as 881
827 well as memory recognition for musical excerpts eliciting four 882
828 different emotions (happiness, sadness, peacefulness, and fear) 883
829 may change with age. To this end, younger and older listeners 884
830 were asked to evaluate the intensity of the emotion felt while their 885
831 facial expressions (i.e., zygomatic and corrugator muscle activity) 886
832 were recorded. They were then instructed to perform an incidental 887
833 recognition task followed by another task in which they had 888
834 to assess for each musical excerpt to what extent they experienced 889
835 each of the four emotions. 890

836 As predicted, the results showed that, when presented with 891
837 happy music, older adults assessed the emotion felt as more 892
838 intense than their younger counterparts. The fact that older adults 893
839 rated their emotional experience as significantly more intense 894
840 for happy music stimuli in comparison to sad and scary music 895
841 stimuli is consistent with the literature showing that aging is asso- 896
842 ciated with a relative preference for positivity over negativity. This 897
843 also supports the view that emotions and motivations cannot 898
844 be disentangled from each other. However, the assumption that 899
845 the stronger emotional experience reported by older adults while 900
846 listening to happy music would be reflected in a greater zygoma- 901
847 tic activity was not supported. Compared to younger adults, 902
848 older adults showed stronger facial expressions for both corruga- 903
849 tor and zygomatic muscles as well as for all intended emotions. 904
850 This suggests that facial expressions are not exclusively aligned 905
851 with the emotional state, thus raising the question of whether 906
852 the general increase of facial expressiveness in older adults would 907
853 simply reflect a deeper engagement in the task. However and 908
854 interestingly, the current results also showed that older adults' 909
855 zygomatic activity, but not for young adults', varied as a func- 910
tion of the intended emotions in such a way that older adults 911
912

showed an increased zygomatic activity for scary excerpts but not 856
for happy or peaceful ones. This is in line with the idea that the 857
reaction of smiling may serve as a defensive goal in inhibiting 858
negative feelings for older adults. It can also be argued that zygoma- 859
tic activity may reflect partial facial expression of fear, but then 860
there we should have observed a greater concomitant activity for 861
the corrugator muscle. However, this was not observed. Taken 862
together, these findings are consistent with the idea that older 863
adults' facial expressions possibly reflect an attempt to regulate 864
emotion. Further research is needed to substantiate the role 865
of voluntary facial expressions in the older adults' response to 866
emotions. 867

Consistent with our expectations, our findings indicated that 868
older adults correctly recognized less musical excerpts than their 869
younger counterparts. Moreover, older adults' range of perfor- 870
mances was quite similar to that found by Kensinger (2008) with 871
emotional words. This suggests that modality has little impact on 872
the strength of the memory decline with aging. The results of the 873
present study also indicated that younger adults as well as older 874
adults better recognized negative and arousing musical excerpts 875
(i.e., scary music) than all other excerpts while producing low 876
false alarms rates for these scary music stimuli. This corroborates 877
the hypothesis of an increased distinctiveness of negative stimuli 878
(e.g., Pesta et al., 2001) and extends previous studies that showed 879
that older adults can visually detect arousing and negative stimuli 880
as well as their younger counterparts (e.g., Magai et al., 2006; 881
Knight et al., 2007). Our findings also gave evidence for increased 882
false recognition for happy music stimuli in older adults but 883
not for young adults. We found a negative relationship between 884
age and the ability to discriminate between true and false happy 885
musical excerpts as well as a positive relationship between the 886
emotion intensity felt in older adults and their rate of false 887
alarms for happy stimuli. Taken together, these findings suggest 888
that positive emotion elicited by happy excerpts may produce 889
an attentional bias in older adults that can lead to confusion 890
between studied and non-studied excerpts and thus enhance 891
the probability of false alarms. Such increase in discrimination 892
threshold is consistent with previous studies showing that aging 893
was associated to a higher false response rate to positive words 894
(Fernandes et al., 2008; Piquet et al., 2008) and corroborates the 895
idea that the reduction of distinctiveness for positive information 896
in older adults would be the result of their liberal bias toward 897
positivity. However, in the present study, the positivity bias is 898
detrimental to memory accuracy. 899

Another main finding of the current research is that, when 900
presented with sad and scary musical excerpts, older adults 901
reported experiencing lower levels of sadness and fear than 902
their younger counterparts. Correlation analyses indicated that 903
the older the people are, the weaker their experience of fear 904
felt while listening to scary music. This fits nicely with pre- 905
vious research demonstrating age-related changes in emotion 906
recognition (Laukka and Juslin, 2007; Lima and Castro, 2011) 907
and emotion perception (Vieillard et al., 2012) in music, and 908
extends these studies by showing these changes also occur when 909
participants are focused on their own emotional experience. 910
Interestingly, compared to Lima and Lima and Castro's (2011) 911
recognition paradigm, the personal engagement involved in the 912
current task seems to facilitate the older adults' ability to process

913 negative emotions. This is consistent with previous findings
 914 demonstrating that older adults benefit more from instructions
 915 encouraging to focus on emotion than on information acquisition
 916 (Mikels et al., 2010) and corroborates the view of an age-related
 917 emphasis on emotion processing. Of course, further research is
 918 needed to compare the older adults' responsiveness to musi-
 919 cal emotions in both contexts of recognition and of emotional
 920 experience.

921 Given the relatively short duration of the musical stimuli used
 922 in the present study, one may argue that this could challenge their
 923 ability to induce emotions, leading participants to rate their per-
 924 ceived emotions rather than their felt emotions. Although studies
 925 aiming to induce felt emotions in listeners tend to use longer
 926 excerpts than those investigating perceived emotions (Eerola and
 927 Vuoskoski, 2013), we believe that the short excerpts used in our
 928 experiment also successfully induced emotions. First of all, our
 929 results indicated that participants reported moderate intensity of
 930 the emotion felt along with significant differences in facial expres-
 931 sivity. Furthermore, previous findings demonstrated that musical
 932 excerpts as short as 13s may recruit neural mechanisms involved
 933 in pleasant/unpleasant emotional responses (Blood et al., 1999).
 934 In the study of Viellard et al. (2008) which used similar 10 s
 935 musical excerpts recorded in a piano timbre, listeners better
 936 recognized some intended emotions when focusing on their emo-
 937 tional experience rather than when focusing on the recognition of
 938 the emotion. This suggests that asking participants to focus on felt
 939 emotions increases the degree of personal engagement in musi-
 940 cal emotion even for short musical excerpts. Taken together, these
 941 data corroborate the hypothesis that the musical emotions were
 942 not only recognized, but indeed felt.

943
 944 **REFERENCES**

945 Allen, R., and Brosgole, L. (1993).
 946 Facial and auditory affect recog-
 947 nition in senile geriatrics, the
 948 normal elderly and young adults.
 949 *Int. J. Neurosci.* 68, 33–42. doi:
 950 10.3109/00207459308994257
 951 Bailey, P. E., Henry, J. D., and Nangle,
 952 M. R. (2009). Electromyographic
 953 evidence for age-related differ-
 954 ences in the mimicry of anger.
 955 *Psychol. Aging* 24, 224–229. doi:
 956 10.1037/a0014112
 957 Bakeman, R. (2005). Recommended
 958 effect size statistics for repeated
 959 measures designs. *Behav. Res.*
 960 *Methods* 37, 379–384. doi:
 961 10.3758/BF03192707
 962 Bayard, S., Erkes, J., and Moroni, C.
 963 (2011). Victoria Stroop Test: nor-
 964 mative data in a sample group of
 965 older people and the study of their
 966 clinical applications in the assess-
 967 ment of inhibition in Alzheimer's
 968 disease. *Arch. Clin. Neuropsychol.*
 969 26, 653–661. doi: 10.1093/arclin/
 970 acr053
 971 Beck, A. T., Steer, R. A., and Brown,
 972 G. K. (1998). *Beck Depression*
 973 *Inventory II*. Paris: Center of
 974 Applied Psychology.

Blood, A. J., Zatorre, R. J., Bermudez, P.,
 and Evans, A. C. (1999). Emotional
 responses to pleasant and unpleas-
 ant music correlate with activity
 in paralimbic brain regions. *Nat.*
Neurosci. 2, 382. doi: 10.1038/7299
 Burriss, L., Powell, D. A., and White,
 J. (2007). Psychophysiological and
 subjective indices of emotion as a
 function of age and gender. *Cogn.*
Emot. 21, 182–210. doi: 10.1080/
 02699930600562235
 Cardebat, D., Doyon, B., Puel, M.,
 Goulet, P., and Joaquette, Y. (1990).
 Evocation lexicale formelle et
 sémantique chez des sujets nor-
 maux. Performances et dynamiques
 de production en fonction du sexe,
 de l'âge et du niveau d'étude. *Acta*
Neurol. Belg. 90, 207–217.
 Carstensen, L. L. (2006). The influ-
 ence of a sense of time on
 human development. *Science*
 312, 1913–1915. doi: 10.1126/
 science.1127488
 Carstensen, L. L., Isaacowitz, D. M.,
 and Charles, S. T. (1999). Taking
 time seriously. A theory of socioe-
 motional selectivity. *Am. Psychol.*
 54, 165–181. doi: 10.1037/0003-
 066X.54.3.165

970 One limitation of this study is that we used a cross-sectional
 971 design. Historical differences in the cultural system and in musical
 972 exposure may have affected young and older adults' performances
 973 differently. The observed age differences in emotional responses
 974 to music might thus reflect a cohort effect rather than an age
 975 effect. Future research would benefit from investigating this issue
 976 more thoroughly. Nevertheless, our study suggests that emo-
 977 tional response to music and memory recognition for musical
 978 excerpts conveying emotions show differences with advancing
 979 age. These age-related differences are characterized by a stronger
 980 emotional reactivity for happiness, an increased zygomatic activ-
 981 ity in response to scary stimuli, an increase in false recognition
 982 for happy musical excerpts, and a decrease in responsiveness to
 983 sad and scary music. This study extends previous findings and
 984 expands them to music, a powerful channel of emotion commu-
 985 nication. Importantly, the findings suggest that aging may cause a
 986 decrease in negative affects and an increase in positive affects even
 987 when these affects are elicited by a more abstract source of emo-
 988 tion that does not refer to specific events. Finally, the current data
 989 are in line with the hypothesis that older adults could use emo-
 990 tional coping skills acquired over the life span in order to avoid
 991 potentially negative events and maintain positive ones (Charles
 992 et al., 2001; Labouvie-Vief et al., 2010).

993
 994 **ACKNOWLEDGMENTS**

995 This research was funded by the ANR "EMCO" program (Project
 996 Strem N ANR 11 EMCO 003 01). We are grateful to Alexandra
 997 Richen for her help in running the experiments as well as to
 998 Alejandra R. Velasquez and Joanna Blatter-Minn for their proof-
 999 reading the manuscript.

Charles, S. T., Mather, M., and
 Carstensen, L. L. (2003). Aging
 and emotional memory: the
 forgettable nature of negative
 images for older adults. *J. Exp.*
Psychol. Gen. 132, 310–324. doi:
 10.1037/0096-3445.132.2.310
 Charles, S. T., and Piazza, J. R. (2007).
 Memories of social interactions: age
 differences in emotional intensity.
Psychol. Aging 22, 300–309. doi:
 10.1037/0882-7974.22.2.300
 Charles, S. T., Reynolds, C. A., and
 Gatz, M. (2001). Age-related dif-
 ferences and change in positive
 and negative affect over 23 years.
J. Pers. Soc. Psychol. 80, 136–151.
 doi: 10.1037/0022-3514.80.1.136
 Cohen, J. (1988). *Statistical Power*
Analysis for the Behavioral Sciences,
2nd Edn. Hillsdale, NJ: Lawrence
 Erlbaum Associates Inc.
 Drapeau, J., Gosselin, N., Gagnon, L.,
 Peretz, I., and Lorrain, D. (2009).
 Emotional recognition from face,
 voice, and music in dementia of the
 Alzheimer type. *Ann. N.Y. Acad. Sci.*
 1169, 342–345. doi: 10.1111/j.1749-
 6632.2009.04768.x
 Eerola, T., and Vuoskoski, J. K.
 (2013). A review of music and
 emotion studies: approaches,
 emotion models, and stimuli.
Music Percept. 30, 307–340. doi:
 10.1525/mp.2012.30.3.307
 Fernandes, M., Ross, M., Wiegand, M.,
 and Schryer, E. (2008). Are the
 memories of older adults positively
 biased. *Psychol. Aging* 23, 297–306.
 doi: 10.1037/0882-7974.23.2.297
 Gosselin, N., Peretz, I., Johnsen,
 E., and Adolphs, R. (2007).
 Amygdala damage impairs emo-
 tion recognition from music.
Neuropsychologia 45, 236–244. doi:
 10.1016/j.neuropsychologia.2006.
 07.012
 Gosselin, N., Peretz, I., Noulhiane,
 M., Hasboun, D., Beckett, C.,
 Baulac, M., et al. (2005). Impaired
 recognition of scary music fol-
 lowing unilateral temporal lobe
 excision. *Brain* 128, 628–640. doi:
 10.1093/brain/awh420
 Green, D. M., and Swets, J. A. (1966).
 Signal detection theory and psy-
 chophysics. New York: Wiley.
 Grünh, D., Smith, J., and Baltes, P.
 B. (2005). No aging bias favoring
 memory for positive material:
 evidence from a heterogeneity-
 homogeneity list paradigm

- 1027 using emotionally toned words. *Psychol. Aging* 20, 579–588. doi: 10.1037/0882-7974.20.4.579
- 1028 Hochhaus, L. (1972). A table for the
1029 calculation of d' and β . *Psychol. Bull.*
1030 77, 375–376. doi: 10.1037/h0032459
- 1031 Isaacowitz, D. M., Wadlinger, H. A.,
1032 Goren, D., and Wilson, H. R. (2006a). Is there an age-related
1033 positivity effect in visual atten-
1034 tion? A comparison of two method-
1035 ologies. *Emotion* 6, 511–516. doi:
1036 10.1037/1528-3542.6.3.511
- 1037 Isaacowitz, D. M., Wadlinger, H. A.,
1038 Goren, D., and Wilson, H. R. (2006b). Selective preference in
1039 visual fixation away from negative
1040 images in old age? An eye-tracking
1041 study. *Psychol. Aging* 21, 40–48. doi:
1042 10.1037/0882-7974.21.1.40
- 1043 Kensinger, E. A. (2008). Age differ-
1044 ences in memory for arousing
1045 and nonarousing emotional
1046 words. *J. Gerontol. B Psychol.*
1047 *Sci. Soc. Sci.* 63, P13–P18. doi:
10.1093/geronb/63.1.P13
- 1048 Kensinger, E. A., Brierley, B., Medford,
1049 N., Growdon, J. H., and Corkin, S. (2002). Effects of normal aging and
1050 Alzheimer's disease on emotional
1051 memory. *Emotion* 2, 118–134. doi:
1052 10.1037/1528-3542.2.2.118
- 1053 Khalfa, S., Peretz, I., Blondin, J.-
1054 P., and Robert, M. (2002). Event-
1055 related skin conductance responses
1056 to musical emotions in humans. *Neurosci. Lett.* 328, 145–149. doi:
1057 10.1016/S0304-3940(02)00462-7
- 1058 Khalfa, S., Roy, M., Rainville, P., Dalla
1059 Bella, S., and Peretz, I. (2008).
1060 Role of tempo entrainment in
1061 psychophysiological differentiation
1062 of happy and sad music? *Int. J. Psychophysiol.* 68, 17–26. doi:
1063 10.1016/j.ijpsycho.2007.12.001
- 1064 Knight, M., Seymour, T. L., Gaunt,
1065 J. T., Baker, C., Nesmith, K., and
1066 Mather, M. (2007). Aging and
1067 goal-directed emotional attention:
1068 distraction reverses emotional
1069 biases. *Emotion* 7, 705–714. doi:
10.1037/1528-3542.7.4.705
- 1070 Labouvie-Vief, G. (2009). Cognition
1071 and equilibrium regulation in devel-
1072 opment and aging. *Restor. Neurol. Neurosci.* 27, 551–565.
- 1073 Labouvie-Vief, G., Grün, D., and
1074 Studer, J. (2010). “Dynamic inte-
1075 gration of emotion and cognition:
1076 equilibrium regulation in develop-
1077 ment and aging,” in *The Handbook of Life-Span Development*, Vol. 2,
1078 eds R. M. Lerner, M. E. Lamb,
1079 and A. M. Freund (Hoboken,
1080 NJ: Wiley), 79–115. doi: 10.1002/
9780470880166.hlsd002004
- 1081 Laukka, P., and Juslin, P. N. (2007).
1082 Similar patterns of age-related dif-
1083 ferences in emotion recognition
1084 from speech and music. *Motiv. Emot.* 31, 182–191. doi: 10.1007/
s11031-007-9063-z
- 1085 Levenson, R. W., Carstensen, L. L.,
1086 Friesen, W. V., and Ekman, P. (1991). Emotion, physiology, and
1087 expression in old age. *Psychol. Aging*
1088 6, 28–35. doi: 10.1037/0882-7974.
6.1.28
- 1089 Lima, C. F., and Castro, S. L. (2011).
1090 Emotion recognition in music
1091 changes across the adult life span. *Cogn. Emot.* 25, 585–598. doi:
1092 10.1080/02699931.2010.502449
- 1093 Magai, C., Consedine, N. S.,
1094 Krivoshekova, Y. S., Kudadjie-
1095 Gyamfi, E., and McPherson, R. (2006). Emotion experience and
1096 expression across the adult life
1097 span: insights from a multimodal
1098 assessment study. *Psychol. Aging* 21,
1099 303–317. doi: 10.1037/0882-7974.
21.2.303
- 1100 Mather, M., Canli, T., English, T.,
1101 Whitfield, S., Wais, P., Ochsner, K.,
1102 et al. (2004). Amygdala responses
1103 to emotionally valenced stimuli in
1104 older and younger adults. *Psychol. Sci.* 15, 259–263. doi: 10.1111/j.
0956-7976.2004.00662.x
- 1105 Mather, M., and Carstensen, L. L. (2005). Aging and motivated cogni-
1106 tion: the positivity effect in atten-
1107 tion and memory. *Trends Cogn. Sci.* 9, 496–502. doi: 10.1016/j.tics.2005.
08.005
- 1108 Mather, M., and Knight, M. (2005).
1109 Goal-directed memory: the role of
1110 cognitive control in older adults' emotional memory. *Psychol. Aging* 20,
1111 554–570. doi: 10.1037/0882-
7974.20.4.554
- 1112 Mikels, J. A., Larkin, G. R., Reuter-
1113 Lorenz, P. A., and Cartensen, L. L. (2005). Divergent trajectories in
1114 the aging mind: changes in work-
1115 ing memory for affective versus
1116 visual information with age. *Psychol. Aging* 20, 542–553. doi: 10.1037/
0882-7974.20.4.542
- 1117 Mikels, J. A., Lockenhoff, C. E.,
1118 Maglio, S. J., Goldstein, M. K.,
1119 Garber, A., and Carstensen, L. L. (2010). Following your heart or
1120 your head: focusing on emotions
1121 versus information differentially
1122 influences the decisions of younger
1123 and older adults. *J. Exp. Psychol. Appl.* 16, 87–95. doi: 10.1037/
a0018500
- 1124 Niedenthal, P. M., Barsalou, L. W.,
1125 Winkielman, P., Krauth-Gruber, S.,
1126 and Ric, F. (2005). Embodiment
1127 in attitudes, social perception,
1128 and emotion. *Pers. Soc. Psychol. Rev.* 9, 184–211. doi: 10.1207/
s15327957pspr0903_1
- 1129 Pesta, B. J., Murphy, M. D., and
1130 Sanders, R. E. (2001). Are emo-
1131 tionally charged lures immune
1132 to false memory? *J. Exp. Psychol. Learn. Mem. Cogn.* 27, 328–338. doi: 10.1037/0278-7393.27.2.328
- 1133 Petit, H., Bakchine, S., Dubois, B.,
1134 Laurent, B., Montagne, B., Touchon,
1135 J., et al. (1998). Consensus state-
1136 ment of an interdisciplinary group
1137 of French experts on modalities of
1138 diagnosis and medical treatment of
1139 Alzheimer's disease at a treatable
1140 stage. *Rev. Neurol.* 154, 432–438.
- 1141 Piguet, O., Connally, E., Krendl, A. C.,
1142 Huot, J. R., and Corkin, S. (2008).
1143 False memory in aging: effects of
1144 emotional valence on word recog-
1145 nition accuracy. *Psychol. Aging* 23,
1146 307–314. doi: 10.1037/0882-7974.
23.2.307
- 1147 Raven, J., Raven, J. C., and Court, J. H. (1998, updated 2003). *Manual for Raven's Progressive Matrices and Vocabulary Scales. Section 1: General Overview*. San Antonio, TX: Harcourt Assessment.
- 1148 Reed, A. E., and Carstensen, L. L. (2012). The theory behind the
1149 age-related positivity effect. *Front. Psychol.* 3:339. doi: 10.3389/fpsyg.2012.00339
- 1150 Reitan, R. M. (1958). Validity of
1151 the Trail Making Test as an indi-
1152 cation of organic brain damage. *Percept. Motor Skills* 8, 271–276. doi: 10.2466/PMS.8.7.271-276
- 1153 Reminger, S. L., Kaszniak, A. W.,
1154 and Dalby, P. R. (2000). Age-
1155 invariance in the asymmetry of
1156 stimulus-evoked emotional facial
1157 muscle activity. *Aging Neuropsychol. Cogn.* 7, 156–168. doi: 10.1076/
1382-5585(200009)7:3;1-Q:FT156
- 1158 Scherer, K. R. (2005). What are emo-
1159 tions? And how can they be mea-
1160 sured? *Soc. Sci. Inf.* 44, 695–729. doi:
10.1177/0539018405058216
- 1161 Schneider, W., Eschman, A., and
1162 Zuccolotto, A. (2002). *E-Prime User's Guide*. Pittsburgh, PA: Psychology Software Tools Inc.
- 1163 Smith, D. P., Hillman, C. H., and Duley,
1164 A. R. (2005). Influence of age on
1165 emotional reactivity during picture
1166 processing. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 60, 49–56. doi: 10.1093/
geronb/60.1.P49
- 1167 Spielberger, C. D. (1993). *State and Trait Anxiety Inventory, Form Y*. Paris: Center of Applied Psychology.
- 1168 Tsai, J. L., Levenson, R. W., and
1169 Carstensen, L. L. (2000). Auto-
1170 nomic, subjective, and
1171 expressive responses to emo-
1172 tional films in older and younger
1173 Chinese Americans and European
1174 Americans. *Psychol. Aging* 15,
1175 684–693. doi: 10.1037/0882-7974.
15.4.684
- 1176 van Tricht, M. J., Smeding, H. M.,
1177 Speelman, J. D., and Schmand,
1178 B. A. (2010). Impaired emotion
1179 recognition in music in Parkinson's
1180 disease. *Brain Cogn.* 74, 58–65. doi: 10.1016/j.bandc.2010.06.005
- 1181 Viellard, S., Didierjean, A., and
1182 Maquestiau, F. (2012). Changes in
1183 the perception and the psychologi-
1184 cal structure of musical emotions
1185 with advancing age. *Exp. Aging Res.* 38, 422–441. doi: 10.1080/
0361073X.2012.699371
- 1186 Viellard, S., Peretz, I., Gosselin,
1187 N., Khalfa, S., Gagnon, L., and
1188 Bouchard, B. (2008). Happy, sad,
1189 scary and peaceful musical excerpts
1190 for research on emotions. *Cogn. Emot.* 22, 720–752. doi: 10.1080/
02699930701503567
- 1191 Wechsler, D. (2000). *WAIS-III Administration and Scoring Manual*. Paris: Center of Applied Psychology.
- 1192 Witvliet, C. V. O., and Vrana, S. R. (2007). Play it again Sam: repeated
1193 exposure to emotionally evocative
1194 music polarises liking and smil-
1195 ing responses, and influences other
1196 affective reports, facial EMG, and
1197 heart rate. *Cogn. Emot.* 21, 3–25. doi: 10.1080/02699930601000672
- 1198 Zimmerman, M. E., Brickman, A. M.,
1199 Paul, R. H., Grieve, S. M., Tate,
1200 D. F., Gunstad, J., et al. (2006).
1201 The relationship between frontal
1202 gray matter volume and cognition
1203 varies across the healthy adult
1204 lifespan. *Am. J. Geriatr. Psychiatry* 14,
1205 823–833. doi: 10.1097/01.JGP.0000238502.40963.ac

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 31 March 2013; accepted: 17 September 2013; published online: xx October 2013.

Citation: Viellard S and Gilet A-L (2013) Age-related differences in affective responses to and memory for emotions conveyed by music: a cross-sectional study. *Front. Psychol.* 4:711. doi: 10.3389/fpsyg.2013.00711

This article was submitted to *Emotion Science*, a section of the journal *Frontiers in Psychology*.

Copyright © 2013 Viellard and Gilet. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.