



**HAL**  
open science

# Asymptotic modelling of multi-dimensional jump boundary conditions at a fluid-porous interface

Philippe Angot, Benoît Goyeau, J. Alberto Ochoa-Tapia

► **To cite this version:**

Philippe Angot, Benoît Goyeau, J. Alberto Ochoa-Tapia. Asymptotic modelling of multi-dimensional jump boundary conditions at a fluid-porous interface. 13èmes Journées d'études des Milieux Poreux 2016, Oct 2016, Anglet, France. hal-01394516

**HAL Id: hal-01394516**

**<https://hal.science/hal-01394516v1>**

Submitted on 9 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Asymptotic modelling of multi-dimensional jump boundary conditions at a fluid-porous interface

Philippe Angot<sup>a</sup>, Benoît Goyeau<sup>b</sup>, J. Alberto Ochoa-Tapia<sup>c</sup>

<sup>a</sup>Aix-Marseille Université, Institut de Mathématiques de Marseille, UMR-CNRS 7373, Centrale Marseille, 39 rue F. Joliot-Curie, 13453 Marseille cedex 13, France.

<sup>b</sup>Ecole Centrale-Supélec, Université Paris-Saclay, EM2C, UPR-CNRS 288, Grande Voie des Vignes, 92295 Châtenay-Malabry, France.

<sup>c</sup>Departamento de Ingeniería de Procesos e Hidráulica, Universidad Autónoma Metropolitana-Iztapalapa, Mexico 09340 D.F., Mexico.

*Keywords:* Asymptotic modelling, coupled fluid-porous transport phenomena, multi-dimensional jump interface conditions, cross-flow pressure jump

## 1. Introduction

We present an original asymptotic modelling [1] for a multi-dimensional unsteady viscous fluid flow and convective transfer at a fluid-porous interface. This asymptotic model is based on the fact that the thickness  $d$  of the interfacial transition region  $\Omega_{fp}$  in the one-domain representation [2, 3] is very small compared to the macroscopic length scale  $L$ . The analysis leads to an equivalent two-domain representation where transport phenomena in the transition layer of the one-domain approach are represented by algebraic jump boundary conditions at a fictive dividing interface  $\Sigma$  between the homogeneous fluid and porous regions. These jump conditions are thus stated in [1] up to first-order in  $O(d/L)$ .

The originality and great interest of this new asymptotic model lies in its general and multi-dimensional character. Indeed, it is shown that all the jump interface conditions [4, 5, 6] derived for the 1D-shear flows are recovered by taking the tangential component of the asymptotic model. In that case, the comparison between the present model and the upscaled models available in the literature gives explicit expressions of the effective jump coefficients and their associated scaling. In addition, for multi-dimensional flows as in [7, 8], the general asymptotic model yields the different components of the jump conditions including a new specific equation for the cross-flow pressure jump on  $\Sigma$ .

## 2. Asymptotic model with multi-dimensional jump interface conditions

We summarize the first-order asymptotic model derived in [1] for the multi-dimensional and unsteady viscous fluid flow with coupled heat or mass transfer inside the fluid-porous system, see Figure 1. *The main assumption is that  $\eta := d/L \ll 1$  with  $d = O(\sqrt{|\mathbf{K}|})$ ,  $\mathbf{K}$  being the permeability tensor in  $\Omega_{fp}$ .*




Figure 1: Fluid-porous configurations – Left: one-domain model with continuous inter-region – Right: two-domain model with a fictive interface  $\Sigma$  at  $\xi := z_\Sigma/d$ .

The jump and mean quantity of any unknown  $\psi$  on  $\Sigma$  are defined as in [9, 7] respectively by :

$$\llbracket \psi \rrbracket_{\Sigma} := \psi^+ - \psi^- = (\psi^f - \psi^p)_{|\Sigma}, \quad \text{and} \quad \bar{\psi}_{\Sigma} := \frac{1}{2} (\psi^+ + \psi^-) = \frac{1}{2} (\psi^f + \psi^p)_{|\Sigma}.$$

For any physical property  $k$ , the arithmetic and harmonic averages over the thickness of  $\Omega_{fp}$  are :

$$\langle k \rangle (x) := \frac{1}{d(x)} \int_{-d/2}^{d/2} k(x, z) dz, \quad \langle k \rangle^h (x) := \left( \frac{1}{d(x)} \int_{-d/2}^{d/2} \frac{dz}{k(x, z)} \right)^{-1} = \langle \frac{1}{k} \rangle.$$

The set of equations corresponds to a Stokes/Darcy-Brinkman transmission problem with source terms inside  $\Omega_f \cup \Omega_p$ , coupled with the advection-diffusion-reaction equation for the temperature  $\theta$  :

$$\left\{ \begin{array}{ll} \nabla \cdot \mathbf{v} = q & \text{in } \Omega_f \cup \Omega_p \times (0, T), \\ \rho \partial_t \mathbf{v} - \mu \Delta \mathbf{v} + \nabla p = \rho \mathbf{f} & \text{in } \Omega_f \times (0, T), \\ \rho \partial_t \mathbf{v} - \tilde{\mu} \Delta \mathbf{v} + \mu \mathbf{K}^{-1} \cdot \mathbf{v} + \nabla p = \rho \phi \mathbf{f} & \text{in } \Omega_p \times (0, T), \\ (\rho c)^* \partial_t \theta + (\rho c)^f \mathbf{v} \cdot \nabla \theta - \nabla \cdot (\mathbf{A} \cdot \nabla \theta) + b \theta = f & \text{in } \Omega_f \cup \Omega_p \times (0, T), \\ \mathbf{v}(t=0) = \mathbf{v}_0 & \text{in } \Omega_f \cup \Omega_p, \\ \theta(t=0) = \theta_0 & \text{in } \Omega_f \cup \Omega_p. \end{array} \right. \quad (1)$$

It must be completed by the *first-order algebraic jump boundary conditions (A.J.B.C.) on the interface*  $\Sigma$  obtained by averaging the transport phenomena inside  $\Omega_{fp}$  from the single-domain model [2, 3], where  $\boldsymbol{\sigma}(\mathbf{v}, p) \cdot \mathbf{n} := -p \mathbf{n} + \tilde{\mu} (\nabla \mathbf{v} + \nabla \mathbf{v}^t) \cdot \mathbf{n}$  denotes the stress vector with  $\tilde{\mu} := \mu/\phi$ ,  $\phi$  being the space variable porosity, and  $\boldsymbol{\varphi}(\theta) := -\mathbf{A} \cdot \nabla \theta + (\rho c)^f \mathbf{v} \theta$  is the convective-diffusive heat flux vector :

$$\left\{ \begin{array}{ll} \llbracket \mathbf{v} \cdot \mathbf{n} \rrbracket_{\Sigma} = d \langle q \rangle & \text{on } \Sigma \times (0, T), \\ -\llbracket \boldsymbol{\sigma}(\mathbf{v}, p) \cdot \mathbf{n} \rrbracket_{\Sigma} + \mu d \mathbf{K}_{\Sigma}^{-1} \cdot \bar{\mathbf{v}}_{\Sigma} = d \langle \rho \phi \mathbf{f} \rangle & \text{on } \Sigma \times (0, T), \\ \overline{\boldsymbol{\sigma}(\mathbf{v}, p) \cdot \mathbf{n}}_{\Sigma} + \bar{p}_{\Sigma} \mathbf{n} - \frac{\mu}{\phi_{\Sigma}} \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} \cdot \frac{\llbracket \mathbf{v} \rrbracket_{\Sigma}}{d} = 0 & \text{on } \Sigma \times (0, T), \\ \llbracket \boldsymbol{\varphi}(\theta) \cdot \mathbf{n} \rrbracket_{\Sigma} + d (\langle b \rangle - (\rho c)^f \langle q \rangle) \bar{\theta}_{\Sigma} = d \langle f \rangle & \text{on } \Sigma \times (0, T), \\ \overline{\boldsymbol{\varphi}(\theta) \cdot \mathbf{n}}_{\Sigma} + \langle A_n \rangle^h \frac{\llbracket \theta \rrbracket_{\Sigma}}{d} - (\rho c)^f \bar{\mathbf{v}} \cdot \bar{\mathbf{n}}_{\Sigma} \bar{\theta}_{\Sigma} = 0 & \text{on } \Sigma \times (0, T). \end{array} \right. \quad (2)$$

Here we have:  $\phi_{\Sigma} := \langle \phi \rangle^h$ ,  $\mathbf{K}_{\Sigma} := \langle \mathbf{K}(\phi) \rangle^h$ ,  $A_n := \mathbf{A}_{nn}$ , and  $(\rho c)^* := \phi (\rho c)^f + (1 - \phi) (\rho c)^p$ .

For a non-centered fictive interface, all the quantities  $\bar{\psi}_{\Sigma}$  are to be replaced by the weighted means :

$$\bar{\psi}_{\Sigma}^w = \bar{\psi}_{\Sigma} + \xi \llbracket \psi \rrbracket_{\Sigma}, \quad \text{with} \quad -\frac{1}{2} \leq \xi := \frac{z_{\Sigma}}{d} \leq \frac{1}{2}. \quad (3)$$

The Stokes/Darcy asymptotic model is obtained by formally letting the effective viscosity  $\tilde{\mu}^p := \tilde{\mu}|_{\Omega_p} = \varepsilon$  tend to zero. We refer to [8] for the theoretical proof when  $\varepsilon \rightarrow 0^+$  with the vanishing viscosity method and [10] with the WKB asymptotic expansion to calculate the associated boundary layer.

The equation for the pressure jump  $\llbracket p \rrbracket_{\Sigma}$  of the cross-flow through  $\Sigma$  will be detailed during the talk.

## References

- [1] PH. ANGOT, B. GOYEAU AND J. A. OCHOA-TAPIA, Asymptotic modelling of transport phenomena at a fluid-porous interface: jump conditions, *J. Fluid Mech.*, submitted (2016).
- [2] PH. ANGOT, Analysis of singular perturbations on the Brinkman problem for fictitious domain models of viscous flows, *Math. Meth. in the Appl. Sci.*, **22**(16), 1395–1412 (1999).
- [3] B. GOYEAU, D. LHULLIER, D. GOBIN AND M. G. VELARDE, Momentum transport at a fluid-porous interface, *Int. J. Heat Mass Transfer*, **46**, 4071–4081 (2003).
- [4] G. S. BEAVERS AND D. D. JOSEPH, Boundary conditions at a naturally permeable wall, *J. Fluid Mech.*, **30**, 197–207 (1967).
- [5] J. A. OCHOA-TAPIA AND S. WHITAKER, Momentum transfer at the boundary between a porous medium and a homogeneous fluid I: theoretical development, *Int. J. Heat Mass Transfer*, **38**, 2635–2646 (1995).
- [6] F. J. VALDÉS-PARADA, C. G. AGUILAR-MADERA, J. A. OCHOA-TAPIA AND B. GOYEAU, Velocity and stress jump conditions between a porous medium and a fluid, *Adv. in Water Res.*, **62**, 327–339 (2013).
- [7] PH. ANGOT, A fictitious domain model for the Stokes/Brinkman problem with jump embedded boundary conditions, *C. R. Math. Acad. Sci. Paris, Ser. I*, **348**(11-12), 697–702 (2010).
- [8] PH. ANGOT, On the well-posed coupling between free fluid and porous viscous flows, *Appl. Math. Lett.*, **24**(6), 803–810 (2011).
- [9] PH. ANGOT, A model of fracture for elliptic problems with flux and solution jumps, *C. R. Acad. Sci. Paris, Série I Math.*, **337**(6), 425–430 (2003).
- [10] PH. ANGOT, G. CARBOU AND V. PÉRON, Asymptotic study for Stokes-Brinkman model with jump embedded transmission conditions, *Asymptotic Analysis*, **96**(3-4), 223–249 (2016).