



HAL
open science

Extension of the Günter derivatives to Lipschitz domains and application to the boundary potentials of elastic waves

Abderrahmane Bendali, Sébastien Tordeux, Yuriy Matveevich Volchkov

► **To cite this version:**

Abderrahmane Bendali, Sébastien Tordeux, Yuriy Matveevich Volchkov. Extension of the Günter derivatives to Lipschitz domains and application to the boundary potentials of elastic waves. Journal of Applied Mechanics and Technical Physics, 2020, 61 (1), pp.21. 10.1134/S0021894420010150 . hal-01395952v2

HAL Id: hal-01395952

<https://hal.science/hal-01395952v2>

Submitted on 7 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Extension of the Günter derivatives to Lipschitz domains and application to the boundary potentials of elastic waves

A. Bendali*, S. Tordeux†

October 13, 2019

Keywords— Boundary integral operators, Günter derivatives, Elastic Waves, Layer potentials, Lipschitz domains.

Abstract

The scalar Günter derivatives of a function defined on the boundary of a three-dimensional domain are expressed as components (or their opposites) of the tangential vector rotational of this function in the canonical orthonormal basis of the ambient space. This in particular implies that these derivatives define bounded operators from H^s into H^{s-1} for $0 \leq s \leq 1$ on the boundary of a Lipschitz domain, and can easily be implemented in boundary element codes. Regularization techniques for the trace and the traction of elastic waves potentials, previously built for a domain of class \mathcal{C}^2 , can in this way be extended to the Lipschitz case. In particular, this yields an elementary way to establish the mapping properties of elastic wave potentials from those of the scalar Helmholtz equation without resorting to the more advanced theory for elliptic systems in Lipschitz domains. Some attention is finally paid to the two-dimensional case.

1 Introduction

All along this paper, Ω^+ and $\Omega^- = \mathbb{R}^3 \setminus \overline{\Omega^+}$ respectively designate a bounded Lipschitz domain of \mathbb{R}^3 , and its exterior. As a result, Ω^+ and Ω^- share a common boundary denoted by $\partial\Omega$. It is well-known that $\partial\Omega$ is endowed with a Lebesgue surface measure s , and that it has an unit normal \mathbf{n} (see figure 1), defined s -almost everywhere, pointing outward from Ω^+ (cf., for example, [21, p. 96]). Vectors with three components a_j ($j = 1, 2, 3$), either real or complex, are identified to column-vectors

$$\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}.$$

The bilinear form underlying the scalar product of two such vectors \mathbf{a} and \mathbf{b} is given by

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{a}^\top \mathbf{b} = \mathbf{b}^\top \mathbf{a} = \sum_{j=1}^3 a_j b_j$$

where \mathbf{a}^\top is the transpose of \mathbf{a} .

Usual notation in the theory of Partial Differential Equations [29] will be used without further comment. We just mention that we make use of the following Fréchet spaces, defined for any integer $m \geq 0$ by

$$H_{\text{loc}}^m(\mathbb{R}^3) = \{v \in \mathcal{D}'(\mathbb{R}^3); \varphi v \in H^m(\mathbb{R}^3), \forall \varphi \in \mathcal{D}(\mathbb{R}^3)\}$$

$$H_{\text{loc}}^m(\overline{\Omega^-}) := \{v \in \mathcal{D}'(\Omega^-); \exists V \in H_{\text{loc}}^m(\mathbb{R}^3), v = V|_{\Omega^-}\},$$

and by

$$H_{\text{comp}}^m(\mathbb{R}^3) = \{v \in H^m(\mathbb{R}^3); \exists R > 0, v|_{|x| \geq R} = 0\}$$

*Université de Toulouse, Institut Mathématique de Toulouse, Département de Génie Mathématique, INSA de Toulouse, 135 avenue de Rangueil, F-31077 Toulouse cedex 4, France, abendali@insa-toulouse.fr

†Project-team Magique-3D, INRIA Bordeaux Sud-Ouest. LMA Pau, CNRS UMR 5142, I.P.R.A., Université de Pau et des Pays de l'Adour, avenue de l'Université BP 1155, 64013 Pau CEDEX, France, sebastien.tordeux@inria.fr




Figure 1: Schematic view of the geometry

$$H_{\text{comp}}^m(\overline{\Omega^-}) = \{v \in H^m(\Omega^-); \exists V \in H_{\text{comp}}^m(\mathbb{R}^3), v = V|_{\Omega^-}\}.$$

With similar definitions, it is trivially true that both $H_{\text{loc}}^m(\overline{\Omega^+})$ and $H_{\text{comp}}^m(\overline{\Omega^+})$ coincide with $H^m(\Omega^+)$. Below, we conveniently use the unified notation $H_{\text{loc}}^m(\overline{\Omega^\pm})$ and $H_{\text{comp}}^m(\overline{\Omega^\pm})$ to refer to both of these spaces. Instead of H^0 , we use the more conventional notation L^2 .

We denote by $u^+ = (u|_{\Omega^+})|_{\partial\Omega}$ (resp. $u^- = (u|_{\Omega^-})|_{\partial\Omega}$) the trace of u on $\partial\Omega$ from the values $u|_{\Omega^+}$ of u in Ω^+ (resp. $u|_{\Omega^-}$ in Ω^-). For simplicity, we omit to explicitly mention the trace when the related function has zero jump across $\partial\Omega$. We also adopt a classical way to denote functional spaces of vector fields having their components in some scalar functional space. For example, $H^s(\partial\Omega; \mathbb{C}^3)$ stands for the space of vector fields \mathbf{u} whose components u_j ($j = 1, 2, 3$) are in $H^s(\partial\Omega)$.

For $1 \leq i, j \leq 3$ and $u \in H_{\text{loc}}^2(\mathbb{R}^3)$, the G unter derivative

$$\mathcal{M}_{ij}^{(\mathbf{n})} u = n_j \partial_{x_i} u - n_i \partial_{x_j} u \quad (1)$$

is well-defined as a function in $L^2(\partial\Omega)$ since the traces of $\partial_{x_i} u$ and $\partial_{x_j} u$ are in $H^{1/2}(\partial\Omega)$ and the components n_i and n_j of the normal \mathbf{n} to $\partial\Omega$ are in $L^\infty(\partial\Omega)$. It is worth recalling that if Ω^+ is a bit more regular, say a $\mathcal{C}^{1,1}$ -domain for example (cf., [21, p. 90] for the definition of a \mathcal{C}^k -domain (resp. $\mathcal{C}^{k,\alpha}$ -domain), also referred to as a domain of class \mathcal{C}^k (resp. $\mathcal{C}^{k,\alpha}$)), $\mathcal{M}_{ij}^{(\mathbf{n})} u$ is in $H^{1/2}(\partial\Omega)$. Seemingly, there is a loss of one-half order of regularity when considering a domain which is only Lipschitz. The purpose of this paper is precisely to show that this one-half order of regularity can be restored for functions in lower order Sobolev spaces.

Let us first recall some well-established properties of the G unter derivatives when Ω^+ is at least a $\mathcal{C}^{1,1}$ -domain. Let

$$\left\{ \mathbf{e}_j = [\delta_{1j}, \delta_{2j}, \delta_{3j}]^\top \right\}_{j=1}^3 \quad (\delta_{ij} = 1 \text{ if } i = j \text{ and } 0 \text{ otherwise})$$

be the canonical basis of \mathbb{R}^3 so that $n_j = \mathbf{n} \cdot \mathbf{e}_j$ for $j = 1, 2, 3$. Define for $1 \leq i \neq j \leq 3$

$$\boldsymbol{\tau}_{ij} = n_j \mathbf{e}_i - n_i \mathbf{e}_j. \quad (2)$$

Clearly

$$\mathcal{M}_{ij}^{(\mathbf{n})} u = \nabla u \cdot \boldsymbol{\tau}_{ij} = \partial_{\boldsymbol{\tau}_{ij}} u$$

with

$$\boldsymbol{\tau}_{ij} \cdot \mathbf{n} = 0. \quad (3)$$

As a result, $\mathcal{M}_{ij}^{(\mathbf{n})}$ is a tangential derivative on $\partial\Omega$, meaning in particular, at least for $u \in \mathcal{C}^1(\mathbb{R}^3)$ and Ω^+ a \mathcal{C}^1 -domain, that $\mathcal{M}_{ij}^{(\mathbf{n})} u$ can be calculated without resorting to interior values of u in Ω^+ or in Ω^- .

These operators were introduced by G unter [12]. It was discovered later [17] that they can be used for bringing out important relations linking the boundary layer potentials of the Lam e system to those of the Laplace equation (see [17, p. 314] and [15, p. 48]). They were then employed to more conveniently express the traction of the double layer elastic potential (see [14] and [15, p. 49]). These approaches were recently

extended to the elastic wave boundary layer potentials by Le Louër [19, 18]. All these results were derived under the assumption that Ω^+ is a \mathcal{C}^2 -domain (actually, $\mathcal{C}^{1,1}$ - is enough). It is the aim of this paper, by defining the G nter derivatives for a Lipschitz domain, that is, a $\mathcal{C}^{0,1}$ -domain, to similarly handle geometries more usual in the applications. More importantly, it is possible in this way to deal with boundary element approximations of the traction of single- and double-layer potentials of Lam  static elasticity and elastic wave systems almost as easily as for the Laplace or the Helmholtz equation.

Actually, in connection with elasticity potential layers, G nter derivatives are involved as entries $\mathcal{M}_{ij}^{(n)}$ ($i, j = 1, 2, 3$) of the skew-symmetric matrix $\mathcal{M}^{(n)}$ acting on vector-valued functions \mathbf{u}

$$\left(\mathcal{M}^{(n)}\mathbf{u}\right)_i = \sum_{j=1}^3 \mathcal{M}_{ij}^{(n)} u_j \quad (i = 1, 2, 3),$$

$(\mathcal{M}^{(n)}\mathbf{u})_i$ and u_j ($j = 1, 2, 3$) being the respective components of $\mathcal{M}^{(n)}\mathbf{u}$ and \mathbf{u} . In [15], this matrix is called the G nter derivatives in matrix form. refer to $\mathcal{M}^{(n)}$ as the G nter derivative matrix.

The G nter derivative matrix actually give rise to a multi-faceted operator, with various expressions, which led to real progresses in the context of Lam  static elasticity boundary layer potentials [17, 15, 19, 14] or in the design of preconditioning techniques for the boundary integral formulations in the scattering of elastic waves [8]. Other ways to write $\mathcal{M}^{(n)}$ do not seem to have been connected with the G nter derivatives [5, 6]. However, all these expressions require either interior values, as for example for above direct definition (1) of $\mathcal{M}_{ij}^{(n)}$, or curvature terms of $\partial\Omega$ as recalled below, making problematic their effective implementation in boundary element codes or in a preconditioning technique. It is among the objectives of this paper to address this issue.

The outline of the paper is as follows. In section 2, we first show that $\mathcal{M}_{ij}^{(n)}u$ corresponds to a component (or its opposite) of the tangential vector rotational $\nabla_{\partial\Omega}u \times \mathbf{n}$ of u in an orthonormal basis of the ambient space. This feature, in addition to some duality properties, enable us to define this derivative as a bounded operator from $H^s(\partial\Omega)$ into $H^{s-1}(\partial\Omega)$ for $0 \leq s \leq 1$. This is actually equivalent to arguing that $u \rightarrow \nabla_{\partial\Omega}u \times \mathbf{n}$ is a bounded operator from $H^s(\partial\Omega)$ into $H^{s-1}(\partial\Omega; \mathbb{C}^3)$ for $0 \leq s \leq 1$, a result which was established for $s = 1/2$ in [2] from a different technique. It then follows that its transpose, yielding the surface rotational $\nabla_{\partial\Omega} \cdot \mathbf{u} \times \mathbf{n}$ of a vector field \mathbf{u} [23, p. 73], defines also a bounded operator from $H^s(\partial\Omega; \mathbb{C}^3)$ into $H^{s-1}(\partial\Omega)$ for $0 \leq s \leq 1$. It is worth noting that in [23, p. 73] the surface rotational was considered for tangential vector fields only. However, the cross-product involved in the expression of this operator makes it possible to extend this definition to a general vector field. We show then that G nter derivatives $\mathcal{M}_{ij}^{(n)}$ can be expressed as differential forms to retrieve an integration by parts formula relatively to these operators on a patch of $\partial\Omega$. Even if this formula was already established by direct calculation in [12], we think that the formalism of differential forms is more appropriate for understanding the basic principle underlying its derivation. It is used here to get explicit expressions for the G nter derivatives of a piecewise smooth function defined on a the boundary of a curved polyhedron. This way to write these derivatives is fundamental in the effective implementations of boundary element codes. In section 3, we begin with some recalls on other previous expressions for G nter derivative matrix $\mathcal{M}^{(n)}$. With the help of a vectorial Green formula, partly introduced in [16] and in a more complete form in [5, 6], we derive a useful volume variational expression for $\mathcal{M}^{(n)}$. As an application in section 4, we extend the regularization techniques (the way for expressing non-integrable kernels involved in boundary layer potentials in terms of integrals converging in the usual meaning) devised by Le Lou r [19, 18] for the elastic wave layer potentials to Lipschitz domains. It is worth recalling that due to its importance in practical implementations of numerical solvers for elastic wave scattering problems, several other regularizations techniques, much more involved in our opinion, have been already proposed (cf., for example, [22, 25, 1, 20, 10, 27] to cite a few). Finally, in Section 5, making use of the connection between two- and three-dimensional Green kernels for the Helmholtz equation, we transpose the regularization techniques in the spatial scale to planar elastic waves.

2 Extension of the G nter derivatives to a Lipschitz domain

We first establish some mapping properties of the G nter derivatives in the framework of a Lipschitz domain. We next show that they can be written as differential 2-forms, up to a Hodge star identification. This will allow us to retrieve an integration by parts formula on the patches of $\partial\Omega$. That yields an expression of these derivatives well suited for boundary element codes.

2.1 Mapping properties of the Günter derivatives for Lipschitz domains

Property (3) ensures that $\mathcal{M}_{ij}^{(\mathbf{n})}$ is a first-order differential operator tangential to $\partial\Omega$ in the sense of [21, p. 147]. This immediately leads to the following first mapping property whose proof is given in Lemma 4.23 of this reference.

Proposition 1 *There exists a constant C independent of $u \in H^1(\partial\Omega)$ such that*

$$\left\| \mathcal{M}_{ij}^{(\mathbf{n})} u \right\|_{L^2(\partial\Omega)} \leq C \|u\|_{H^1(\partial\Omega)}. \quad (4)$$

To go further, we make the following observation which, surprisingly enough, does not seem to have been done before. It consists in noting that vector $\boldsymbol{\tau}_{ij}$, defined in (2), can be written under the following form using the elementary double product formula

$$\boldsymbol{\tau}_{ij} = (\mathbf{n} \cdot \mathbf{e}_j) \mathbf{e}_i - (\mathbf{n} \cdot \mathbf{e}_i) \mathbf{e}_j = \mathbf{n} \times (\mathbf{e}_i \times \mathbf{e}_j) \quad (1 \leq i, j \leq 3).$$

In this way, using the properties of the mixed product, we can also put Günter derivative $\mathcal{M}_{ij}^{(\mathbf{n})} u$ in the following form

$$\mathcal{M}_{ij}^{(\mathbf{n})} u = \nabla u \cdot \mathbf{n} \times (\mathbf{e}_i \times \mathbf{e}_j) = \nabla u \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j. \quad (5)$$

Indeed, formula (5) expresses $\mathcal{M}_{ij}^{(\mathbf{n})} u$ as a component (or its opposite) of the tangential vector rotational of u in the canonical basis of \mathbb{R}^3

$$\mathcal{M}_{ij}^{(\mathbf{n})} u = \nabla_{\partial\Omega} u \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j. \quad (6)$$

(See [23, p. 69] for the definition and properties of the tangential gradient $\nabla_{\partial\Omega} u$ and the tangential vector rotational of a function when, for example, Ω^+ is a \mathcal{C}^2 -domain.)

Remark 1 *Formula (6) was already given in [17, Formulae (1.12) in Chap. V] but written in terms of the components of $\nabla_{\partial\Omega} u$, \mathbf{n} , and the Levi-Civita symbol ε_{ijk} ($\varepsilon_{ijk} = \pm 1$ if $\{i, j, k\}$ is an even or odd permutation of $\{1, 2, 3\}$ respectively, and 0 otherwise) as follows*

$$\mathcal{M}_{jk}^{(\mathbf{n})} u = - \sum_{i=1}^3 \varepsilon_{ijk} \partial_{s_i} u, \quad (7)$$

with

$$\partial_{s_k} u = \sum_{i,j=1}^3 \varepsilon_{ijk} n_i (\nabla_{\partial\Omega} u)_j.$$

Actually, it is necessary to remark that

$$\partial_{s_i} u = (\mathbf{n} \times \nabla_{\partial\Omega} u)_i$$

and to use the connection of the Levi-Civita symbol to the components of $\mathbf{e}_i \times \mathbf{e}_j$

$$\mathbf{e}_i \times \mathbf{e}_j = \sum_{k=1}^3 \varepsilon_{ijk} \mathbf{e}_k \quad (8)$$

to be able to write (7) in the form

$$\mathcal{M}_{jk}^{(\mathbf{n})} u = \sum_{i=1}^3 (\nabla_{\partial\Omega} u \times \mathbf{n})_i \mathbf{e}_i \cdot \mathbf{e}_j \times \mathbf{e}_k$$

which readily gives (6). It must be noticed that neither the compact form (6), nor more importantly formula (5), expressing Günter derivative $\mathcal{M}_{jk}^{(\mathbf{n})} u$ as a component of the rotational vector $\nabla_{\partial\Omega} u \times \mathbf{n}$ have been stated explicitly in [17] or elsewhere. This is more than just two different ways of writing the same formula as illustrated in the proof of the next lemma.

We have next the following lemma which is established in a less straightforward way in [12] for a $\mathcal{C}^{1,\alpha}$ -domain ($0 < \alpha \leq 1$).

Lemma 1 For u and v in $\mathcal{C}_{comp}^1(\mathbb{R}^3)$, the following integration by parts formula

$$\int_{\partial\Omega} v \mathcal{M}_{ij}^{(\mathbf{n})} u \, ds = \int_{\partial\Omega} u \mathcal{M}_{ji}^{(\mathbf{n})} v \, ds \quad (9)$$

holds true.

Proof. The proof directly follows from the following simple observation

$$\nabla \times (u \mathbf{e}_i \times \mathbf{e}_j) = v \nabla \times (\mathbf{e}_i \times \mathbf{e}_j) - u \nabla \times (\mathbf{e}_j \times \mathbf{e}_i)$$

and Green's formula in Lipschitz domains [21, Th. 3.34]

$$\begin{aligned} \int_{\Omega^\pm} \underbrace{\nabla \cdot \nabla \times (u \mathbf{e}_i \times \mathbf{e}_j)}_{=0} \, dx &= \pm \int_{\partial\Omega} (\nabla \times (u \mathbf{e}_i \times \mathbf{e}_j)) \cdot \mathbf{n} \, ds \\ &= \mp \left(\int_{\partial\Omega} v \mathcal{M}_{ij}^{(\mathbf{n})} u \, ds - \int_{\partial\Omega} u \mathcal{M}_{ji}^{(\mathbf{n})} v \, ds \right). \end{aligned}$$

■

We then come to the following theorem embodying optimal mapping properties of the G nter derivatives.

Theorem 1 Under the above assumption that Ω^+ is a bounded Lipschitz domain, G nter derivative $\mathcal{M}_{ij}^{(\mathbf{n})}$ can be extended in a bounded linear operator from $H^s(\partial\Omega)$ into $H^{s-1}(\partial\Omega)$ for $0 \leq s \leq 1$.

Proof. It is a straightforward consequence of estimate (4) and symmetry property (9) by duality and interpolation techniques. ■

Corollary 1 Under the general assumptions of the above theorem, the tangential vector rotational defines a bounded linear operator $u \rightarrow \nabla_{\partial\Omega} u \times \mathbf{n}$ from $H^s(\partial\Omega)$ into $H^{s-1}(\partial\Omega; \mathbb{C}^3)$ for $0 \leq s \leq 1$. Consequently, the surface rotational gives rise to a bounded operator $\mathbf{u} \in H^s(\partial\Omega; \mathbb{C}^3) \rightarrow \nabla_{\partial\Omega} \cdot \mathbf{u} \times \mathbf{n} \in H^{s-1}(\partial\Omega)$ for $0 \leq s \leq 1$.

Proof. Immediate since the components of $\nabla_{\partial\Omega} u \times \mathbf{n}$ are nothing else but G nter derivatives and the surface rotational is the transpose of the tangential vector rotational. ■

Remark 2 When u and v are the respective traces of functions in $H^1(\mathbb{R}^3)$, it is established in [2, p. 855] that $\nabla u \times \mathbf{n}$ is well-defined in $H_{\parallel}^{-1/2}(\partial\Omega; \mathbb{C}^3)$, the dual space of

$$\begin{aligned} H_{\parallel}^{1/2}(\partial\Omega; \mathbb{C}^3) &= \{ \mathbf{v} \in L^2(\partial\Omega; \mathbb{C}^3); \\ &\quad \mathbf{v} = \mathbf{n} \times (\mathbf{w} \times \mathbf{n}), \mathbf{w} \in H^1(\Omega^\pm; \mathbb{C}^3) \} \end{aligned}$$

equipped with the graph norm and that $\nabla u \times \mathbf{n}$ depends on the trace $u|_{\partial\Omega}$ of u on $\partial\Omega$ [2, p. 855] only. It is also proved in this paper [2, formulae (15) p. 850 and Lemma 2.3 p. 851] that $H_{\parallel}^{-1/2}(\partial\Omega; \mathbb{C}^3)$ can be identified to a closed subspace of $H^{-1/2}(\partial\Omega; \mathbb{C}^3)$. This is the particular case corresponding to $s = 1/2$ which has been previously mentioned.

The following symmetry result is known for a long time in the case of smoother domains and more regular functions [17, p. 284] and is an immediate consequence of the definition of the G nter derivative matrix and the symmetry and mapping properties of $\mathcal{M}_{ij}^{(\mathbf{n})}$.

Corollary 2 G nter derivative matrix $\mathcal{M}^{(\mathbf{n})}$ defines a bounded linear operator from $H^s(\partial\Omega; \mathbb{C}^3)$ into $H^{s-1}(\partial\Omega; \mathbb{C}^3)$ for $0 \leq s \leq 1$ with the following symmetry property

$$\begin{aligned} \langle \mathbf{v}, \mathcal{M}^{(\mathbf{n})} \mathbf{u} \rangle_{1-s, \partial\Omega} &= \langle \mathbf{u}, \mathcal{M}^{(\mathbf{n})} \mathbf{v} \rangle_{s, \partial\Omega}, \\ \mathbf{u} \in H^s(\partial\Omega; \mathbb{C}^3), \mathbf{v} \in H^{1-s}(\partial\Omega; \mathbb{C}^3). \end{aligned} \quad (10)$$

For simplicity, we keep the same notation for the bilinear form underlying the duality product between $H^s(\partial\Omega; \mathbb{C}^3)$ and $H^{-s}(\partial\Omega; \mathbb{C}^3)$ and that $\langle \cdot, \cdot \rangle_{s, \partial\Omega}$ between $H^s(\partial\Omega)$ and $H^{-s}(\partial\Omega)$

$$\langle \mathbf{v}, \boldsymbol{\ell} \rangle_{s, \partial\Omega} = \sum_{i=1}^3 \langle v_i, \ell_i \rangle_{s, \partial\Omega}, \quad \boldsymbol{\ell} \in H^{-s}(\partial\Omega; \mathbb{C}^3), \mathbf{v} \in H^s(\partial\Omega; \mathbb{C}^3).$$

Remark 3 The duality $H^s(\partial\Omega)$, $H^{-s}(\partial\Omega)$ is usually denoted by $\langle \ell, v \rangle_{s, \partial\Omega}$ for $\ell \in H^{-s}(\partial\Omega)$ and $v \in H^s(\partial\Omega)$. The transposition used here is convenient for the notation of the single-layer potential of elastic waves given below.

2.2 Explicit expression for the Günter derivatives

Up to now, we have defined the Günter derivatives just in the distributional sense: $\mathcal{M}_{ij}^{(n)}u \in H^{s-1}(\partial\Omega)$ for $u \in H^s(\partial\Omega)$, $0 \leq s \leq 1$. In concrete applications, $\partial\Omega$ must be considered as the boundary of a curved polyhedron. This is the case of course when $\partial\Omega$ presents curved faces and edges, and vertices, but also once the geometry has been effectively approximated (cf., for example, [28, p. 15]). This means that $\partial\Omega$ can be covered by a non-overlapping decomposition \mathcal{T}

$$\partial\Omega = \cup_{\omega \in \mathcal{T}} \bar{\omega}$$

where \mathcal{T} is a finite family of open domains ω of $\partial\Omega$ such that for all $\omega, v \in \mathcal{T}$, $\omega \cap v = \emptyset$ when $\omega \neq v$. Each ω is assumed to be a “surface polygonal domain” in the meaning that $\bar{\omega} \subset U_\omega$ (U_ω being an open C^∞ -parametrized surface of \mathbb{R}^3), that its boundary $\partial\omega$ is a piecewise smooth curve, and that ω is a Lipschitz domain of U_ω . Lipschitz domains of smooth manifolds are defined similarly to Lipschitz domains of \mathbb{R}^N replacing “rigid motions” in [21, Definition 3.28] by local C^∞ -diffeomorphisms onto domains of \mathbb{R}^2 . Recall that Ω^+ is globally a Lipschitz domain, hence preventing $\partial\Omega$ to present cusp points. Simple and widespread examples of such boundaries are given by triangular meshes of surfaces of \mathbb{R}^3 . Figure 2 depicts a surface triangular mesh of a $C^{1,1}$ -domain. The geometry and the mesh have been designed using the free software *Gmsh* [9]. For the exact surface, ω and U_ω are obtained by local coordinate systems (local charts) (see, for example, [3]). For the approximate surface, ω is a triangle of \mathbb{R}^3 and U_ω is the plane supporting this triangle.




Figure 2: Polyhedral domain obtained from the surface mesh of $C^{1,1}$ -domain.

Boundary element spaces are generally subspaces of the following one

$$\mathcal{P}_{m,\mathcal{T}}(\partial\Omega) = \left\{ u \in L^\infty(\partial\Omega); u|_\omega \circ \Phi_\omega \in \mathbb{P}_m^{(2)}, \forall \omega \in \mathcal{T} \right\}$$

where $\Phi_\omega : D_\omega \subset \mathbb{R}^2 \rightarrow U_\omega$ is a local coordinate system on U_ω (cf., for example, [3, p. 111]), and $\mathbb{P}_m^{(2)}$ is the space of polynomials of degree $\leq m$ in two variables with complex coefficients. Such kinds of spaces are contained in $H^s(\partial\Omega)$ for $1/2 \leq s \leq 1$ if and only if they are contained in

$$\mathcal{C}_\mathcal{T}(\partial\Omega) = \left\{ u \in C^0(\partial\Omega); u|_\omega \in C^\infty(\bar{\omega}), \forall \omega \in \mathcal{T} \right\}$$

(see, for example, [11] when $\partial\Omega = \mathbb{R}^2$).

For $u \in \mathcal{C}_\mathcal{T}(\partial\Omega)$, we can define $\mathcal{M}_{ij,\mathcal{T}}^{(n)}u$ almost everywhere on $\partial\Omega$ by

$$\left(\mathcal{M}_{ij,\mathcal{T}}^{(n)}u \right)|_\omega = \nabla_\omega u|_\omega \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j, \forall \omega \in \mathcal{T}$$

where ∇_ω is the tangential gradient on ω and \mathbf{n} is the unit normal on ω pointing outward from Ω^+ . Our objective is to show that

$$\mathcal{M}_{ij}^{(n)}u = \mathcal{M}_{ij,\mathcal{T}}^{(n)}u \text{ for all } u \in \mathcal{C}_\mathcal{T}(\partial\Omega). \quad (11)$$

This identification requires some preliminaries to be established.

First, we can assume that $u|_\omega$ is the trace of a function u_ω which is C^∞ in a neighborhood in \mathbb{R}^3 of U_ω . We can hence write

$$\begin{aligned}\nabla_\omega u|_\omega \times \mathbf{n} \cdot (\mathbf{e}_i \times \mathbf{e}_j) &= (\nabla u_\omega)|_\omega \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j \\ &= (\mathbf{e}_i \times \mathbf{e}_j) \times (\nabla u_\omega)|_\omega \cdot \mathbf{n}\end{aligned}$$

Using (8), the canonical identification of vector fields to 1-forms on \mathbb{R}^3 and the Hodge star operator on \mathbb{R}^3 , $(\mathbf{e}_i \times \mathbf{e}_j) \times (\nabla u_\omega)|_\omega$ can be written as follows

$$\begin{aligned}(\mathbf{e}_i \times \mathbf{e}_j) \times (\nabla u)|_\omega &= \sum_{k=1}^3 \varepsilon_{ijk} * (dx_k \wedge du_\omega)|_\omega \\ &= \sum_{k=1}^3 \varepsilon_{ijk} * (d(-u_\omega dx_k))|_\omega\end{aligned}$$

We thus retrieve the following result established component by component in [12] without the formalism of differential forms.

Lemma 2 For $u \in C^\infty(\bar{\omega})$ and $v \in C_{comp}^1(\mathbb{R}^3)$, the following integration by parts formula

$$\begin{aligned}\int_\omega v \nabla_\omega u \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j ds &= - \sum_{k=1}^3 \varepsilon_{ijk} \int_{\partial\omega_\circ} uv dx_k \\ &\quad - \int_\omega u \nabla_\omega v \times \mathbf{n} \cdot \mathbf{e}_i \times \mathbf{e}_j ds\end{aligned}$$

holds true. The orientation $\partial\omega_\circ$ is that induced by \mathbf{n} .

Proof. The lemma results from the following observations

$$\nabla_\omega uv = v \nabla_\omega u + u \nabla_\omega v$$

$$\begin{aligned}\int_\omega \nabla_\omega uv \times \mathbf{n} \cdot (\mathbf{e}_i \times \mathbf{e}_j) ds &= \sum_{k=1}^3 \varepsilon_{ijk} \int_\omega * (d(-uv dx_k)) \cdot \mathbf{n} ds \\ &= \sum_{k=1}^3 \varepsilon_{ijk} \int_\omega d(-uv dx_k)\end{aligned}$$

and Stokes' formula. ■

The following theorem gives a simple way to calculate the Günter derivatives when dealing with a boundary element method.

Theorem 2 Formula (11) holds true for any $u \in \mathcal{C}_\mathcal{T}(\partial\Omega)$.

Proof. Clearly, $\mathcal{C}_\mathcal{T}(\partial\Omega) \subset H^1(\partial\Omega)$. Hence, for $u \in \mathcal{C}_\mathcal{T}(\partial\Omega)$ and $v \in C_{comp}^\infty(\mathbb{R}^3)$, symmetry property (10) yields

$$\left\langle v, \mathcal{M}_{ij}^{(\mathbf{n})} u \right\rangle_{1-s, \partial\Omega} = \left\langle u, \mathcal{M}_{ji}^{(\mathbf{n})} v \right\rangle_{s, \partial\Omega} = \int_{\partial\Omega} u \mathcal{M}_{ji, \mathcal{T}}^{(\mathbf{n})} v ds.$$

Integrating by parts, we can write

$$\int_{\partial\Omega} u \mathcal{M}_{ji, \mathcal{T}}^{(\mathbf{n})} v ds = - \sum_{\omega \in \mathcal{T}} \sum_{k=1}^3 \varepsilon_{ijk} \int_{\partial\omega_\circ} uv dx_k + \int_{\partial\Omega} v \mathcal{M}_{ij, \mathcal{T}}^{(\mathbf{n})} u ds.$$

Since

$$\sum_{\omega \in \mathcal{T}} \int_{\partial\omega_\circ} uv dx_k = 0,$$

due to the opposite orientation on each curved edge of the non-overlapping decomposition \mathcal{T} of $\partial\Omega$, we get

$$\int_{\partial\Omega} v \mathcal{M}_{ij, \mathcal{T}}^{(\mathbf{n})} u ds = \left\langle v, \mathcal{M}_{ij}^{(\mathbf{n})} u \right\rangle_{1-s, \partial\Omega}.$$

Formula (11) then results from the density of $C_{comp}^\infty(\mathbb{R}^3)$ in $H^{1-s}(\partial\Omega)$. ■

Remark 4 The density of $C_{comp}^\infty(\mathbb{R}^3)$ in $H^s(\partial\Omega)$ ($0 \leq s \leq 1$), for a Lipschitz domain Ω^+ , can be established along the same lines than that of $C_{comp}^\infty(\mathbb{R}^3)$ in $L^2(\partial\Omega)$, which is proved in [24, Th. 4.9].

3 Other expressions of the G unter derivative matrix

We first examine previous ways to write the G unter derivative matrix when Ω^+ is of class $\mathcal{C}^{1,1}$. We then show whether or not these expressions can be extended to a Lipschitz domain. In particular, we recall a way to write $\mathcal{M}^{(n)}$ variationally by means of a volume integral, already considered elsewhere but not in the present context.

3.1 Previous equivalent expressions for the G unter derivative matrix

We begin with the following compact expression for the G unter derivative matrix given in [19]

$$\mathcal{M}^{(n)}\mathbf{u} = \nabla\mathbf{u}\mathbf{n} - \mathbf{n}\nabla\cdot\mathbf{u}, \quad \mathbf{u} \in H_{\text{loc}}^2(\mathbb{R}^3; \mathbb{C}^3) \quad (12)$$

which can be obtained by observing that

$$\sum_{j=1}^3 \mathcal{M}_{ij}^{(n)} u_j = \sum_{j=1}^3 \partial_{x_i} u_j n_j - n_i \sum_{j=1}^3 \partial_{x_j} u_j.$$

Recall that gradient $\nabla\mathbf{u}$ of vector \mathbf{u} is the matrix whose column j is ∇u_j for $j = 1, 2, 3$.

Probably to more clearly bring out that expression (12) depends on $\mathbf{u}|_{\partial\Omega}$ only, Le Lou er [19] used the following way to write the gradient on $\partial\Omega$

$$\nabla u_j = \nabla_{\partial\Omega} u_j + \mathbf{n} \partial_{\mathbf{n}} u_j \quad (j = 1, 2, 3)$$

to get

$$\mathcal{M}^{(n)}\mathbf{u} = \nabla_{\partial\Omega}\mathbf{u}\mathbf{n} - \text{tr} \nabla_{\partial\Omega}\mathbf{u} \mathbf{n}, \quad (13)$$

$\nabla_{\partial\Omega}\mathbf{u}$ being defined similarly to $\nabla\mathbf{u}$ and $\text{tr} \nabla_{\partial\Omega}\mathbf{u}$ being the algebraic trace of $\nabla_{\partial\Omega}\mathbf{u}$. The reader should be aware that in the formula stating the above expression of $\mathcal{M}^{(n)}\mathbf{u}$ in [19] $\nabla_{\partial\Omega}\cdot\mathbf{u}$ stands for $\text{tr} \nabla_{\partial\Omega}\mathbf{u}$.

Actually, $\text{tr} \nabla_{\partial\Omega}\mathbf{u}$ can be put in the following form

$$\text{tr} \nabla_{\partial\Omega}\mathbf{u} = \text{tr} \nabla\mathbf{u} - \text{tr} \mathbf{n} (\partial\mathbf{u})^\top = \nabla\cdot\mathbf{u} - \mathbf{n}\cdot\partial_{\mathbf{n}}\mathbf{u}$$

and since

$$\nabla\cdot\mathbf{u} = \nabla_{\partial\Omega}\cdot\mathbf{n} \times (\mathbf{u} \times \mathbf{n}) + 2\mathcal{H}\mathbf{u}\cdot\mathbf{n} + \mathbf{n}\cdot\partial_{\mathbf{n}}\mathbf{u} \quad \text{on } \partial\Omega, \quad (14)$$

where $\nabla_{\partial\Omega}\cdot$ denotes the surface divergence (see, for example, [23, p. 72 and 75]), this yields

$$\mathcal{M}^{(n)}\mathbf{u} = \nabla_{\partial\Omega}\mathbf{u}\mathbf{n} - \mathbf{n} (\nabla_{\partial\Omega}\cdot\mathbf{n} \times (\mathbf{u} \times \mathbf{n}) + 2\mathcal{H}\mathbf{u}\cdot\mathbf{n}). \quad (15)$$

We have denoted by $2\mathcal{H} = \text{tr} \mathcal{C}$ the mean Gaussian curvature of $\partial\Omega$, defined as the algebraic trace of the Gauss curvature operator $\mathcal{C} = \nabla\mathbf{n}$. Formula (14) requires a domain of class $\mathcal{C}^{1,1}$ at least to be stated.

There is one concern with both expressions (13) and (15): they do not clearly express that $\mathcal{M}^{(n)}$ is a symmetric operator as stated in (10). With regard to this point, one can first observe that

$$\nabla_{\partial\Omega}\mathbf{v} \mathbf{n} = \sum_{j=1}^3 n_j \nabla_{\partial\Omega} v_j = \sum_{j=1}^3 \nabla_{\partial\Omega} (n_j v_j) - \sum_{j=1}^3 v_j \nabla_{\partial\Omega} n_j$$

Since $\nabla_{\partial\Omega} n_j = \nabla n_j = \mathcal{C}_{*j}$, the j -th column of \mathcal{C} , and $\mathcal{C}\mathbf{n} = 0$, we can write

$$\nabla_{\partial\Omega}\mathbf{v} \mathbf{n} = \nabla_{\partial\Omega}\mathbf{v}\cdot\mathbf{n} - \mathcal{C}\mathbf{n} \times (\mathbf{v} \times \mathbf{n})$$

coming, at least when Ω^+ is $\mathcal{C}^{1,1}$ -domain and $\mathbf{u} \in H^2(\mathbb{R}^3; \mathbb{C}^3)$, to the following way to write the G unter derivative matrix

$$\begin{aligned} \mathcal{M}^{(n)}\mathbf{u} = & \nabla_{\partial\Omega}\mathbf{u}\cdot\mathbf{n} - \mathbf{n}\nabla_{\partial\Omega}\cdot\mathbf{n} \times (\mathbf{u} \times \mathbf{n}) \\ & - \mathcal{C}(\mathbf{n} \times (\mathbf{u} \times \mathbf{n})) - 2\mathcal{H}\mathbf{u}\cdot\mathbf{n} \mathbf{n} \end{aligned} \quad (16)$$

more clearly expressing the symmetry properties stated above. However, since this expression involves the curvature operator \mathcal{C} and the mean curvature $2\mathcal{H}$ in an explicit way, it becomes meaningless for a Lipschitz domain even when not taking care of its derivation.

We now come to the expression of the G unter derivative matrix most often used to express the traction in Lam e static elasticity [17, formula (1.14) p. 282]

$$\mathcal{M}^{(n)}\mathbf{u} = \partial_{\mathbf{n}}\mathbf{u} + \mathbf{n} \times \nabla \times \mathbf{u} - \mathbf{n}\nabla\cdot\mathbf{u}. \quad (17)$$

Since the derivation of this formula does not seem to have been explicitly carried out before, for the convenience of the reader, we show how it can be established from the above compact expression of $\mathcal{M}^{(\mathbf{n})}$. Writing

$$\mathcal{M}^{(\mathbf{n})}\mathbf{u} = \left(\nabla\mathbf{u} - \nabla\mathbf{u}^\top\right)\mathbf{n} + \nabla\mathbf{u}^\top\mathbf{n} - \mathbf{n}\nabla\cdot\mathbf{u},$$

we get

$$\mathcal{M}^{(\mathbf{n})}\mathbf{u} = \partial_{\mathbf{n}}\mathbf{u} + \left(\nabla\mathbf{u} - \nabla\mathbf{u}^\top\right)\mathbf{n} - \mathbf{n}\nabla\cdot\mathbf{u}.$$

Now

$$\left(\nabla\mathbf{u} - \nabla\mathbf{u}^\top\right)_{ij} = \partial_{x_i}u_j - \partial_{x_j}u_i = \sum_{l,m=1}^3 (\delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}) \partial_{x_l}u_m.$$

Using the elementary writing of $\delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}$ in terms of the Levi-Civita symbol

$$\delta_{il}\delta_{jm} - \delta_{im}\delta_{jl} = \sum_{k=1}^3 \varepsilon_{ijk}\varepsilon_{lmk}$$

we come to

$$\begin{aligned} \left(\nabla\mathbf{u} - \nabla\mathbf{u}^\top\right)_{ij} &= \sum_{l,m,k=1}^3 \varepsilon_{ijk}\varepsilon_{lmk}\partial_{x_l}u_m \\ &= \sum_{k=1}^3 \varepsilon_{ijk}(\nabla\times\mathbf{u})_k, \end{aligned}$$

and thus to

$$\begin{aligned} \left(\left(\nabla\mathbf{u} - \nabla\mathbf{u}^\top\right)\mathbf{n}\right)_i &= \sum_{j,k=1}^3 \varepsilon_{ijk}n_j(\nabla\times\mathbf{u})_k \\ &= (\mathbf{n}\times\nabla\times\mathbf{u})_i \quad (i = 1, 2, 3). \end{aligned}$$

Form (17) of $\mathcal{M}^{(\mathbf{n})}\mathbf{u}$ gives rise to two concerns also:

- At least, in a direct way, it can not be evaluated from $\mathbf{u}|_{\partial\Omega}$ only;
- Contrary to (16), it keeps a meaning when Ω^+ is only a Lipschitz domain but requires that $\mathbf{u} \in H^2(\mathbb{R}^3; \mathbb{C}^3)$ to be defined.

With regard to the first of the above two points, Darbas and Le Louër [8] used expression (14) for the divergence [23, Formula (2.5.215)] together with the following one for the curl

$$\mathbf{n}\times\nabla\times\mathbf{u} = \nabla_{\partial\Omega}\mathbf{u}\cdot\mathbf{n} - \mathcal{C}\mathbf{n}\times(\mathbf{u}\times\mathbf{n}) - \mathbf{n}\times(\partial_{\mathbf{n}}\mathbf{u}\times\mathbf{n})$$

[23, Formula (2.5.225)] to get formula (16) from formula (17).

3.2 Expression of the G unter derivative matrix by a volume integral

The trace $\partial_{\mathbf{n}}\mathbf{u} + \mathbf{n}\times\nabla\times\mathbf{u} - \mathbf{n}\nabla\cdot\mathbf{u}$ has been considered in [7, Proof of Lemma 2.1 p. 248] without any reference to the G unter derivatives. More particularly, collecting some formulae in this paper, we readily come to the following Green formula

$$\begin{aligned} \int_{\Omega^\pm} \nabla\mathbf{u}\cdot\nabla\mathbf{v} - \nabla\times\mathbf{u}\cdot\nabla\times\mathbf{v} - \nabla\cdot\mathbf{u}\nabla\cdot\mathbf{v} \, dx = \\ \pm \int_{\partial\Omega} (\partial_{\mathbf{n}}\mathbf{u} + \mathbf{n}\times\nabla\times\mathbf{u} - \mathbf{n}\nabla\cdot\mathbf{u})\cdot\mathbf{v} \, ds \end{aligned} \quad (18)$$

for \mathbf{u} and \mathbf{v} in $H^2(\mathbb{R}^3; \mathbb{C}^3)$ where the bilinear form underlying the scalar product of two 3×3 matrices is defined by

$$\nabla\mathbf{u}\cdot\nabla\mathbf{v} = \sum_{j=1}^3 \nabla u_j\cdot\nabla v_j = \sum_{i,j=1}^3 \partial_{x_i}u_j\partial_{x_i}v_j.$$

It is assumed there that Ω^+ is a curved polyhedron but the derivation remains valid when Ω^+ is a Lipschitz domain and for \mathbf{v} in $H^1(\Omega^\pm; \mathbb{C}^3)$. In the same way, the above Green formula is still holding true for $\mathbf{u} \in H_{loc}^2(\mathbb{R}^3; \mathbb{C}^3)$ and $\mathbf{v} \in H_{comp}^1(\mathbb{R}^3; \mathbb{C}^3)$ or $\mathbf{u} \in H_{comp}^2(\mathbb{R}^3; \mathbb{C}^3)$ and $\mathbf{v} \in H_{loc}^1(\mathbb{R}^3; \mathbb{C}^3)$. Actually, formula (18) can also be directly deduced from an older Green formula considered in [16, p. 220]

$$\int_{\Omega^\pm} \Delta \mathbf{u} \cdot \mathbf{v} + \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} + \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} \, dx = \pm \int_{\partial\Omega} (\nabla \times \mathbf{u} \times \mathbf{n} + \mathbf{n} \nabla \cdot \mathbf{u}) \cdot \mathbf{v} \, ds. \quad (19)$$

We then directly come to the following theorem giving the expression of the Günter derivative matrix in terms of a volume integral.

Theorem 3 *Let Ω^+ be a bounded Lipschitz domain of \mathbb{R}^3 . Using the general notation introduced above, we have*

$$\langle \mathbf{v}, \mathcal{M}^{(n)} \mathbf{u} \rangle_{1/2, \partial\Omega} = \pm \int_{\Omega^\pm} \nabla \mathbf{u} \cdot \nabla \mathbf{v} - \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} - \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} \, dx \quad (20)$$

for $\mathbf{u} \in H_{loc}^1(\mathbb{R}^3; \mathbb{C}^3)$ and $\mathbf{v} \in H_{comp}^1(\mathbb{R}^3; \mathbb{C}^3)$.

Proof. In view of (18), assuming that $\mathbf{v} \in H_{comp}^2(\mathbb{R}^3; \mathbb{C}^3)$, we can write

$$\int_{\partial\Omega} \mathbf{u} \cdot \mathcal{M}^{(n)} \mathbf{v} \, ds = \pm \int_{\Omega^\pm} \nabla \mathbf{u} \cdot \nabla \mathbf{v} - \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} - \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} \, dx.$$

Noting then that

$$\int_{\partial\Omega} \mathbf{u} \cdot \mathcal{M}^{(n)} \mathbf{v} \, ds = \langle \mathbf{u}, \mathcal{M}^{(n)} \mathbf{v} \rangle_{1/2, \partial\Omega} = \langle \mathbf{v}, \mathcal{M}^{(n)} \mathbf{u} \rangle_{1/2, \partial\Omega}$$

we get (20) for $\mathbf{v} \in H_{comp}^2(\mathbb{R}^3; \mathbb{C}^3)$. The proof can then be readily completed from the density of $H_{comp}^2(\mathbb{R}^3; \mathbb{C}^3)$ in $H_{comp}^1(\mathbb{R}^3; \mathbb{C}^3)$. ■

4 Application to the elastic wave boundary-layer potentials

In this section, we extend the regularization of elastic wave boundary-layer potentials devised by Le Louër [19, 18] for a geometry of class \mathcal{C}^2 to the case of a Lipschitz domain. This extension is straightforward for the traces of the single- and the double-layer potentials. We just more explicitly bring out an intermediary expression for the double-layer potential and an identity linking the elastic wave boundary-layer potentials to those related to the Helmholtz equation. We focus on the traction of the double-layer potential which requires a different technique of proof. Meanwhile, as an application of these regularization techniques, we show how the mapping properties of the elastic waves potentials easily reduce to those related to the Helmholtz equation without resorting to the general theory of boundary layer potentials for elliptic systems.

4.1 Layer potentials of elastic waves

For $\mathbf{p} \in H^{-1/2}(\partial\Omega; \mathbb{C}^3)$, the elastic wave single-layer potential can be expressed as follows

$$S\mathbf{p}(x) = \langle \Gamma(x, y), \mathbf{p}_y \rangle_{1/2, \partial\Omega} \quad (x \in \Omega^+ \cup \Omega^-),$$

in terms of the Kupradze matrix $\Gamma(x, y)$ whose entries are given by [17, p. 85]

$$\Gamma_{kl}(x, y) = \frac{1}{\omega^2 \varrho} (\kappa_s^2 G_{\kappa_s}(x, y) \delta_{kl} + \partial_{x_k} \partial_{x_l} (G_{\kappa_p} - G_{\kappa_s})(x, y)) \quad (k, l = 1, 2, 3).$$

Dummy variable y is used to indicate that the duality brackets link \mathbf{p} to the function $y \rightarrow \Gamma(x, y)$ indexed by parameter x . The notation $\langle \Gamma(x, y), \mathbf{p}_y \rangle_{1/2, \partial\Omega}$ refers to the vector whose component k is given by

$$\sum_{l=1}^3 \langle \Gamma_{kl}(x, y), (p_l)_y \rangle_{1/2, \partial\Omega}$$

where $(p_l)_y$ is component l of \mathbf{p}_y . It is this formula that motivates the transposition in the duality brackets $H^{1/2}(\partial\Omega)$, $H^{-1/2}(\partial\Omega)$ adopted above. As usual

$$\kappa_p = \omega\sqrt{\varrho/(2\mu + \lambda)} \text{ and } \kappa_s = \omega\sqrt{\varrho/\mu}$$

are the wavenumbers corresponding to compression or P-waves and shear or S-waves respectively. The constants ω , ϱ , $\mu > 0$ and $\lambda \geq 0$ characterize the angular frequency of the wave, the density and the Lamé coefficients of the elastic medium respectively. Finally, $G_\kappa(x, y) = \exp(i\kappa|x - y|)/4\pi|x - y|$ is the Green kernel characterizing the solutions of the Helmholtz equation

$$\Delta_y G_\kappa(x, y) + \kappa^2 G_\kappa(x, y) = -\delta_x \text{ in } \mathcal{D}'(\mathbb{R}^3),$$

satisfying the Sommerfeld radiation condition

$$\lim_{|y| \rightarrow \infty} |y| (\partial_{|y|} G_\kappa(x, y) - i\kappa G_\kappa(x, y)) = 0,$$

δ_x being the Dirac mass at x .

Actually, we think that it is more convenient to express $S\mathbf{p}$ in terms of the Helmholtz equation single-layer potentials $V_{\kappa_p}\mathbf{p}$ and $V_{\kappa_s}\mathbf{p}$ characterizing the P- and the S-waves respectively

$$S\mathbf{p} = \frac{1}{\omega^2 \varrho} (\kappa_s^2 V_{\kappa_s}\mathbf{p} + \nabla \nabla \cdot (V_{\kappa_s} - V_{\kappa_p})\mathbf{p}), \quad (21)$$

where generically the single-layer potential related to the Helmholtz equation corresponding to the wave number $\kappa > 0$ is defined by

$$(V_\kappa \mathbf{p}(x))_\ell = \left\langle G_\kappa(x, y), (p_\ell)_y \right\rangle_{1/2, \partial\Omega}, \quad x \in \Omega^+ \cup \Omega^-,$$

$(V_\kappa \mathbf{p}(x))_\ell$ ($\ell = 1, 2, 3$) being the ℓ -th component of $V_\kappa \mathbf{p}(x)$.

The following proposition recalls some important properties of these potentials.

Proposition 2 For $p \in H^{-1/2+s}(\partial\Omega)$, $V_\kappa p \in H_{loc}^{1+s}(\mathbb{R}^3)$, $-1/2 \leq s \leq 1/2$. It satisfies the Helmholtz equation $\Delta V_\kappa p + \kappa^2 V_\kappa p = 0$ in $\Omega^+ \cup \Omega^-$ and the Sommerfeld radiation condition. Moreover

$$(V_{\kappa_s} - V_{\kappa_p})p \in H_{loc}^{3+s}(\mathbb{R}^3). \quad (22)$$

Proof. The mapping property of V_κ is a particular case of that of single-layer potentials of more general elliptic equations (cf., for example, [4, Th. 1] or [21, Th. 6.11]). The fact that it satisfies the Helmholtz equation and the Sommerfeld radiation condition is stated for example in [23, p. 117]. The final property is well-known. For the convenience of the reader, we prove it below. From the definition of V_κ (cf., for example, [21, p. 201]), we can write

$$\Delta V_\kappa p + \kappa^2 V_\kappa p = -p\delta_{\partial\Omega}$$

where $p\delta_{\partial\Omega}$ is the single-layer distribution defined by

$$\langle \varphi, p\delta_{\partial\Omega} \rangle_{\mathcal{D}, \mathcal{D}'} = \langle \varphi|_{\partial\Omega}, p \rangle_{1/2, \partial\Omega}, \quad \varphi \in \mathcal{D}(\mathbb{R}^3)$$

where $\langle \cdot, \cdot \rangle_{\mathcal{D}, \mathcal{D}'}$ is the bilinear form underlying the duality brackets $\mathcal{D}(\mathbb{R}^3)$, $\mathcal{D}'(\mathbb{R}^3)$. Thus

$$\Delta (V_{\kappa_s} - V_{\kappa_p})p = \kappa_p^2 V_{\kappa_p} p - \kappa_s^2 V_{\kappa_s} p.$$

Property (22) is then a direct consequence of the interior regularity for the solutions of the elliptic equations (see, for example, [21, Th. 4.16]). ■

The double-layer potential for elastic waves is defined for $\boldsymbol{\psi} \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$ by [17, p. 301]

$$K\boldsymbol{\psi}(x) = - \int_{\partial\Omega} \left(T_y^{(n)} \Gamma(x, y) \right)^\top \boldsymbol{\psi}(y) ds_y, \quad x \in \Omega^+ \cup \Omega^-,$$

where $T_y^{(n)}$ denotes the traction operator defined for $\mathbf{u} \in H_{loc}^2(\mathbb{R}^3; \mathbb{C}^3)$ by

$$T^{(n)}\mathbf{u} = 2\mu\partial_n \mathbf{u} + \lambda n \nabla \cdot \mathbf{u} + \mu n \times \nabla \times \mathbf{u},$$

$T_y^{(n)}\Gamma(x, y)$ being the matrix whose column j is obtained by applying $T_y^{(n)}$ to column j of $\Gamma(x, y)$. The reader must take care of the fact that the above double-layer as well as the one associated with the Helmholtz equation

$$N_\kappa \lambda(x) = - \int_{\partial\Omega} \partial_{\mathbf{n}_y} G_\kappa(x, y) \lambda(y) ds_y \quad x \in \Omega^+ \cup \Omega^-,$$

are of the opposite sign of those considered in the literature (cf. [17, p. 301] and [15, Formulae (2.2.19) and (1.2.2)]). We find this notation more compatible with the formulae expressing the jump of the traction of the single-layer potential for elastic waves and the normal derivative of the Helmholtz equation single-layer potential.

The above extension of $\mathcal{M}^{(n)}$ to a Lipschitz domain allows us to do the same for the expressions of the double-layer potential devised by Le Louër [19] for \mathcal{C}^2 -domains.

Proposition 3 *The double-layer potential can be expressed as*

$$K\psi = \nabla V_{\kappa_p} \mathbf{n} \cdot \psi - \nabla \times V_{\kappa_s} \mathbf{n} \times \psi - 2\mu S \mathcal{M}^{(n)} \psi \text{ in } \Omega^+ \cup \Omega^-. \quad (23)$$

Moreover, in view of the following identity

$$\nabla V_{\kappa_s} \mathbf{n} \cdot \psi - \nabla \times V_{\kappa_s} \mathbf{n} \times \psi = N_{\kappa_s} \psi + V_{\kappa_s} \mathcal{M}^{(n)} \psi \text{ in } \Omega^+ \cup \Omega^- \quad (24)$$

it can be put also in the following form

$$K\psi = N_{\kappa_s} \psi + (V_{\kappa_s} - 2\mu S) \mathcal{M}^{(n)} \psi + \nabla (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \psi \text{ in } \Omega^+ \cup \Omega^-. \quad (25)$$

Proof. Both the above expressions of $K\psi$ are straightforward extensions of calculations carried out in [19]. Formulae (23) and (24) are stated here in their own right instead of being parts of the calculations. ■

The following theorem can then be proved in an elementary fashion from the properties of the Helmholtz equation layer potentials.

Theorem 4 *The elastic wave layer potentials have the following mapping properties:*

$$\begin{aligned} S &: H^{-1/2+s}(\partial\Omega; \mathbb{C}^3) \rightarrow H_{loc}^{1+s}(\mathbb{R}^3; \mathbb{C}^3) \\ K &: H^{1/2+s}(\partial\Omega; \mathbb{C}^3) \rightarrow H_{loc}^{1+s}(\overline{\Omega^\pm}; \mathbb{C}^3) \end{aligned} \quad \text{for } -1/2 < s < 1/2;$$

The potentials $\mathbf{u} = S\mathbf{p}$ or $\mathbf{u} = K\psi$ satisfy

$$\begin{cases} \Delta^* \mathbf{u} + \omega^2 \varrho \mathbf{u} = 0 \text{ in } \Omega^+ \cup \Omega^-, \\ \mathbf{u} \text{ fulfils the Kupradze radiation conditions [17, p. 124]} \end{cases}$$

where Δ^* is the elastic laplacian given by

$$\Delta^* \mathbf{u} = \mu \Delta \mathbf{u} + (\lambda + \mu) \nabla \nabla \cdot \mathbf{u}.$$

Proof. The first part of the proof follows from Costabel's results on mapping properties of scalar elliptic operators [4]. The second one is obtained by straightforward calculations from (21) and (23). ■

4.2 Traces of elastic wave layer potentials

The traces of the single- and double-layer potentials S and K and their mapping properties can also be deduced from the traces of the layer potentials of the Helmholtz equation.

Theorem 5 *The operators defined by $(S\mathbf{p})^\pm$ for $\mathbf{p} \in H^{-1/2}(\partial\Omega; \mathbb{C}^3)$ and $(K\psi)^\pm$ for $\psi \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$ have the following expressions*

$$\begin{aligned} (S\mathbf{p})^\pm &= \frac{1}{\omega^2 \varrho} (\kappa_s^2 V_{\kappa_s} \mathbf{p} + \nabla \nabla \cdot (V_{\kappa_s} - V_{\kappa_p}) \mathbf{p}), \\ (K\psi)^\pm &= (N_{\kappa_s} \psi)^\pm + (V_{\kappa_s} - 2\mu S) \mathcal{M}^{(n)} \psi + \nabla (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \psi. \end{aligned}$$

In particular, the jumps of the related potentials are given by

$$[S\mathbf{p}] = (S\mathbf{p})^+ - (S\mathbf{p})^- = 0, \quad [K\psi] = \psi.$$

As a result, we simply refer to $(S\mathbf{p})^\pm$ by $S\mathbf{p}$ below.

The mapping properties of these operators are given, for $-1/2 < s < 1/2$, by

$$\begin{aligned} S\mathbf{p} &\in H^{1/2+s}(\partial\Omega; \mathbb{C}^3), \text{ for } \mathbf{p} \in H^{-1/2+s}(\partial\Omega; \mathbb{C}^3); \\ (K\psi)^\pm &\in H^{1/2+s}(\partial\Omega; \mathbb{C}^3), \text{ for } \psi \in H^{1/2+s}(\partial\Omega; \mathbb{C}^3). \end{aligned}$$

Proof. The only point requiring some care concerns the term $\nabla(V_{\kappa_s} - V_{\kappa_p})\mathbf{n} \cdot \boldsymbol{\psi}$. But since $\boldsymbol{\psi} \in H^{1/2+s}(\partial\Omega; \mathbb{C}^3)$, $\mathbf{n} \cdot \boldsymbol{\psi}$ is well-defined in $L^2(\partial\Omega)$ and thus belongs to $H^{-1/2+s}(\partial\Omega)$ for $-1/2 < s < 1/2$. Regularity property (22) then yields

$$\nabla(V_{\kappa_s} - V_{\kappa_p})\mathbf{n} \cdot \boldsymbol{\psi} \in H_{\text{loc}}^{s+2}(\mathbb{R}^3; \mathbb{C}^3) \text{ for } -1/2 < s < 1/2.$$

This is enough to define its trace in $H^{1/2+s}(\partial\Omega; \mathbb{C}^3)$ from Costabel's extension of the trace theorem for Lipschitz domains (see [4, Lemma 3.6] and [21, Th. 3.38]). ■

Remark 5 *The point preventing the extension of the above mapping properties to the end-points, $s = \pm 1/2$, concerns Costabel's extension of the trace theorem from $H_{\text{loc}}^s(\overline{\Omega^\pm})$ onto $H^{s-1/2}(\partial\Omega)$, valid only for $1/2 < s < 3/2$.*

4.3 Traction of the elastic waves layer potentials

We begin with the following classical lemma which defines the traction $T^{(n)}\mathbf{u}$ for \mathbf{u} in the following space

$$H_{\text{loc}}^1(\Delta^*, \overline{\Omega^\pm}) = \left\{ \mathbf{v} \in H_{\text{loc}}^1(\overline{\Omega^\pm}; \mathbb{C}^3); \Delta^* \mathbf{v} \in L_{\text{loc}}^2(\overline{\Omega^\pm}; \mathbb{C}^3) \right\}.$$

Meanwhile, we adapt previous expressions of this operator, written in terms of the G\"unter derivative matrix, to the present context of a Lipschitz geometry.

Lemma 3 *For $\mathbf{u} \in H_{\text{loc}}^1(\Delta^*, \overline{\Omega^\pm})$ and $\mathbf{v} \in H_{\text{comp}}^1(\mathbb{R}^3; \mathbb{C}^3)$, the following formula defines $(T^{(n)}\mathbf{u})^\pm$ in $H^{-1/2}(\partial\Omega; \mathbb{C}^3)$*

$$\int_{\Omega^\pm} 2\mu \nabla \mathbf{u} \cdot \nabla \mathbf{v} - \mu \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} + \lambda \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} + \Delta^* \mathbf{u} \cdot \mathbf{v} \, dx = \left\langle \mathbf{v}, \pm (T^{(n)}\mathbf{u})^\pm \right\rangle_{1/2, -1/2}. \quad (26)$$

Moreover, the traction $(T^{(n)}\mathbf{u})^\pm$ can also be expressed in either of the two following forms

$$\int_{\Omega^\pm} \mu \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} + (\lambda + 2\mu) \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} + \Delta^* \mathbf{u} \cdot \mathbf{v} \, dx + \left\langle \mathbf{v}, \pm 2\mu \mathcal{M}^{(n)} \mathbf{u}^\pm \right\rangle_{1/2, \partial\Omega} = \left\langle \mathbf{v}, \pm (T^{(n)}\mathbf{u})^\pm \right\rangle_{1/2, \partial\Omega}, \quad (27)$$

$$\int_{\Omega^\pm} \mu \nabla \mathbf{u} \cdot \nabla \mathbf{v} + (\lambda + \mu) \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} + \Delta^* \mathbf{u} \cdot \mathbf{v} \, dx + \left\langle \mathbf{v}, \pm \mu \mathcal{M}^{(n)} \mathbf{u}^\pm \right\rangle_{1/2, \partial\Omega} = \left\langle \mathbf{v}, \pm (T^{(n)}\mathbf{u})^\pm \right\rangle_{1/2, \partial\Omega}. \quad (28)$$

Proof. Identity (26) is obtained from identity (19) and usual Green formula for $\mathbf{u} \in H_{\text{loc}}^2(\overline{\Omega^\pm}; \mathbb{C}^3)$ by putting the left-hand side in the form

$$\begin{aligned} \int_{\Omega^\pm} 2\mu \nabla \mathbf{u} \cdot \nabla \mathbf{v} - \mu \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} + \lambda \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v} + \Delta^* \mathbf{u} \cdot \mathbf{v} \, dx = \\ \int_{\Omega^\pm} 2\mu (\nabla \mathbf{u} \cdot \nabla \mathbf{v} + \Delta \mathbf{u} \cdot \mathbf{v}) \, dx \\ + \int_{\Omega^\pm} (\lambda + \mu) (\nabla \nabla \cdot \mathbf{u} \cdot \mathbf{v} + \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v}) \, dx \\ - \int_{\Omega^\pm} \mu (\Delta \mathbf{u} \cdot \mathbf{v} + \nabla \times \mathbf{u} \cdot \nabla \times \mathbf{v} + \nabla \cdot \mathbf{u} \nabla \cdot \mathbf{v}) \, dx. \end{aligned}$$

It is extended to $\mathbf{u} \in H_{\text{loc}}^1(\Delta^*, \overline{\Omega^\pm})$ by usual density, continuity and duality arguments (cf., for example, [11] for the case of the Laplace operator and [4, 21] for more general elliptic problems). Formulae (27) and (28) are then a simple recast of this identity from volume expression (20) of $\mathcal{M}^{(n)}$. ■

Remark 6 *It is worth noting the following two important features:*

1. A first part of the integrand in (26) is exactly the (opposite) of the density of virtual work

$$2\mu\nabla\mathbf{u}\cdot\nabla\mathbf{v}-\mu\nabla\times\mathbf{u}\cdot\nabla\times\mathbf{v}+\lambda\nabla\cdot\mathbf{u}\nabla\cdot\mathbf{v}=\boldsymbol{\Sigma}\mathbf{u}\cdot\mathbf{E}\mathbf{v}$$

done by the internal stresses

$$\boldsymbol{\Sigma}\mathbf{u}=2\mu\mathbf{E}\mathbf{u}+\lambda\nabla\cdot\mathbf{u}\mathbb{I}_3$$

under the virtual displacement \mathbf{v} ; $\mathbf{E}\mathbf{u}=(1/2)(\nabla\mathbf{u}+\nabla\mathbf{u}^\top)$ and \mathbb{I}_3 are the strain tensor and the 3×3 unit matrix respectively;

2. When $\mathbf{u}\in H_{loc}^2(\overline{\Omega^\pm};\mathbb{C}^3)$, usual Green formula yields the well-known representation formulae of the traction in terms of $\mathcal{M}^{(n)}$ (cf. [17, Formula (V, 1.16)] and [15, Formula (2.2.35)])

$$T^{(n)}\mathbf{u}=2\mu\mathcal{M}^{(n)}\mathbf{u}-\mu\mathbf{n}\times\nabla\times\mathbf{u}+(\lambda+2\mu)\mathbf{n}\nabla\cdot\mathbf{u}\quad (29)$$

$$T^{(n)}\mathbf{u}=\mu\mathcal{M}^{(n)}\mathbf{u}+\mu\partial_n\mathbf{u}+(\lambda+\mu)\mathbf{n}\nabla\cdot\mathbf{u}.\quad (30)$$

We can thus establish the representation of the traction of the single-layer potential in terms of the traces of the Helmholtz equation potentials.

Theorem 6 *The operators defined by $(T^{(n)}S\mathbf{p})^\pm$ for $\mathbf{p}\in H^{-1/2}(\partial\Omega;\mathbb{C}^3)$ have the following representation*

$$\left(T^{(n)}S\mathbf{p}\right)^\pm=(\partial_n V_{\kappa_s}\mathbf{p})^\pm+\mathbf{n}\nabla\cdot(V_{\kappa_p}-V_{\kappa_s})\mathbf{p}-\mathcal{M}^{(n)}(V_{\kappa_s}-2\mu S)\mathbf{p}.\quad (31)$$

In particular, the jump of the traction of the single-layer potential is given by

$$\left[T^{(n)}S\mathbf{p}\right]=\mathbf{p}.$$

The mapping properties of these operators can be stated as follows

$$\left(T^{(n)}S\mathbf{p}\right)^\pm\in H^{-1/2+s}(\partial\Omega;\mathbb{C}^3),$$

for $\mathbf{p}\in H^{-1/2+s}(\partial\Omega;\mathbb{C}^3)$ and $-1/2<s<1/2$.

Proof. Keeping the general notation of Lemma 3, we use representation (27) of the traction to write

$$\begin{aligned}\left\langle\mathbf{v},\left(T^{(n)}S\mathbf{p}\right)^\pm\right\rangle_{1/2,\partial\Omega}&=\left\langle\mathbf{v},2\mu\mathcal{M}^{(n)}S\mathbf{p}\right\rangle_{1/2,\partial\Omega} \\ &\pm\int_{\Omega^\pm}\mu\nabla\times S\mathbf{p}\cdot\nabla\times\mathbf{v}+(\lambda+2\mu)\nabla\cdot S\mathbf{p}\nabla\cdot\mathbf{v}+\Delta^*S\mathbf{p}\cdot\mathbf{v}\,dx.\end{aligned}$$

Noting that $\mu\nabla\times S\mathbf{p}=\nabla\times V_{\kappa_s}\mathbf{p}$, $(\lambda+2\mu)\nabla\cdot S\mathbf{p}=\nabla\cdot V_{\kappa_p}\mathbf{p}$, and $\Delta^*S\mathbf{p}=-\omega^2\rho S\mathbf{p}$ in Ω^\pm , we get

$$\begin{aligned}\left\langle\mathbf{v},\left(T^{(n)}S\mathbf{p}\right)^\pm\right\rangle_{1/2,\partial\Omega}&=\left\langle\mathbf{v},2\mu\mathcal{M}^{(n)}S\mathbf{p}\right\rangle_{1/2,\partial\Omega} \\ &\pm\int_{\Omega^\pm}\nabla\times V_{\kappa_s}\mathbf{p}\cdot\nabla\times\mathbf{v}+\nabla\cdot V_{\kappa_p}\mathbf{p}\nabla\cdot\mathbf{v}-\omega^2\rho S\mathbf{p}\cdot\mathbf{v}\,dx.\end{aligned}$$

or in a more explicit form

$$\begin{aligned}\left\langle\mathbf{v},\left(T^{(n)}S\mathbf{p}\right)^\pm\right\rangle_{1/2,\partial\Omega}&=\left\langle\mathbf{v},2\mu\mathcal{M}^{(n)}S\mathbf{p}\right\rangle_{1/2,\partial\Omega} \\ &\pm\int_{\Omega^\pm}\nabla\times V_{\kappa_s}\mathbf{p}\cdot\nabla\times\mathbf{v}+\nabla\cdot V_{\kappa_p}\mathbf{p}\nabla\cdot\mathbf{v}\,dx \\ &\pm\int_{\Omega^\pm}(-\kappa_s^2V_{\kappa_s}\mathbf{p}-\nabla\nabla\cdot V_{\kappa_s}\mathbf{p}+\nabla\nabla\cdot V_{\kappa_p}\mathbf{p})\cdot\mathbf{v}\,dx.\end{aligned}$$

Reorganizing the integrands, we come to

$$\begin{aligned}\left\langle\mathbf{v},\left(T^{(n)}S\mathbf{p}\right)^\pm\right\rangle_{1/2,\partial\Omega}&=\left\langle\mathbf{v},2\mu\mathcal{M}^{(n)}S\mathbf{p}\right\rangle_{1/2,\partial\Omega} \\ &\pm\int_{\Omega^\pm}\nabla\times V_{\kappa_s}\mathbf{p}\cdot\nabla\times\mathbf{v}+\nabla\cdot V_{\kappa_s}\mathbf{p}\nabla\cdot\mathbf{v}+\Delta V_{\kappa_s}\mathbf{p}\cdot\mathbf{v}\,dx \\ &\pm\int_{\Omega^\pm}\nabla\nabla\cdot(V_{\kappa_p}-V_{\kappa_s})\mathbf{p}\cdot\mathbf{v}+\nabla\cdot(V_{\kappa_p}-V_{\kappa_s})\mathbf{p}\nabla\cdot\mathbf{v}\,dx,\end{aligned}$$

which can also be written as

$$\begin{aligned}
\left\langle \mathbf{v}, (T^{(n)}S\mathbf{p})^\pm \right\rangle_{1/2, \partial\Omega} &= \left\langle \mathbf{v}, 2\mu\mathcal{M}^{(n)}S\mathbf{p} \right\rangle_{1/2, \partial\Omega} \\
&\pm \int_{\Omega^\pm} \nabla \times V_{\kappa_s}\mathbf{p} \cdot \nabla \times \mathbf{v} + \nabla \cdot V_{\kappa_s}\mathbf{p} \nabla \cdot \mathbf{v} - \nabla V_{\kappa_s}\mathbf{p} \cdot \nabla \mathbf{v} \, dx \\
&\pm \int_{\Omega^\pm} \Delta V_{\kappa_s}\mathbf{p} \cdot \mathbf{v} + \nabla V_{\kappa_s}\mathbf{p} \cdot \nabla \mathbf{v} \, dx \\
&\pm \int_{\Omega^\pm} \nabla \nabla \cdot (V_{\kappa_p} - V_{\kappa_s})\mathbf{p} \cdot \mathbf{v} + \nabla \cdot (V_{\kappa_p} - V_{\kappa_s})\mathbf{p} \nabla \cdot \mathbf{v} \, dx.
\end{aligned}$$

Volume expression (20) of $\mathcal{M}^{(n)}$ and usual Green formula directly yield (31). The jump of $T^{(n)}S\mathbf{p}$ directly follows from that of the normal derivative of the single-layer potential of the Helmholtz equation. The mapping properties are obtained in the same way than those related to the traces of the double-layer potential. ■

Remark 7 Representation formula (31) establishes the duality identity

$$\left\langle \boldsymbol{\psi}, (T^{(n)}S\mathbf{p})^\pm \right\rangle_{1/2, \partial\Omega} = - \left\langle (K\boldsymbol{\psi})^\mp, \mathbf{p} \right\rangle_{1/2, \partial\Omega},$$

for $\boldsymbol{\psi} \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$ and $\mathbf{p} \in H^{-1/2}(\partial\Omega; \mathbb{C}^3)$, from the corresponding formula for the potentials of the Helmholtz equation without resorting to the general theory for elliptic systems [21, p. 211].

Now we address the perhaps most important issue in this paper: a suitable regularization of the hyper-singular kernels arising in the representation of the traction of the double-layer potential. As said above, we here extend two regularizations, devised by Le Louër [19, 18] for a geometry of class \mathcal{C}^2 , to a Lipschitz domain.

The first regularization is based on formula (25), and can be viewed, at some extent, as a generalization of the static elasticity case derived by Han (cf. [14] and [15, Lemma 2.3.3]).

Theorem 7 For $\boldsymbol{\psi} \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$, the traction of the double-layer potential on each side of $\partial\Omega$ is given by

$$\begin{aligned}
(T^{(n)}K\boldsymbol{\psi})^\pm &= \mu \left((\partial_{\mathbf{n}} N_{\kappa_s}\boldsymbol{\psi}) + \mathcal{M}^{(n)}(N_{\kappa_s}\boldsymbol{\psi})^\pm - (\partial_{\mathbf{n}} V_{\kappa_s}\mathcal{M}^{(n)}\boldsymbol{\psi})^\pm \right) \\
&+ 2\mu \left(\mathcal{M}^{(n)}\nabla(V_{\kappa_p} - V_{\kappa_s})\mathbf{n} \cdot \boldsymbol{\psi} - \mathbf{n}\nabla \cdot (V_{\kappa_p} - V_{\kappa_s})\mathcal{M}^{(n)}\boldsymbol{\psi} \right) \\
&+ \mathcal{M}^{(n)}(3\mu V_{\kappa_s} - 4\mu^2 S)\mathcal{M}^{(n)}\boldsymbol{\psi} - \omega^2 \rho \mathbf{n} (V_{\kappa_p} - V_{\kappa_s})\mathbf{n} \cdot \boldsymbol{\psi}.
\end{aligned} \tag{32}$$

In particular, $[T^{(n)}K\boldsymbol{\psi}] = 0$ and $(T^{(n)}K\boldsymbol{\psi})^\pm = T^{(n)}K\boldsymbol{\psi}$ defines a bounded operator from $H^{1/2+s}(\partial\Omega; \mathbb{C}^3)$ into $H^{-1/2+s}(\partial\Omega; \mathbb{C}^3)$ for $-1/2 < s < 1/2$.

Proof. The calculations follow those in [18, Lemma 2.3]. They are however carried out here on the potentials instead on the kernels. The approach in [18, Lemma 2.3], more or less explicitly, requires a smooth extension of the unit normal in a neighborhood of $\partial\Omega$, which, of course, is not available for a Lipschitz geometry. The derivation is based on the decomposition of the double-layer potential in three terms

$$K\boldsymbol{\psi} = \underbrace{N_{\kappa_s}\boldsymbol{\psi} + V_{\kappa_s}\mathcal{M}^{(n)}\boldsymbol{\psi}}_{\mathbf{w}_0} + \underbrace{\nabla(V_{\kappa_p} - V_{\kappa_s})\mathbf{n} \cdot \boldsymbol{\psi} - 2\mu S\mathcal{M}^{(n)}\boldsymbol{\psi}}_{\mathbf{w}_1}.$$

The last term is just a multiple of the single-layer potential created by the density $\mathcal{M}^{(n)}\boldsymbol{\psi}$: thus the corresponding tractions are given by (31)

$$\begin{aligned}
\left(T^{(n)}(-2\mu S\mathcal{M}^{(n)}\boldsymbol{\psi}) \right)^\pm &= -2\mu \left(\partial_{\mathbf{n}} V_{\kappa_s}\mathcal{M}^{(n)}\boldsymbol{\psi} \right)^\pm \\
&- 2\mu \left(\mathbf{n}\nabla \cdot (V_{\kappa_p} - V_{\kappa_s})\mathcal{M}^{(n)}\boldsymbol{\psi} - \mathcal{M}^{(n)}(V_{\kappa_p} - V_{\kappa_s})\mathcal{M}^{(n)}\boldsymbol{\psi} \right).
\end{aligned}$$

The second term is in $H_{\text{loc}}^2(\overline{\Omega^\pm}; \mathbb{C}^3)$. The corresponding traction can be calculated using the direct definition and the fact that $\nabla \times \mathbf{w}_1 = 0$ and $\nabla \cdot \mathbf{w}_1 = (-\kappa_p^2 V_{\kappa_p} + \kappa_s^2 V_{\kappa_s})\mathbf{n} \cdot \boldsymbol{\psi}$

$$T^{(n)}\mathbf{w}_1 = 2\mu\mathcal{M}^{(n)}\mathbf{w}_1 + (\lambda + 2\mu)(-\kappa_p^2 V_{\kappa_p} + \kappa_s^2 V_{\kappa_s})\mathbf{n} \cdot \boldsymbol{\psi}.$$

For the last term, we first observe that $\mathbf{w}_0 \in H_{\text{loc}}^1(\overline{\Omega^\pm}; \mathbb{C}^3)$ and $\Delta \mathbf{w}_0 = -\kappa_s^2 \mathbf{w}_0$ in Ω^\pm since \mathbf{w}_0 is a combination of layer potentials of the Helmholtz equation corresponding to the wavenumber κ_s with respective densities $\boldsymbol{\psi} \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$ and $\mathcal{M}^{(\mathbf{n})}\boldsymbol{\psi} \in H^{-1/2}(\partial\Omega; \mathbb{C}^3)$. Next using (24), we can write

$$\nabla \cdot \mathbf{w}_0 = \Delta V_{\kappa_s} \mathbf{n} \cdot \boldsymbol{\psi} = -\kappa_s^2 V_{\kappa_s} \mathbf{n} \cdot \boldsymbol{\psi} \in H_{\text{loc}}^1(\overline{\Omega^\pm})$$

so that by Green's formula we readily get that $(T^{(\mathbf{n})}\mathbf{w}_0)^\pm$ can be expressed by (30) from (28) so arriving to

$$\begin{aligned} (T^{(\mathbf{n})}\mathbf{w}_0)^\pm &= \mu \mathcal{M}^{(\mathbf{n})}\mathbf{w}_0 + \mu (\partial_{\mathbf{n}} N_{\kappa_s} \boldsymbol{\psi})|_{\partial\Omega} + \mu \left(\partial_{\mathbf{n}} V_{\kappa_s} \mathcal{M}^{(\mathbf{n})}\boldsymbol{\psi} \right)^\pm \\ &\quad - \kappa_s^2 (\lambda + \mu) \mathbf{n} V_{\kappa_s} \mathbf{n} \cdot \boldsymbol{\psi}. \end{aligned}$$

It is enough to collect the above three terms to obtain (32). The rest of the proof is obtained from the jump and mapping properties of the layer potentials of the Helmholtz equation (cf. [4] or [21, p. 202]). ■

Remark 8 *Actually, representation formula (32) leads to an expression of $T^{(\mathbf{n})}K\boldsymbol{\psi}$ where the integrals are converging in the usual meaning, in other words with no need for Cauchy principal values or Hadamard finite parts to be defined. This property is provided by the fact that the term $\partial_{\mathbf{n}} N_{\kappa_s} \boldsymbol{\psi}$ can be represented in a variational form using Hamdi's regularization formula [13]*

$$\begin{aligned} \langle \boldsymbol{\varphi}, \partial_{\mathbf{n}} N_{\kappa_s} \boldsymbol{\psi} \rangle_{1/2, \partial\Omega} &= \sum_{j=1}^3 \langle \mathbf{n} \times \nabla \varphi_j, V_{\kappa_s} \mathbf{n} \times \nabla \psi_j \rangle_{1/2, \partial\Omega} \\ &\quad - \int_{\partial\Omega} \varphi_j \mathbf{n} \cdot V_{\kappa_s} (\psi_j \mathbf{n}) ds \end{aligned}$$

with $\boldsymbol{\varphi} \in H^{1/2}(\partial\Omega; \mathbb{C}^3)$, ψ_j and φ_j being the components of $\boldsymbol{\psi}$ and $\boldsymbol{\varphi}$ respectively (see [21, p. 289] for a comprehensive proof).

Remark 9 *Le Louër [19, Lemma 2.3] gave a second representation formula for $T^{(\mathbf{n})}K\boldsymbol{\psi}$*

$$\begin{aligned} T^{(\mathbf{n})}K\boldsymbol{\psi} &= \mu \nabla_{\partial\Omega} (V_{\kappa_s} \nabla_{\partial\Omega} \cdot \boldsymbol{\psi} \times \mathbf{n}) \times \mathbf{n} \\ &\quad + 2\mu \left(\mathcal{M}^{(\mathbf{n})} (N_{\kappa_s} \boldsymbol{\psi})^\pm - (\partial_{\mathbf{n}} V_{\kappa_s} \mathcal{M}^{(\mathbf{n})}\boldsymbol{\psi})^\pm \right) \\ &\quad + 2\mu \left(\mathcal{M}^{(\mathbf{n})} \nabla (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \boldsymbol{\psi} - \mathbf{n} \nabla \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathcal{M}^{(\mathbf{n})}\boldsymbol{\psi} \right) \\ &\quad + (4/\kappa_s^2) \mathcal{M}^{(\mathbf{n})} \nabla \nabla \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathcal{M}^{(\mathbf{n})}\boldsymbol{\psi} \\ &\quad - \omega^2 \varrho (\mathbf{n} \times V_{\kappa_s} (\boldsymbol{\psi} \times \mathbf{n}) + \mathbf{n} V_{\kappa_p} \mathbf{n} \cdot \boldsymbol{\psi}). \end{aligned} \tag{33}$$

The derivation of this author can similarly be adapted to deal with a Lipschitz geometry starting this once from representation formula (23) and using variational form (27) for the traction. The mapping properties of the related operator result as above from those of the layer potentials of the Helmholtz equation and of those of the tangential vector rotational and the surface rotational given in Corollary 1.

5 The two-dimensional case

We limit ourselves here to the case where both the geometry and the mechanical characteristics of the elastic medium are invariant to translations along the x_3 -axis. We first examine what happens to the Günter derivatives when applied to a function independent of the variable x_3 . We next use the relation linking the 2D and 3D Green kernels of the Helmholtz equation to express the two-dimensional elastic wave potentials similarly as above in \mathbb{R}^3 .

5.1 Two-dimensional Günter derivatives

In this part, we assume that the geometry is described as follows: $\Omega^\pm = \Omega_\perp^\pm \times (-\infty, +\infty)$ where Ω_\perp^\pm is a bounded 2D Lipschitz domain of the plane and $\Omega_\perp^- = \mathbb{R}^2 \setminus \overline{\Omega_\perp^+}$ is its complement. Any vector field \mathbf{u} , depending only on the transverse variable (x_1, x_2) , can be written as the superposition of a plane vector field \mathbf{u}_\perp and a scalar field u_3 , respectively called the plane and the anti-plane components of \mathbf{u} , according to the decomposition

$$\mathbf{u}(x_1, x_2) = \mathbf{u}_\perp(x_1, x_2) + u_3(x_1, x_2) \mathbf{e}_3.$$

Recall that $\{\mathbf{e}_j\}_{j=1}^3$ denotes the canonical basis of the space. The unit normal \mathbf{n} to $\partial\Omega$ is independent of x_3 , and verifies $n_3 = 0$. As a result, we do not distinguish between \mathbf{n} and its plane component \mathbf{n}_\perp . Subscript \perp is used to denote 2D analogs of 3D symbols. Let us just mention that $\nabla_\perp \times \mathbf{u}_\perp$ and $\nabla_\perp \times u_3$ are the scalar curl and the vector curl and are defined by

$$\nabla_\perp \times \mathbf{u}_\perp = \partial_{x_1} u_2 - \partial_{x_2} u_1, \quad \nabla_\perp \times u_3 = \partial_{x_2} u_3 \mathbf{e}_1 - \partial_{x_1} u_3 \mathbf{e}_2.$$

Let u be a function independent of x_3 . We readily get that

$$\mathcal{M}_{i3}^{(\mathbf{n})} u = n_3 \partial_{x_i} u - n_i \partial_{x_3} u = 0.$$

As a result, only two G unter derivatives are not zero

$$\mathcal{M}_{21}^{(\mathbf{n})} u = -\mathcal{M}_{12}^{(\mathbf{n})} u = n_1 \partial_{x_2} u - n_2 \partial_{x_1} u = \partial_\tau u$$

with $\tau = R_{\pi/2} \mathbf{n}$, R_θ being the counterclockwise rotation around the x_3 -axis by θ . In other words,

$$\mathcal{M}_{21}^{(\mathbf{n})} u = -\mathcal{M}_{12}^{(\mathbf{n})} u = \partial_s u$$

where s is the curvilinear abscissa of $\partial\Omega_\perp$ growing in the counterclockwise direction. The following version of Theorem 1 is more usual.

Theorem 8 *Under the above general assumptions, operator ∂_s is bounded from $H^s(\partial\Omega_\perp)$ into $H^{s-1}(\partial\Omega_\perp)$ for $0 \leq s \leq 1$.*

Remark 10 *G unter derivative matrix $\mathcal{M}^{(\mathbf{n})}$ reduces to an operator of a particularly simple form*

$$\mathcal{M}^{(\mathbf{n})} \mathbf{u} = \mathcal{M}_\perp^{(\mathbf{n})} \mathbf{u}_\perp = R_{\pi/2} \partial_s \mathbf{u}_\perp = \mathbf{e}_3 \times \partial_s \mathbf{u}.$$

5.2 Two-dimensional elastic waves layer potentials

Noting that

$$\mu \Delta \mathbf{u} + (\mu + \lambda) \nabla \nabla \cdot \mathbf{u} = \begin{bmatrix} \mu \Delta_\perp \mathbf{u}_\perp + (\mu + \lambda) \nabla_\perp \nabla_\perp \cdot \mathbf{u}_\perp \\ \mu \Delta_\perp u_3 \end{bmatrix},$$

we readily get that the plane \mathbf{u}_\perp and the antiplane u_3 components of \mathbf{u} are uncoupled at the level of the propagation equations.

Finally, the plane component $(T^{(\mathbf{n})} \mathbf{u})_\perp$ and the antiplane $(T^{(\mathbf{n})} \mathbf{u})_3$ one of the traction, corresponding to a field \mathbf{u} independent of x_3 , respectively depend on the plane displacement \mathbf{u}_\perp and the antiplane one u_3 only

$$\begin{aligned} (T^{(\mathbf{n})} \mathbf{u})_\perp &= T_\perp^{(\mathbf{n})} \mathbf{u}_\perp = 2\mu \partial_n \mathbf{u}_\perp + \mathbf{n} \nabla_\perp \cdot \mathbf{u}_\perp + \tau \nabla_\perp \times \mathbf{u}_\perp \\ (T^{(\mathbf{n})} \mathbf{u})_3 &= T_3^{(\mathbf{n})} u_3 = \mu \partial_n u_3, \end{aligned}$$

The expressions of the layer potentials and the related boundary integral operators can thus be obtained in a simple way using the following integral representation of the 2D fundamental solution of the Helmholtz equation

$$\frac{i}{4} H_0^{(1)}(\kappa r) = \int_{-\infty}^{+\infty} \frac{\exp\left(i\kappa \sqrt{r^2 + x_3^2}\right)}{4\pi \sqrt{r^2 + x_3^2}} dx_3 \text{ for } r > 0,$$

which is classically obtained by the change of variable $x_3 = \sinh t$ from the Mehler-Sonine integrals [26, Formulae 10.9.9]

$$\frac{i}{4} H_0^{(1)}(\kappa r) = \frac{1}{4\pi} \int_{-\infty}^{+\infty} \exp(i\kappa r \cosh t) dt.$$

For simplicity, we avoid to distinguish by subscript \perp the single-layer and the double-layer potentials related to the Helmholtz equation in 2D

$$V_\kappa p(x_1, x_2) = \int_{\partial\Omega_\perp} \frac{i}{4} H_0^{(1)}\left(\kappa \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}\right) p(y_1, y_2) ds_{(y_1, y_2)},$$

$$N_\kappa \varphi(x_1, x_2) =$$

$$- \int_{\partial\Omega_\perp} \frac{i}{4} \partial_{\mathbf{n}_{y_1, y_2}} H_0^{(1)} \left(\kappa \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2} \right) \varphi(y_1, y_2) ds_{(y_1, y_2)},$$

leaving the context to define whether it is the 2D case or the 3D one which is considered.

Each potential or boundary integral operator related to 2D elastic waves is decomposed in its plane and antiplane parts:

- Single-layer potential

$$\begin{aligned} S\mathbf{p} &= S_\perp \mathbf{p}_\perp + (S_3 p_3) \mathbf{e}_3, \\ S_\perp \mathbf{p}_\perp &= \frac{1}{\omega^2 \varrho} (\kappa_s^2 V_{\kappa_s} \mathbf{p}_\perp + \nabla_\perp \nabla_\perp \cdot (V_{\kappa_s} - V_{\kappa_p}) \mathbf{p}_\perp), \\ S_3 p_3 &= \frac{1}{\mu} V_{\kappa_s} p_3, \end{aligned}$$
- Double-layer potential

$$\begin{aligned} K\boldsymbol{\psi} &= K_\perp \boldsymbol{\psi}_\perp + (K_3 \psi_3) \mathbf{e}_3, \\ K_\perp \boldsymbol{\psi}_\perp &= N_{\kappa_s} \boldsymbol{\psi}_\perp \\ &\quad + (V_{\kappa_s} - 2\mu S_\perp) \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp + \nabla_\perp (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \boldsymbol{\psi}_\perp, \\ K_3 \psi_3 &= N_{\kappa_s} \psi_3, \end{aligned}$$
- Traction of the single-layer potential

$$\begin{aligned} T^{(\mathbf{n})} S\mathbf{p} &= T_\perp^{(\mathbf{n})} S_\perp \mathbf{p}_\perp + \left(T_3^{(\mathbf{n})} S_3 p_3 \right) \mathbf{e}_3, \\ T_\perp^{(\mathbf{n})} S_\perp \mathbf{p}_\perp &= (\partial_{\mathbf{n}} V_{\kappa_s} \mathbf{p})^\pm + \mathbf{n} \nabla_\perp \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathbf{p}_\perp \\ &\quad - \mathcal{M}_\perp^{(\mathbf{n})} (V_{\kappa_s} - 2\mu S_\perp) \mathbf{p}_\perp, \\ T_3^{(\mathbf{n})} S_3 p_3 &= (\partial_{\mathbf{n}} V_{\kappa_s} p_3)^\pm. \end{aligned}$$
- Traction of the double-layer potential

$$\begin{aligned} T^{(\mathbf{n})} K\boldsymbol{\psi} &= T_\perp^{(\mathbf{n})} K_\perp \boldsymbol{\psi}_\perp + \left(T_3^{(\mathbf{n})} K_3 \psi_3 \right) \mathbf{e}_3, \\ T_\perp^{(\mathbf{n})} K_\perp \boldsymbol{\psi}_\perp &= \\ &\quad \mu \left(\partial_{\mathbf{n}} N_{\kappa_s} \boldsymbol{\psi}_\perp + \mathcal{M}_\perp^{(\mathbf{n})} (N_{\kappa_s} \boldsymbol{\psi}_\perp)^\pm - \left(\partial_{\mathbf{n}} V_{\kappa_s} \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp \right)^\pm \right) \\ &\quad + 2\mu \left(\mathcal{M}_\perp^{(\mathbf{n})} \nabla_\perp (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \boldsymbol{\psi}_\perp - \mathbf{n} \nabla_\perp \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp \right) \\ &\quad + \left(\mathcal{M}_\perp^{(\mathbf{n})} (3\mu V_{\kappa_s} - 4\mu^2 S_\perp) \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi} - \omega^2 \varrho \mathbf{n} (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \boldsymbol{\psi}_\perp \right), \\ T_3^{(\mathbf{n})} K_3 \psi_3 &= \mu \partial_{\mathbf{n}} N_{\kappa_s} \psi_3. \end{aligned}$$

Remark 11 Another expression for the traction of the double-layer potential

$$\begin{aligned} T^{(\mathbf{n})} K\boldsymbol{\psi} &= T_\perp^{(\mathbf{n})} K_\perp \boldsymbol{\psi}_\perp + \left(T_3^{(\mathbf{n})} K_3 \psi_3 \right) \mathbf{e}_3, \\ T_\perp^{(\mathbf{n})} K_\perp \boldsymbol{\psi}_\perp &= 2\mu \left(\mathcal{M}_\perp^{(\mathbf{n})} (N_{\kappa_s} \boldsymbol{\psi}_\perp)^\pm - \left(\partial_{\mathbf{n}} V_{\kappa_s} \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp \right)^\pm \right) \\ &\quad + 2\mu \left(\mathcal{M}_\perp^{(\mathbf{n})} \nabla_\perp (V_{\kappa_p} - V_{\kappa_s}) \mathbf{n} \cdot \boldsymbol{\psi}_\perp - \mathbf{n} \nabla_\perp \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp \right) \\ &\quad + (4/\kappa_s^2) \mathcal{M}_\perp^{(\mathbf{n})} \nabla_\perp \nabla_\perp \cdot (V_{\kappa_p} - V_{\kappa_s}) \mathcal{M}_\perp^{(\mathbf{n})} \boldsymbol{\psi}_\perp \\ &\quad - \omega^2 \varrho (\boldsymbol{\tau} V_{\kappa_s} (\boldsymbol{\psi}_\perp \cdot \boldsymbol{\tau}) + \mathbf{n} V_{\kappa_p} \mathbf{n} \cdot \boldsymbol{\psi}_\perp), \\ T_3^{(\mathbf{n})} K_3 \psi_3 &= -\mu \partial_s V_{\kappa_s} \partial_s \psi_3 - \omega^2 \varrho \boldsymbol{\tau} \cdot V_{\kappa_s} (\psi_3 \boldsymbol{\tau}). \end{aligned}$$

can also be obtained from (33).

References

- [1] E. BECACHE, J.-C. NÉDÉLEC, AND N. NISHIMURA, *Regularization in 3d of anisotropic elastodynamic crack and obstacle problems*, Journal of Elasticity, 31 (1993), pp. 25–46.
- [2] A. BUFFA, M. COSTABEL, AND D. SHEEN, *On traces for $h(\text{curl}, \omega)$ in Lipschitz domains*, J. Math. Anal. Appl., 276 (2002), pp. 845–867.
- [3] Y. CHOQUET-BRUHAT AND C. DEWITT MORETTE, *Analysis, Manifolds and Physics - Part I: Basics*, Elsevier, London, New York, etc., 1991. 2nd revised edition.

- [4] M. COSTABEL, *Boundary integral operators on lipschitz domains: elementary results*, SIAM J. Math. Anal., 19 (1988), pp. 613–625.
- [5] ———, *A coercive bilinear form for Maxwell's equations*, Journal of Mathematical Analysis and Application, 157 (1991), pp. 527–541.
- [6] M. COSTABEL AND M. DAUGE, *A singularly perturbed mixed boundary value problem*, Comm. Partial Differential Equations, 21 (1996), p. 16671703.
- [7] ———, *Maxwell and Lamé eigenvalues on polyhedra*, Math. Methods Appl. Sci., 22 (1999), pp. 243–258.
- [8] M. DARBAS AND F. LE LOUËR, *Well-conditioned boundary integral formulations for high-frequency elastic scattering problems in three dimensions*, Mathematical Methods in the Applied Sciences, 38 (2015), pp. 1705–1733.
- [9] C. GEUZAINÉ AND J.-F. REMACLE, *Gmsh: a three-dimensional finite element mesh generator with built-in pre- and post-processing facilities*, International Journal for Numerical Methods in Engineering, 79 (2009), pp. 1309–1331.
- [10] L. J. GRAY AND S. J. CHANG, *Hypersingular integral formulation of elastic wave scattering*, Engineering Analysis with Boundary Elements, 10 (1992), pp. 337–343.
- [11] P. GRISVARD, *Elliptic Equations in Non-Smooth Domains.*, Monographs and Studies in Mathematics., Pitman, Toronto, 1982.
- [12] N. M. GUNTHER, *La théorie du potentiel et ses applications aux problèmes fondamentaux de la physique mathématique*, Gauthiers-Villars, Paris, 1934.
- [13] M. HAMDI, *Une formulation variationnelle par équations intégrales pour la résolution de l'équation d'Helmholtz avec des conditions aux limites mixtes*, C. R. Acad. Sci. Paris Sér II, 292 (1981), pp. 17–20.
- [14] H. HAN, *The boundary integro-differential equations of three-dimensional Neumann problem in linear elasticity*, Numer. Math., 68 (1994), pp. 269–281.
- [15] G. C. HSIAO AND W. L. WENDLAND, *Boundary Integral Equations*, Springer, Berlin-Heidelberg, 2008.
- [16] W. KNAUFF AND R. KRESS, *On the exterior boundary-value problem for the time-harmonic Maxwell equations*, Journal of Mathematical Analysis and Application, 72 (1979), pp. 215–235.
- [17] V. D. KUPRADZE, T. G. GEGELIA, M. O. BASHELEISHVILI, AND T. V. BURCHULADZE, *Three-Dimensional Problems of the Mathematical Theory of Elasticity and Thermoelasticity.*, North Holland, Amsterdam New-York Oxford, 1979.
- [18] F. LE LOUËR, *A high-order spectral algorithm for elastic obstacle scattering in three dimensions*. <hal-00700779v1>, 2012.
- [19] ———, *A high-order spectral algorithm for elastic obstacle scattering in three dimensions*, Journal of Computational Physics, 279 (2014), pp. 1–17.
- [20] Y. LIU AND F. J. RIZZO, *Hypersingular boundary integral equations for radiation and scattering of elastic waves in three dimensions*, Comput. Meth. Appl. Mech. Engrg., 107 (1993), pp. 131–144.
- [21] W. MCLEAN, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, Cambridge, UK, and New York, USA, 2000.
- [22] J. C. NÉDÉLEC, *The double layer potential for periodic elastic waves in \mathbf{R}^3* , in Boundary elements, D. Quinghua, ed., Pergamon Press, 1986, pp. 439–448.
- [23] J.-C. NÉDÉLEC, *Acoustic and Electromagnetic Equations: Integral Representations for Harmonic Problems*, Springer, Berlin, 2001.
- [24] J. NEČAS, *Direct Methods in the Theory of Elliptic Equations*, Springer, Heidelberg Dordrecht London New-York, 2012.
- [25] N. NISHUMURA AND S. KOBAYASHI, *A regularized boundary integral equation method for elastodynamic crack problems*, Computational Mechanics, 4 (1989), pp. 319–328.

- [26] F. W. OLVER, D. W. LOZIER, R. F. BOISVERT, AND C. W. CLARK, *NIST Handbook of Mathematical Functions*, NIST U.S. Department of Commerce and Cambridge University Press, New-York, 2010.
- [27] F. J. RIZZO, D. J. SHIPPY, AND M. REZAYAT, *A boundary integral equation method for radiation and scattering of elastic waves in three dimensions*, *International Journal for Numerical Methods in Engineering*, 21 (1985), pp. 115–129.
- [28] S. A. SAUTER AND C. SCHWAB, *Boundary Element Methods*, Springer-Verlag, Berlin-Heidelberg, 2011.
- [29] M. TAYLOR, *Partial Differential Equations I, Basic Theory*, Springer-Verlag, New York, 1996.