



**HAL**  
open science

# BOUNDARY LAYER ANALYSIS FOR THE FAST HORIZONTAL ROTATING FLUIDS

Wei-Xi Li, Van-Sang Ngo, Chao-Jiang Xu

► **To cite this version:**

Wei-Xi Li, Van-Sang Ngo, Chao-Jiang Xu. BOUNDARY LAYER ANALYSIS FOR THE FAST HORIZONTAL ROTATING FLUIDS. *Communications in Mathematical Sciences*, 2019, 17, pp.299-338. hal-01396779

**HAL Id: hal-01396779**

**<https://hal.science/hal-01396779>**

Submitted on 15 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# BOUNDARY LAYER ANALYSIS FOR THE FAST HORIZONTAL ROTATING FLUIDS

WEI-XI LI, VAN-SANG NGO AND CHAO-JIANG XU

ABSTRACT. It is well known that, for fast rotating fluids with the axis of rotation being perpendicular to the boundary, the boundary layer is of Ekman-type, described by a linear ODE system. In this paper we consider fast rotating fluids, with the axis of rotation being parallel to the boundary. We show that the corresponding boundary layer is describe by a nonlinear, degenerated PDE system which is similar to the 2-D Prandtl system. Finally, we prove the well-posedness of the governing system of the boundary layer in the space of analytic functions with respect to tangential variable.

## 1. INTRODUCTION

The incompressible Navier-Stokes equation coupled with a large Coriolis term reads

$$\begin{cases} \partial_t u^\varepsilon - \nu \Delta u^\varepsilon + u^\varepsilon \cdot \nabla u^\varepsilon + \frac{\omega \times u^\varepsilon}{\varepsilon} + \nabla p^\varepsilon = 0, \\ \operatorname{div} u^\varepsilon = 0, \\ u^\varepsilon|_{t=0} = u_0^\varepsilon, \end{cases}$$

with Dirichlet boundary condition, where  $\frac{\omega \times u^\varepsilon}{\varepsilon}$  stands for the Coriolis force and  $\omega$  is the rotation vector,  $\varepsilon^{-1}$  the rescaled speed of rotation,  $\nu$  the viscosity coefficients. The above system is sufficient to describe the rotation fluids which is a significant part of geophysics. Due to the earth's self-rotation, we can't neglect the Coriolis force in order to model the oceanography and meteorology dealing with large-scale magnitude. When the fluid is between a strip and the direction of rotation is not parallel to the boundary, we have the well-developed Ekman layers to match the interior flow with Dirichlet boundary condition, cf. [4, 5, 15, 24] and the references therein. The situation will be more complicated when the direction of rotation is parallel to the boundary, considering cylinder for instance and letting the fluid rotate around the vertical axis. Then we will have two types of boundaries, the horizontal boundary layer which is Ekman layers and the vertical boundary layers for which much less is known, despite various studies [5, 33, 35]. We refer to [5] for detailed discussions on the problem of vertical boundary layers.

In this paper, we consider the fast rotating viscous fluids where the the axe of rotation is horizontal with respect to the boundary. We prove that the governing equation for boundary layer is nonlinear PDE system which is similar to classical 2-D Prandtl boundary layer system, and we also obtain the well-posedness of this vertical boundary layers in the space of analytic functions.

As a preliminary step we first consider the half space case  $\mathbb{R}_+^3 = \mathbb{R}_h^2 \times \mathbb{R}_+$ . More precisely, we consider the following system

$$(N-S_\varepsilon) \quad \begin{cases} \partial_t u^\varepsilon - \nu \Delta u^\varepsilon + u^\varepsilon \cdot \nabla u^\varepsilon + \frac{e_2 \times u^\varepsilon}{\varepsilon} + \nabla p^\varepsilon = 0 & \text{in } \mathbb{R}_h^2 \times \mathbb{R}_+, \forall t \geq 0 \\ \operatorname{div} u^\varepsilon = 0 & \text{in } \mathbb{R}_h^2 \times \mathbb{R}_+, \forall t \geq 0 \\ u^\varepsilon|_{x_3=0} = 0 & \text{on } \mathbb{R}_h^2 \\ u^\varepsilon|_{t=0} = u_0^\varepsilon, & \text{in } \mathbb{R}_h^2 \times \mathbb{R}_+. \end{cases}$$

where  $e_2 = (0, 1, 0)$  is the unit horizontal vector,  $\nu > 0$  the coefficient of viscosity of fluids and  $\varepsilon$  the Rossby number. These equations describe the evolution of an incompressible three-dimensional viscous fluid in a rotating frame,  $\frac{e_2 \times u^\varepsilon}{\varepsilon}$  being the Coriolis force due to the rotation at high frequency  $\varepsilon^{-1}$ . According to the Taylor-Proudman theorem [34], the fast rotation penalize the movement of the fluid in the direction of the rotation axis. As a consequence, the fluid has tendency to move in columns, parallel to the rotation

*Date:* November 14, 2016.

*2000 Mathematics Subject Classification.* 35M13, 35Q30, 35Q35, 76U05.

*Key words and phrases.* Incompressible Navier Stokes equation, boundary layer, rotating fluids.

axis, which are widely known as the Taylor columns. This phenomenon is well-known in oceanography and meteorology, which is observed in many large-scale atmospheric and oceanic flows. In mathematical point of view, when  $\varepsilon$  goes to zero, the rotation term  $\frac{e_2 \times u^\varepsilon}{\varepsilon}$  becomes large and can only be balanced by the pressure. This means that, if  $u$  is the (formal) limit of  $u^\varepsilon$ , as  $\varepsilon \rightarrow 0$ , then  $e_2 \times u$  need to be a gradient term, which implies that  $u$  is independent of  $x_2$  (more explanations will be found in Section 2). In this paper, we will consider the case where the initial data are well prepared, *i.e.*  $u_0^\varepsilon$  do not depend on  $x_2$ .

When there is no Coriolis force, the zero-viscosity limit for the Navier-Stokes equations for incompressible fluids in a domain with boundary, with non-slip boundary conditions, is a challenging problem due to the formation of a boundary layer which is governed by the Prandtl equations ([29]). The mathematical analysis theory of Prandtl equation is also a challenging problem, see [1, 8, 9, 12, 26] and references therein. Far from the boundary, the inviscid limit problem was treated by several authors; we can refer, for instance, to Swann [32] and Kato [20]. In another work, Kato [19] gives some equivalent formulations of this problem in the case of bounded domains, showing that the convergence to the Euler system is equivalent to the fact that the  $L^2$  strength of the boundary layer goes to 0. Caffisch & Sammartino [31] solved the problem for analytic solutions on a half space by solving the Prandtl equations via abstract Cauchy-Kowaleskaya theorem. We also refer to [13, 16, 23] and the references therein for the recent progress on the inviscid limit of the Navier-Stokes equations when the initial vorticity is located away from the boundary. On the other hand, another commonly used boundary conditions are Navier-type slip boundary conditions, in which case the vanishing viscosity limit is rigorously justified; cf. [22, 36, 37, 38] and references therein.

We want to say a few words to compare the system (N-S $_\varepsilon$ ) with the case where the rotation axis is vertical with respect to the boundary (the rotation axis is in the direction of  $e_3 = (0, 0, 1)$  instead of  $e_2$ ). If the domain considered is between two parallel plates ( $\mathbb{T}^2 \times [0, 1]$  or  $\mathbb{R}^2 \times [0, 1]$ ), it was proved in Grenier & Masmoudi [15], Masmoudi [24, 25] and Chemin *et al.* [4] that for the rotating fluids with anisotropic viscosity  $-\nu \Delta_h - \varepsilon \partial_{x_3}^2$ , all the weak solutions of Navier-Stokes equation converge to the solution of the 2D Euler or 2D Navier-Stokes system (with damping term - effect of the Ekman pumping). The vertical rotation and the specific form of the domain (between two parallel plates) permit to explicitly construct the boundary layer velocity term from the interior velocity term (which satisfies a 2D damped Euler system), without using the Prandtl equations. We also want to mention the work of Dalibard and Gérard-Varet [7] in the case of fast rotating fluids on a rough domain with non-slip boundary conditions. The boundary layer is also proved to be of size  $\varepsilon$  (contrary to the case of Prandtl equations where the boundary layer is of size  $\sqrt{\varepsilon}$ ). We also refer to a series of work for the rotating fluids with anisotropic viscosity (see for exemple [2], [3], [10], [11], [14],[18], [27], [28]).

We want to emphasize that the formation of the boundary layers in the case of vertical rotation axis is due to the incompatibility of the Dirichlet boundary conditions with the columnar movement of the limit fluid (as  $\varepsilon \rightarrow 0$ ). Indeed, as the rotation axis is  $e_3$ , the limiting velocity of the fluid is independent of  $x_3$ , and so, the Dirichlet boundary conditions imply that the limit velocity should be zero. This incompatibility leads to the fact that a thin layer (Ekman's layer) is formed near the boundary, and the fluid's evolution is violent in this small scale zone, in a way that stops the fluid on the boundary.

In the case of horizontal rotation axis (in the direction of  $e_2$ ), the incompatibility of boundary conditions will be more complicated, because of the fact that the limit velocity is independent of  $x_2$  instead of  $x_3$ . In Section 2, we prove that the limit system is a 2D Euler-like system. This means that we are no longer in the case considered by Ekman. The techniques of [15] and [4] do not work and we can not explicitly calculate the boundary layer. The fast rotation only penalizes the fluid motion in the  $x_2$  direction, and leads to a problem very close to the inviscid limit of two-dimensional Navier-Stokes system. It is then relevant to look for a boundary layer of size  $\sqrt{\varepsilon}$  and we will show in Section 2 that in this boundary layer of size  $\sqrt{\varepsilon}$ , the fluid velocity actually satisfies a two-dimensional Prandtl-like system. Finally, we remark that in this paper, we only consider the case where  $\nu = \varepsilon$ . Indeed, as explained in [15] and also in [5], if the ratio  $\nu/\varepsilon$  goes to infinity, the fluid rapidly stops after a few evolutions. It is then more interesting to consider the case where  $\nu \lesssim \varepsilon$ , which moreover better fits physical observations.

In this work, we study the formation of the boundary layer when  $\nu = \varepsilon \rightarrow 0$ . We suppose the existence of a boundary layer of size  $\sqrt{\varepsilon}$  near the boundary  $\{x_3 = 0\}$  of  $\mathbb{R}_+^3$ . We will derive the limit equation and the boundary layer equation by using a formal asymptotic expansion in the Section 2. We refer to the book of Pedlovsky [30] for more detail about this formal expansion. To this end, we suppose that the solution of

(N-S $_{\varepsilon}$ ) accepts the following asymptotic expansion

$$(1.1) \quad u^{\varepsilon}(t, x_1, x_2, x_3) = \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} \left[ u^{I,j}(t, x_1, x_2, x_3) + u^{B,j} \left( t, x_1, x_2, \frac{x_3}{\sqrt{\varepsilon}} \right) \right] + \dots,$$

$$(1.2) \quad p^{\varepsilon}(t, x_1, x_2, x_3) = \sum_{j=-2}^1 \varepsilon^{\frac{j}{2}} p^{I,j}(t, x_1, x_2, x_3) + \sum_{j=-2}^0 \varepsilon^{\frac{j}{2}} p^{B,j} \left( t, x_1, x_2, \frac{x_3}{\sqrt{\varepsilon}} \right) + \dots,$$

where  $u^{B,j}(t, x_1, x_2, y)$  and  $p^{B,j}(t, x_1, x_2, y)$  exponentially go to zero as  $y \stackrel{\text{def}}{=} \frac{x_3}{\sqrt{\varepsilon}} \rightarrow +\infty$ . The remaining terms is supposed to be very small (at least of order 3).

Throughout this paper, we will always use  $\partial_t$ ,  $\partial_i$  (or  $\partial_{x_i}$ ),  $i = 1, 2, 3$ , and  $\partial_y$  to respectively denote the derivatives with respect to the time variable  $t$ , the space variables  $x_i$ ,  $i = 1, 2, 3$ , and the boundary layer variable  $y = \frac{x_3}{\sqrt{\varepsilon}}$ . Using the above asymptotic expansion, we first deduce that the behavior of the fluid near the boundary is governed by the following 2D Prandtl-like equation

$$(P1) \quad \begin{cases} \partial_t \mathcal{U}_1^{p,0} - \partial_y^2 \mathcal{U}_1^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_1^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0} + \partial_1 p^{B,0} + \overline{\partial_1 p^{I,0}} = 0, \\ \partial_t \mathcal{U}_3^{p,1} - \partial_y^2 \mathcal{U}_3^{p,1} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_3^{p,1} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_3^{p,1} + \overline{\partial_3 p^{I,1}} + y \overline{\partial_3^2 p^{I,0}} = 0, \\ \partial_1 \mathcal{U}_1^{p,0} + \partial_y \mathcal{U}_3^{p,1} = 0, \\ \mathcal{U}_1^{p,0}|_{y=0} = 0, \quad \lim_{y \rightarrow +\infty} \mathcal{U}_1^{p,0}(t, x_1, y) = \overline{u_1^{I,0}}, \\ \mathcal{U}_3^{p,1}|_{y=0} = 0, \quad \partial_y \mathcal{U}_3^{p,1}|_{y=0} = 0, \\ (\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1})|_{t=0} = (\mathcal{U}_{1,0}^{p,0}, \mathcal{U}_{3,0}^{p,1}), \end{cases}$$

with the unknown functions  $(\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1}, p^{B,0})$ , and the horizontal second component satisfies a parabolic type equation

$$(P2) \quad \begin{cases} \partial_t \mathcal{U}_2^{p,0} - \partial_y^2 \mathcal{U}_2^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_2^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_2^{p,0} = 0 \\ \mathcal{U}_2^{p,0}|_{y=0} = 0, \quad \lim_{y \rightarrow +\infty} \mathcal{U}_2^{p,0}(t, x_1, y) = \overline{u_2^{I,0}} \\ \mathcal{U}_2^{p,0}|_{t=0} = \mathcal{U}_{2,0}^{p,0}. \end{cases}$$

Here

$$\partial_2 \mathcal{U}_1^{p,0} = \partial_2 \mathcal{U}_2^{p,0} = \partial_2 \mathcal{U}_3^{p,1} = 0.$$

Here, we emphasize the ‘‘Prandtl-like’’ property of our system by using the new unknown functions

$$\begin{aligned} \mathcal{U}_j^{p,0} &= u_j^{B,0} + \overline{u_j^{I,0}}, \quad j = 1, 2 \\ \mathcal{U}_3^{p,1} &= u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \end{aligned}$$

where  $\overline{u^{I,j}}, \overline{p^{I,j}}$ ,  $j = 1, 2$  are the values on the boundary of the tangential velocity and pressure of the outflow satisfying the Bernoulli-type law

$$\begin{cases} \partial_t \overline{u_1^{I,0}} + \overline{u_1^{I,0}} \partial_1 \overline{u_1^{I,0}} + \overline{\partial_1 p^{I,0}} = 0 \\ \partial_t \overline{u_2^{I,0}} + \overline{u_1^{I,0}} \partial_1 \overline{u_2^{I,0}} + \overline{\partial_2 p^{I,0}} = 0 \\ \partial_t \overline{u_3^{I,1}} + \overline{u_1^{I,0}} \partial_1 \overline{u_3^{I,1}} + \overline{u_3^{I,1}} \partial_3 \overline{u_3^{I,0}} + \overline{\partial_3 p^{I,1}} = 0 \end{cases}$$

which is the restriction of the Euler system and linearized Euler system on the boundary  $x_3 = 0$ , so that they depend only on the variables  $(t, x_1)$ . More precise description will be found in Section 2.

Note that the boundary layer equation (P1) look very close to that of classical 2D Prandtl equation, but the fast rotating produces the boundary layer pressure for the first components, so that the boundary layer equation (P1) is now really a system of 3 equations with both the velocity  $(\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1})$  and the boundary pressure  $p^{B,0}$  to determined. We remark that on one side, the first equation in (P1) admits the similar structure of Prandtl equation, i.e., the degeneracy in  $x_1$  coupled with the nonlocal property arising from the term  $\mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0}$ , so that the system (P1) is quite similar to Prandtl equation. Therefore we can only expect the local well-posedness for analytic initial data if no additional assumptions are imposed. On the

other hand, there is a crucial difference between Prandtl equation and the first equation in (P1), due to the unknown pressure  $p^{B,0}$ . Recall the pressure term in Prandtl equation is from outflow and can be defined by the Bernoulli law, so that the pressure therein is a given function and therefore Prandtl equation is a kind of degenerate parabolic equation. But here the situation is quite complicated since we have the unknown pressure  $p^{B,0}$  in (P1), which arises because of the fast rotation parallel to the boundary, and can't be defined by the Bernoulli law anymore. So the classical theory for Prandtl equation is not applicable directly to our case and moreover we can't follow the same strategy as in Prandtl equation to treat the the first equation in (P1). To overcome the difficulty due to the unknown pressure term in the first equation of (P1), we will firstly solve the second equation for  $U_3^{p,1}$ , and then use the divergence-free property to find  $U_1^{p,0}$  (see Section 3 for detail). Finally we mention that the mathematical justification of the inviscid limit for solutions to (N-S $_\varepsilon$ ), is also complicated as classical Prandtl boundary layer theory. We only concentrate in this work on the well-posedness of boundary layer and will investigate this inviscid limit problem in the future work.

On the other hand, we will prove in Section 2 that the limiting velocity of the outer flow satisfies a classical 2D Euler-type equation, which is,

$$(1.3) \quad \begin{cases} \partial_t u_1^{I,0} + u_1^{I,0} \partial_1 u_1^{I,0} + u_3^{I,0} \partial_3 u_1^{I,0} + \partial_1 p^{I,0} = 0 \\ \partial_t u_2^{I,0} + u_1^{I,0} \partial_1 u_2^{I,0} + u_3^{I,0} \partial_3 u_2^{I,0} = 0 \\ \partial_t u_3^{I,0} + u_1^{I,0} \partial_1 u_3^{I,0} + u_3^{I,0} \partial_3 u_3^{I,0} + \partial_3 p^{I,0} = 0 \\ \partial_2 u_1^{I,0} = \partial_2 u_2^{I,0} = \partial_2 u_3^{I,0} = \partial_2 p^{I,0} = 0 \\ \partial_1 u_1^{I,0} + \partial_3 u_3^{I,0} = 0 \\ u_3^{I,0}|_{x_3=0} = 0 \\ u^{I,0}|_{t=0} = u_0^{I,0}(x_1, x_3). \end{cases}$$

In the system (1.3), the components  $(u_1^{I,0}, u_3^{I,0}, p^{I,0})$  satisfy exactly a 2-D incompressible Euler equation on the half-plane, so that the existence and regularity in Gevery class of local in time solution is well know, (see Vicol [21] and references therein), but in the study of boundary layer equation, we need some weighted on the tangential variables, we cite in particular the results of [6].

**Definition 1.1.** Let  $\frac{1}{2} < \ell \leq 1$  be given. We denote by  $\mathcal{A}_\tau$  the space of analytic functions with analytic radius  $\tau > 0$ , which is consist of all functions  $f \in L^2(\mathbb{R}_+^2)$  such that

$$\|f\|_{\mathcal{A}_\tau} \stackrel{\text{def}}{=} \sup_{|\alpha| \geq 0} \frac{\tau^{|\alpha|}}{|\alpha|!} \|\langle z \rangle^\ell \partial_z^\alpha f\|_{L^2(\mathbb{R}_+^2)} < +\infty.$$

**Theorem 1.2** ([6]). Suppose that the initial data  $u_0^{I,0} = (u_{1,0}^{I,0}, u_{2,0}^{I,0}, u_{3,0}^{I,0})$  in (1.3) satisfies

$$u_{1,0}^{I,0}, u_{2,0}^{I,0}, u_{3,0}^{I,0} \in \mathcal{A}_{\tau_0}$$

for some  $\tau_0 > 0$ , the divergence-free condition and the compatibility condition. Then Euler-type system (1.3) admits a unique solution  $(u_1^{I,0}, u_2^{I,0}, u_3^{I,0}) \in L^\infty([0, T]; \mathcal{A}_\tau)$  for some  $T > 0$  and  $\tau > 0$ .

The construction of the components  $(u_1^{I,0}, u_3^{I,0}, p^{I,0})$  is given in [6]. The construction of  $u_2^{I,0}$  is standard, using the classical theory of transport equation.

Now we list several estimates, which are just immediate consequences of the definition of  $\|\cdot\|_{\mathcal{A}_\tau}$  and Sobolev inequalities. For  $u_1^{I,0} \in L^\infty([0, T]; \mathcal{A}_\tau)$ , we have, for all  $p, q \geq 0$ ,

$$(1.4) \quad \|\langle x_1 \rangle^\ell \partial_1^p \partial_3^q u_3^{I,0}(x_1, x_3)\|_{L^\infty(\mathbb{R}_+; L^2(\mathbb{R}_{x_1}))} \leq C \|u_3^{I,0}\|_{\mathcal{A}_\tau} \frac{(p+q+3)!}{\tau^{p+q+3}}.$$

Using the equation

$$\partial_t u_1^{I,0} + u_1^{I,0} \partial_1 u_1^{I,0} + u_3^{I,0} \partial_3 u_1^{I,0} + \partial_1 p^{I,0} = 0,$$

we can calculate, by virtue of Leibniz formula,

$$(1.5) \quad \begin{aligned} & \|\langle x_1 \rangle^\ell \partial_t \partial_1^p \partial_3^q u_3^{I,0}\|_{L^\infty(\mathbb{R}_+; L^2(\mathbb{R}_{x_1}))} \\ & \leq C_\tau \left( \|u_1^{I,0}\|_{\mathcal{A}_\tau}^2 + \|u_1^{I,0}\|_{\mathcal{A}_\tau} \|u_3^{I,0}\|_{\mathcal{A}_\tau} + \|p^{I,0}\|_{\mathcal{A}_\tau} \right) \frac{2^{p+q}(p+q)!}{\tau^{p+q}}. \end{aligned}$$

In order to completely give the solutions of the systems (P1) and (P2), we also need the following linearized Euler system, which describes the evolution of the fluids in the interior part of the domain, far from the boundary, at the order  $\sqrt{\varepsilon}$ .

$$(1.6) \quad \begin{cases} \partial_t u_1^{I,1} + u_1^{I,0} \partial_1 u_1^{I,1} + u_3^{I,0} \partial_3 u_1^{I,1} + u_1^{I,1} \partial_1 u_1^{I,0} + u_3^{I,1} \partial_3 u_1^{I,0} + \partial_1 p^{I,1} = 0 \\ \partial_t u_2^{I,1} + u_1^{I,0} \partial_1 u_2^{I,1} + u_3^{I,0} \partial_3 u_2^{I,1} + u_1^{I,1} \partial_1 u_2^{I,0} + u_3^{I,1} \partial_3 u_2^{I,0} = 0 \\ \partial_t u_3^{I,1} + u_1^{I,0} \partial_1 u_3^{I,1} + u_3^{I,0} \partial_3 u_3^{I,1} + u_1^{I,1} \partial_1 u_3^{I,0} + u_3^{I,1} \partial_3 u_3^{I,0} + \partial_3 p^{I,1} = 0 \\ \partial_2 u_1^{I,1} = \partial_2 u_2^{I,1} = \partial_2 u_3^{I,1} = \partial_2 p^{I,1} = 0 \\ \partial_1 u_1^{I,1} + \partial_3 u_3^{I,1} = 0 \\ u_3^{I,1}|_{x_3=0} = -u_3^{B,1}(t, x_1, 0) \\ u^{I,1}|_{t=0} = u_0^{I,1}(x_1, x_3). \end{cases}$$

For this linearized Euler system (1.6), we have

**Theorem 1.3.** *Let  $\ell > 1/2$ ,  $\tau_0 > 0$  and  $u_3^{B,1}(t, x_1, 0)$  a given function such that*

$$\sum_{m \leq 2} \|\langle x_1 \rangle^\ell \partial_1^m u_3^{B,1}(t, x_1, 0)\|_{L^2(\mathbb{R}_{x_1})}^2 + \sum_{m \geq 3} \left[ \frac{\tau_0^{m-1}}{(m-3)!} \right]^2 \|\langle x_1 \rangle^\ell \partial_1^m u_3^{B,1}(t, x_1, 0)\|_{L^2(\mathbb{R}_{x_1})}^2 < +\infty.$$

Suppose that the initial data  $u_0^{I,1} = (u_{1,0}^{I,1}, u_{2,0}^{I,1}, u_{3,0}^{I,1})$  in (1.6) satisfies the divergence-free condition, the compatibility condition and

$$u_{1,0}^{I,1}, u_{2,0}^{I,1}, u_{3,0}^{I,1} \in \mathcal{A}_{\tau_0}.$$

Then the linearized Euler system (1.6) admits a unique solution  $(u_1^{I,1}, u_2^{I,1}, u_3^{I,1}) \in L^\infty([0, T]; \mathcal{A}_\tau)$  for some  $T > 0$  and  $\tau > 0$ .

We remark that, the compatibility condition ask

$$u_{3,0}^{I,1}(x_1, 0) = -u_{3,0}^{B,1}(x_1, 0).$$

It is exactly the non-slip condition of (N-S $_\varepsilon$ ) at order 1. Because of its linearity, treating the system (1.6) is still much easier than treating the system (1.3), even with the presence of the given boundary function  $u_3^{B,1}(t, x_1, 0)$ . So, to prove Theorem 1.3, we can simply follow the lines of the proof of Theorem 1.2 as in [6].

Before giving the well-posedness results on (P1) and (P2), we need the following weighted analytic function spaces in tangential variable. We also remark that there is no coupling between  $(\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1})$  and  $\mathcal{U}_2^{p,0}$ . Then, the strategy consists in separately solving the systems (P1) and (P2).

**Definition 1.4.** *Let  $1/2 < \ell \leq 1$  be given throughout the paper. With each pair  $(\rho, a)$  with  $\rho > 0$  and  $a > 0$  we associate a space  $X_{\rho,a}$  of all functions  $u(x_1, y) \in H^\infty(\mathbb{R}_{x_1}; H^2(\mathbb{R}_+))$  such that*

$$\sum_{m \leq 2} \left( \sum_{0 \leq j \leq 1} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}^2 \right) + \sum_{m \geq 3} \left( \sum_{0 \leq j \leq 1} \left[ \frac{\rho^{m-1}}{(m-3)!} \right]^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}^2 \right) < +\infty,$$

where we use the convention  $0! = 1$ . We endow  $X_{\rho,a}$  with the norm

$$|u|_{X_{\rho,a}}^2 = \sum_{m \leq 2} \left( \sum_{0 \leq j \leq 1} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}^2 \right) + \sum_{m \geq 3} \left( \sum_{0 \leq j \leq 1} \left[ \frac{\rho^{m-1}}{(m-3)!} \right]^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}^2 \right).$$

The well-posedness of the system (P1) can be stated as follows.

**Theorem 1.5.** *Suppose that the initial data*

$$\mathcal{U}_{3,0}^{p,1} = u_{3,0}^{B,1} + \overline{u_{3,0}^{I,1}} + \overline{y \partial_3 u_{3,0}^{I,0}}$$

in (P1) satisfies that

$$u_{3,0}^{B,1} \in X_{\rho_0, a_0}, \quad u_{3,0}^{I,1}, u_{3,0}^{I,0} \in \mathcal{A}_{\tau_0}$$

for some  $a_0 > 0$ ,  $\rho_0 > 0$  and  $\tau_0 > 0$  and

$$\mathcal{U}_{1,0}^{p,0}(x_1, y) = - \int_{-\infty}^{x_1} \partial_y u_{3,0}^{B,1}(z, y) dz + \overline{u_{1,0}^{I,0}}(x_1).$$

Then there exist  $T > 0$ ,  $\tau > 0$  and a pair  $(\rho, a)$  with  $\rho, a > 0$ , such that the system (P1) admits a unique solution  $(\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1}, \partial_1 p^{B,0})$ , and moreover

$$\begin{aligned} \mathcal{U}_3^{p,1} &= u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \\ \mathcal{U}_1^{p,0}(t, x_1, y) &= - \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, y) dz + \overline{u_1^{I,0}}(t, x_1), \end{aligned}$$

with  $u_3^{B,1} \in L^\infty([0, T]; X_{\rho, a})$  and  $u_1^{I,0}, u_3^{I,0}, u_3^{I,1} \in L^\infty([0, T]; \mathcal{A}_\tau)$ .

**Remark 1.6.** (i) Here we consider the well prepared initial data, that is the initial data are independent of  $x_2$ .

(ii) We want to remark that once we find  $\mathcal{U}_3^{p,1}$ , we can obtain  $\mathcal{U}_1^{p,0}$  using the divergence-free property in the third equation of the system (P1).

Let  $\mathcal{U}_1^{p,0}, \mathcal{U}_3^{p,1}$  be the solutions to the system (P1) given by the theorem above. Then we see (P2) is a linear parabolic equation, and we have the following theorem concerned with its well-posedness.

**Theorem 1.7.** Let  $\rho_0 > 0$ ,  $a_0 > 0$ ,  $\tau_0 > 0$  be given. For any initial data

$$\mathcal{U}_{2,0}^{p,0} = u_{2,0}^{B,0} + \overline{u_{2,0}^{I,0}}$$

where  $u_{2,0}^{B,0} \in X_{\rho_0, a_0}$  and  $u_{2,0}^{I,0} \in \mathcal{A}_{\tau_0}$ , there exist  $T > 0$ ,  $0 < \tau < \tau_0$  and  $0 < a < a_0$ , such that the equation (P2) admits a unique solution  $\mathcal{U}_2^{p,0}$  satisfying  $\mathcal{U}_2^{p,0} = u_2^{B,0} + \overline{u_2^{I,0}}$  with

$$u_2^{B,0} \in L^\infty([0, T], X_{\rho_0, a}), \quad u_2^{I,0} \in L^\infty([0, T], \mathcal{A}_\tau).$$

By the two theorems above we obtain the well-posedness for the boundary layer equation of the system (N-S $_\varepsilon$ ) in the frame of analytic space in tangential variable.

The paper is organized as follows. In section 2, we formally derive the governing equations of the outer flow inside the domain and the systems (P1) and (P2) which describe the fluid motion inside the boundary layer. The sections 3-4 are devoted to prove the well-posedness of the system (P1). Finally, we give some brief ideas of the proof of Theorem 1.7 for the well-posedness of equation (P2) in the section 5.

## 2. FORMAL ASYMPTOTIC EXPANSION

First of all, we want to give a few words to explain our special choice of the order of the expansions of the velocity and the pressure. Indeed, we remark that as for the formulation of Prandtl boundary layer equations, we are only interested in the leading orders which are necessary to allow us to formally obtain the governing equations of the evolution of the boundary layer. By using the asymptotic expansions (1.1) and (1.2), we have the following asymptotic identities for the leading terms up to order  $\varepsilon^{1/2}$  and all the

remaining terms are of higher order in  $\varepsilon$ .

$$(2.1) \quad \left\{ \begin{array}{l} \partial_t u^\varepsilon = \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} (\partial_t u^{I,j} + \partial_t u^{B,j}) + \dots \\ -\varepsilon \Delta u^\varepsilon = -\partial_y^2 u^{B,0} - \varepsilon^{\frac{1}{2}} \partial_y^2 u^{B,1} - \sum_{j=0}^1 \varepsilon^{1+\frac{j}{2}} (\Delta u^{I,j} + \Delta_h u^{B,j}) + \dots \\ u^\varepsilon \cdot \nabla u^\varepsilon = \sum_{j=0}^1 \varepsilon^{\frac{j-1}{2}} \left[ \sum_{k=0}^j (u_3^{B,k} + u_3^{I,k}) \partial_y u^{B,j-k} \right] + \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} \left[ \sum_{k=0}^j (u_h^{B,k} + u_h^{I,k}) \cdot \nabla_h u^{B,j-k} \right] \\ \quad + \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} \left[ \sum_{k=0}^j (u^{B,k} + u^{I,k}) \cdot \nabla u^{I,j-k} \right] + \dots \\ \frac{e_2 \times u^\varepsilon}{\varepsilon} = \sum_{j=0}^1 \varepsilon^{\frac{j}{2}-1} \left[ \begin{pmatrix} u_3^{B,j} \\ 0 \\ -u_1^{B,j} \end{pmatrix} + \begin{pmatrix} u_3^{I,j} \\ 0 \\ -u_1^{I,j} \end{pmatrix} \right] + \dots \\ \nabla p^\varepsilon = \varepsilon^{-\frac{3}{2}} \begin{pmatrix} 0 \\ 0 \\ \partial_y p^{B,-1} \end{pmatrix} + \sum_{j=-2}^{-1} \varepsilon^j \begin{pmatrix} \partial_1 p^{B,j} \\ \partial_2 p^{B,j} \\ \partial_y p^{B,j+1} \end{pmatrix} + \begin{pmatrix} \partial_1 p^{B,0} \\ \partial_2 p^{B,0} \\ 0 \end{pmatrix} + \sum_{j=-2}^1 \varepsilon^{\frac{j}{2}} \nabla p^{I,j} + \dots \end{array} \right.$$

**2.1. Formal derivation of the fluid behavior far from the boundary.** We put all the asymptotic identities (2.1) into the system (N-S $_\varepsilon$ ) and we deduce that

$$(2.2) \quad \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} \partial_t u^{I,j} - \sum_{j=0}^1 \varepsilon^{1+\frac{j}{2}} \Delta u^{I,j} + \sum_{j=0}^1 \varepsilon^{\frac{j}{2}} \sum_{k=0}^j u^{I,k} \cdot \nabla u^{I,j-k} + \sum_{j=0}^1 \varepsilon^{\frac{j}{2}-1} \begin{pmatrix} u_3^{I,j} \\ 0 \\ -u_1^{I,j} \end{pmatrix} + \sum_{j=-2}^1 \varepsilon^{\frac{j}{2}} \nabla p^{I,j} = 0(\varepsilon).$$

Taking the limit  $y = \frac{x_3}{\sqrt{\varepsilon}} \rightarrow +\infty$  ( $\varepsilon \rightarrow 0$ ), the divergence-free property writes

$$(2.3) \quad \operatorname{div} u^{I,j} = 0, \quad \forall j \geq 0.$$

At the leading term of  $\varepsilon^{-1}$  in (2.2), we simply have

$$(2.4) \quad \begin{pmatrix} u_3^{I,0} \\ 0 \\ -u_1^{I,0} \end{pmatrix} + \begin{pmatrix} \partial_1 p^{I,-2} \\ \partial_2 p^{I,-2} \\ \partial_3 p^{I,-2} \end{pmatrix} = 0.$$

Then, classical calculations (see Grenier-Masmoudi [15] or Chemin *et al.* [5]) give

$$(2.5) \quad \partial_2 p^{I,-2} = \partial_2 u_1^{I,0} = \partial_2 u_2^{I,0} = \partial_2 u_3^{I,0} = 0.$$

At the order  $\varepsilon^{-1/2}$  in (2.2), we have

$$(2.6) \quad \begin{pmatrix} u_3^{I,1} \\ 0 \\ -u_1^{I,1} \end{pmatrix} + \begin{pmatrix} \partial_1 p^{I,-1} \\ \partial_2 p^{I,-1} \\ \partial_3 p^{I,-1} \end{pmatrix} = 0,$$

which imply

$$(2.7) \quad \partial_2 p^{I,-1} = \partial_2 u_1^{I,1} = \partial_2 u_2^{I,1} = \partial_2 u_3^{I,1} = 0.$$

**Remark 2.1.** Identities (2.5) and (2.7) mean that the limit behaviour of the outer flow is two-dimensional, as predicts the Taylor-Proudman theorem.



**At the order**  $\varepsilon^0$  in (2.2), taking into account (2.5) and the divergence-free condition (2.3), we obtain

$$(2.8) \quad \begin{cases} \partial_t u_1^{I,0} + u_1^{I,0} \partial_1 u_1^{I,0} + u_3^{I,0} \partial_3 u_1^{I,0} + \partial_1 p^{I,0} = 0 \\ \partial_t u_2^{I,0} + u_1^{I,0} \partial_1 u_2^{I,0} + u_3^{I,0} \partial_3 u_2^{I,0} + \partial_2 p^{I,0} = 0 \\ \partial_t u_3^{I,0} + u_1^{I,0} \partial_1 u_3^{I,0} + u_3^{I,0} \partial_3 u_3^{I,0} + \partial_3 p^{I,0} = 0 \\ \partial_2 u_1^{I,0} = \partial_2 u_2^{I,0} = \partial_2 u_3^{I,0} = 0 \\ \partial_1 u_1^{I,0} + \partial_3 u_3^{I,0} = 0 \end{cases}$$

Now, by applying  $\partial_2$  to the second equation of the system (2.8), we obtain

$$\partial_2^2 p^{I,0} = 0,$$

which means that there exist  $g_1(x_1, x_3)$  and  $g_2(x_1, x_3)$  such that

$$p^{I,0} = x_2 g_1 + g_2.$$

Now, differentiating the first and third equations of (2.8) with respect to  $x_2$ , we obtain

$$\partial_1 g_1 = \partial_3 g_1 = 0.$$

By taking  $|x| \rightarrow +\infty$  in the second equation of (2.8), we conclude that  $g_1 \equiv 0$ . Thus, the system (2.8) becomes the following 2D Euler-type system with three components in the half-plane, which is the formal limiting system of (N-S $_\varepsilon$ ) far from the boundary as  $\varepsilon \rightarrow 0$

$$\begin{cases} \partial_t u_1^{I,0} + u_1^{I,0} \partial_1 u_1^{I,0} + u_3^{I,0} \partial_3 u_1^{I,0} + \partial_1 p^{I,0} = 0 \\ \partial_t u_2^{I,0} + u_1^{I,0} \partial_1 u_2^{I,0} + u_3^{I,0} \partial_3 u_2^{I,0} = 0 \\ \partial_t u_3^{I,0} + u_1^{I,0} \partial_1 u_3^{I,0} + u_3^{I,0} \partial_3 u_3^{I,0} + \partial_3 p^{I,0} = 0 \\ \partial_2 u_1^{I,0} = \partial_2 u_2^{I,0} = \partial_2 u_3^{I,0} = \partial_2 p^{I,0} = 0 \\ \partial_1 u_1^{I,0} + \partial_3 u_3^{I,0} = 0 \\ u_3^{I,0}|_{x_3=0} = 0. \end{cases}$$

Since this system is independent of  $x_2$ , for the compatibility, we need to impose the well prepared initial data, which means that

$$u^{I,0}(0, x_1, x_3) = u_0^{I,0}(x_1, x_3).$$

The boundary condition will be discussed in (2.17).

The system (1.3) will be completed with a boundary condition for the second component  $u_2^{I,0}$ . In fact, the trace function  $\overline{u_2^{I,0}}(t, x_1)$  on the boundary  $\{x_3 = 0\}$  satisfies the following system

$$\begin{cases} \partial_t \overline{u_2^{I,0}} + u_1^{I,0} \partial_1 \overline{u_2^{I,0}} = 0 \\ \overline{u_2^{I,0}}(0, x_1) = u_{0,2}^{I,0}(x_1, 0). \end{cases}$$

**At the order**  $\varepsilon^{1/2}$  in (2.2), using (2.7) and the divergence-free condition (2.3), we obtain the system

$$\begin{cases} \partial_t u_1^{I,1} + u_1^{I,0} \partial_1 u_1^{I,1} + u_3^{I,0} \partial_3 u_1^{I,1} + u_1^{I,1} \partial_1 u_1^{I,0} + u_3^{I,1} \partial_3 u_1^{I,0} + \partial_1 p^{I,1} = 0 \\ \partial_t u_2^{I,1} + u_1^{I,0} \partial_1 u_2^{I,1} + u_3^{I,0} \partial_3 u_2^{I,1} + u_1^{I,1} \partial_1 u_2^{I,0} + u_3^{I,1} \partial_3 u_2^{I,0} + \partial_2 p^{I,1} = 0 \\ \partial_t u_3^{I,1} + u_1^{I,0} \partial_1 u_3^{I,1} + u_3^{I,0} \partial_3 u_3^{I,1} + u_1^{I,1} \partial_1 u_3^{I,0} + u_3^{I,1} \partial_3 u_3^{I,0} + \partial_3 p^{I,1} = 0 \\ \partial_2 u_1^{I,1} = \partial_2 u_2^{I,1} = \partial_2 u_3^{I,1} = \partial_2 p^{I,1} = 0 \\ \partial_1 u_1^{I,1} + \partial_3 u_3^{I,1} = 0 \end{cases}$$

We also remark that we can not obtain any determined boundary condition for  $u^{I,1}$ , but only a condition depending on the boundary condition of  $u^{B,1}$ . Indeed, on the boundary, we recall the value of  $u_i^{I,j}$  is related to the value of  $u_i^{B,j}$  by the equation

$$u_i^{I,j}(t, x_1, 0) + u_i^{B,j}(t, x_1, 0) = 0 \quad j = 0, 1; \quad i = 1, 2, 3.$$

Using the same argument, we can prove that  $\partial_2 p^{I,1} = 0$ , and we obtain the following 2D linearized Euler-type system with three components in the half-plane

$$\begin{cases} \partial_t u_1^{I,1} + u_1^{I,0} \partial_1 u_1^{I,1} + u_3^{I,0} \partial_3 u_1^{I,1} + u_1^{I,1} \partial_1 u_1^{I,0} + u_3^{I,1} \partial_3 u_1^{I,0} + \partial_1 p^{I,1} = 0 \\ \partial_t u_2^{I,1} + u_1^{I,0} \partial_1 u_2^{I,1} + u_3^{I,0} \partial_3 u_2^{I,1} + u_1^{I,1} \partial_1 u_2^{I,0} + u_3^{I,1} \partial_3 u_2^{I,0} = 0 \\ \partial_t u_3^{I,1} + u_1^{I,0} \partial_1 u_3^{I,1} + u_3^{I,0} \partial_3 u_3^{I,1} + u_1^{I,1} \partial_1 u_3^{I,0} + u_3^{I,1} \partial_3 u_3^{I,0} + \partial_3 p^{I,1} = 0 \\ \partial_2 u_1^{I,1} = \partial_2 u_2^{I,1} = \partial_2 u_3^{I,1} = \partial_2 p^{I,1} = 0 \\ \partial_1 u_1^{I,1} + \partial_3 u_3^{I,1} = 0 \\ u_3^{I,1}(t, x_1, 0) = -u_3^{B,1}(t, x_1, 0) \\ u^{I,1}(0, x_1, x_3) = u_0^{I,1}(x_1, x_3). \end{cases}$$

Here, we also suppose that the initial data are well prepared, *i.e.* independent of  $x_2$ .

**2.2. Formal asymptotic expansions inside the boundary layer.** Inside the boundary layer (in the domain  $0 < x_3 \leq \sqrt{\varepsilon}$ ), we consider the Taylor expansions

$$\begin{aligned} u_i^{I,j}(t, x_h, x_3) &= u_i^{I,j}(t, x_h, 0) + x_3 \partial_3 u_i^{I,j}(t, x_h, 0) + \frac{x_3^2}{2} \partial_3^2 u_i^{I,j}(t, x_h, 0) + \dots \\ p^{I,j}(t, x_h, x_3) &= p^{I,j}(t, x_h, 0) + x_3 \partial_3 p^{I,j}(t, x_h, 0) + \frac{x_3^2}{2} \partial_3^2 p^{I,j}(t, x_h, 0) + \dots \end{aligned}$$

Performing the change of variable  $y = \frac{x_3}{\sqrt{\varepsilon}}$ , we have

$$(2.9) \quad \begin{cases} u_i^{I,j}(t, x_h, x_3) = \overline{u_i^{I,j}} + \varepsilon^{\frac{1}{2}} y \overline{\partial_3 u_i^{I,j}} + \frac{\varepsilon y^2}{2!} \overline{\partial_3^2 u_i^{I,j}} + \dots \\ p^{I,j}(t, x_h, x_3) = \overline{p^{I,j}} + \varepsilon^{\frac{1}{2}} y \overline{\partial_3 p^{I,j}} + \frac{\varepsilon y^2}{2!} \overline{\partial_3^2 p^{I,j}} + \dots \end{cases}$$

where  $\overline{f} = f(t, x_1, x_2, 0)$  is the trace of  $f$  on  $\{x_3 = 0\}$ . Now, we will rewrite the identities (2.1), taking into account the expansion (2.9). First, we have

$$(2.10) \quad \begin{aligned} u^\varepsilon &= \left( u^{B,0} + \overline{u^{I,0}} \right) + \varepsilon^{\frac{1}{2}} \left( u^{B,1} + \overline{u^{I,1}} + y \overline{\partial_3 u^{I,0}} \right) + \sum_{k=2}^3 \varepsilon^{\frac{k}{2}} \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^{k-1} u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^k u^{I,0}} \right) + \dots \\ &= \mathcal{U}^{p,0} + \varepsilon^{\frac{1}{2}} \mathcal{U}^{p,1} + \sum_{k=2}^3 \varepsilon^{\frac{k}{2}} \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^{k-1} u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^k u^{I,0}} \right) + \dots \end{aligned}$$

where we note

$$(2.11) \quad \mathcal{U}^{p,0} = u^{B,0} + \overline{u^{I,0}}, \quad \mathcal{U}^{p,1} = u^{B,1} + \overline{u^{I,1}} + y \overline{\partial_3 u^{I,0}}.$$

The derivatives of  $u^\varepsilon$  with respect to tangential variables write

$$\partial_{t,1,2}^m u^\varepsilon = \partial_{t,1,2}^m \mathcal{U}^{p,0} + \varepsilon^{\frac{1}{2}} \partial_{t,1,2}^m \mathcal{U}^{p,1} + \sum_{k=2}^3 \varepsilon^{\frac{k}{2}} \partial_{t,1,2}^m \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^{k-1} u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^k u^{I,0}} \right) + \dots$$

where  $m = 1, 2$ . For the normal variable, we have

$$\partial_3 u^\varepsilon = \varepsilon^{-\frac{1}{2}} \partial_y u^{B,0} + \left( \partial_y u^{B,1} + \overline{\partial_3 u^{I,0}} \right) + \sum_{k=1}^3 \varepsilon^{\frac{k}{2}} \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^k u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^{k+1} u^{I,0}} \right) + \dots$$

and

$$\partial_3^2 u^\varepsilon = \varepsilon^{-1} \partial_y^2 u^{B,0} + \varepsilon^{-\frac{1}{2}} \partial_y^2 u^{B,1} + \overline{\partial_3^2 u^{I,0}} + \sum_{k=1}^3 \varepsilon^{\frac{k}{2}} \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^{k+1} u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^{k+2} u^{I,0}} \right) + \dots$$

Thus,

$$\begin{aligned} -\varepsilon \Delta u^\varepsilon &= -\varepsilon \Delta_h \mathcal{U}^{p,0} - \varepsilon^{\frac{3}{2}} \Delta_h \mathcal{U}^{p,1} - \partial_y^2 \mathcal{U}^{p,0} - \varepsilon^{\frac{1}{2}} \partial_y^2 \mathcal{U}^{p,1} - \varepsilon \overline{\partial_3^2 u^{I,0}} \\ &\quad - \varepsilon^{\frac{3}{2}} \left( \frac{y^{k-1}}{(k-1)!} \overline{\partial_3^{k+1} u^{I,1}} + \frac{y^k}{k!} \overline{\partial_3^{k+2} u^{I,0}} \right) + \dots \end{aligned}$$

For the non-linear term, we only give the explicit calculations for the first orders of its expansion. We write

$$u^\varepsilon \cdot \nabla u^\varepsilon = u_h^\varepsilon \cdot \nabla_h u^\varepsilon + u_3^\varepsilon \partial_3 u^\varepsilon.$$

Then, we have

$$\begin{aligned} u_h^\varepsilon \cdot \nabla_h u^\varepsilon &= \mathcal{U}_h^{p,0} \cdot \nabla_h \mathcal{U}_h^{p,0} + \varepsilon^{\frac{1}{2}} \mathcal{U}_h^{p,0} \cdot \nabla_h \mathcal{U}_h^{p,1} + \varepsilon^{\frac{1}{2}} \mathcal{U}_h^{p,1} \cdot \nabla_h \mathcal{U}_h^{p,0} + \dots \\ u_3^\varepsilon \partial_3 u^\varepsilon &= \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_h^{p,0} + \varepsilon^{\frac{1}{2}} \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_h^{p,1} + \varepsilon^{\frac{1}{2}} \left( \overline{y \partial_3 u_3^{I,1}} + \frac{y^2}{2} \overline{\partial_3^2 u_3^{I,0}} \right) \partial_y \mathcal{U}_h^{p,0} + \dots \end{aligned}$$

For the Coriolis forcing term (the rotation term), we have

$$\begin{aligned} &\frac{e_2 \times u^\varepsilon}{\varepsilon} \\ &= \varepsilon^{-1} \begin{pmatrix} 0 \\ 0 \\ -\mathcal{U}_1^{p,0} \end{pmatrix} + \varepsilon^{-\frac{1}{2}} \begin{pmatrix} \mathcal{U}_3^{p,1} \\ 0 \\ -\mathcal{U}_1^{p,1} \end{pmatrix} + \sum_{k=2}^3 \varepsilon^{\frac{k}{2}-1} \left[ \frac{y^{k-1}}{(k-1)!} \begin{pmatrix} \overline{\partial_3^{k-1} u_3^{I,1}} \\ 0 \\ -\overline{\partial_3^{k-1} u_1^{I,1}} \end{pmatrix} + \frac{y^k}{k!} \begin{pmatrix} \overline{\partial_3^k u_3^{I,0}} \\ 0 \\ -\overline{\partial_3^k u_1^{I,0}} \end{pmatrix} \right] + \dots \end{aligned}$$

Finally, the pressure term is

$$\begin{aligned} (2.12) \quad \begin{pmatrix} \partial_{x_1} p^\varepsilon \\ \partial_{x_2} p^\varepsilon \\ \partial_{x_3} p^\varepsilon \end{pmatrix} &= \varepsilon^{-\frac{3}{2}} \begin{pmatrix} 0 \\ 0 \\ \partial_y \mathcal{P}^{p,-2} \end{pmatrix} + \sum_{j=-2}^{-1} \varepsilon^{\frac{j}{2}} \begin{pmatrix} \partial_1 \mathcal{P}^{p,j} \\ \partial_2 \mathcal{P}^{p,j} \\ \partial_y \mathcal{P}^{p,j+1} \end{pmatrix} + \left[ \begin{pmatrix} \partial_1 \mathcal{P}^{p,0} \\ \partial_2 \mathcal{P}^{p,0} \\ 0 \end{pmatrix} + \sum_{k=0}^2 \frac{y^k}{k!} \begin{pmatrix} 0 \\ 0 \\ \overline{\partial_3^{k+1} p^{I,-k}} \end{pmatrix} \right] \\ &+ \varepsilon^{\frac{1}{2}} \sum_{j=-2}^1 \frac{y^{1-j}}{(1-j)!} \begin{pmatrix} \overline{\partial_1 \partial_3^{1-j} p^{I,j}} \\ \overline{\partial_2 \partial_3^{1-j} p^{I,j}} \\ \overline{\partial_3^{2-j} p^{I,j}} \end{pmatrix} + \dots \end{aligned}$$

where

$$(2.13) \quad \mathcal{P}^{p,-2} = p^{B,-2} + \overline{p^{I,-2}}, \quad \mathcal{P}^{p,-1} = p^{B,-1} + \overline{p^{I,-1}} + \overline{y \partial_3 p^{I,-2}}$$

$$(2.14) \quad \mathcal{P}^{p,0} = p^{B,0} + \overline{p^{I,0}} + \overline{y \partial_3 p^{I,-1}} + \frac{y}{2} \overline{\partial_3 p^{I,-2}}.$$

**2.3. Incompressibility and boundary conditions.** The divergence-free property of the velocity field is rewritten as follows

$$\begin{aligned} 0 = \operatorname{div} u^\varepsilon &= \varepsilon^{-\frac{1}{2}} \partial_y u_3^{B,0} \left( t, x_h, \frac{x_3}{\sqrt{\varepsilon}} \right) + \left[ \operatorname{div} u^{I,0} + \partial_1 u_1^{B,0} + \partial_2 u_2^{B,0} + \partial_y u_3^{B,1} \left( t, x_h, \frac{x_3}{\sqrt{\varepsilon}} \right) \right] \\ &+ \varepsilon^{\frac{1}{2}} \left[ \operatorname{div} u^{I,1} + \partial_1 u_1^{B,1} + \partial_2 u_2^{B,1} \right] + \dots \end{aligned}$$

Inside the boundary layer, using the expansion (2.3) and (2.9), we deduce the following divergence-free condition

$$\varepsilon^{-\frac{1}{2}} \partial_y u_3^{B,0} + \left( \partial_1 u_1^{B,0} + \partial_2 u_2^{B,0} + \partial_y u_3^{B,1} \right) + \varepsilon^{\frac{1}{2}} \left( \partial_1 u_1^{B,1} + \partial_2 u_2^{B,1} \right) = 0.$$

Thus, we obtain the incompressibility of the boundary layer

$$(2.15) \quad \begin{aligned} \partial_1 u_1^{B,0} + \partial_2 u_2^{B,0} + \partial_y u_3^{B,1} &= 0, \\ \partial_1 u_1^{B,1} + \partial_2 u_2^{B,1} &= 0. \end{aligned}$$

Moreover, we have

$$\partial_y u_3^{B,0} = 0,$$

which, by taking  $y \rightarrow +\infty$ , gives

$$u_3^{B,0} = 0.$$

For the boundary condition in (N-S $_\varepsilon$ ) on  $\{x_3 = 0\}$ , we have

$$\sum_{j=0}^1 \varepsilon^{\frac{j}{2}} [u^{I,j}(t, x_h, 0) + u^{B,j}(t, x_h, 0)] = 0,$$

which implies that

$$\overline{u^{I,0}(t)} + u^{B,0}(t, x_h, 0) = 0,$$

$$(2.16) \quad \overline{u^{I,1}(t)} + u^{B,1}(t, x_h, 0) = 0.$$

In particular,  $u_3^{B,0} = 0$  imply

$$(2.17) \quad u_3^{I,0}|_{x_3=0} = \overline{u_3^{I,0}} = 0,$$

which is the boundary condition for Euler equation in (1.3), and the third components in (2.16) gives the boundary condition of linearized Euler equation in (1.6).

**2.4. Formal derivation of the governing equations of the fluid in the boundary layer.** Now, we consider the system (N-S $_\varepsilon$ ) near  $\{x_3 = 0\}$ , using the asymptotic formal (2.10) - (2.12).

**At the order  $\varepsilon^{-\frac{3}{2}}$ ,** we have

$$\partial_y p^{B,-2} = 0,$$

which implies that  $p^{B,-2} = 0$  because  $p^{B,-2}$  goes to zero as  $y \rightarrow +\infty$ . Using the new notation of the pressure defined in (2.13), we get

$$\partial_y \mathcal{P}^{p,-2} = 0.$$

**At the order  $\varepsilon^{-1}$ ,** using the fact that  $\overline{u_3^{I,0}} = 0$ ,  $u_3^{B,0} = 0$  and  $p^{B,-2} = 0$ , we get

$$\begin{pmatrix} 0 \\ 0 \\ -u_1^{B,0} \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ -u_1^{I,0} \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \partial_y p^{B,-1} \end{pmatrix} + \overline{\nabla p^{I,-2}} = 0,$$

which implies that  $\overline{\partial_1 p^{I,-1}} = \overline{\partial_2 p^{I,-1}} = 0$  and

$$(2.18) \quad -u_1^{B,0} - \overline{u_1^{I,0}} + \partial_y p^{B,-1} + \overline{\partial_3 p^{I,-2}} = 0.$$

Using the new velocity and pressure defined in (2.13) and taking into account the fact that  $\mathcal{U}_3^{p,0} = 0$ , we can also write

$$(2.19) \quad \begin{pmatrix} 0 \\ 0 \\ -\mathcal{U}_1^{p,0} \end{pmatrix} + \begin{pmatrix} \partial_1 \mathcal{P}^{p,-2} \\ \partial_2 \mathcal{P}^{p,-2} \\ \partial_y \mathcal{P}^{p,-1} \end{pmatrix} = 0.$$

**At the order  $\varepsilon^{-1/2}$ ,** we have

$$\begin{pmatrix} u_3^{B,1} \\ 0 \\ -u_1^{B,1} \end{pmatrix} + \begin{pmatrix} \overline{u_3^{I,1}} \\ 0 \\ -\overline{u_1^{I,1}} \end{pmatrix} + y \begin{pmatrix} \overline{\partial_3 u_3^{I,0}} \\ 0 \\ -\overline{\partial_3 u_1^{I,0}} \end{pmatrix} + \begin{pmatrix} \partial_1 p^{B,-1} \\ \partial_2 p^{B,-1} \\ \partial_y p^{B,0} \end{pmatrix} + \begin{pmatrix} \overline{\partial_1 p^{I,-1}} \\ \overline{\partial_2 p^{I,-1}} \\ \overline{\partial_3 p^{I,-1}} \end{pmatrix} + y \begin{pmatrix} \overline{\partial_1 \partial_3 p^{I,-2}} \\ \overline{\partial_2 \partial_3 p^{I,-2}} \\ \overline{\partial_3^2 p^{I,-2}} \end{pmatrix} = 0,$$

or in a equivalent way, using the new velocity and pressure defined in (2.14),

$$(2.20) \quad \begin{pmatrix} \mathcal{U}_3^{p,1} \\ 0 \\ -\mathcal{U}_1^{p,1} \end{pmatrix} + \begin{pmatrix} \partial_1 \mathcal{P}^{p,-1} \\ \partial_2 \mathcal{P}^{p,-1} \\ \partial_y \mathcal{P}^{p,0} \end{pmatrix} = 0.$$

then

$$\partial_2 \mathcal{P}^{p,-1} = 0.$$

and

$$\begin{aligned} \partial_2 \mathcal{U}_1^{p,0} &= \partial_2 \partial_y \mathcal{P}^{p,-1} = \partial_y \partial_2 \mathcal{P}^{p,-1} = 0 \\ \partial_2 \mathcal{U}_3^{p,1} &= -\partial_2 \partial_1 \mathcal{P}^{p,-1} = -\partial_1 \partial_2 \mathcal{P}^{p,-1} = 0. \end{aligned}$$

Using the divergence-free properties (2.3) and (2.15), we also have

$$\partial_2 \mathcal{U}_2^{p,0} = -\partial_1 \mathcal{U}_1^{p,0} - \partial_y \mathcal{U}_3^{p,1} = -\partial_1 \partial_y \mathcal{P}^{p,-1} - (-\partial_y \partial_1 \mathcal{P}^{p,-1}) = 0.$$

We deduce that  $(\mathcal{U}_1^{p,0}, \mathcal{U}_2^{p,0}, \mathcal{U}_3^{p,1})$  is a divergence-free vector field which is independent on  $x_2$ . The fact that  $\partial_2 u^{I,0} = \partial_2 u^{I,1} = 0$  implies that

$$(2.21) \quad \partial_2 u_1^{B,0} = \partial_2 u_2^{B,0} = \partial_2 u_3^{B,1} = 0.$$

**Remark 2.2.** *The leading order of the velocity of the fluid inside the boundary layer also obeys the Taylor-Proudman theorem.*

**At the order**  $\varepsilon^0$ , recalling that  $u_3^{B,0} = \overline{u_3^{I,0}} = 0$ , we get the following equation

$$\begin{aligned} & \partial_t \left( u_h^{B,0} + \overline{u_h^{I,0}} \right) - \partial_y^2 u_h^{B,0} + \left( u_h^{B,0} + \overline{u_h^{I,0}} \right) \cdot \nabla_h \left( u_h^{B,0} + \overline{u_h^{I,0}} \right) + \left( u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \right) \partial_y u_h^{B,0} \\ & + y \left( \frac{\overline{\partial_3 u_3^{I,1}}}{-\partial_3 u_1^{I,1}} \right) + \frac{y^2}{2} \left( \frac{\overline{\partial_3^2 u_3^{I,0}}}{-\partial_3^2 u_1^{I,0}} \right) + \begin{pmatrix} \overline{\partial_1 p^{B,0}} \\ \overline{\partial_2 p^{B,0}} \\ 0 \end{pmatrix} + \begin{pmatrix} \overline{\partial_1 p^{I,0}} \\ \overline{\partial_2 p^{I,0}} \\ \overline{\partial_3 p^{I,0}} \end{pmatrix} + y \begin{pmatrix} \overline{\partial_1 \partial_3 p^{I,-1}} \\ \overline{\partial_2 \partial_3 p^{I,-1}} \\ \overline{\partial_3^2 p^{I,-1}} \end{pmatrix} + \frac{y^2}{2} \begin{pmatrix} \overline{\partial_1 \partial_3^2 p^{I,-2}} \\ \overline{\partial_2 \partial_3^2 p^{I,-2}} \\ \overline{\partial_3^3 p^{I,-2}} \end{pmatrix} = 0. \end{aligned}$$

From (2.4) and (2.6), we deduce that

$$-y \overline{\partial_3 u_1^{I,1}} - \frac{y^2}{2} \overline{\partial_3^2 u_1^{I,0}} + y \overline{\partial_3^2 p^{I,-1}} + \frac{y^2}{2} \overline{\partial_3^3 p^{I,-2}} = 0.$$

We also remark that the boundary condition applying to the third equation of the Euler system implies that

$$\overline{\partial_3 p^{I,0}} = 0.$$

Then, using the new velocity and pressure defined in (2.11) and (2.13), we get

$$\partial_t \mathcal{U}_h^{p,0} - \partial_y^2 \mathcal{U}_h^{p,0} + \mathcal{U}_h^{p,0} \cdot \nabla_h \mathcal{U}_h^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_h^{p,0} + \begin{pmatrix} \overline{\partial_1 p^{B,0}} + \overline{\partial_1 p^{I,0}} \\ \overline{\partial_2 p^{p,0}} \end{pmatrix} = 0.$$

Taking into account the divergence-free condition (2.15), the identities (2.18) and (2.19), and  $(\mathcal{U}_1^{p,0}, \mathcal{U}_2^{p,0}, \mathcal{U}_3^{p,1})$  is independent on  $x_2$ , we deduce that  $(\mathcal{U}_1^{p,0}, \mathcal{U}_2^{p,0}, \mathcal{U}_3^{p,1})$  satisfies the following system

$$\begin{cases} \partial_t \mathcal{U}_1^{p,0} - \partial_y^2 \mathcal{U}_1^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_1^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0} + \partial_1 p^{B,0} + \overline{\partial_1 p^{I,0}} = 0 \\ \partial_t \mathcal{U}_2^{p,0} - \partial_y^2 \mathcal{U}_2^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_2^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_2^{p,0} + \partial_2 p^{p,0} = 0 \\ \partial_1 \mathcal{U}_1^{p,0} + \partial_y \mathcal{U}_3^{p,1} = 0, \\ \partial_2 \mathcal{U}_1^{p,0} = \partial_2 \mathcal{U}_2^{p,0} = \partial_2 \mathcal{U}_3^{p,1} = 0. \end{cases}$$

We remark that the above system is not complete, since we need another equation for the component  $\mathcal{U}_3^{p,1}$ .

**At the order**  $\varepsilon^{1/2}$ , we have

$$\begin{aligned} & \partial_t \mathcal{U}^{p,1} - \partial_y^2 \mathcal{U}^{p,1} + \mathcal{U}_h^{p,0} \cdot \nabla_h \mathcal{U}^{p,1} + \mathcal{U}_h^{p,1} \cdot \nabla_h \mathcal{U}_h^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}^{p,1} + \left( y \overline{\partial_3 u_3^{I,1}} + \frac{y^2}{2} \overline{\partial_3^2 u_3^{I,0}} \right) \partial_y \mathcal{U}_h^{p,0} \\ & + \left[ \frac{y^2}{2} \begin{pmatrix} \overline{\partial_3^2 u_3^{I,1}} \\ 0 \\ -\partial_3^2 u_1^{I,1} \end{pmatrix} + \frac{y^3}{6} \begin{pmatrix} \overline{\partial_3^3 u_3^{I,0}} \\ 0 \\ -\partial_3^3 u_1^{I,0} \end{pmatrix} \right] + \sum_{k=0}^3 \frac{y^k}{k!} \begin{pmatrix} \overline{\partial_1 \partial_3^k p^{I,1-k}} \\ \overline{\partial_2 \partial_3^k p^{I,1-k}} \\ \overline{\partial_3^{k+1} p^{I,1-k}} \end{pmatrix} = 0. \end{aligned}$$

Here, we are only interested in the component  $\mathcal{U}_3^{p,1}$ . Using the fact that  $\partial_2 \mathcal{U}_3^{p,1} = 0$ , we obtain

$$\partial_t \mathcal{U}_3^{p,1} - \partial_y^2 \mathcal{U}_3^{p,1} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_3^{p,1} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_3^{p,1} + \overline{\partial_3 p^{I,1}} + y \overline{\partial_3^2 p^{I,0}} = 0.$$

Collect all the above formal calculations, we deduce the following governing equations of the boundary layer

$$(P1) \quad \begin{cases} \partial_t \mathcal{U}_1^{p,0} - \partial_y^2 \mathcal{U}_1^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_1^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0} + \partial_1 p^{B,0} + \overline{\partial_1 p^{I,0}} = 0 \\ \partial_t \mathcal{U}_3^{p,1} - \partial_y^2 \mathcal{U}_3^{p,1} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_3^{p,1} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_3^{p,1} + \overline{\partial_3 p^{I,1}} + y \overline{\partial_3^2 p^{I,0}} = 0 \\ \partial_1 \mathcal{U}_1^{p,0} + \partial_y \mathcal{U}_3^{p,1} = 0 \\ \partial_2 \mathcal{U}_1^{p,0} = \partial_2 \mathcal{U}_3^{p,1} = 0 \\ \mathcal{U}_1^{p,0}(t, x_1, 0) = 0, \quad \lim_{y \rightarrow +\infty} \mathcal{U}_1^{p,0}(t, x_1, y) = \overline{u_1^{I,0}}(x_1) \\ \mathcal{U}_3^{p,1}(t, x_1, 0) = 0, \quad \partial_y \mathcal{U}_3^{p,1}(t, x_1, 0) = 0 \\ \mathcal{U}_1^{p,0}(0, x_1, y) = u_{0,1}^{B,0}(x_1, y) + \overline{u_{0,1}^{I,0}}(x_1) \\ \mathcal{U}_3^{p,1}(0, x_1, y) = u_{0,3}^{B,1}(x_1, y) + \overline{u_{0,3}^{I,1}}(x_1) + y \overline{\partial_3 u_{0,3}^{I,0}}(x_1). \end{cases}$$

and

$$(P2) \quad \begin{cases} \partial_t \mathcal{U}_2^{p,0} - \partial_y^2 \mathcal{U}_2^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_2^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_2^{p,0} + \partial_2 \mathcal{P}^{p,0} = 0 \\ \mathcal{U}_2^{p,0}(0, x_1, y) = u_{0,2}^{B,0}(x_1, y) + \overline{u_{0,2}^{I,0}}(x_1) \\ \mathcal{U}_2^{p,0}(t, x_1, 0) = 0, \quad \lim_{y \rightarrow +\infty} \mathcal{U}_2^{p,0}(t, x_1, y) = \overline{u_2^{I,0}}(x_1) \\ \mathcal{U}_2^{p,0}(0, x_1, y) = u_{0,2}^{B,0}(x_1, y) + \overline{u_{0,2}^{I,0}}(x_1). \end{cases}$$

**Claim:** *The pressure term of the (P2) satisfies  $\partial_2 \mathcal{P}^{p,0} = 0$ .*

Indeed, applying  $\partial_2$  to the first equation of the systems (P1) and (P2), and using the fact that

$$\partial_2 \mathcal{U}_1^{p,0} = \partial_2 \mathcal{U}_2^{p,0} = \partial_2 \mathcal{U}_3^{p,1} = 0,$$

we deduce that

$$\partial_1 \partial_2 \mathcal{P}^{p,0} = \partial_2^2 \mathcal{P}^{p,0} = 0.$$

This means that, modulo a constant, we have

$$\mathcal{P}^{p,0} = x_2 G_1(t, y) + \int_{-\infty}^{x_1} \tilde{f}(t, x, y) dx,$$

where

$$G_1 = - \left( \partial_t \mathcal{U}_2^{p,0} - \partial_y^2 \mathcal{U}_2^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_2^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_2^{p,0} \right)$$

is to be determined and

$$\tilde{f} = \partial_1 \mathcal{P}^{p,0} = -\partial_t \mathcal{U}_1^{p,0} + \partial_y^2 \mathcal{U}_1^{p,0} - \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_1^{p,0} - \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0} - \left( \frac{y^2}{2} \overline{\partial_3^2 u_3^{I,0}} + y \overline{\partial_3 u_3^{I,1}} \right).$$

We recall that, from (2.20), we have

$$\partial_y \mathcal{P}^{p,0} = \mathcal{U}_1^{p,1},$$

where  $\mathcal{U}_1^{p,1}$  is the solution of the system

$$\begin{cases} \partial_t \mathcal{U}_1^{p,1} - \partial_y^2 \mathcal{U}_1^{p,1} + \mathcal{U}_1^{p,0} \cdot \partial_1 \mathcal{U}_1^{p,1} + \mathcal{U}_2^{p,0} \cdot \partial_2 \mathcal{U}_1^{p,1} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,1} + \mathcal{U}_1^{p,1} \partial_1 \mathcal{U}_1^{p,0} \\ \quad + \left( \overline{y \partial_3 u_3^{I,1}} + \frac{y^2}{2} \overline{\partial_3^2 u_3^{I,0}} \right) \partial_y \mathcal{U}_1^{p,0} + \left[ \frac{y^2}{2} \overline{\partial_3^2 u_3^{I,1}} + \frac{y^3}{6} \overline{\partial_3^3 u_3^{I,0}} \right] + \sum_{k=0}^3 \frac{y^k}{k!} \overline{\partial_1 \partial_3^k p^{I,1-k}} = 0 \\ \mathcal{U}_1^{p,1}(0, x_1, x_2, y) = u_{0,1}^{B,1}(x_1, y) + \overline{u_{0,1}^{I,1}}(x_1) + y \overline{\partial_3 u_{0,1}^{I,0}}(x_1) + \alpha_1(y) x_2 \\ \mathcal{U}_1^{p,1}(t, x_1, 0) = 0. \end{cases}$$

We remark that  $\partial_y G_1(t, y) = \partial_2 \mathcal{U}_1^{p,1}$  and we recall that  $\partial_1 \partial_2 \mathcal{U}_1^{p,1} = \partial_2^2 \mathcal{U}_1^{p,1}$ . So, in fact, we will find  $\partial_y G_1$  by solving the following system

$$(2.22) \quad \begin{cases} \partial_t (\partial_y G_1) - \partial_y^2 (\partial_y G_1) + (\partial_1 \mathcal{U}_1^{p,0}) (\partial_y G_1) + \mathcal{U}_3^{p,1} \partial_y (\partial_y G_1) = 0 \\ \partial_y G_1(0, y) = \alpha_1(y) \\ \partial_y G_1(t, 0) = 0. \end{cases}$$

where  $\alpha_1$  is a given function, with  $\alpha_1(0) = 0$ . For the case of well prepared data, we consider the initial data to be independent of  $x_2$ , so  $\alpha_1 \equiv 0$  and it is easy to see that the system (2.22) admits 0 as a trivial solution. Then, the uniqueness of this solution implies  $\partial_y G_1(t, \cdot) \equiv 0$ . Replacing  $y = 0$  in (2.22), we obtain  $G_1(t, 0) = 0$ , and so  $G_1(t, \cdot) \equiv 0$ , for any  $t \in \mathbb{R}_+$ .

## 3. WELL-POSEDNESS OF THE BOUNDARY LAYER SYSTEM

In this section we will prove the well-posedness for system (P1). Since the pressure term in the first equation of (P1) is unknown, we begin with handling the second one to prove the existence of  $u_3^{I,1}$  and then use the divergence-free property to find  $u_1^{I,0}$ . To do so we insert the representations

$$u_1^{p,0} = u_1^{B,0} + \overline{u_1^{I,0}}, \quad u_3^{p,1} = u_3^{B,1} + \overline{u_3^{I,1}} + \overline{y\partial_3 u_3^{I,0}}$$

into the second equation of (P1), and then make use of the equations (1.3) and (1.6) of  $u_3^{I,0}$  and  $u_3^{I,1}$ . It then follows that the unknowns  $u_3^{B,1}$ ,  $u_1^{B,0}$  and  $u_3^{I,1}$  satisfy the equation

$$\begin{aligned} & \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) u_3^{B,1} + \left( u_1^{B,0} + \overline{u_1^{I,0}} \right) \partial_1 u_3^{B,1} \\ & + \left( u_3^{B,1} + \overline{u_3^{I,1}} \right) \partial_y u_3^{B,1} + \overline{\partial_3 u_3^{I,0}} u_3^{B,1} + \left( -\partial_1 \overline{u_3^{I,1}} + \overline{y\partial_1 \partial_3 u_3^{I,0}} \right) u_1^{B,0} = 0, \end{aligned}$$

and the divergence-free properties (2.15) and (2.21) yield

$$u_1^{B,0} = - \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, y) dz.$$

Thus the above is just a equation for  $u_3^{B,1}$ . To solve the system (P1), we consider the following nonlinear initial-boundary problem,

$$(3.1) \quad \begin{cases} \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) u + \left( v + \overline{u_1^{I,0}} \right) \partial_1 u \\ \quad + (u - u(t, x_1, 0)) \partial_y u + \overline{\partial_3 u_3^{I,0}} u + \left( \partial_1 u(t, x_1, 0) + \overline{y\partial_1 \partial_3 u_3^{I,0}} \right) v = 0, \\ \partial_y u|_{y=0} = -\overline{\partial_3 u_3^{I,0}}(t, x_1), \quad \lim_{y \rightarrow +\infty} u(t, x_1, y) = 0, \\ u|_{t=0} = u_0(x_1, y), \end{cases}$$

where the unknown functions  $u$  and  $v$  are linked by the relation

$$(3.2) \quad v(t, x_1, y) = - \int_{-\infty}^{x_1} \partial_y u(t, z, y) dz.$$

Recall the functions  $u_1^{I,0}, u_3^{I,0}$  are the solutions to the Euler-type system (1.3). By Theorem 1.2, we see  $u_1^{I,0}, u_3^{I,0} \in \mathcal{A}_\tau$  for some  $\tau > 0$ .

The main result of this section can be stated as follows.

**Theorem 3.1.** *Suppose the initial data  $u_0 \in X_{\rho_0, a_0}$  for some  $\rho_0 > 0$  and  $a_0 > 0$  and satisfies the compatibility conditions. Then the system (3.1) admits a unique solution*

$$u \in L^\infty([0, T_*]; X_{\rho_*, a})$$

for some  $\rho_* > 0$ ,  $a > 0$  and  $T_* > 0$ .

We now proceed the proof of the theorem 3.1 through the following parabolic approximations.

**The approximate solutions.** Consider the following regularized system, for  $\varepsilon > 0$ ,

$$(3.3) \quad \begin{cases} \left( \partial_t - \varepsilon \partial_1^2 - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) u^\varepsilon + \left( v^\varepsilon + \overline{u_1^{I,0}} \right) \partial_1 u^\varepsilon \\ \quad + (u^\varepsilon - u^\varepsilon(t, x_1, 0)) \partial_y u^\varepsilon + \overline{\partial_3 u_3^{I,0}} u^\varepsilon + \left( \partial_1 u(t, x_1, 0) + \overline{y\partial_1 \partial_3 u_3^{I,0}} \right) v = 0, \\ \partial_y u^\varepsilon(t, x_1, 0) = \overline{\partial_3 u_3^{I,0}}(t, x_1), \quad \lim_{y \rightarrow +\infty} u^\varepsilon(t, x_1, y) = 0, \\ u^\varepsilon|_{t=0} = u_0(x_1, y). \end{cases}$$

The above is a nonlinear parabolic equation, and from classical theory we can deduce the following local well-posedness result.

**Theorem 3.2.** *Suppose the initial data  $u_0 \in X_{2\rho_0, a_0}$  for some  $\rho_0 > 0$ ,  $a_0 > 0$  and satisfies the compatibility conditions. Then the system (3.3) admits a unique solution*

$$u^\varepsilon \in L^\infty([0, T_\varepsilon]; X_{\rho_0, a})$$

for some  $0 < a < a_0$  independent of  $\varepsilon$  and  $T_\varepsilon > 0$  depends on  $\varepsilon$ .

**Uniform estimates for the approximate solutions.** We will perform the uniform estimate with respect to  $\varepsilon$  for the approximate solutions  $u^\varepsilon$  given in the previous Theorem. The main result here can be stated as follows.

**Proposition 3.3.** *Suppose  $u^\varepsilon \in L^\infty([0, T_\varepsilon]; X_{\rho_0, a})$  is a solution to the initial-boundary problem (3.3). Then there exists  $0 < \rho_* \leq \rho_0$ , depending only on  $|u_0|_{X_{\rho_0, a_0}}$ , such that  $u^\varepsilon \in L^\infty([0, T_\varepsilon]; X_{\rho_*, a})$  for all  $\varepsilon > 0$ . Moreover*

$$(3.4) \quad \|u^\varepsilon\|_{L^\infty([0, T_\varepsilon]; X_{\rho_*, a})} \leq C |u_0|_{X_{\rho_0, a_0}},$$

where  $C$  is a constant depending only on  $a_0, \rho_0, \tau, \|u_3^{I,0}\|_{\mathcal{A}_\tau}$  and  $\|u_1^{I,0}\|_{\mathcal{A}_\tau}$ , but independent of  $\varepsilon$ .

To prove the above proposition, we need another two *auxiliary norms*  $|\cdot|_{Y_{\rho, a}}$  and  $|\cdot|_{Z_{\rho, a}}$  which are defined by

$$(3.5) \quad |u|_{Y_{\rho, a}}^2 = \sum_{m \leq 2} \left( \sum_{0 \leq j \leq 1} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)} \right)^2 \\ + \sum_{m \geq 3} \left( \sum_{0 \leq j \leq 1} (m-1)^{1/2} \rho^{-1/2} \frac{\rho^{m-1}}{(m-3)!} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)} \right)^2,$$

and

$$|u|_{Z_{\rho, a}}^2 = \sum_{m \leq 2} \left( \sum_{1 \leq j \leq 2} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)} \right)^2 \\ + \sum_{m \geq 3} \left( \sum_{1 \leq j \leq 2} \frac{\rho^{m-1}}{(m-3)!} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)} \right)^2.$$

The following energy estimate is a key part to prove Proposition 3.3.

**Proposition 3.4.** *Let  $u^\varepsilon \in L^\infty([0, T_\varepsilon]; X_{\rho_0, a})$  be a solution to the initial-boundary problem (3.3) and  $0 < \rho(t) \leq \min\{\rho_0/2, \tau/3\}$  a smooth function. Then for any  $t \in [0, T_\varepsilon]$ ,*

$$(3.6) \quad |u^\varepsilon(t)|_{X_{\rho(t), a}}^2 + \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho(t), a}}^2 dt - \int_0^T \rho'(t) |u^\varepsilon(t)|_{Y_{\rho(t), a}}^2 dt \\ \leq |u_0|_{X_{\rho_0, a_0}}^2 + C \int_0^{T_\varepsilon} \left( |\rho'(t)| \rho(t)^{-2} |u^\varepsilon(t)|_{X_{\rho(t), a}} + |u^\varepsilon(t)|_{X_{\rho(t), a}}^2 + |u^\varepsilon(t)|_{X_{\rho(t), a}}^4 \right) dt \\ + C \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho(t), a}} |u^\varepsilon(t)|_{Y_{\rho(t), a}}^2 dt.$$

The proof of the proposition above is postponed to the next section, and we now use it to prove Proposition 3.3.

**Proof of Proposition 3.3.** To simplify the notations we will use  $C$  in the following discussion to denote different suitable constants, which depend only on  $a_0, \rho_0, \tau, \|u_3^{I,0}\|_{\mathcal{A}_\tau}$  and  $\|u_1^{I,0}\|_{\mathcal{A}_\tau}$ , but independent of  $\varepsilon$ .

Let  $\rho_\varepsilon$  be the solution to the differential equation:

$$(3.7) \quad \begin{cases} \rho'_\varepsilon(t) = -|u^\varepsilon(t)|_{Z_{\rho_\varepsilon(t), a}}, \\ \rho|_{t=0} = \min\{\rho_0/2, \tau/3\}, \end{cases}$$

or equivalently

$$(3.8) \quad \rho_\varepsilon(t) = \min\{\rho_0/2, \tau/3\} - \int_0^t |u^\varepsilon(s)|_{Z_{\rho_\varepsilon(s), a}} ds.$$

Observe, for any  $0 < \rho, \tilde{\rho} \leq \rho_0/2$ , we have

$$\left| |u^\varepsilon|_{Z_{\rho, a}} - |u^\varepsilon|_{Z_{\tilde{\rho}, a}} \right| \leq C |u^\varepsilon|_{Z_{\rho_0, a}} |\rho - \tilde{\rho}|,$$



which along with Cauchy-Lipschitz Theorem gives the existence of  $\rho_\varepsilon$  to equation (3.7). Now choosing  $\rho(t) = \rho_\varepsilon(t)$  in (3.6) and observing (3.7), we can rewrite (3.6) as

$$\begin{aligned} & |u^\varepsilon(t)|_{X_{\rho_\varepsilon, a}}^2 + \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho_\varepsilon, a}}^2 dt \\ \leq & |u_0|_{X_{\rho_0, a_0}}^2 + C \int_0^{T_\varepsilon} \left( |\rho'_\varepsilon(t)| \rho_\varepsilon^{-2} |u^\varepsilon|_{X_{\rho_\varepsilon, a}} + |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^2 + |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^4 \right) dt. \end{aligned}$$

Thus, using (3.7),

$$\begin{aligned} (3.9) \quad & |u^\varepsilon(t)|_{X_{\rho_\varepsilon, a}}^2 + \frac{1}{2} \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho_\varepsilon, a}}^2 dt \\ & \leq |u_0|_{X_{\rho_0, a_0}}^2 + C \int_0^{T_\varepsilon} \left( \rho_\varepsilon^{-4} |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^2 + |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^2 + |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^4 \right) dt. \end{aligned}$$

In view of (3.8) for  $T_\varepsilon$  be small sufficiently we have

$$\forall t \in [0, T_\varepsilon], \quad \rho_\varepsilon(t) \geq \frac{1}{8} \min \{ \rho_0, \tau/3 \},$$

and thus it follows from (3.9) that, for any  $t \in [0, T_\varepsilon]$ ,

$$|u^\varepsilon(t)|_{X_{\rho_\varepsilon, a}}^2 + \frac{1}{2} \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho_\varepsilon, a}}^2 dt \leq |u_0|_{X_{\rho_0, a_0}}^2 + C \int_0^{T_\varepsilon} \left( |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^2 + |u^\varepsilon|_{X_{\rho_\varepsilon, a}}^4 \right) dt,$$

with  $C$  depending only on  $a_0, \rho_0, \tau, \|u_3^{I,0}\|_{\mathcal{A}_\tau}$  and  $\|u_1^{I,0}\|_{\mathcal{A}_\tau}$ , but independent of  $\varepsilon$ . Thus by general Gronwall inequality, we conclude

$$(3.10) \quad |u^\varepsilon(t)|_{X_{\rho_\varepsilon, a}}^2 \leq C |u_0|_{X_{\rho_0, a_0}}^2,$$

and

$$\int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho_\varepsilon, a}}^2 dt \leq 3 |u_0|_{X_{\rho_0, a_0}}^2 + |u_0|_{X_{\rho_0, a_0}}^4.$$

As a result, in view of (3.8) we see

$$\begin{aligned} \rho_\varepsilon(t) &= \min \{ \rho_0/2, \tau/3 \} - \int_0^t |u^\varepsilon(s)|_{Z_{\rho_\varepsilon(s), a}} ds \\ &\geq \min \{ \rho_0/2, \tau/3 \} - t^{1/2} \left( \int_0^{T_\varepsilon} |u^\varepsilon(t)|_{Z_{\rho_\varepsilon, a}}^2 dt \right)^{1/2} \\ &\geq \min \{ \rho_0/2, \tau/3 \} - t^{1/2} \left( 2 |u_0|_{X_{\rho_0, a_0}}^2 + |u_0|_{X_{\rho_0, a_0}}^4 \right)^{1/2}. \end{aligned}$$

So if we choose  $T_*$  such that

$$(3.11) \quad T_* = 4^{-1} \left( 3 |u_0|_{X_{\rho_0, a_0}}^2 + |u_0|_{X_{\rho_0, a_0}}^4 \right)^{-1} \left( \min \{ \rho_0/2, \tau/3 \} \right)^2$$

Then

$$\forall t \in [0, T_\varepsilon] \subset [0, T_*], \quad \rho_\varepsilon(t) \geq \rho_* \stackrel{\text{def}}{=} \frac{1}{4} \min \{ \rho_0, \tau/3 \}.$$

By (3.10), it follows that

$$\forall t \in [0, T_\varepsilon] \subset [0, T_*], \quad |u^\varepsilon(t)|_{X_{\rho_*, a}}^2 \leq C |u_0|_{X_{\rho_0, a_0}}^2.$$

This completes the proof of Proposition 3.3.  $\square$

**Completion of the proof of Theorem 3.1.** Due to the uniform estimate (3.4), we can extend the lifespan  $T_\varepsilon$  to  $T_*$  with  $T_*$  defined in (3.11), following the standard bootstrap arguments. Thus we see for any  $\varepsilon > 0$  the system (3.3) admits a unique solution  $u^\varepsilon \in L^\infty([0, T_*]; X_{\rho_*, a})$  such that

$$\|u^\varepsilon\|_{L^\infty([0, T_*]; X_{\rho_*, a})} \leq C |u_0|_{X_{\rho_0, a_0}},$$

with  $T_*, \rho_*, a, C$  independent of  $\varepsilon$ . Thus letting  $\varepsilon \rightarrow 0$ , the compactness arguments show that the limit  $u \in L^\infty([0, T_*]; X_{\rho_*, a})$  solves the system (3.1), proving Theorem 3.1.  $\square$

**Proof of Theorem 1.5 .** Taking

$$u = u_3^{B,1}, \quad v = - \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, y) dz,$$

the system (3.1) implies that the function

$$\mathcal{U}_3^{p,1} = u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}}$$

satisfies

$$\begin{cases} \partial_t \mathcal{U}_3^{p,1} - \partial_y^2 \mathcal{U}_3^{p,1} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_3^{p,1} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_3^{p,1} + \overline{\partial_3 p^{I,1}} + y \overline{\partial_3^2 p^{I,0}} = 0, \\ \partial_y \mathcal{U}_3^{p,1}(t, x_1, 0) = 0, \\ \mathcal{U}_3^{p,1}(0, x_1, y) = u_{0,3}^{B,1}(x_1, y) + \overline{u_{0,3}^{I,1}}(x_1) + y \overline{\partial_3 u_{0,3}^{I,0}}(x_1), \end{cases}$$

with

$$\mathcal{U}_1^{p,0} = - \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, y) dz + \overline{u_1^{I,0}}.$$

So we need to check the boundary condition

$$(3.12) \quad \mathcal{U}_3^{p,1}|_{y=0} = u_3^{B,1}(t, x_1, 0) + \overline{u_3^{I,1}}(t, x_1) = 0.$$

For this purpose, we first use Theorem 3.1 to determine  $u_3^{B,1}$ , then use Theorem 1.3 to solve the linearized Euler system (1.6) with the boundary condition

$$u_3^{I,1}|_{x_3=0} = -u_3^{B,1}(t, x_1, 0).$$

For the component  $\mathcal{U}_1^{p,0}$ , using the divergence-free properties of  $u^{I,0}$ , we have firstly

$$\begin{aligned} \mathcal{U}_1^{p,0}|_{y=0} &= - \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, 0) dz + \overline{u_1^{I,0}}(t, x_1) \\ &= \int_{-\infty}^{x_1} \overline{\partial_3 u_3^{I,0}}(t, z) dz + \overline{u_1^{I,0}}(t, x_1) \\ &= - \int_{-\infty}^{x_1} \overline{\partial_1 u_1^{I,0}}(t, z) dz + \overline{u_1^{I,0}}(t, x_1) \\ &= 0. \end{aligned}$$

On the other hand, since  $u_3^{B,1} \in L^\infty([0, T_*]; X_{\rho_*, a})$ , we have the limit

$$\begin{aligned} \lim_{y \rightarrow +\infty} \mathcal{U}_1^{p,0}(t, x_1, y) &= - \lim_{y \rightarrow +\infty} \int_{-\infty}^{x_1} \partial_y u_3^{B,1}(t, z, y) dz + \overline{u_1^{I,0}}(t, x_1) \\ &= \overline{u_1^{I,0}}(t, x_1). \end{aligned}$$

So the boundary conditions for  $\mathcal{U}_1^{p,0}$  are satisfied. Finally, for the pressure term of the first equation in (P1), once we obtain  $\mathcal{U}_3^{p,1}$ ,  $\mathcal{U}_1^{p,0}$  and  $\overline{\partial_1 p^{I,0}}$ , it is enough to put

$$\partial_1 p^{B,0} = -\partial_t \mathcal{U}_1^{p,0} + \partial_y^2 \mathcal{U}_1^{p,0} - \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_1^{p,0} - \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_1^{p,0} - \overline{\partial_1 p^{I,0}}.$$

We then complete the proof of Theorem 1.5.

#### 4. UNIFORM ENERGY ESTIMATES

In this section we proceed through the following lemmas to prove Proposition 3.4. To simplify the notations in the following proof we will write  $u$  instead of  $u^\varepsilon$ , omitting the superscript  $\varepsilon$ , and use  $C$  in the following discussion to denote different suitable constants, which depend only on  $a_0, \rho_0, \tau, \|u_3^{I,0}\|_{\mathcal{A}_\tau}$  and  $\|u_1^{I,0}\|_{\mathcal{A}_\tau}$ .

In view of the definition of  $|\cdot|_{X_{\rho, a}}$  it suffices to estimate terms

$$(4.1) \quad \sum_{m \leq 2} \left( \| \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m u \|_{L^2(\mathbb{R}_+^2)} \right) + \sum_{m \geq 3} \left( \frac{\rho^{m-1}}{(m-3)!} \| \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m u \|_{L^2(\mathbb{R}_+^2)} \right)$$

and

$$(4.2) \quad \sum_{m \leq 2} \left( \| \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u \|_{L^2(\mathbb{R}_+^2)} \right) + \sum_{m \geq 3} \left( \frac{\rho^{m-1}}{(m-3)!} \| \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u \|_{L^2(\mathbb{R}_+^2)} \right)$$

Here we first treat the terms in (4.2), and the ones in (4.1) can be deduced similarly with simpler arguments. To do so, we use the notation  $\omega = \partial_y u$ . Then it follows from (3.1) that

$$(4.3) \quad \left\{ \begin{array}{l} \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) \omega + \left( v + \overline{u_1^{I,0}} \right) \partial_1 \omega \\ \quad + (u - u(t, x_1, 0)) \partial_y \omega + 2 \overline{\partial_3 u_3^{I,0}} \omega + \left( \partial_1 u(t, x_1, 0) + y \overline{\partial_1 \partial_3 u_3^{I,0}} \right) \partial_y v \\ \quad + (\partial_y v) \partial_1 u + \omega^2 + \overline{\partial_1 \partial_3 u_3^{I,0}} v = 0, \\ \omega|_{y=0} = \overline{\partial_1 u_1^{I,0}}(t, x_1), \\ \omega|_{t=0} = \partial_y u_{3,0}^{B,1}. \end{array} \right.$$

Thus the function, defined by

$$(4.4) \quad \varphi_m = \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \omega(t) = \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u(t),$$

solves the equation

$$\left\{ \begin{array}{l} \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) \varphi_m - a'(t) y^2 \varphi_m + \left( v + \overline{u_1^{I,0}} \right) \partial_1 \varphi_m + (u - u(t, x_1, 0)) \partial_y \varphi_m = \mathcal{R}^m(t), \\ \varphi_m|_{y=0} = \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1), \\ \varphi_m|_{t=0} = \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u_{3,0}^{B,1}, \end{array} \right.$$

where

$$\mathcal{R}^m(t) = \sum_{j=1}^{11} \mathcal{R}_j^m(t)$$

with

$$\begin{aligned} \mathcal{R}_1^m &= -4ay \partial_y \varphi_m + 4a^2 y^2 \varphi_m - 2a \varphi_m, \\ \mathcal{R}_2^m &= 2ay^2 \overline{\partial_3 u_3^{I,0}} \varphi_m + 2ay (u - u(t, x_1, 0)) \varphi_m, \\ \mathcal{R}_3^m &= \left( \partial_1 \langle x_1 \rangle^\ell \right) e^{ay^2} \left( v + \overline{u_1^{I,0}} \right) \partial_1^m \omega, \\ \mathcal{R}_4^m &= - \sum_{k=1}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \overline{\partial_1^k \partial_3 u_3^{I,0}} \right) y \partial_y \partial_1^{m-k} \omega, \\ \mathcal{R}_5^m &= - \sum_{k=1}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \partial_1^k v + \overline{\partial_1^k u_1^{I,0}} \right) \partial_1^{m-k+1} \omega, \\ \mathcal{R}_6^m &= - \sum_{k=1}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \partial_1^k u - \partial_1^k u(t, x_1, 0) \right) \partial_1^{m-k} \partial_y \omega, \\ \mathcal{R}_7^m &= - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \overline{\partial_1^k \partial_3 u_3^{I,0}} \right) \partial_1^{m-k} \omega, \\ \mathcal{R}_8^m &= - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \partial_1^{k+1} u(t, x_1, 0) + y \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}} \right) \partial_1^{m-k} \partial_y v, \\ \mathcal{R}_9^m &= - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( \partial_1^{k+1} u \right) \partial_1^{m-k} \partial_y v, \\ \mathcal{R}_{10}^m &= - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( (\partial_1^k \omega) \partial_1^{m-k} \omega + \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}} \partial_1^{m-k} v \right). \end{aligned}$$

From the first equation in (4.3), it follows that

$$(4.5) \quad \begin{aligned} & \left( \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} - (a'(t) y^2 \varphi_m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \\ & + \left( \left( v + \overline{u_1^{I,0}} \right) \partial_1 \varphi_m + (u - u(t, x_1, 0)) \partial_y \varphi_m(t), \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \\ & = (\mathcal{R}^m(t), \varphi_m(t))_{L^2(\mathbb{R}_+^2)}, \end{aligned}$$

with  $\mathcal{R}^m$  given above.

In the following lemmas, let  $0 < a(t) < a_0$  to be determined later, and let  $0 < \rho = \rho(t) \leq \min\{\rho_0/2, \tau/3\}$  be an arbitrary smooth function of  $t$ .

**Lemma 4.1.** *A constants  $C$  exists such that for any  $N \geq 3$ ,*

$$\sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_1^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \leq C \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2,$$

and

$$\sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_2^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \leq C \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{X_{\rho,a}}^4.$$

*Proof.* We have, integrating by parts,

$$(\mathcal{R}_1^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} = (4a^2 y^2 \varphi_m, \varphi_m)_{L^2(\mathbb{R}_+^2)} = 4a^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2$$

Direct verification shows

$$(\mathcal{R}_2^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \leq \left( 2a \|\overline{\partial_3 u_3^{I,0}}\|_{L^\infty} + a^2 \right) \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + 4 \|u\|_{L^\infty(\mathbb{R}_+^2)}^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2.$$

Observe

$$\|\overline{\partial_3 u_3^{I,0}}\|_{L^\infty} \leq C \|u_3^{I,0}\|_{G_\tau}$$

and

$$\sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \leq \sum_{m=3}^{\infty} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \leq |u|_{X_{\rho,a}},$$

and thus the desired results follow, completing the proof.  $\square$

**Lemma 4.2.** *There exists a constant  $C$  such that for any  $\rho$  with  $0 < \rho \leq \tau/3$ , we have*

$$\sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_8^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \leq \frac{1}{8} |u|_{Z_{\rho,a}}^2 + C \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2.$$

*Proof.* Recall  $\mathcal{R}_8^{m,j}$  can be written as, for any  $\tilde{\varepsilon} > 0$ ,

$$\begin{aligned} (\mathcal{R}_8^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} & = \left( - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} (\partial_1^{k+1} u(x_1, 0)) \partial_1^{m-k} \partial_y v, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \\ & + \left( - \sum_{k=0}^m \binom{m}{k} \langle x_1 \rangle^\ell e^{ay^2} \left( y \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}} \right) \partial_1^{m-k} \partial_y v, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \\ & \leq \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \\ & + \tilde{\varepsilon} \left[ \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}}\|_{L_{x_1}^2(\mathbb{R})} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \right]^2 \\ & + C_{\tilde{\varepsilon}} \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2. \end{aligned}$$

Then it suffices to show that

$$(4.6) \quad \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \\ \leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2,$$

and

$$(4.7) \quad \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left[ \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}}\|_{L_{x_1}^2(\mathbb{R})} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \right]^2 \\ \leq C |u|_{Z_{\rho,a}}^2.$$

We will proceed to prove the above estimate through the following steps.

**Step (a)** We begin with several estimates to be used later in the proof. Firstly in view of the definition of  $|\cdot|_{Y_{\rho,a}}$  given in (3.5), we may write

$$|u|_{Y_{\rho,a}}^2 = \sum_{m=0}^{+\infty} |u|_{Y_{\rho,a,m}}^2$$

where  $|u|_{Y_{\rho,a,m}}$  is defined by

$$|u|_{Y_{\rho,a,m}} = \begin{cases} \sum_{0 \leq j \leq 1} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}, & 0 \leq m \leq 2 \\ \sum_{0 \leq j \leq 1} (m-1)^{1/2} \rho^{-1/2} \frac{\rho^{m-1}}{(m-3)!} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}, & m \geq 3. \end{cases}$$

Thus

$$(4.8) \quad \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \leq \begin{cases} |u|_{Y_{\rho,a}}, & 0 \leq m \leq 2, \\ |u|_{Y_{\rho,a,m}} m^{-1/2} \rho^{1/2} \frac{(m-3)!}{\rho^{m-1}}, & m \geq 3, \end{cases}$$

Next from the relations (3.2) it follows that

$$\|e^{ay^2} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \leq C \|\langle x_1 \rangle^\ell e^{ay^2} \partial_y^2 u\|_{L^2(\mathbb{R}_+^2)} \leq C |u|_{Z_{\rho,a}},$$

and that for  $j \geq 1$ ,

$$\|e^{ay^2} \partial_1^j \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} = \|e^{ay^2} \partial_1^{j-1} \partial_y^2 u\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \\ \leq C \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^j \partial_y^2 u\|_{L^2(\mathbb{R}_+^2)} \\ \leq \begin{cases} C |u|_{Z_{\rho,a}}, & 1 \leq j \leq 2, \\ C |u|_{Z_{\rho,a,j}} \frac{(j-3)!}{\rho^{j-1}}, & j \geq 3, \end{cases}$$

where  $|u|_{Z_{\rho,a,k}}$  is defined by the relation  $|u|_{Z_{\rho,a}} = \sum_{k \geq 0} |u|_{Z_{\rho,a,k}}$ , so that

$$|u|_{Z_{\rho,a,k}} = \begin{cases} \sum_{1 \leq j \leq 2} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^k \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}, & 0 \leq k \leq 2 \\ \sum_{1 \leq j \leq 2} \frac{\rho^{k-1}}{(k-3)!} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^k \partial_y^j u\|_{L^2(\mathbb{R}_+^2)}, & k \geq 3. \end{cases}$$

Thus we conclude

$$(4.9) \quad \|e^{ay^2} \partial_1^j \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \leq \begin{cases} C |u|_{Z_{\rho,a}}, & 0 \leq j \leq 2, \\ C |u|_{Z_{\rho,a,j}} \frac{(j-3)!}{\rho^{j-1}}, & j \geq 3. \end{cases}$$

Using the Sobolev inequality

$$\|\langle x_1 \rangle^\ell \partial_1^j u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \leq C \|\langle x_1 \rangle^\ell \partial_1^j u\|_{L^2(\mathbb{R}_+)} + C \|\langle x_1 \rangle^\ell \partial_1^j \partial_y u\|_{L^2(\mathbb{R}_+)},$$

gives

$$(4.10) \quad \|\langle x_1 \rangle^\ell \partial_1^j u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \leq \begin{cases} C |u|_{Y_{\rho,a}}, & \text{if } 0 \leq j \leq 2, \\ C |u|_{Y_{\rho,a,j}} j^{-1/2} \rho^{1/2} \frac{(j-3)!}{\rho^{j-1}}, & \text{if } j \geq 3. \end{cases}$$

Finally,

$$(4.11) \quad \forall k \geq 0, \quad \|\langle x_1 \rangle^\ell \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}}\|_{L_{x_1}^2(\mathbb{R})} \leq C \|u_3^{I,0}\|_{\mathcal{A}_\tau} \frac{(k+3)!}{\tau^{k+3}}$$

due to (1.4).

**Step (b).** We now prove (4.7). For this purpose we use (4.11) and (4.9) to calculate

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}}\|_{L_{x_1}^2(\mathbb{R})} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \\ \leq & C \|u_3^{I,0}\|_{\mathcal{A}_\tau} \sum_{k=0}^{m-3} \frac{m!}{k!(m-k)!} \frac{(k+3)!}{\tau^{k+3}} \frac{(m-k-3)!}{\rho^{m-k-1}} |u|_{Z_{\rho,a,m-k}} \\ & + C \|u_3^{I,0}\|_{\mathcal{A}_\tau} \sum_{k=m-2}^m \frac{m!}{k!(m-k)!} \frac{(k+3)!}{\tau^{k+3}} |u|_{Z_{\rho,a,m-k}} \\ \leq & C \frac{(m-3)!}{\rho^{m-1}} \sum_{k=0}^{m-3} \frac{m^3}{k^3(m-k-2)^3} \frac{2^k \rho^k}{\tau^{k+3}} |u|_{Z_{\rho,a,m-k}} + C \frac{(m-3)!}{\rho^{m-1}} \sum_{k=m-2}^m \frac{2^k \rho^{m-1}}{\tau^{k+3}} |u|_{Z_{\rho,a,m-k}} \\ \leq & C \tau^{-3} \frac{(m-3)!}{\rho^{m-1}} \sum_{k=0}^{m-3} \frac{2^k \rho^k}{\tau^k} |u|_{Z_{\rho,a,m-k}} + C \tau^{-3} \frac{(m-3)!}{\rho^{m-1}} \sum_{k=m-2}^m \frac{2^k \rho^{m-1}}{\tau^k} |u|_{Z_{\rho,a,m-k}}, \end{aligned}$$

which yields

$$\begin{aligned} & \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left[ \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \overline{\partial_1^{k+1} \partial_3 u_3^{I,0}}\|_{L_{x_1}^2(\mathbb{R})} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \right]^2 \\ \leq & C \sum_{m=3}^N \left( \sum_{k=0}^{m-3} \frac{2^k \rho^k}{\tau^k} |u|_{Z_{\rho,a,m-k}} \right)^2 + C \sum_{m=3}^N \left( \sum_{k=m-2}^m \frac{2^k \rho^{m-1}}{\tau^k} |u|_{Z_{\rho,a,m-k}} \right)^2 \end{aligned}$$

On the other hand, by virtue of Young's inequality for discrete convolution (cf. [17, Theorem 20.18]) we have

$$\sum_{m=3}^N \left( \sum_{k=0}^{m-3} \frac{2^k \rho^k}{\tau^k} |u|_{Z_{\rho,a,m-k}} \right)^2 \leq C \left( \sum_{k=0}^N \frac{2^k \rho^k}{\tau^k} \right)^2 \sum_{k=0}^N |u|_{Z_{\rho,a,k}}^2 \leq C |u|_{Z_{\rho,a}}^2,$$

since  $\rho \leq \tau/3$ . And direct computation yields

$$\sum_{m=3}^N \left( \sum_{k=m-2}^m \frac{2^k \rho^{m-1}}{\tau^k} |u|_{Z_{\rho,a,m-k}} \right)^2 \leq C |u|_{Z_{\rho,a}}^2.$$

Then we obtain (4.7), combining the above inequalities.

**Step (c).** Now we check (4.6) and write

$$\begin{aligned} & \sum_{k=0}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \\ \leq & S_1 + S_2 + S_3 \end{aligned}$$

with

$$\begin{aligned} S_1 &= \sum_{k=0}^2 \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}, \\ S_2 &= \sum_{k=3}^{m-3} \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \end{aligned}$$

and

$$S_3 = \sum_{k=m-2}^m \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}.$$

For the term  $S_{2,m}$ , we use (4.8), (4.9) and (4.10) to compute

$$\begin{aligned}
S_{2,m} &= \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \sum_{k=3}^{m-3} \binom{m}{k} \|\langle x_1 \rangle^\ell \partial_1^{k+1} u\|_{L_y^\infty(\mathbb{R}_+; L_{x_1}^2(\mathbb{R}))} \|e^{ay^2} \partial_1^{m-k} \partial_y v\|_{L_y^2(\mathbb{R}_+; L_{x_1}^\infty(\mathbb{R}))} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \\
&\leq C \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \sum_{k=3}^{m-3} \frac{m!}{k!(m-k)!} \left[ k^{-1/2} \rho^{1/2} \frac{(k-2)!}{\rho^k} |u|_{Y_{\rho,a,k+1}} \right] \frac{(m-k-3)!}{\rho^{m-k-1}} |u|_{Z_{\rho,a,m-k}} \\
&\quad \times m^{-1/2} \rho^{1/2} \frac{(m-3)!}{\rho^{m-1}} |u|_{Y_{\rho,a,m}} \\
&\leq C \rho |u|_{Y_{\rho,a,m}} \sum_{k=3}^{m-3} \frac{m^3}{k^2(m-k-2)^3} k^{-1/2} m^{-1/2} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \\
&\leq C \rho |u|_{Y_{\rho,a,m}} \left( \sum_{k=3}^{m-3} \frac{m^3}{k^2(m-k-2)^3} k^{-1/2} m^{-1/2} |u|_{Y_{\rho,a,k+1}}^2 \right)^{1/2} \\
&\quad \times \left( \sum_{k=3}^{m-3} \frac{m^3}{k^2(m-k-2)^3} k^{-1/2} m^{-1/2} |u|_{Z_{\rho,a,m-k}}^2 \right)^{1/2} \\
&\leq C \rho |u|_{Z_{\rho,a,m}} |u|_{Y_{\rho,a,m}}^2,
\end{aligned}$$

and thus

$$\begin{aligned}
\sum_{m=3}^N S_{2,m} &\leq C \rho \left( \sum_{m=3}^N |u|_{Y_{\rho,a,m}}^2 \right)^{1/2} \left( \sum_{m=3}^N \left[ \sum_{k=3}^{m-3} \frac{m^3}{k^2(m-k-2)^3} k^{-1/2} m^{-1/2} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \right]^2 \right)^{1/2} \\
&\leq C \rho |u|_{Y_{\rho,a}} \left( \sum_{m=3}^N \left[ \sum_{k=3}^{m-3} \frac{1}{k^2} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \right]^2 \right)^{1/2} \\
&\quad + C \rho |u|_{Y_{\rho,a}} \left( \sum_{m=3}^N \left[ \sum_{k=3}^{m-3} \frac{1}{(m-k-2)^3} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \right]^2 \right)^{1/2}
\end{aligned}$$

the last inequality following from the fact that

$$\forall 3 \leq k \leq m-3, \quad \frac{m^3}{k^2(m-k-2)^3} k^{-1/2} m^{-1/2} \leq C \left( \frac{1}{k^2} + \frac{1}{(m-k-2)^3} \right).$$

Moreover, by virtue of Young's inequality for discrete convolution (cf. [17, Theorem 20.18]) we obtain

$$\begin{aligned}
\left( \sum_{m=3}^N \left[ \sum_{k=3}^{m-3} \frac{1}{k^2} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \right]^2 \right)^{1/2} &\leq C \left( \sum_{m=3}^{+\infty} |u|_{Z_{\rho,a,m}}^2 \right)^{1/2} \sum_{k=3}^{+\infty} \frac{1}{k^2} |u|_{Y_{\rho,a,k}} \\
&\leq C |u|_{Z_{\rho,a}} \left( \sum_{k=1}^{+\infty} |u|_{Y_{\rho,a,k}}^2 \right)^{1/2} \left( \sum_{k=1}^{+\infty} \frac{1}{k^4} \right)^{1/2} \\
&\leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}.
\end{aligned}$$

Similarly

$$\left( \sum_{m=3}^N \left[ \sum_{k=3}^{m-3} \frac{1}{(m-k-2)^3} |u|_{Y_{\rho,a,k+1}} |u|_{Z_{\rho,a,m-k}} \right]^2 \right)^{1/2} \leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}.$$

Combining these inequality we conclude

$$\sum_{m=3}^N S_{2,m} \leq C \rho |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2 \leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2.$$

The estimates on the rest two terms  $S_1$  and  $S_3$  can be deduced similarly and directly, and we have

$$\sum_{m=3}^N (S_{1,m} + S_{3,m}) \leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2,$$

proving (4.6). The proof of Lemma 4.2 is complete.  $\square$

**Lemma 4.3.** *A constant  $C$  exists such that*

$$\begin{aligned} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_3^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C \left( |u|_{X_{\rho,a,m}}^3 + |u|_{X_{\rho,a,m}}^2 \right), \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_4^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq \frac{1}{8} |u|_{Z_{\rho,a}}^2 + C \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2, \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_5^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C \left( |u|_{X_{\rho,a,m}}^3 + |u|_{X_{\rho,a,m}}^2 \right), \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_6^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2, \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_7^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C |u|_{X_{\rho,a}}^2, \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_9^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2, \\ \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 (\mathcal{R}_{10}^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} &\leq C \left( |u|_{X_{\rho,a}}^3 + |u|_{X_{\rho,a}}^2 \right). \end{aligned}$$

*Proof.* The treatment of  $\mathcal{R}_6, \mathcal{R}_9$  is exactly the same as in the proof of (4.6). The other terms can be deduced similarly by following the proof in Lemma 4.2 with slightly changes, and the arguments here will be simpler since there is no the highest derivative  $\partial_1^{m+1}$  involved. This means we can perform the estimates with the norm  $Y_{\rho,a}$  in Lemma 4.2 replaced by  $X_{\rho,a}$  here. So we omit the proof for brevity.  $\square$

Combining the estimates in Lemma 4.1-Lemma 4.3, we have

**Corollary 4.4.** *There are two constants  $C, C_0$  such that*

$$\begin{aligned} &\sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left( \sum_{k=1}^{10} (\mathcal{R}_k^m, \varphi_m)_{L^2(\mathbb{R}_+^2)} \right) \\ &\leq \frac{1}{4} |u|_{Z_{\rho,a}}^2 + C_0 \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2 + C \left( |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^4 \right). \end{aligned}$$

**Lemma 4.5.** *We have*

$$\begin{aligned} &\left( \left( \partial_t - \partial_y^2 + \overline{\partial_3 u_3^{I,0}} y \partial_y \right) \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} - \left( a'(t) y^2 \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \\ &\quad + \left( \left( v + \overline{u_1^{I,0}} \right) \partial_1 \varphi_m + (u - u(t, x_1, 0)) \partial_y \varphi_m(t), \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \\ &\geq \frac{1}{2} \frac{d}{dt} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 - a'(t) \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \\ &\quad + \frac{d}{dt} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 - C \left( \frac{(m-3)!}{\rho^{m-1}} \right)^2 |u|_{X_{\rho,a,m}}^2. \end{aligned}$$



*Proof.* Firstly we calculate, integrating by parts and using the relation (3.2),

$$(4.12) \quad \left| \left( \overline{\partial_3 u_3^{I,0}} y \partial_y \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \right| + \left| \left( (v + \overline{u_1^{I,0}}) \partial_1 \varphi_m + (u - u(t, x_1, 0)) \partial_y \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} \right| \\ \leq \frac{1}{2} \left( \left\| \overline{\partial_3 u_3^{I,0}} \right\|_{L^\infty} + \left\| \overline{\partial_3 u_1^{I,0}} \right\|_{L^\infty} \right) \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2.$$

Integrating by parts and using the boundary condition in (4.3), we have

$$(4.13) \quad \left( (\partial_t - \partial_y^2) \varphi_m, \varphi_m \right)_{L^2(\mathbb{R}_+^2)} = \frac{1}{2} \frac{d}{dt} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \\ + \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) (\partial_y \varphi_m)(t, x_1, 0) dx_1.$$

Now we check the boundary value of  $\partial_y \varphi$ . In view of (4.4) we see

$$\partial_y \varphi_m|_{y=0} = \langle x_1 \rangle^\ell \partial_1^m \partial_y^2 u|_{y=0}.$$

And moreover, using the relation

$$\langle x_1 \rangle^\ell \partial_y^2 u|_{y=0} = \partial_t u(x_1, 0) - \overline{\partial_3 u_3^{I,0}} u(x_1, 0) + \overline{u_1^{I,0}} (\partial_1 u)(x_1, 0)$$

which follows from (3.1), we conclude

$$\partial_y \varphi_m|_{y=0} = \partial_t \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) - \langle x_1 \rangle^\ell \partial_1^m \left( \overline{\partial_3 u_3^{I,0}} u(t, x_1, 0) \right) + \langle x_1 \rangle^\ell \partial_1^m \left( \overline{u_1^{I,0}} (\partial_1 u)(t, x_1, 0) \right).$$

As a result,

$$\int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell (\partial_y \varphi_m)(t, x_1, 0) dx_1 \\ = \frac{d}{dt} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ - \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_t \partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ - \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \partial_1^m \left( \overline{\partial_3 u_3^{I,0}} u(x_1, 0) \right) dx_1 \\ + \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \partial_1^m \left( \overline{u_1^{I,0}} (\partial_1 u)(t, x_1, 0) \right) dx_1.$$

Moreover, In view of (1.5), we can repeat the arguments in Lemma 4.2 and Lemma 4.3, to obtain, observing  $\rho < \tau/4$ ,

$$\left| \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_t \partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \right| \leq C \left( \frac{(m-3)!}{\rho^{m-1}} \right)^2 \|u_1^{I,0}\|_{\mathcal{A}_\tau} |u|_{X_{\rho,a,m}},$$

$$\left| \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \partial_1^m \left( \overline{\partial_3 u_3^{I,0}} u(x_1, 0) \right) dx_1 \right| \leq C \left( \frac{(m-3)!}{\rho^{m-1}} \right)^2 \|u_1^{I,0}\|_{\mathcal{A}_\tau} \|u_3^{I,0}\|_{\mathcal{A}_\tau} |u|_{X_{\rho,a,m}}$$

and

$$\left| \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \partial_1^m \left( \overline{u_1^{I,0}} (\partial_1 u)(t, x_1, 0) \right) dx_1 \right| \leq C \left( \frac{(m-3)!}{\rho^{m-1}} \right)^2 \|u_1^{I,0}\|_{\mathcal{A}_\tau}^2 |u|_{X_{\rho,a,m}}.$$

Combing these inequalities above, we conclude

$$\int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell (\partial_y \varphi_m)(t, x_1, 0) dx_1 \\ \geq \frac{d}{dt} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 - C \left( \frac{(m-3)!}{\rho^{m-1}} \right)^2 |u|_{X_{\rho,a,m}},$$

which, along with (4.12) and (4.13), yields the conclusion, completing the proof.  $\square$

**Lemma 4.6.** *Let  $a(t) = a_0 - (2a_0^2 + C_0)t$  with  $C_0$  the constants given in Corollary 4.4. Then for any  $N$ ,*

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \frac{1}{2} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^2 u(t)\|_{L^2(\mathbb{R}_+^2)}^2 \\ & \quad - \rho'(t) \sum_{m=3}^N \left( (m-1)^{1/2} \rho^{-1/2} \frac{\rho^{m-1}}{(m-3)!} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \right)^2 \\ & \leq \frac{1}{4} |u|_{Z_{\rho,a}}^2 - \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \quad + C \left( |\rho'| \rho^{-2} |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^4 \right). \end{aligned}$$

*Proof.* Using the equality (4.5) and Lemma 4.5, we obtain

$$\begin{aligned} & \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left[ \frac{1}{2} \frac{d}{dt} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 - a'(t) \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \right] \\ & \leq - \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \quad + \sum_{m=3}^N (2m-2) \frac{\rho'(t) \rho^{2m-3}}{[(m-3)!]^2} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 + |u|_{\rho,a} \\ & \quad + \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left( \sum_{k=1}^{10} (\mathcal{R}_k^m(t), \varphi_m(t))_{L^2(\mathbb{R}_+^2)} \right) \\ & \leq - \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \quad + \sum_{m=3}^N (2m-2) \frac{\rho'(t) \rho^{2m-3}}{[(m-3)!]^2} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 + |u|_{\rho,a} \\ & \quad + \frac{1}{4} |u|_{Z_{\rho,a}}^2 + C_0 \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2 + C \left( |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^4 \right), \end{aligned}$$

the last inequality following from Corollary 4.4. On the other hand,

$$\begin{aligned} & \sum_{m=3}^N (2m-2) \frac{\rho'(t) \rho^{2m-3}}{[(m-3)!]^2} \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \leq C \sum_{m=3}^N \frac{2^m \rho'(t) \rho^{2m-3}}{[(m-3)!]^2} \|\langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}\|_{L^2(\mathbb{R}_{x_1})} \left( \|\langle x_1 \rangle^\ell \partial_1^m u\|_{L^2} + \|\langle x_1 \rangle^\ell \partial_1^m \partial_y u\|_{L^2} \right) \\ & \leq C \sum_{m=3}^N \frac{2^m \rho'(t) \rho^{m-2}}{\tau^{m+3}} \|u_1^{I,0}\|_{G_\tau} \left[ \frac{\rho^{m-1}}{(m-3)!} \left( \|\langle x_1 \rangle^\ell \partial_1^m u\|_{L^2} + \|\langle x_1 \rangle^\ell \partial_1^m \partial_y u\|_{L^2} \right) \right] \\ & \leq C |\rho'| \rho^{-2} |u|_{X_{\rho,a}} \end{aligned}$$

the last inequality using the fact that  $\rho < \tau/3$ . As a result, combining the equalities above yields

$$\begin{aligned} & \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left[ \frac{1}{2} \frac{d}{dt} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 - a'(t) \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \right] \\ & \leq \frac{1}{4} |u|_{Z_{\rho,a}}^2 - \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \quad + C_0 \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2 + C \left( |\rho'| \rho^{-2} |u|_{X_{\rho,a}} + |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^4 \right). \end{aligned}$$

Moreover from the relations

$$\|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \geq \frac{1}{2} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^2 u(t)\|_{L^2(\mathbb{R}_+^2)}^2 - 2a_0^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2$$

and

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 - a'(t) \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \\ & \quad - \rho'(t) \left( (m-1)^{1/2} \rho^{-1/2} \frac{\rho^{m-1}}{(m-3)!} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \right)^2 \\ &= \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \left[ \frac{1}{2} \frac{d}{dt} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \|\partial_y \varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 - a'(t) \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \right], \end{aligned}$$

it follows that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + \frac{1}{2} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^2 u(t)\|_{L^2(\mathbb{R}_+^2)}^2 \\ & \quad - (a'(t) + 2a_0^2) \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 \\ & \quad - \rho'(t) \sum_{m=3}^N \left( (m-1)^{1/2} \rho^{-1/2} \frac{\rho^{m-1}}{(m-3)!} \|\varphi_m\|_{L^2(\mathbb{R}_+^2)} \right)^2 \\ & \leq \frac{1}{4} |u|_{Z_{\rho,a}}^2 - \frac{d}{dt} \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \int_{\mathbb{R}_{x_1}} \langle x_1 \rangle^\ell \overline{\partial_1^{m+1} u_1^{I,0}}(t, x_1) \langle x_1 \rangle^\ell \partial_1^m u(t, x_1, 0) dx_1 \\ & \quad + C_0 \sum_{m=3}^N \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|y\varphi_m\|_{L^2(\mathbb{R}_+^2)}^2 + C |u|_{Z_{\rho,a}} |u|_{Y_{\rho,a}}^2 + C \left( |\rho'| \rho^{-2} |u|_{X_{\rho,a}} + |u|_{X_{\rho,a}}^2 + |u|_{X_{\rho,a}}^4 \right). \end{aligned}$$

Now observing  $a(t) = a_0 - (2a_0^2 + C_0)t$ , we complete the proof.  $\square$

**Completion of the proof of Proposition 3.4.** By Lemma 4.6, we integrate both sides over  $[0, t] \subset [0, T]$  and then let  $N \rightarrow +\infty$ , to obtain that for any  $t \in [0, T]$ ,

$$\begin{aligned} & \sum_{m=3}^{+\infty} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m(t)\|_{L^2(\mathbb{R}_+^2)}^2 + \int_0^t \sum_{m=3}^{+\infty} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^2 u(t)\|_{L^2(\mathbb{R}_+^2)}^2 dt \\ & \quad - \int_0^t \rho'(t) \sum_{m=3}^{+\infty} \frac{m-1}{\rho} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\varphi_m(t)\|_{L^2(\mathbb{R}_+^2)}^2 dt \\ & \leq |u_0|_{X_{\rho_0, a_0}}^2 + \frac{1}{2} \int_0^t |u(t)|_{Z_{\rho,a}}^2 dt + C \int_0^t \left( |\rho'(t)| \rho^{-2} |u(t)|_{X_{\rho,a}} + |u(t)|_{X_{\rho,a}}^2 + |u(t)|_{X_{\rho,a}}^4 \right) dt \\ & \quad + C \int_0^t |u(t)|_{Z_{\rho,a}} |u(t)|_{Y_{\rho,a}}^2 dt. \end{aligned}$$

Direct computation also gives

$$\begin{aligned} & \sum_{m \leq 2} \|\varphi_m(t)\|_{L^2(\mathbb{R}_+^2)}^2 + \int_0^t \sum_{m \leq 2} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y^2 u(t)\|_{L^2(\mathbb{R}_+^2)}^2 dt - \int_0^t \rho'(t) \sum_{m \leq 2} \|\varphi_m(t)\|_{L^2(\mathbb{R}_+^2)}^2 dt \\ & \leq |u_0|_{X_{\rho_0, a_0}}^2 + \frac{1}{2} \int_0^t |u(t)|_{Z_{\rho,a}}^2 dt + C \int_0^t \left( |\rho'(t)| \rho^{-2} |u(t)|_{X_{\rho,a}} + |u(t)|_{X_{\rho,a}}^2 + |u(t)|_{X_{\rho,a}}^4 \right) dt \\ & \quad + C \int_0^t |u(t)|_{Z_{\rho,a}} |u(t)|_{Y_{\rho,a}}^2 dt. \end{aligned}$$

Similarly, using the notation

$$\psi_m = \langle x_1 \rangle^\ell e^{ay^2} \partial_1^m u(t),$$

we can deduce, following the proof of the above two inequalities with slight modification and simpler arguments,

$$\begin{aligned}
& \sum_{m \leq 2} \|\psi_m(t)\|_{L^2}^2 + \sum_{m=3}^{+\infty} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\psi_m(t)\|_{L^2(\mathbb{R}_+^2)}^2 \\
& + \int_0^T \left( \sum_{m \leq 2} \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u(t)\|_{L^2(\mathbb{R}_+^2)}^2 + \sum_{m=3}^{+\infty} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\langle x_1 \rangle^\ell e^{ay^2} \partial_1^m \partial_y u(t)\|_{L^2(\mathbb{R}_+^2)}^2 \right) dt \\
& - \int_0^T \rho'(t) \left( \sum_{m \leq 2} \|\psi_m\|_{L^2}^2 + \sum_{m=3}^{+\infty} \frac{m-1}{\rho} \left( \frac{\rho^{m-1}}{(m-3)!} \right)^2 \|\psi_m\|_{L^2(\mathbb{R}_+^2)}^2 \right) dt \\
\leq & |u_0|_{X_{\rho_0, a_0}}^2 + \frac{1}{2} \int_0^T |u(t)|_{Z_{\rho, a}}^2 dt + C \int_0^T \left( |\rho'(t)| \rho^{-2} |u(t)|_{X_{\rho, a}} + |u(t)|_{X_{\rho, a}}^2 + |u(t)|_{X_{\rho, a}}^4 \right) dt \\
& + C \int_0^T |u(t)|_{Z_{\rho, a}} |u(t)|_{Y_{\rho, a}}^2 dt
\end{aligned}$$

Combining these inequalities we conclude, observing the definition of  $|\cdot|_{X_{\rho, a}}$ ,  $|\cdot|_{Y_{\rho, a}}$  and  $|\cdot|_{Z_{\rho, a}}$  and any  $\rho \leq \min\{\rho_0, \tau/3\}$ ,

$$\begin{aligned}
& |u(t)|_{X_{\rho, a}}^2 + \int_0^T |u(t)|_{Z_{\rho, a}}^2 dt - \int_0^T \rho'(t) |u(t)|_{Y_{\rho, a}}^2 dt \\
\leq & |u_0|_{X_{\rho_0, a_0}}^2 + \frac{1}{2} \int_0^T |u(t)|_{Z_{\rho, a}}^2 dt + C \int_0^T \left( |\rho'(t)| \rho^{-2} |u(t)|_{X_{\rho, a}} + |u(t)|_{X_{\rho, a}}^2 + |u(t)|_{X_{\rho, a}}^4 \right) dt \\
& + C \int_0^T |u(t)|_{Z_{\rho, a}} |u(t)|_{Y_{\rho, a}}^2 dt
\end{aligned}$$

Thus Claim (3.6) follows and the proof is complete.  $\square$

## 5. EXISTENCE OF SOLUTION FOR SECOND COMPONENT

In this section, we determine the second component  $\mathcal{U}_2^{p,0}$  by solving the parabolic-type equation

$$\begin{cases} \partial_t \mathcal{U}_2^{p,0} - \partial_y^2 \mathcal{U}_2^{p,0} + \mathcal{U}_1^{p,0} \partial_1 \mathcal{U}_2^{p,0} + \mathcal{U}_3^{p,1} \partial_y \mathcal{U}_2^{p,0} = 0, \\ \mathcal{U}_2^{p,0}(t, x_1, 0) = 0, \quad \lim_{y \rightarrow +\infty} \mathcal{U}_2^{p,0}(t, x_1, y) = \overline{u_2^{I,0}}(x_1), \\ \mathcal{U}_2^{p,0}(0, x_1, y) = u_{0,2}^{B,0}(x_1, y) + \overline{u_{0,2}^{I,0}}(x_1). \end{cases}$$

We recall that

$$(5.1) \quad \overline{\partial_t u_2^{I,0}} + \overline{u_1^{I,0}} \partial_1 \overline{u_2^{I,0}} = 0.$$

Then, the system (P2) becomes

$$(P2bis) \quad \begin{cases} \partial_t u_2^{B,0} - \partial_y^2 u_2^{B,0} + \left( u_1^{B,0} + \overline{u_1^{I,0}} \right) \partial_1 u_2^{B,0} + \left( u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \right) \partial_y u_2^{B,0} + \overline{\partial_1 u_2^{I,0}} u_1^{B,0} = 0, \\ \partial_2 u_2^{B,0} = 0, \\ u_2^{B,0}(t, x_1, 0) = -\overline{u_2^{I,0}}, \quad \lim_{y \rightarrow +\infty} u_2^{B,0}(t, x_1, y) = 0, \\ u_2^{B,0}(0, x_1, y) = u_{0,2}^{B,0}(x_1, y). \end{cases}$$

We have the following results

**Theorem 5.1.** *Let  $\rho_0 > 0$ ,  $a_0 > 0$ . For any initial data  $u_{2,0}^{B,0} \in X_{\rho_0, a_0}$ , there exists  $T > 0$ ,  $\tau > 0$  and  $0 < a < a_0$ , such that the system (P2bis) admits a unique solution  $u_2^{B,0} \in L^\infty([0, T], X_{\rho_0, a})$ .*

**Proof.** In order to prove Theorem 5.1, the idea is to define an auxiliary function

$$v = u_2^{B,0} + e^{-2a_0 y^2} \overline{u_2^{I,0}},$$

which satisfies the following boundary conditions

$$v(t, x_1, 0) = \lim_{y \rightarrow +\infty} v(t, x_1, y) = 0.$$

Then, the first equation of the system (P2bis) becomes

$$\begin{aligned} \partial_t \left( v - e^{-2a_0 y^2} \overline{u_2^{I,0}} \right) - \partial_y^2 \left( v - e^{-2a_0 y^2} \overline{u_2^{I,0}} \right) + \left( u_1^{B,0} + \overline{u_1^{I,0}} \right) \partial_1 \left( v - e^{-2a_0 y^2} \overline{u_2^{I,0}} \right) \\ + \left( u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \right) \partial_y \left( v - e^{-2a_0 y^2} \overline{u_2^{I,0}} \right) + \overline{\partial_1 u_2^{I,0}} u_1^{B,0} = 0. \end{aligned}$$

Using (5.1), we can rewrite the system (P2bis) as

$$(P2_v) \quad \begin{cases} \partial_t v - \partial_y^2 v + \left( u_1^{B,0} + \overline{u_1^{I,0}} \right) \partial_1 v + \left( u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \right) \partial_y v + R = 0, \\ \partial_2 v = 0, \\ v(t, x_1, 0) = 0, \quad \lim_{y \rightarrow +\infty} u_2^{B,0}(t, x_1, y) = 0, \\ v(0, x_1, y) = u_{0,2}^{B,0}(x_1, y) + e^{-2a_0 y^2} \overline{u_{0,2}^{I,0}}(x_1), \end{cases}$$

where

$$R = (16a_0^2 y^2 - 4a_0) e^{-2a_0 y^2} \overline{u_2^{I,0}} + 4a_0 \left( u_3^{B,1} + \overline{u_3^{I,1}} + y \overline{\partial_3 u_3^{I,0}} \right) y e^{-2a_0 y^2} \overline{u_2^{I,0}} + \left( 1 - e^{-2a_0 y^2} \right) \overline{\partial_1 u_2^{I,0}} u_1^{B,0}.$$

We remark that the system  $(P2_v)$  is in the same form as the system (3.1) with Dirichlet boundary conditions. Thus, we can prove Theorem 5.1 in the same way (with a lot of simplifications) as we did to prove Theorem 3.1.

**Acknowledgements.** The research of the first author was supported by NSF of China(11422106) and Fok Ying Tung Education Foundation (151001), and he would like to thank the invitation of “laboratoire de mathématiques Raphaël Salem” of the Université de Rouen. The second author would like to express his sincere thanks to School of mathematics and statistics of Wuhan University for the invitations. The research of the last author is supported partially by “The Fundamental Research Funds for Central Universities of China”.

#### REFERENCES

- [1] Alexandre R., Wang Y., Xu C.-J. and Yang T., Well-posedness of The Prandtl Equation in Sobolev Spaces, *J. Amer. Math. Soc.*, **28** (2015), 745-784,2015.
- [2] J.-Y. Chemin, B. Desjardins, I. Gallagher and E. Grenier, Anisotropy and dispersion in rotating fluids, *Nonlinear Partial Differential Equations and their application, Collège de France Seminar, Studies in Mathematics and its Applications*, **31** (2002) 171-191.
- [3] J.-Y. Chemin, B. Desjardins, I. Gallagher and E. Grenier, Fluids with anisotropic viscosity, Special issue for R. Temam’s 60th birthday, *M2AN. Mathematical Modelling and Numerical Analysis*, **34** (2000) 315-335.
- [4] J.-Y. Chemin, B. Desjardins, I. Gallagher and E. Grenier, Ekman boundary layers in rotating fluids, *ESAIM Controle optimal et calcul des variations, A tribute to J.-L. Lions*, **8** (2002) 441-466.
- [5] J.-Y. Chemin, B. Desjardins, I. Gallagher and E. Grenier, *Mathematical Geophysics: An introduction to rotating fluids and to the Navier-Stokes equations, Oxford University Press*, 2006.
- [6] Cheng F. and Li W.-X., Gevrey regularity in the weighted function space for incompressible Euler equation on half-space. *to appear Acta Math Scien.*
- [7] A.-L. Dalibard, D. Gérard-Varet, Nonlinear boundary layers for rotating fluids (submitted).
- [8] E, W.: Boundary layer theory and the zero-viscosity limit of the Navier-Stokes equation. *Acta Math. Sin. (Engl. Ser.)* **16** (2000) 207-218.
- [9] E, W. & Enquist, B.: Blow up of solutions of the unsteady Prandtl’s equation, *Comm. Pure Appl. Math.*, **50** (1997) 1287-1293.
- [10] I. Gallagher, Applications of Schochet’s Methods to Parabolic Equation, *Journal de Mathématiques Pures et Appliquées*, **77** (1998) 989-1054.
- [11] I. Gallagher and L. Saint-Raymond, Weak convergence results for inhomogeneous rotating fluid equations, *Journal d’Analyse Mathématique*, **99** (2006) 1-34.
- [12] D. Gérard-Varet, & E. Dormy, On the ill-posedness of the Prandtl equation, *J. Amer. Math. Soc.*, **23** (2010) 591-609.
- [13] D. Gérard-Varet, Y. Maekawa, N. Masmoudi. Gevrey Stability of Prandtl Expansions for 2D Navier-Stokes. Preprint. arXiv:1607.06434.
- [14] S.Gong, Y. Guo, Y.-G.Wang, Ekman boundary layer expansions of Navier-Stokes equations with rotation. *Bull. Inst. Math. Acad. Sin. (N.S.)* **10** (2015), no. 3, 375-392.
- [15] E. Grenier, & N. Masmoudi, Ekman layers of rotating fluids, the case of well prepared initial data. *Comm. Partial Differential Equations* **22** (1997) 953-975.

- [16] Y. Guo, Toan T. Nguyen. Prandtl boundary layer expansions of steady Navier-Stokes flows over a moving plate. Preprint. arXiv:1411.6984
- [17] E. Hewitt, K. Ross, Abstract harmonic analysis. Vol. I. Structure of topological groups, integration theory, group representations. Second edition. Grundlehren der Mathematischen Wissenschaften, 115. Springer-Verlag, Berlin-New York, 1979
- [18] D. Iftimie, A uniqueness result for the Navier-Stokes equations with vanishing vertical viscosity, *SIAM Journal on Mathematical Analysis*, **33** (2002) 1483-1493.
- [19] T. Kato, Remarks on zero viscosity limit for non-stationary Navier-Stokes flows with boundary, in Seminar on Nonlinear PDE, ed. S. S. Chern, MSRI, 1984.
- [20] T. Kato, Non-stationary flows of viscous and ideal fluids in  $\mathbb{R}^3$ , *J. Func. Anal.* **9** (1972) 296305.
- [21] I. Kukavica, V. Vicol, On the analyticity and Gevrey class regularity up to the boundary for the Euler equation, *Nonlinearity* **24** (2011) 765-796.
- [22] J.-L. Lions, Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod; Gauthier-Villars, Paris, 1969
- [23] Y. Maekawa, On the Inviscid Limit Problem of the Vorticity Equations for Viscous Incompressible Flows in the Half-Plane, *Comm. Pure Appl. Math.*, **67** (2014) 1045-1128.
- [24] N. Masmoudi, Ekman layers of rotating fluids: The case of general initial data. *Comm. Pure Appl. Math.* **53** (2000) 432-483.
- [25] N. Masmoudi, The Euler Limit of the Navier-Stokes Equations, and Rotating Fluids with Boundary. *Arch. Rational Mech. Anal.* **142** (1998) 375394.
- [26] N. Masmoudi and T. K. Wong. Local-in-time existence and uniqueness of solutions to the Prandtl equations by energy methods. *Comm. Pure Appl. Math.*, **68** (2015) 1683-1741.
- [27] V.-S. Ngo, Rotating Fluids with small viscosity, *International Mathematics Research Notices IMRN*, **10** (2009) 1860-1890.
- [28] M. Paicu, Étude asymptotique pour les fluides anisotropes en rotation rapide dans le cas périodique, *Journal de Mathématiques Pures et Appliquées*, **83** (2004) 163-242.
- [29] O. A. Oleinik, V. N. Samokhin, *Mathematical Models in Boundary Layers Theory*. Chapman & Hall/CRC, 1999.
- [30] J. Pedlovsky, Geophysical Fluid Dynamics. Springer-Verlag (1979).
- [31] M. Sammartino, R. E. Caflisch, Zero viscosity limit for analytic solutions of the Navier-Stokes equations on a half-space, I. Existence for Euler and Prandtl equations. *Comm. Math. Phys.*, 192(1998) 433-461; II. Construction of the Navier-Stokes solution. *Comm. Math. Phys.*, **192** (1998) 463-491.
- [32] H. Swann, The convergence with vanishing viscosity of non-stationary Navier-Stokes flow to ideal flow in  $\mathbb{R}^3$ , *Trans. Amer. Math. Soc.* **157** (1971) 373397.
- [33] K. Stewartson, On almost rigid rotations, Part 2, *J. Fluid Mech.*, **3** (1957) 17-26.
- [34] G. I. Taylor, Experiments on the motion of solid bodies in rotating fluids, *Proc. Roy. Soc. A* **104** (1923) 213218.
- [35] A.I. Van de Vooren, The connection between Ekman and Stewartson layers for a rotating disk, *J. of Engineering Mathematics*, **27** (1993) 189-207.
- [36] X.-P. Wang, Y.-G. Wang, Z. Xin, Boundary layers in incompressible Navier-Stokes equations with Navier boundary conditions for the vanishing viscosity limit. *Commun. Math. Sci.* 8 (2010), no. 4, 965-998.
- [37] Y.L. Xiao, Z.P. Xin, On the vanishing viscosity limit for the 3D Navier-Stokes equations with a slip boundary condition. *Comm. Pure Appl. Math.* 60 (2007), no. 7, 1027-1055
- [38] V. I. Yudovich, A two-dimensional non-stationary problem on the flow of an ideal incompressible fluid through a given region. *Mat. Sb.* 4 (1964), no. 64, 562-588

(W.-X. Li) SCHOOL OF MATHEMATICS AND STATISTICS, AND COMPUTATIONAL SCIENCE HUBEI KEY LABORATORY, WUHAN UNIVERSITY, WUHAN 430072, CHINA

*E-mail address:* wei-xi.li@whu.edu.cn

(V.-S. Ngo) UNIVERSITÉ DE ROUEN, CNRS UMR 6085, LABORATOIRE DE MATHÉMATIQUES, 76801 SAINT-ETIENNE DU ROUVRAY, FRANCE

*E-mail address:* van-sang.ngo@univ-rouen.fr

(C.-J. Xu) UNIVERSITÉ DE ROUEN, CNRS UMR 6085, LABORATOIRE DE MATHÉMATIQUES, 76801 SAINT-ETIENNE DU ROUVRAY, FRANCE, AND, SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, CHINA

*E-mail address:* Chao-Jiang.Xu@univ-rouen.fr