



HAL
open science

Pigments produced by the bacteria belonging to the genus *Arthrobacter*

Nuthathai Sutthiwong, Yanis Caro, Mireille Fouillaud, Philippe Laurent, A.
Valla, Laurent Dufossé

► **To cite this version:**

Nuthathai Sutthiwong, Yanis Caro, Mireille Fouillaud, Philippe Laurent, A. Valla, et al.. Pigments produced by the bacteria belonging to the genus *Arthrobacter*. 7th International Congress of Pigments in Food – New technologies towards health, through colors, Jun 2013, Novara, Italy. 2016. hal-01397507

HAL Id: hal-01397507

<https://hal.science/hal-01397507v1>

Submitted on 16 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

Pigments produced by the bacteria belonging to the genus *Arthrobacter*

Sutthiwong N.^{1, 2}, Caro Y.², Fouillaud M.², Laurent P.³, Valla A.⁴, Dufossé L.^{2, 5}

¹ Agricultural Technology Department, Thailand Institute of Scientific and Technological Research, Pathum Thani, Thailand

² Laboratoire de Chimie des Substances Naturelles et des Sciences des Aliments, Université de La Réunion, ESIROI Agroalimentaire, Sainte-Clotilde, Ile de La Réunion, France

³ Laboratoire de Chimie des Substances Naturelles et des Sciences des Aliments, Département de Génie Biologique, IUT, Université de La Réunion, Saint-Pierre, Ile de La Réunion, France

⁴ FRE 2125 CNRS, Chimie et Biologie des Substances Naturelles, 6 rue de l'Université, Quimper, France

⁵ Laboratoire ANTIQX, Université de Bretagne Occidentale, Pôle Universitaire Pierre-Jakez Hélias, Quimper, France E-mail: laurent.dufosse@univ-reunion.fr



INTRODUCTION

For several decades, pigments have been used as a taxonomic tool for the identification and classification of bacteria. Nowadays, pigment producing microorganisms attract wide interest in many scientific disciplines because of their biotechnological potential. With the growing concern in microbial pigments because of factors such as production independent from seasons and geographical conditions, novel combinations of microorganisms and pigments that can be extracted from the biomass or the culture medium are being evaluated.

The genus *Arthrobacter* is one among the most diverse microbial groups which have been found to produce pigments. Most of bacteria in this genus produce a range of pigments with orange, yellow, blue, green or red hues. At the present time, 80 species in this genus have been accepted by taxonomists. However, the purification and characterization of pigments produced by bacteria belonging to the genus *Arthrobacter* have not been frequently conducted up to the complete description of the chemical structures and the role(s) of pigments in these strains.

Carotenoids

Two psychrophilic bacteria, *Arthrobacter glacialis* and *Arthrobacter flavus* sp. nov., have been discovered as yellow pigments producers. Their pigments were characterized as three C₅₀-carotenoids with molecular formula C₅₀H₇₈O₂ (Fig. 2). More recently, *Arthrobacter arilaitensis*, one of the major bacterial species found at the surface of smear-ripened cheeses, has been reported as a yellow pigment producer, pigments which were tentatively identified as carotenoids. Furthermore, the carotenoids excreted by this strain may also belong to the C₅₀-subfamily.




Fig. 2. Chemical structures of carotenoids produced by *Arthrobacter glacialis*. 1, a: R = H; b: R = Ac; Derivatized 2, a: R = H; b: R = SiMe₃; Derivatized 3, a: R = R¹; b: R = Ac; R² = H; c: R = Ac; R³ = SiMe₃.

Riboflavin

Arthrobacter globiformis isolated from soil excreted a yellow pigment during exponential growth, pigment which was identified as riboflavin, C₁₇H₂₀N₄O₆, also known as vitamin B2 (Fig. 1)



Fig. 1. Chemical structure of riboflavin



Indigoidine

Brilliant blue in color and water-insoluble pigments produced by *Arthrobacter atrocyaneus* and *Arthrobacter polychromogenes* were identified as indigoidine and its derivatives (Fig. 3)

Another strain which produces a blue pigment related to indigoidine is *Arthrobacter crystallopoietes*, although colonies of this *Arthrobacter* appear brilliant green in color. The strain *Arthrobacter oxydans* has been also reported that it produces the blue pigment which is also related to indigoidine.




Fig. 3. Chemical structures of indigoidine and its derivatives produced by *Arthrobacter* species. 1, a: R = H; b: R = CH₂CO₂; 2, a: R¹ = R² = H; b: R¹ = R² = CH₂CO₂; c: R¹ = R² = CH₂; d: R¹ = H; R² = CH₂CO₂; 3, R = CH₂CO₂.



Porphyrins

Many bacteria in the genus *Arthrobacter* produce red pigment porphyrins. A compound belonging to the family of red extracellular pigments porphyrin was isolated in *A. globiformis*, *A. photogenimus* and *A. aureus* and identified as coproporphyrin III, C₃₄H₃₆N₄O₈ (Fig. 5). Another form of porphyrin was also described from pigment excreted by *A. hyalinus*. This pigment was identified as uroporphyrin III, C₄₀H₃₈N₄O₁₀ (Fig. 6).




Fig. 5. Chemical structure of coproporphyrin III




Fig. 6. Chemical structure of uroporphyrin III

Carotenoids

Red-carotenoids accumulation in *Arthrobacter agilis*, a psychrotrophic bacterium isolated from Antarctic sea ice, has been investigated. The pigments were identified as a series of geometrical isomers of the C₅₀ carotenoid bacterioruberin (Fig.7). Another *Arthrobacter*, *Arthrobacter roseus* sp. nov., has been also reported to produce red-carotenoids.





Fig. 7. Chemical structures of carotenoids produced by *Arthrobacter agilis*. 1, tetra-anhydrobacterioruberin; 2, Bacterioruberin; R=H; a series of glycosylated derivatives: R=H, hexose or dihexose



Indochrome

Apart from indigoidine, other chromophores of the water-soluble pigments produced by *A. atrocyaneus* and *A. polychromogenes* were identified as indochromes with chemical formula C₂₉H₂₃N₃O₁₂ (Fig. 4). These pigments were released into the culture liquid only by indigoidine-producing bacteria.




Fig. 4. Chemical structures of indochrome produced by *Arthrobacter polychromogenes*

CONCLUSION

Pigments produced by microorganisms gain interest from the scientific community not only as a taxonomic tool to identify classify the microorganisms but also for a commercial purpose. The utilization of natural pigments in manufacturing has been increasing since the nineties due to the consumer awareness of the toxicity problems linked to synthetic pigments. Microorganisms seem to be a reasonable choice for colorant production due to biotechnological advantages e.g. production

regardless of season and geographical conditions; controllable and predictable yield. The genus *Arthrobacter* is one among diverse microorganisms which has been found to produce pigments with several hues. Furthermore, these bacteria have been commonly found in various environments. By these advantageous points, the study of the bacteria belonging to the genus *Arthrobacter* might lead to the discovery a novel source of natural colorants.

REFERENCES

- Imanari, Y. et al. Overproduction of riboflavin by an *Arthrobacter* sp. mutant resistant to 5-Fluorouracil. *Appl Microbiol Biotech.* **1995**, *50*, 317-322.
- Caro, Y. et al. Bacterial carotenoids. XIV.1. C50-Carotenoid, 14'-** C50-Carotenoids from *Arthrobacter glacialis*. *Acta Chem Scand B*, **1975**, *29*, 921-926.
- Reddy, G.S.N. et al. *Arthrobacter flavus* sp. nov., a psychrotrophic bacterium isolated from a pond in McMurdo Dry Valley, Antarctica. *Int J Syst Evol Microbiol.* **2000**, *50*, 1553-1561.
- Galaup, P. et al. First pigment fingerprints from the rind of French PDO red-smear ripened soft cheeses Epoisses, Mont d'Or and Maroilles. *Inn Food Sci Emerg Technol.* **2007**, *8*, 373-378.
- Kuhn, R. et al. Indigoidine and other bacterial pigments related to 3,3'-bipyridyl. *Arch Microbiol.* **1965**, *51*, 71-84.
- Ensign, J.C. et al. A crystalline pigment produced from 2-hydroxypyridine by *Arthrobacter crystallopoietes* n. sp. *Arch Microbiol.* **1963**, *47*, 137-153.
- Kajiwara, M. et al. Properties of zinc uroporphyrin III produced from isopropanol by *Arthrobacter hyalinus*. *J Ferment Bioeng.* **1995**, *79*(2), 174-176.
- Fong, N.J.C. et al. Carotenoids accumulation in the psychrotrophic bacterium *Arthrobacter agilis* in response to thermal and salt stress. *Appl Microbiol Biotech.* **2001**, *56*, 750-756.
- Reddy, G.S.N. et al. *Arthrobacter roseus* sp. nov., a psychrotrophic bacterium isolated from an Antarctic cyanobacterial mat sample. *Int J Syst Evol Microbiol.* **2003**, *52*, 1017-1021.



PIGMENTS PRODUCED BY THE BACTERIA BELONGING TO THE GENUS *ARTHROBACTER*

Sutthiwong N.^{1,2}, Caro Y.², Fouillaud M.², Laurent P.³, Valla A.⁴, Dufossé L.^{2,5}

¹ Agricultural Technology Department, Thailand Institute of Scientific and Technological Research, Pathum Thani, Thailand

² Laboratoire de Chimie des Substances Naturelles et des Sciences des Aliments, ESIROI, Université de La Réunion, Sainte-Clotilde, Ile de La Réunion, France

³ Département Génie Biologique, IUT, Université de La Réunion, Saint-Pierre, Ile de La Réunion, France

⁴ FRE 2125 CNRS, Chimie et Biologie des Substances Naturelles, Thiverval-Grignon, France

⁵ Laboratoire ANTiOX, Université de Bretagne Occidentale, Pôle Université Pierre-Jakez Hélias, Quimper, France

E-mail: Laurent.Dufosse@univ-reunion.fr

Abstract

Since several decades, pigments have been used as a taxonomic tool for the identification and classification of bacteria. Nowadays, pigment producing microorganisms have been also widely interested in scientific disciplines because of their biotechnological potential. With the growing interest in microbial pigments because of factors such as production regardless of season and geographical conditions, novel microorganisms which their pigments can be extracted are being evaluated. In the nature, a numerous number of microorganisms e.g. yeast, fungi, algae and bacteria produce pigments. The genus *Arthrobacter* is one among diverse microorganisms which has been found to produce pigments. Most of bacteria in this genus produce a range of pigments. Several previous studies show that pigments produced by bacteria belonging to the genus *Arthrobacter* have various hues depending on the chromophore which is present, e.g. yellow by carotenoid and riboflavin, green and blue by indigoidine and indochrome, and red by porphyrins and carotenoids. Since long time numerous strains in this genus have been reported that their colonies are colored; however, the purification and characterization of their pigments were not frequently conducted until well know chemical structures and role in these strains. Consequently, a study of pigments produced by the genus *Arthrobacter* may be worthy to play attention for discovering a novel source of natural colourants.

References

[1] Dufossé, L. Microbial production of food grade pigments. *Food Technol Biotechnol.* **2006**, *44*(3), 313-321.

[2] Fong, N.J.C. et al. Carotenoid accumulation in the psychrotrophic bacteria *Arthrobacter agilis* in response to thermal and salt stress. *Appl Microbiol Biotechnol.* **2001**, *56*, 750-756.

[3] <http://www.bacterio.cict.fr/a/arthrobacter.html>



PROGRAM

7th International Congress on Pigments in Food

June 18-21, 2013
Novara, Italy



Scientific Committee

Øyvind M. Andersen (Norway)
Marco Arlorio (Italy - Chair)
George Britton (United Kingdom)
Reinhold Carle (Germany)
Laurent Dufossé (Réunion Island)
José Empis (Portugal)
Vincenzo Fogliano (Italy)
Nicola Galaffu (Switzerland)
Vural Gökmen (Turkey)
Marina Heinonen (Finland)
Adriana Mercadante (Brazil)
Maria Roca Lopez Cepero (Spain)
Steven Schwartz (USA)
Livia Simon-Sarkadi (Hungary)
Carmen Socaciu (Romania)

Organizing Committee

Marco Arlorio (Chair)
Jean Daniel Coisson
Vincenzo Fogliano
Daniele Giuffrida
Aldo Martelli
Fabiano Travaglia

Secretariat

Matteo Bordiga
Elisabetta Cereti
Cristiano Garino
Monica Locatelli

e-mail: pif2013@pif2013.org

Phone: +39 0321 375873

www.pif2013.com

7th International Congress on Pigments in Food



18-21 June 2013, Novara, Italy

SCIENTIFIC PROGRAM

18 JUNE

16.30 - 17.00 Opening ceremony and welcome

SESSION 1 CHEMISTRY AND BIOCHEMISTRY

17.00 - 17.40

Plenary lecture Prof. G. BRITTON (University of Liverpool, U.K.)
"NATURAL CAROTENOIDS: A STUDY IN OILS AND WATER COLOURS"

17.40 - 18.00

CAROTENOID ESTER PROFILES IN *SOLANUM TUBEROSUM* AND *SOLANUM PHUREJA* CULTIVARS
Burmeister A., Bondiek S., Jerz G., Fleischmann P.
Institute of Food Chemistry, TU Braunschweig, Braunschweig, Germany

18.00 - 18.20

INTRAMOLECULAR AND INTERMOLECULAR FACTORS AFFECTING THE DEGRADATION KINETICS OF XANTHOPHYLL ESTERS
Jarén-Galán M., Hornero-Méndez D., Pérez-Gálvez A.
Food Biotechnology Department, Instituto de la Grasa (CSIC), Sevilla, Spain.

18.20 - 18.40

ANALYTICAL AND TECHNOLOGICAL ASPECT OF CAROTENOIDS FROM RED-BELL PEPPER
Daoud H.G., Palotas G., Palotas G., Pek Z.; Helyes L.
Szent Istvan University & Univer Products PSI



19 JUNE

SESSION 1 CHEMISTRY AND BIOCHEMISTRY

9.00 - 9.40	<p>Plenary lecture Prof. Ø.M. ANDERSEN (University of Bergen, Norway)</p> <p>“DIFFERENCES IN ANTHOCYANIN CONTENT OF FOOD AND NATURAL SOURCES CORRELATED WITH DIFFERENCES IN ANTHOCYANIN CHEMISTRY AND PROPERTIES”</p>
9.40 - 10.00	<p>ANTHOCYANIN-SYNTHESIZING TOMATO GENOTYPE ‘SUN BLACK™’ AS PRINCIPAL INGREDIENT FOR A NEW FUNCTIONAL TOMATO SAUCE</p> <p>Blando F.¹, Albano C.¹, Gerardi C.¹, Mita G.¹, Mazzucato A.²</p> <p>¹Institute of Sciences of Food Production, CNR, Lecce, Italy; ² Department of Sciences and Technologies for Agriculture, Forestry, Nature and Energy, Tuscia University, Viterbo, Italy</p>
10.00 - 10.20	<p>STUDIES ON COUPLING REACTIONS OF PROANTHOCYANIDINS AND MALVIDIN-3-O-GLUCOSIDE IN A WINE-LIKE MODEL SOLUTION SYSTEM</p> <p>Nickolaus P., Weber F., <u>Durner D.</u></p> <p>Competence Center for Viticulture & Enology, Neustadt an der Weinstraße, Germany.</p>
10.20 - 10.40	<p>POST-HARVEST MODIFICATIONS ENHANCE THE ZEAXANTHIN CONTENT IN VEGETABLES</p> <p>Esteban R.¹, Fleta E.¹, Buezo J.¹, Miguez F.¹, Becerril J.M.¹, García-Plazaola J.I.¹</p> <p>¹ Department of Plant Biology and Ecology, University of Basque Country, UPV/EHU, Bilbao, Spain</p>
10.40 - 11.00	Coffee break
11.00 - 12.00	TIME FOR POSTER SESSION
12.00 - 12.20	<p>DESCRIPTION OF A NEW CHLOROPHYLL CATABOLITE IN RIPENED FRUITS OF QUINCE (<i>Cydonia Oblonga</i>, Mill.)</p> <p><u>Roca M.</u>, Ríos J.J., Pérez-Gálvez A.</p> <p>Food Biotechnology Department, Instituto de la Grasa (CSIC), Sevilla, Spain.</p>
12.20 - 12.40	<p>RELATIONSHIPS AMONG FLAG LEAF CHLOROPHYLL CONTENT, AGRONOMICAL TRAITS, AND SOME PHYSIOLOGICAL TRAITS OF WINTER BREAD WHEAT GENOTYPES</p> <p>Bahar B., Sirat A., Kilic R., Aydin I.</p> <p>(Siran Vocational School, Gumushane University, Gumushane, Turkey)</p>
12.40 - 13.00	<p>OXIDATION ROUTES FOR BETACYANINS</p> <p>Wybraniec S.¹, Szot D.¹, Nemzer B.², Pietrzowski Z.³</p> <p>¹Department of Analytical Chemistry, Cracow University of Technology, Cracow, Poland; ²Chemistry Research, FutureCeuticals Inc., Mommence, IL, USA; ³Applied BioClinical Inc., Irvine, CA, USA.</p>

19 JUNE

**SESSION 2 (part 1) TECHNOLOGY, BIOTECHNOLOGY,
AND PROCESSING**

14.00 - 14.40

Plenary lecture prof. Vural GÖKMEN (Hacettepe University, Ankara, Turkey)
"ARTIFICIAL INTELLIGENCE: IMPROVING THE COLOR MEASUREMENT"

14.40 - 15.00

INFLUENCE OF SOME OAK WOOD COMPONENTS ON STABILITY OF MALVIDIN-3-GLUCOSIDE AND CHROMATIC CHARACTERISTICS IN MODEL WINE SOLUTIONS

Correia A.C., Jordão A.M.
Agrarian Higher School, Polytechnic Institute of Viseu (CI&DETS), Viseu, Portugal.

15.00 - 15.20

STABILIZATION OF ANTHOCYANIN-METAL CHELATES WITH HYDROCOLLOIDS FOR THEIR APPLICATION AS BLUE FOOD COLORANTS

Buchweitz M., Kammerer D. R., Carle R.
Institute of Food Science and Biotechnology, Chair Plant Foodstuff Technology, Hohenheim University, Stuttgart, Germany.

15.20 - 15.40

STABILISATION OF BEETROOT DERIVED BETANIN THROUGH INTERACTION WITH AN EXTRACT FROM BARBADOS CHERRY

Kendrick A.
Diana Food Division, Rennes, France.

15.40 - 16.20

Plenary LECTURE Prof. G. CRAVOTTO (University of Turin, Italy)

"MICROWAVE AND ULTRASOUND ASSISTED FOOD PIGMENTS EXTRACTION: HIGHLY EFFICIENT REACTORS FOR GREEN, SUSTAINABLE PROCESSES"

20 JUNE

SESSION 3 PIGMENTS FROM MICROALGAE

9.00 - 9.40	<p>Plenary lecture Prof. Eriksen N.T. (University of Aalborg, Denmark)</p> <p>PIGMENTS FROM MICROALGAE: A NEW PERSPECTIVE WITH EMPHASIS ON PHYCOCYANIN</p>
9.40 - 10.00	<p>Invited lecture</p> <p>ALGAL CAROTENOIDS AS NOVEL PIGMENTS IN NUTRITION</p> <p><u>Christaki E.</u> Laboratory of Nutrition, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece.</p>
10.00 - 10.20	<p>FUNCTIONAL FOOD DEVELOPMENT USING AQUEOUS EXTRACT OF ARTROSPIRA (SPIRULINA) MAXIMA RICH IN PHYCOBILIPROTEINS</p> <p><u>Langellotti A.L.</u>, Buono S., Vargas I., Martello A., Fogliano V. CRIAq Research Centre, University of Naples "Federico II", Portici, Italy</p>
10.20 - 10.40	Coffee break
10.40 - 11.40	TIME FOR POSTER SESSION

SESSION 2 (part 2) TECHNOLOGY, BIOTECHNOLOGY, AND PROCESSING

11.40 - 12.00	<p>NATURAL HYDROXYANTHRAQUINOID PIGMENTS: CURRENT SITUATION AND FUTURE OPPORTUNITIES IN FOOD</p> <p>Caro Y., Fouillaud M., Laurent P., <u>Dufossé L.</u> Laboratoire de Chimie des Substances Naturelles et des Sciences des Aliments, Université de la Réunion, Sainte-Clotilde, Ile de la Réunion, France</p>
12.00 - 12.20	<p>DEGRADATION OF ANTHOCYANINS IN PROCESSED STRAWBERRY FRUIT</p> <p><u>Kermasha S.</u>, Borgomano S. Department of Food Science and Agricultural Chemistry, McGill University, Ste-Anne de Bellevue, Canada</p>
12.20 - 13.00	"Short communications" from selected posters

21 JUNE

SESSION 4 HEALTH AND NUTRITION

9.00 - 9.40

Plenary lecture Dr. Schweiggert (University of Hohenheim, Germany)

ENHANCED BIOAVAILABILITY OF CAROTENOIDS: THE INFLUENCE OF CHROMOPLAST MORPHOLOGY DIETARY LIPID, AND THERMAL PROCESSING

Schweiggert R.M.^{1,2}, Kopec R.E.^{2,6}, Cooperstone J.L.², Villalobos-Gutierrez M.G.³, Högel J.⁴, Young G.S.⁵, Francis D.M.⁷, Quesada S.³, Esquivel P.³, Schwartz S.J.², Carle R.¹

9.40 - 10.00

BIOACCESSIBILITY AND CHANGES IN THE CAROTENOID PROFILE FROM MURICI FRUIT AFTER *IN VITRO* GASTROINTESTINAL DIGESTION

Mariutti L., Rodrigues E., Mandelli F., Mercadante A.

Department of Food Science, University of Campinas, Campinas, Brazil.

10.00 - 10.20

A MINI REVIEW ON THE COLOURLESS CAROTENOIDS PHYTOENE AND PHYTOFLUENE. ARE THEY INVISIBLE BIOACTIVE COMPOUNDS?

Meléndez-Martínez A. J.^{1,2}, Mapelli Brahm P.², Stinco C.M.² and Wang X-D.^{1,3}

¹J. Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA; ²Food Colour & Quality Laboratory, Department of Nutrition and Food Science, Universidad de Sevilla, Sevilla, Spain; ³Department of Nutritional Science, Gerald J. and Dorothy R. Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA.

10.20 - 10.40

DISSECTING THE PHARMACOPHORE OF CURCUMIN: TWO CASE STUDIES

Minassi A., Appendino G.

Department of Pharmaceutical Sciences, University of Piemonte Orientale "A. Avogadro", Novara, Italy.

10.40 - 11.00

Coffee break

11.00 - 12.30

ROUND TABLE: New natural pigments from foods: technical applicability and regulatory affairs

12.30 - 13.00

Closure ceremony and advertisement of next PIF

Poster Session

P 01: Synthesis of water-soluble carotenoids via click-reaction

Agócs A., Háda M., Nagy V., Deli J.

P 02: Thermal and light stability of β -cryptoxanthin esters

Bunea A., Andrei S., Rugină D., Pinteau A.

P 03: Effect of esterification on thermal stability and antioxidant activity of zeaxanthin

Pinteau A., Bunea A., Socaciu C.

P 04: Measurement of enzymatic hydrolysis of lutein esters from dairy products during *in vitro* digestion

Xavier A.A.O., Garrido-Fernández J., Mercadante A.Z., Pérez-Gálvez A..

P 05: Oil bodies as a potential microencapsulation carrier for astaxanthin stabilization and safe delivery

Acevedo F., Rubilar M., Villarroel M., Navarrete P., Jofré I., Romero F., Acevedo V., Shene C..

P 06: Microencapsulation of astaxanthin oleoresin from *Phaffia rhodozyma*

Villalobos-Castillejos F., Yáñez-Fernández J., Barragán-Huerta B.E.

P 07: Effect of genotype and growing conditions on lutein and β -carotene content of green leafy *Brassica* species

Arrigoni E., Reif C., Berger F., Baumgartner D., Nyström L.

P 08: Effect of processing on content of vital carotenoids in new vegetable puree

Palotás Gábor, Palotás Gabriella, Daood H., Pék Z., Helyes L.

P 09: Effect of addition of sodium erythorbate and urucum on the lipid oxidation in pork meat

Figueiredo B., Bragagnolo N.

P 10: Identification of *Cionosicyos macranthus* carotenoids

Murillo E., Watts M., Reyna G.

P 11: Bioactive compounds in supercritical CO₂-extracted pumpkin oil

Durante M., Lenucci M.S., D'Amico L., Dalessandro G., Mita G.

P 12: Evaluation of carotenoids and capsaicinoids content in powder of chilli peppers during one year of shelf-life

Giuffrida D., Cavazza A., Dugo P., Torre G., Corradini C., Bignardi C., Dugo G.mo

P 13: Carotenoids in red fleshed sweet oranges

Merussi G.D., Latado R.R., Rossi E.A., Sylos C.M.

P 14: Colour changes in heat-treated orange juice during ambient storage

Wibowo S., Vervoort L., Lemmens L., Hendrickx M., Van Loey A.

P 15: Carotenoid deposition and profiles in peach palm (*Bactris gasipaes* Kunth) fruits, and their implication on its nutritional potential

Hempel J., Esquivel P., Carle R., Schweiggert R.M.

P 16: Deposition of lycopene, β -carotene, and β -cryptoxanthin in different chromoplast substructures in papaya fruits

Schweiggert R.M., Steingass C.B., Heller A., Esquivel P., Carle R.

P 17: Evaluation of quality parameters and carotenoid content of three cultivars of mango (*Mangifera indica* L.) from Réunion island

Rosalie R., Chillet M., Joas J., Lechaudel M., Payet B., Vulcain E., Dufossé L.

P 18: Genuine profiles and bioaccessibilities of carotenoids from red- and yellow-fleshed Mamey sapote (*Pouteria sapota*) fruits

Chacón-Ordóñez T., Jiménez V.M., Esquivel P., Carle R., Schweiggert R.M.

P 19: Transgenic tomatoes and their carotenoid and flavour profiles

Höfelmeier H., Burmeister A., Schwab W., Fleischmann P.

P 20: Study of the time-course *cis/trans* isomerisation of lycopene, phytoene and phytofluene from tomato

Meléndez-Martínez A.J., Paulino M., Stinco C.M., Wang X.-D.

P 21: Carotenoid composition of three Hungarian algae species

Deli J., Vasas G., Parizsa P., Hajdú G., Szabó I., Lambert N.

P 22: HPLC method validation for the determination of fucoxanthin

Travaglia F., Bordiga M., Locatelli M., Coisson J.D., Arlorio M.

P 23: Carotenoids stabilisation for use in beverages: two different approaches

Mesnier X., Boukobza F., Bily A., Roller M.

P 24: Effect of heat processing on the profile of pigments and antioxidant capacity of Jalapeño peppers at intermediate ripening stages

Cervantes-Paz B., Ornelas-Paz J. de J., Yahia E.M.

P 25: Micellarization and digestive stability of pigments from Jalapeño peppers at intermediate ripening stages

Victoria-Campos C.I., Ornelas-Paz J. de J.

P 26: Changes in lutein, chlorophylls and chlorophyll degradation products in pistachio kernels (*Pistacia vera*) during roasting

Pumilia G., Schwartz S.J., Cichon M.J., Giuffrida D., Dugo G.mo

P 27: Decolouration processes under non-oxygen thermal auto-oxidation of chlorophyll and carotenoids fractions in virgin olive oils

Aparicio-Ruiz R., Gandul-Rojas B.

P 28: Pigment changes during processing of green table olive specialities treated with alkali and without fermentation

Gallardo-Guerrero L., Gandul-Rojas B.

P 29: Polyphenols and volatile compounds in Ogliarola and Cellina olive

Romani A., Banelli L., Fierini E., Mancuso S., Masi E., Haimler D.

P 30: Chlorophyllian pigments in extra virgin olive oils

Rovellini P., Venturini S., Fusari P.

P 31: Subcellular distribution in olive fruit of peroxidase activity on chlorophyll substrate

Vergara-Dominguez H., Roca M., Gandul-Rojas B.

P 32: Chlorophyll and carotenoid pigments in a survey of marketed apple varieties

Delgado-Pelayo R., Gallardo-Guerrero L., Hornero-Mendez D.

P 33: Quantitation of polyphenols in different apple varieties cultivated in Aosta valley

Valentini S., Sado A., Chasseur M., Thedy L., Lale Murix H., Barrel I., Chatel A.

P 34: Anthocyanins, phenolic acids and antioxidant activity in yellow, red and purple-fleshed potatoes after steam cooking

Bellumori M., Innocenti M., Cerretani L., Mulinacci N.

P 35: Chemical characterization and antioxidant activity of six rice cultivars grown in Piedmont (pigmented and non-pigmented)

Bordiga M., Coisson J.D., Locatelli M., Travaglia F., Arlorio M.

P 36: Effect of the use of enzymatic preparations on extraction of phenolic compounds from blue maize (*Zea mays* L.), from the region of Tlaxcala, Mexico

Martínez de Santos M.L., Conteras-Llano L.E., Lozada-Ramírez J.D., Ortega-Regules A.E.

P 37: Techno-functional properties of tomato sauce fortified with anthocyanin pigments

Blando F., Biasco M., Albano C., Gerardi C., Dal Porto L., Mita G.

P 38: Effect of post-harvest treatment on anthocyanin content and total phenolics in mango (*Mangifera indica* L.) peels

Geerkens C.H., Müller-Maatsch J.T.L., Geissler M., Carle R.

P 39: Maqui (*Aristotelia chilensis* (Mol.) Stuntz) – Detailed analysis of the highly pigmented “superfruit”

Brauch J., Buchweitz M., Carle R.

P 40: *Prunus mahaleb* L. fruit extracts: a novel source for natural food pigments

Gerardi C., Albano C., Blando F., Pinthus E., Rescio L., Mita G.

P 41: Red orange anthocyanins in colored juices and drinks: analytical method validation

Scordino M., Sabatino L., Gargano M., Lazzaro F., Borzi M.A., Traulo P., Gagliano G.

P 42: Anthocyanins extraction from mulberry by a combination of high hydrostatic pressure and enzymatic hydrolysis as emerging technology

Kim C.-T., Maeng J.-S., Kim C.-J., Cho Y.-J., Kim N., Oh H.-J., Kwon S.-J., Sung G.B.

P 43: Anthocyanins and bioactives content in healthy red fruit drinks

Castellar M.R., Díaz-García M.C., Obón J.M., Vicente-Castillo A.

- P 44: Bioactive compounds and antioxidant activity in fruits from Atlantic rainforest, Southeast Brazil**
Azevedo-Silva N., Rodrigues E., Mercadante A.Z., Oyama L.M., De Rosso V.V.
- P 45: Phenolic composition of Nebbiolo grapes from Piedmont: changes during ripening and identification of geographic origin**
Locatelli M., Travaglia F., Bordiga M., Coisson J.D., Arlorio M.
- P 46: Antioxidant pigments in red grape juices (*Vitis vinifera* L. cv Aglianico N.): *in vitro* bioaccessibility, bioavailability and plasma protein interaction**
Tenore G.C., Ritieni A., Campiglia P., Manfra M., Coppola L., Novellino E.
- P 47: Stability of naturally coloured food plant extracts**
Papetti A., Gazzani G.
- P 48: Color diversity and antioxidant activity in cactus pear fruits from Southern Italy genotypes**
Albano C., Aprile A., Negro C., Gerardi C., Mita G., Miceli A., De Bellis L., Blando F.
- P 49: Betanin stability in selected aqueous-organic solutions influenced by heavy metals**
Wybraniec S., Szot D., Nemzer B., Pietrzowski Z.
- P 50: Teaching food biotechnology in secondary schools using riboflavin as example**
Pietzner V., Zorn H.
- P 51: Application and stability of the natural pigment neocandentone in candy products in comparison with a commercial anthocyanin**
Gutierrez Zúñiga C., Yáñez-Fernández J., Barragán-Huerta B.E.
- P 52: Characterization and genetic fingerprint of saffron**
Vignolini P., Pinelli P., Albertini E., Romani R.
- P 53: Extraction methods of natural pigments from stamen of saffron flower**
Einafshar S., Rohani R., Khorsand Beheshti H.
- P 54: Effect of salt-stress on the production of pigments by *Chlorella vulgaris* under heterotrophic culture**
Benavente-Valdés J.R., Montañez J.C., Aguilar C.N., Méndez-Zavala A.
- P 55: Carotenoids profile of ultrasound-assisted extract *Phormidium* sp.**
Rodrigues D.B., Weis G.C.C., Schio K.L., Jacob-Lopes E., Zepka L.Q.
- P 56: Microorganisms used as pigment source**
Sarıçoban C., Battal S.
- P 57: Pigmented filamentous fungi isolated from tropical marine environments around Réunion island**
Fouillaud M., Boyer E., Fel A., Caro Y., Dufossé L.
- P 58: Valorisation of vinasse, a rum distillery effluent, by the production of carotenoid pigments using filamentous fungi**
Dorla E., Caro Y., Fouillaud M., Dufossé L., Laurent P.
- P 59: Pigments produced by the bacteria belonging to the genus *Arthrobacter***
Sutthiwong N., Caro Y., Fouillaud M., Laurent P., Valla A., Dufossé L.
- P 60: Characterization of *Arthrobacter arilaitensis* pigmentation using spectrophotometry**
Sutthiwong N., Caro Y., Fouillaud M., Dufossé L.
- P 61: Modeling thermal stability of red pigments produced by *Penicillium purpurogenum* GH2**
Morales-Oyervides L., Oliveira J., Souza-Gallagher M.J., Méndez-Zavala A., Montanez J.C.
- P 62: pH stability of red pigments produced by *Penicillium purpurogenum* GH2**
Morales-Oyervides L., Oliveira J., Souza-Gallagher M.J., Méndez-Zavala A., Montanez J.C.
- P 63: Preparation of brown-coloured submicron-sized hazelnut skin fiber with high antioxidant capacity using high shear homogenization**
Özdemir K.S., Yılmaz C., Gökmen V.
- P 64: Survey on occurrence of aminocarmine acid in E120 (carmine)-labeled food additives and beverages**
Sabatino L., Scordino M., Gargano M., Lazzaro F., Borzi M.A., Traulo P., Gagliano G.

The Organizing Committee expresses sincere thanks to

