



**HAL**  
open science

# Linear time algorithms for finding independent spanning trees on pyramid networks †

Shuo-I Wang, Fu-Hsing Wang

► **To cite this version:**

Shuo-I Wang, Fu-Hsing Wang. Linear time algorithms for finding independent spanning trees on pyramid networks †. 2016. hal-01402110

**HAL Id: hal-01402110**

**<https://hal.science/hal-01402110>**

Preprint submitted on 24 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Linear time algorithms for finding independent spanning trees on pyramid networks<sup>†</sup>

Shuo-I Wang<sup>1</sup>, Fu-Hsing Wang<sup>2‡</sup>

<sup>1</sup>Department of Maritime Policing, Taiwan Police College  
Taipei, Taiwan, R.O.C.

<sup>2</sup>Department of Information Management, Chinese Culture University  
Taipei, Taiwan, R.O.C.

## Abstract

The use of independent spanning trees (ISTs) has scientific applications in fault-tolerant requirement in network protocols and secure message distributions. Most of the designs of ISTs are for those interconnection networks with vertex symmetric property, implying that one can find ISTs rooted on a designated vertex, and, by the vertex symmetry property of the given network, hence have solved the ISTs problem on any arbitrary vertex. The existence of asymmetry makes the ISTs problem even harder than its symmetric counterpart. In this paper, we propose linear time algorithms that solved 3-ISTs rooted at an arbitrary vertex of pyramid networks.

**Keywords:** independent spanning trees; interconnection networks; pyramid networks; graph algorithms.

---

<sup>†</sup>This work was supported in part by the Ministry of Science and Technology of the Republic of China under the contract MOST 104-2221-E-034-001.

<sup>‡</sup>Corresponding Author: Professor Fu-Hsing Wang (wang.fuhsing@gmail.com).

# 1 Introduction

The vertex set and the edge set of a graph  $G$  are denoted by  $V(G)$  and  $E(G)$ , respectively. Let  $P = \langle v_1, v_2, \dots, v_j \rangle$  be a *path* from  $v_1$  to  $v_j$  in  $G$ . We use  $I(P)$  to denote the set  $V(P) \setminus \{v_1, v_j\}$ . Two paths  $P$  and  $Q$  are called *disjoint*, denoted by  $P \parallel Q$ , if  $V(P) \cap V(Q) = \emptyset$ . We say that  $P$  and  $Q$  are *internally disjoint* if  $I(P) \cap I(Q) = \emptyset$ . A *spanning tree* of a graph  $G$  is a tree  $T$  such that  $V(T) = V(G)$ . Given a designated vertex  $r \in V$  as a root vertex and a natural number  $k$ , spanning trees  $T_1, T_2, \dots, T_k$  of  $G$  are called *k-independent spanning trees* of  $G$  rooted at  $r$  (referred as *k-ISTs* for short) if the  $k$  paths connecting  $r$  and  $v$ , for any vertex  $v$ , in  $T_1, T_2, \dots, T_k$  are pairwise internally disjoint in  $G$ .

The use of independent spanning trees has scientific applications in fault-tolerant requirement in network protocols and secure message distributions [1]. For example, an efficient and reliable broadcasting can be achieved by sending  $k$  copies of the message rooted at the source vertex along  $k$ -ISTs such that the message can be broadcasted to all the other vertices in a network with at most  $k - 1$  faulty vertices. Although the ISTs problem is hard for general graphs, several results are known for some special classes of graphs, especially on interconnection networks, such as odd graphs [5],  $n$ -dimensional torus [9], product graphs [6], chordal rings [4], recursive circulant graphs [11], parity cubes [10], and hypercubes [8]. Most of the above results are for symmetric graphs implying that one can find ISTs rooted on a designated vertex, and, by the vertex symmetry property of the given graph, hence have solved the ISTs problem on any arbitrary vertex. However, the investigation into the ISTs problem on interconnection networks with no vertex symmetric property, to the best of our knowledge, are poor.

We focus attention on the construction of ISTs of a given pyramid network. Pyramid networks have potentially powerful architecture for many applications such as image processing, visualization, and data mining [3]. The major advantage of pyramids which is important for image processing systems is hierarchical abstracting and transferring the data from different directions and forward them toward the apex of a pyramid [7]. A pyramid network is a 3-connected graph. In [2], Cheriyan and Maheshwari showed that, for any 3-connected graph, 3-ISTs rooted at any vertex can be found in  $O(|V||E|)$  time. In this paper, we propose linear time algorithms for finding 3-ISTs of a pyramid network rooted at an arbitrary vertex.

## 2 Preliminaries

All graphs considered here are finite, undirected, without loops and multiple edges. A *path* in a graph  $G$  is a single vertex or a sequence of distinct vertices  $\langle v_1, v_2, \dots, v_n \rangle$  such that  $(v_1, v_2), (v_2, v_3), \dots, (v_{n-1}, v_n)$  are edges of  $G$  and the  $v_i$  are distinct. A vertex of degree 0 is considered as an *isolated* vertex.

A  $2^k \times 2^k$  *square mesh*, denoted by  $M_k$ , has the vertex set  $V(M_k) = \{(x, y) \mid 0 \leq x, y \leq 2^k - 1\}$  where any two vertices  $(x_1, y_1)$  and  $(x_2, y_2)$  are connected by an edge iff  $|x_1 - x_2| + |y_1 - y_2| = 1$ . Each  $M_k, k \geq 2$ , can be partitioned into  $2^{k-2} \times 2^{k-2}$  square submeshes  $M_2$  and each  $M_2$  is called a *cluster*. Every cluster can be further decomposed into  $2 \times 2$   $M_1$  and each  $M_1$  is said to be a *block*. Let  $v, v^+, v^{++}, v^-$  denote the four vertices of a block in a clockwise order starting from  $v$ , and  $B, B^+, B^{++}, B^-$  denote the four blocks of a cluster in a clockwise order starting from  $B$ .

Let  $\mathbb{P}_n$  be an  $n$ -dimensional *pyramid* with the vertex set  $\bigcup_{k=0}^n V_k$ , where  $V_k = \{(k; x, y) \mid 0 \leq x, y \leq 2^k - 1\}$ . Vertex  $(0; 0, 0)$  is the *apex* of  $\mathbb{P}_n$  and also denoted as  $\Delta$ . The subgraph induced by  $V_k$  is connected as an  $M_k$  and called *layer  $k$*  of  $\mathbb{P}_n$ . Let  $S_0$  and  $S_1$  denote the layers 0 and 1 of  $\mathbb{P}_n$  with the vertex set  $(0; 0, 0)$  and  $\{(1; x, y) \mid 0 \leq x, y \leq 3\}$ , respectively. Let  $S(k; a, b)$  denote the cluster on layer  $k$  with the vertex set  $\{(k; x, y) \mid 4a \leq x \leq 4a + 3 \text{ and } 4b \leq y \leq 4b + 3\}$ , where  $2 \leq k \leq n$  and  $0 \leq a, b \leq 2^{k-2} - 1$ . Notice that layer  $k$  has  $2^{k-2}$  clusters. Vertex  $(k; x, y)$  has exactly four *children*  $(k + 1; 2x, 2y), (k + 1; 2x, 2y + 1), (k + 1; 2x + 1, 2y), (k + 1; 2x + 1, 2y + 1)$  in  $V_{k+1}$  and a *parent vertex*  $(k - 1; \lfloor \frac{x}{2} \rfloor, \lfloor \frac{y}{2} \rfloor)$  in  $V_{k-1}$ . For simplicity, we shall use the abbreviated notation  $S_k$  for  $S(k; a, b)$ , whenever no confusion can arise. Let  $p(v)$  denote the parent vertex of  $v$  and  $l(v)$  the layer number of  $v$ . The block containing the four children of  $v$ , denoted  $\mathcal{C}_{\mathcal{B}}(v)$ , is called the *child block* of  $v$ . In an  $S_k$ , the block with the vertex set  $\{p(v) \mid v \in V(S_{k+1})\}$  is called the *parent block* of  $S_{k+1}$  and denoted by  $\mathcal{P}_{\mathcal{B}}(S_{k+1})$ . An edge between  $v$  and  $p(v)$  is called a *layer edge*, while every edge of a cluster is called a *cluster edge*. Every vertex on the shortest path from  $v$  to  $\Delta$  is called an *ancestor* of  $v$ .

In a cluster  $S$ , every vertex of degree 4 is called an *inner vertex*, while the remaining vertices of  $S$  are said to be *outer vertices*. An outer vertex of degree 2 is also called a *corner vertex*. The subgraph induced by all inner vertices of  $S$  is an *i-cycle*, while the cycle induced by all of the outer vertices is an *o-cycle*. For any two vertices  $u$  and  $v$  in an *o-cycle* (respectively, *i-cycle*)  $O$ , each path from  $u$  to

$v$  is called an  $o$ -path (respectively,  $i$ -path). Vertex  $v$  is also labeled as  $u^{+d}$  (respectively,  $u^{-d}$ ) if  $v$  can be visited from  $u$  via  $d$  edges of  $O$  in a clockwise order (respectively, a counterclockwise order). In particular,  $u = v$  if  $d = 0$ . Figure 1 depicts an example of a 2-layered pyramid  $\mathbb{P}_2$ . The dash lines indicate layer edges, while the solid lines are cluster edges. Each bold 4-cycle is a block. The top-right block of  $S(2;0,0)$  has a corner vertex  $(2;0,3)$ , an inner vertex  $(2;1,2)$  and outer vertices  $(2;0,3)$ ,  $(2;1,3)$  and  $(2;0,2)$ . The subgraph induced by the set  $X = \{(2;1,1), (2;1,2), (2;2,2), (2;2,1)\}$  is the  $i$ -cycle of  $S(2;0,0)$ , and the  $o$ -cycle of  $S(2;0,0)$  is induced by  $V(S(2;0,0)) \setminus V(X)$ .

### 3 Constructions of 3-ISTs Rooted at Different Layers

For a given root vertex  $r$ , we shall construct 3-ISTs of  $\mathbb{P}_n$  rooted at  $r$  by considering the position of  $r$  to fulfill the vertex asymmetry of  $\mathbb{P}_n$ .

#### 3.1 Rooted at the Apex

Before the construction of 3-ISTs of  $\mathbb{P}_n$  rooted at an arbitrary vertex, we first define a vertex coloring on every vertex of  $\mathbb{P}_n$  for selecting edges to be tree edges of a spanning tree. Let  $\pi_0 : V \rightarrow \{0, 1, 2, 3\}$  be a coloring on  $\mathbb{P}_n$ , where for each vertex  $v \in V(\mathbb{P}_n - r)$ ,  $\pi_0(v) \in \{0, 1, 2, 3\}$ , and the four vertices of each block are assigned distinct colors. For convenience, let  $\rho_1, \rho_2$  and  $\rho_3$  be three distinct color numbers in  $\{1, 2, 3\}$  in the remaining text. In  $S_1$ , we arbitrarily select a vertex  $v$  as the corner vertex and then assign colors  $0, \rho_1, \rho_2$  and  $\rho_3$  to vertices  $v, v^+, v^-$  and  $v^{++}$ , respectively. Let  $\hat{H}_{1,\rho_i}$  be the subgraph with the vertex set  $V(S_1) \cup \{r\}$ , where  $1 \leq i \leq 3$ . And let  $E(\hat{H}_{1,\rho_1}) = E(\langle v^{++}, v^+, v, v^- \rangle) \cup \{(r, v^+)\}$ ,  $E(\hat{H}_{1,\rho_2}) = E(\langle v^{++}, v^-, v, v^+ \rangle) \cup \{(r, v^-)\}$  and  $E(\hat{H}_{1,\rho_3}) = E(\langle v^+, v^{++}, v^- \rangle) \cup \{(r, v), (r, v^{++})\}$ . The  $\pi_0$  coloring of a cluster is determined by the coloring of its parent block. For a cluster  $S_k, k \geq 2$ , let  $x, x^+, x^-$  and  $x^{++}$  be the four parent vertices of  $S_k$  such that  $\pi_0(x) = 0, \pi_0(x^+) = \rho_1, \pi_0(x^-) = \rho_2$  and  $\pi_0(x^{++}) = \rho_3$ . We assign color  $\pi_0(x^+)$  to the corner vertex of  $\mathcal{C}_{\mathcal{B}}(x^+)$ , to the two outer vertices incident to  $\mathcal{C}_{\mathcal{B}}(x^+)$ , and to the outer vertex incident to  $\mathcal{C}_{\mathcal{B}}(x)$  in  $\mathcal{C}_{\mathcal{B}}(x^-)$  (see Statement 2 of **HFinding Algorithm**). Statement 3 assigns color  $\pi_0(x^-)$  to the corner vertex of  $\mathcal{C}_{\mathcal{B}}(x^-)$ , to the two outer vertices incident to  $\mathcal{C}_{\mathcal{B}}(x^-)$ , and to the outer vertex incident to  $\mathcal{C}_{\mathcal{B}}(x^{++})$  in  $\mathcal{C}_{\mathcal{B}}(x^+)$ . In Statement 4, we assign color  $\pi_0(x^{++})$  to each inner vertex of  $S_k$ . The remaining uncolored vertices of  $S_k$  are assigned color 0 in Statement 5.

We now define three graphs for any cluster, and then each subgraph will be selected to be

---

**Input:** Given a coloring on the vertices  $x, x^+, x^-, x^{++}$  of  $\mathcal{P}_{\mathcal{B}}(S_k)$  as:  $\pi_0(x) = 0, \pi_0(x^+) = \rho_1,$   
 $\pi_0(x^-) = \rho_2, \pi_0(x^{++}) = \rho_3.$

**Output:** The colorings of blocks  $\mathcal{C}_{\mathcal{B}}(x), \mathcal{C}_{\mathcal{B}}(x^+), \mathcal{C}_{\mathcal{B}}(x^-)$  and  $\mathcal{C}_{\mathcal{B}}(x^{++})$  in  $S_k.$

**begin**

Let the inner vertex of  $\mathcal{C}_{\mathcal{B}}(x^{++})$  be  $\hat{h}_{k,\rho_3}$  and the corner vertices of  $\mathcal{C}_{\mathcal{B}}(x^+)$  and  $\mathcal{C}_{\mathcal{B}}(x^-)$  be  
 $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$ , respectively;

Assign color  $\rho_1$  to every vertex of  $\{\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_1}^{-5}\};$

Assign color  $\rho_2$  to every vertex of  $\{\hat{h}_{k,\rho_2}, \hat{h}_{k,\rho_2}^{+2}, \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-5}\};$

Assign color  $\rho_3$  to every inner vertex of  $S_k;$

Assign color 0 to every uncolored outer vertex of  $S_k;$

$E(\hat{H}_{k,\rho_1}) = E(P_{k,\rho_1}) \cup \{(\hat{h}_{k,\rho_3}, \hat{h}_{k,\rho_1}^{+2}), (\hat{h}_{k,\rho_3}^{-1}, \hat{h}_{k,\rho_1}^{-1}), (\hat{h}_{k,\rho_3}^{+2}, \hat{h}_{k,\rho_1}^{-2}), (\hat{h}_{k,\rho_3}^{+1}, \hat{h}_{k,\rho_1}^{-5}), (\hat{h}_{k,\rho_1}, p(\hat{h}_{k,\rho_1}))\};$

$E(\hat{H}_{k,\rho_2}) = E(P_{k,\rho_2}) \cup \{(\hat{h}_{k,\rho_3}, \hat{h}_{k,\rho_2}^{-2}), (\hat{h}_{k,\rho_3}^{-1}, \hat{h}_{k,\rho_2}^{-5}), (\hat{h}_{k,\rho_3}^{+2}, \hat{h}_{k,\rho_2}^{+2}), (\hat{h}_{k,\rho_3}^{+1}, \hat{h}_{k,\rho_2}^{-1}), (\hat{h}_{k,\rho_2}, p(\hat{h}_{k,\rho_2}))\};$

$E(\hat{H}_{k,\rho_3}) = E(P_{k,\rho_3}) \cup \{(u, v) \in E(S_k) | \pi_0(u) = \rho_3, \pi_0(v) \neq$   
 $\rho_3\} \cup \{(\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_1}^{-1}), (\hat{h}_{k,\rho_2}, \hat{h}_{k,\rho_2}^{-1}), (\hat{h}_{k,\rho_1}^{+3}, p(\hat{h}_{k,\rho_1}^{+3})), (\hat{h}_{k,\rho_1}^{-3}, p(\hat{h}_{k,\rho_1}^{-3})), (\hat{h}_{k,\rho_3}, p(\hat{h}_{k,\rho_3}))\};$

**end**

**HFinding Algorithm**

---

a subgraph of one of the 3-ISTs of  $\mathbb{P}_n$ . Let  $\hat{H}_i(k; a, b)$  denote the subgraph with the vertex set  $V(S(k; a, b) + V(\mathcal{P}_{\mathcal{B}}(S(k; a, b))))$ , where  $1 \leq i \leq 3, 2 \leq k \leq n$  and  $0 \leq a, b \leq 2^{k-2} - 1$ . A vertex  $v \in V(\hat{H}_i(k; a, b))$ , denoted  $\hat{h}_i(k; a, b)$ , is called a *junction vertex* of  $\hat{H}_i(k; a, b)$  if  $\pi_0(v) = \pi_0(p(v)) = i$ . The vertex of color  $\rho_i$  in  $S_1$  is denoted by  $\hat{h}_{1,\rho_i}$  for each  $i$  and the vertex of color 0 in  $S_1$  is  $\hat{h}_{1,0}$ . For simplicity, we shall also use the abbreviated notation  $\hat{h}_{k,i}$  and  $\hat{H}_{k,i}$  for  $\hat{h}_i(k; a, b)$  and  $\hat{H}_i(k; a, b)$ , respectively, and let  $\hat{h}_k = \{\hat{h}_{k,1}, \hat{h}_{k,2}, \hat{h}_{k,3}\}.$

By the first statement of **HFinding Algorithm**, there are exactly two junction vertices on the  $o$ -cycle of  $S_k$ , while one junction vertex belongs to the  $i$ -cycle of  $S_k$ . In Statements 6-8, we define three subgraphs  $\hat{H}_{k,\rho_1}, \hat{H}_{k,\rho_2}$  and  $\hat{H}_{k,\rho_3}$  on  $S_k$ .  $\hat{H}_{k,\rho_1}$  has the outer path  $P_{k,\rho_1} = \langle \hat{h}_{k,\rho_2}, \hat{h}_{k,\rho_1}^{-5}, \dots, \hat{h}_{k,\rho_1}^{-1}, \hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_1}^{+1}, \hat{h}_{k,\rho_1}^{+2}, \dots, \hat{h}_{k,\rho_1}^{+5} \rangle$ , and every inner vertex of  $S_k$  has exactly one edge incident to  $P_{k,\rho_1}$  in  $\hat{H}_{k,\rho_1}$  (See Statement 6).  $\hat{H}_{k,\rho_2}$  contains the outer path  $P_{k,\rho_2} = \langle \hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_2}^{-5}, \dots, \hat{h}_{k,\rho_2}^{-1}, \hat{h}_{k,\rho_2}, \hat{h}_{k,\rho_2}^{+1}, \hat{h}_{k,\rho_2}^{+2}, \dots, \hat{h}_{k,\rho_2}^{+5} \rangle$ , and every inner vertex has an edge incident to  $P_{k,\rho_2}$  (See Statement 7).  $\hat{H}_{k,\rho_3}$  has the inner path  $P_{k,\rho_3} = \langle \hat{h}_{k,\rho_3}^{-3}, \hat{h}_{k,\rho_3}^{-2}, \hat{h}_{k,\rho_3}^{-1}, \hat{h}_{k,\rho_3} \rangle$  and every outer vertex  $v \in V(S_k)$  has an edge incident to  $P_{k,\rho_3}$  if  $v$  is not a corner vertex. Among the four corner vertices of  $S_k$ ,  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$  are adjacent to  $\hat{h}_{k,\rho_1}^{-1}$  and  $\hat{h}_{k,\rho_2}^{-1}$ , respectively, in  $\hat{H}_{k,\rho_3}$ . And the two corner vertices of color 0 have layer edges  $(\hat{h}_{k,\rho_1}^{+3}, p(\hat{h}_{k,\rho_1}^{+3}))$  and  $(\hat{h}_{k,\rho_1}^{-3}, p(\hat{h}_{k,\rho_1}^{-3}))$  in  $\hat{H}_{k,\rho_3}$  (See Statement 8). For the connectivity of layers, the edge  $(\hat{h}_{k,\rho_i}, p(\hat{h}_{k,\rho_i}))$  is used to connect  $\hat{H}_{k,\rho_i}$  and

$\hat{H}_{k-1,\rho_i}$ , where  $1 \leq i \leq 3$  (See Statements 6-8). It can be seen that  $P_{k,\rho_1}, P_{k,\rho_2}$  and  $P_{k,\rho_3}$  are pairwise disjoint. Figure 2 illustrates subgraphs  $\hat{H}_{k,\rho_1}, \hat{H}_{k,\rho_2}$  and  $\hat{H}_{k,\rho_3}$  of a cluster  $S_k$  on the condition that the parent vertex of the top-left block of  $S_k$  is of color 0. The bold edges are edges of subgraphs and the grey vertices represent junction vertices. The formal definitions of  $\hat{H}_{k,\rho_1}, \hat{H}_{k,\rho_2}$  and  $\hat{H}_{k,\rho_3}$  refer to **HFinding Algorithm**.

The following observation gives some properties of  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$ :

**Observation 3.1.** *Let  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$  be on the  $o$ -cycle of  $S_k$ , where  $2 \leq k \leq n$ .*

$$(1) \hat{h}_{k,\rho_1}^{+d} = \hat{h}_{k,\rho_2}^{-(6-d)} \text{ and } \hat{h}_{k,\rho_2}^{+d} = \hat{h}_{k,\rho_1}^{-(6-d)}.$$

(2) *If  $v = \hat{h}_{k,\rho_1}^{+d}$ , then  $v$  is an end-vertex of  $\langle \hat{h}_{k,\rho_1}^{+d}, \hat{h}_{k,\rho_1}^{+(d-1)}, \dots, \hat{h}_{k,\rho_1} \rangle$  and  $\langle \hat{h}_{k,\rho_2}^{-(6-d)}, \hat{h}_{k,\rho_2}^{-(6-d-1)}, \dots, \hat{h}_{k,\rho_2} \rangle$  in  $\hat{H}_{k,\rho_1}$  and  $\hat{H}_{k,\rho_2}$ , respectively.*

(3) *If  $v = \hat{h}_{k,\rho_1}^{-d}$ , then  $v$  is an end-vertex of  $\langle \hat{h}_{k,\rho_1}^{-d}, \hat{h}_{k,\rho_1}^{-(d-1)}, \dots, \hat{h}_{k,\rho_1} \rangle$  and  $\langle \hat{h}_{k,\rho_2}^{+(6-d)}, \hat{h}_{k,\rho_2}^{+(6-d-1)}, \dots, \hat{h}_{k,\rho_2} \rangle$  in  $\hat{H}_{k,\rho_1}$  and  $\hat{H}_{k,\rho_2}$ , respectively.*

**Proof.** Let  $B_1$  and  $B_2$  be the block containing  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$ , respectively. According to the first statement of **HFinding Algorithm**, both  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$  are corner vertices and  $B_1$  is in fact  $B_2^{++}$ . Thus,  $\hat{h}_{k,\rho_1} = \hat{h}_{k,\rho_2}^{+6}$  and  $\hat{h}_{k,\rho_2} = \hat{h}_{k,\rho_1}^{+6}$ . Besides, both (2) and (3) immediately follow (1).

Q. E. D.

Since the  $\pi_0$  coloring of a cluster can be decided by its parent block, the coloring of  $S_2$  is determined by the coloring of  $S_1$ . After that, the coloring and subgraphs of every cluster in  $\mathbb{P}_n$  can then be established. We now continue with the derivation of building spanning subgraphs  $T_{\rho_i}^0, i = 1, 2, 3$ , on  $\mathbb{P}_n$ . Each  $T_{\rho_i}^0$  is regarded as an  $n$ -layered graph. Let  $V(T_{\rho_i}^0) = V(\mathbb{P}_n)$  and  $E(T_{\rho_i}^0) = E(\hat{H}_{1,\rho_i}) \bigcup_{k=2}^n \bigcup_{a,b=0}^{2^{k-2}-1} E(\hat{H}_{\rho_i}(k; a, b))$  for  $i = 1, 2, 3$ .

To show that there are three pairwise internally disjoint paths from any vertex  $u_k$  to the apex  $r$ , we first find three pairwise internally disjoint paths from  $u_k$  to the block  $B$  containing  $p(u_k)$ .

**Lemma 3.1.** *If  $u_k$  is a vertex of  $S_k, 2 \leq k \leq n$ , and  $u'_{k-1}$  is a dummy vertex with dummy edges incident to every vertex of  $\mathcal{P}_B(S_k)$ , then paths from  $u_k$  to  $u'_{k-1}$  in  $T_1^0, T_2^0$  and  $T_3^0$  are pairwise internally disjoint.*

**Proof.** Let  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$  be outer vertices and  $\hat{h}_{k,\rho_3}$  be an inner vertex. Since  $\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_2}$  and

$\hat{h}_{k,\rho_3}$  belong to three different blocks,  $p(\hat{h}_{k,\rho_1}), p(\hat{h}_{k,\rho_2})$  and  $p(\hat{h}_{k,\rho_3})$  are distinct. Let  $P_{\rho_i}$  be the path from  $u_k$  to  $u'_{k-1}$  in  $T_{\rho_i}^0$ , where  $1 \leq i \leq 3$ . Each  $P_{\rho_i}$  can be regarded as a concatenation of subpaths  $\langle u_k \rangle Q_{\rho_i} \langle t_i, u'_{k-1} \rangle$ , where  $t_i \in V(\mathcal{P}_{\mathcal{B}}(S_k))$ . Let  $v_i$  be the neighbor of  $u_k$  in  $\hat{H}_{k,\rho_i}$  for each  $i$ . Note that  $v_i \neq v_j$  and  $t_i \neq t_j$  if  $i \neq j$ . We shall prove  $Q_{\rho_1}, Q_{\rho_2}, Q_{\rho_3}$  pairwise disjoint instead of showing  $P_{\rho_1}, P_{\rho_2}, P_{\rho_3}$  pairwise internally disjoint.

*Case 1.*  $u_k$  is a corner vertex and  $\pi_0(u_k) = 0$ .

By **HFinding Algorithm**,  $u_k$  is either  $\hat{h}_{k,\rho_1}^{+3}$  or  $\hat{h}_{k,\rho_1}^{-3}$ . From the definition of  $E(\hat{H}_{k,\rho_3})$ ,  $(u_k, p(u_k)) \in E(\hat{H}_{k,\rho_3})$ . Then  $P_{\rho_3} = \langle u_k, p(u_k), u'_{k-1} \rangle$  and thus  $Q_{\rho_3}$  is empty. So,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$ . If  $u_k = \hat{h}_{k,\rho_1}^{+3}$ , by Observation 3.1 (2),  $u_k$  is an end-vertex of  $\langle \hat{h}_{k,\rho_1}^{+3}, \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{+1}, \hat{h}_{k,\rho_1} \rangle$  in  $\hat{H}_{k,\rho_1}$  and is also an end-vertex of  $\langle \hat{h}_{k,\rho_2}^{-3}, \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-1}, \hat{h}_{k,\rho_2} \rangle$  in  $\hat{H}_{k,\rho_2}$ . It means that  $Q_{\rho_1} = \langle \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{+1}, \hat{h}_{k,\rho_1} \rangle$  and  $Q_{\rho_2} = \langle \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-1}, \hat{h}_{k,\rho_2} \rangle$ . For the case  $u_k = \hat{h}_{k,\rho_1}^{-3}$ , by Observation 3.1 (3),  $u_k$  is an end-vertex of  $\langle \hat{h}_{k,\rho_1}^{-3}, \hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_1}^{-1}, \hat{h}_{k,\rho_1} \rangle$  in  $\hat{H}_{k,\rho_1}$  and an end-vertex of  $\langle \hat{h}_{k,\rho_2}^{+3}, \hat{h}_{k,\rho_2}^{+2}, \hat{h}_{k,\rho_2}^{+1}, \hat{h}_{k,\rho_2} \rangle$  in  $\hat{H}_{k,\rho_2}$ . Then  $Q_{\rho_1} = \langle \hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_1}^{-1}, \hat{h}_{k,\rho_1} \rangle$  and  $Q_{\rho_2} = \langle \hat{h}_{k,\rho_2}^{+2}, \hat{h}_{k,\rho_2}^{+1}, \hat{h}_{k,\rho_2} \rangle$ . It concludes that  $Q_{\rho_1} \parallel Q_{\rho_2}$  for cases  $u_k = \hat{h}_{k,\rho_1}^{+3}$  and  $u_k = \hat{h}_{k,\rho_1}^{-3}$ .

*Case 2.*  $u_k = \hat{h}_{k,\rho_1}$ .

Obviously,  $P_{\rho_1} = \langle u_k, p(u_k), u'_{k-1} \rangle$  and thus  $Q_{\rho_1}$  is empty. So,  $Q_{\rho_1} \parallel Q_{\rho_2}$  and  $Q_{\rho_1} \parallel Q_{\rho_3}$ . Furthermore, since  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}^{+5}$  are the two end-vertices of the path  $P_{k,\rho_2}$  and  $\hat{h}_{k,\rho_2} \in V(P_{k,\rho_2})$ ,  $Q_{\rho_2}$  is a subpath of  $P_{k,\rho_2}$  and  $Q_{\rho_2}$  does not contain the vertex  $\hat{h}_{k,\rho_2}^{+5}$ . From the definition of  $\hat{H}_{k,\rho_3}$ ,  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_1}^{-1}$  are adjacent and  $\hat{h}_{k,\rho_1}^{-1}$  is adjacent to an inner vertex  $v$  in  $E(\hat{H}_{k,\rho_3})$ . Since  $v \in V(P_{k,\rho_3})$ ,  $Q_{\rho_3} - \hat{h}_{k,\rho_1}^{-1}$  is a subpath of  $P_{k,\rho_3}$ . That is,  $Q_{\rho_3} - \hat{h}_{k,\rho_2}^{+5}$  is a subpath of  $P_{k,\rho_3}$  because  $\hat{h}_{k,\rho_1}^{-1} = \hat{h}_{k,\rho_2}^{+5}$ . Moreover, since  $P_{k,\rho_3} \parallel P_{k,\rho_2}$  and  $\hat{h}_{k,\rho_2}^{+5} \notin V(Q_{\rho_2})$ ,  $Q_{\rho_3} \parallel Q_{\rho_2}$ .

*Case 3.*  $u_k = \hat{h}_{k,\rho_2}$ .

The proof is similar to *Case 1* and is left to the reader.

*Case 4.*  $u_k$  is an inner vertex.

Notably,  $v_1$  and  $v_2$  belong to the same block. If  $u_k = \hat{h}_{k,\rho_3}$ , then  $Q_{\rho_3}$  is empty. By the definitions of  $\hat{H}_{k,\rho_1}$  and  $\hat{H}_{k,\rho_2}$ ,  $v_1$  and  $v_2$  are  $\hat{h}_{k,\rho_1}^{+2}$  and  $\hat{h}_{k,\rho_2}^{-2}$ , respectively. Then  $Q_{\rho_1} = \langle \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{+1}, \hat{h}_{k,\rho_1} \rangle$  and  $Q_{\rho_2} = \langle \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-1}, \hat{h}_{k,\rho_2} \rangle$ . So,  $Q_{\rho_1}, Q_{\rho_2}$  and  $Q_{\rho_3}$  are pairwise disjoint. Similarly, we get pairwise disjoint paths  $Q_{\rho_1}, Q_{\rho_2}$  and  $Q_{\rho_3}$  for cases  $u_k = \hat{h}_{k,\rho_3}^{-j}$ , where  $j = 1, 2, 3$ .



Case 5.  $u_k$  is adjacent to a corner vertex.

Clearly,  $v_i \in V(P_{k,\rho_i})$  for each  $i$ . That is,  $Q_{\rho_3}$  is an  $i$ -path, while both  $Q_{\rho_1}$  and  $Q_{\rho_2}$  are  $o$ -paths. So,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$ . Vertex  $u_k$  is either  $\hat{h}_{k,\rho_1}^{+d_1}$  or  $\hat{h}_{k,\rho_1}^{-d_2}$ , where  $1 \leq d_1, d_2 \leq 5$  and  $d_1, d_2 \neq 3$ . For the former case, by Observation 3.1 (1),  $Q_{\rho_1} = \langle \hat{h}_{k,\rho_1}^{+(d_1-1)}, \dots, \hat{h}_{k,\rho_1} \rangle$  and  $Q_{\rho_2} = \langle \hat{h}_{k,\rho_2}^{-(6-d_1-1)}, \dots, \hat{h}_{k,\rho_2} \rangle$ . Thus,  $Q_{\rho_1} \parallel Q_{\rho_2}$ . For the latter case, by Observation 3.1 (2),  $Q_{\rho_1} = \langle \hat{h}_{k,\rho_1}^{-(d_2-1)}, \dots, \hat{h}_{k,\rho_1} \rangle$  and  $Q_{\rho_2} = \langle \hat{h}_{k,\rho_2}^{+(6-d_2-1)}, \dots, \hat{h}_{k,\rho_2} \rangle$ . Therefore,  $Q_{\rho_1} \parallel Q_{\rho_2}$ .

Q. E. D.

We further present three pairwise internally disjoint paths from a block  $B$  on a cluster  $S_k$  to  $\mathcal{P}_B(S_k)$ .

**Lemma 3.2.** *Let  $B$  be a block of  $S_k$ ,  $2 \leq k \leq n$ . If  $u'_k$  and  $u'_{k-1}$  are dummy vertices with dummy edges incident to every vertex of  $B$  and  $\mathcal{P}_B(S_k)$ , respectively, then the paths from  $u'_k$  to  $u'_{k-1}$  in  $T_i^0$ , for  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Proof.** Let  $V(B) = \{v_0, v_{\rho_1}, v_{\rho_2}, v_{\rho_3}\}$  and  $V(\mathcal{P}_B(S_k)) = \{t_0, t_{\rho_1}, t_{\rho_2}, t_{\rho_3}\}$ , where  $\pi_0(v_0) = \pi_0(t_0) = 0$  and  $\pi_0(v_{\rho_i}) = \pi_0(t_{\rho_i}) = \rho_i$  for each  $i$ . Let  $P_{\rho_i}$  denote the path from  $u'_k$  to  $u'_{k-1}$  in  $T_{\rho_i}^0$  and  $Q_{\rho_j}$  be the subpath of  $P_{\rho_j}$  on  $S_k$  for each  $i$ . Let  $\hat{h}_{k,\rho_1}$  and  $\hat{h}_{k,\rho_2}$  be outer vertices and  $\hat{h}_{k,\rho_3}$  be an inner vertex. By the definition of  $E(\hat{H}_{k,\rho_j})$ ,  $P_{\rho_j}$  must visit  $v_{\rho_j}$  and  $t_{\rho_j}$ , where  $j = 1, 2$ . We shall prove  $Q_{\rho_1}, Q_{\rho_2}, Q_{\rho_3}$  pairwise disjoint instead of showing  $P_{\rho_1}, P_{\rho_2}, P_{\rho_3}$  pairwise internally disjoint.

According to the **HFinding Algorithm**,  $v_{\rho_1} \in \{\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_1}^{-5}\}$  and  $v_{\rho_2} \in \{\hat{h}_{k,\rho_2}, \hat{h}_{k,\rho_2}^{+2}, \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-5}\}$ . Actually, the ordered pair  $(v_{\rho_1}, v_{\rho_2}) \in \{(\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_2}^{-5}), (\hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_2}^{-2}), (\hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_2}^{+2}), (\hat{h}_{k,\rho_1}^{-5}, \hat{h}_{k,\rho_2})\}$ . Since  $p(\hat{h}_{k,\rho_j})$  is  $t_{\rho_j}$  for each  $j$ , we get the ordered pair  $(Q_{\rho_1}, Q_{\rho_2})$  as follows:

$$(Q_{\rho_1}, Q_{\rho_2}) = \begin{cases} (\langle \hat{h}_{k,\rho_1} \rangle, \langle \hat{h}_{k,\rho_2}^{-5}, \hat{h}_{k,\rho_2}^{-4}, \dots, \hat{h}_{k,\rho_2} \rangle) & \text{if } (v_{\rho_1}, v_{\rho_2}) = (\hat{h}_{k,\rho_1}, \hat{h}_{k,\rho_2}^{-5}), \\ (\langle \hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_1}^{+1}, \hat{h}_{k,\rho_1} \rangle, \langle \hat{h}_{k,\rho_2}^{-2}, \hat{h}_{k,\rho_2}^{-1}, \hat{h}_{k,\rho_2} \rangle) & \text{if } (v_{\rho_1}, v_{\rho_2}) = (\hat{h}_{k,\rho_1}^{+2}, \hat{h}_{k,\rho_2}^{-2}), \\ (\langle \hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_1}^{-1}, \hat{h}_{k,\rho_1} \rangle, \langle \hat{h}_{k,\rho_2}^{+2}, \hat{h}_{k,\rho_2}^{+1}, \hat{h}_{k,\rho_2} \rangle) & \text{if } (v_{\rho_1}, v_{\rho_2}) = (\hat{h}_{k,\rho_1}^{-2}, \hat{h}_{k,\rho_2}^{+2}), \\ (\langle \hat{h}_{k,\rho_1}^{-5}, \hat{h}_{k,\rho_1}^{-4}, \dots, \hat{h}_{k,\rho_1} \rangle, \langle \hat{h}_{k,\rho_2} \rangle) & \text{if } (v_{\rho_1}, v_{\rho_2}) = (\hat{h}_{k,\rho_1}^{-5}, \hat{h}_{k,\rho_2}). \end{cases}$$

It is clear that  $Q_{\rho_1} \parallel Q_{\rho_2}$  for each case. Consider  $P_{\rho_3}$ . By the definition of  $E(\hat{H}_{k,\rho_3})$ ,  $P_{\rho_3}$  visits either  $t_0$  or  $t_{\rho_3}$ .

Case 1.  $P_{\rho_3}$  visits  $t_0$ .

By the definition of  $E(\hat{H}_{k,\rho_3})$ ,  $P_{\rho_3} = \langle u'_k, v_0, t_0, u'_{k-1} \rangle$  and thus  $Q_{\rho_3} = \langle v_0 \rangle$ . Clearly,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$  for every ordered pair  $(v_{\rho_1}, v_{\rho_2})$ .

*Case 2.*  $P_{\rho_3}$  visits both  $v_{\rho_3}$  and  $t_{\rho_3}$ .

Since every vertex of color  $\rho_3$  is an inner vertex,  $Q_{\rho_3}$  is a subpath of  $P_{k,\rho_3}$  and is in fact an  $i$ -path. However, both  $Q_{\rho_1}$  and  $Q_{\rho_2}$  are  $o$ -paths. So,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$ .

*Case 3.*  $P_{\rho_3}$  visits both  $v_0$  and  $t_{\rho_3}$ .

If  $v_0$  is a corner vertex, then  $v_0$  must be  $\hat{h}_{k,\rho_1}^{+3}$  and  $t_{\rho_3} = p(v_0)$ . Since, by the definition of  $E(\hat{H}_{k,\rho_3})$ ,  $(\hat{h}_{k,\rho_1}^{+3}, p(\hat{h}_{k,\rho_1}^{+3})) \in E(\hat{H}_{k,\rho_3})$ ,  $P_{\rho_3} = \langle u'_k, v_0, t_{\rho_3}, u'_{k-1} \rangle$ . That is,  $Q_{\rho_3} = \langle v_0 \rangle$ . So,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$  for every ordered pair  $(v_{\rho_1}, v_{\rho_2})$ . Otherwise,  $v_0$  is not a corner vertex and so  $v_0$  is adjacent to an inner vertex, say  $v$ . Since  $v \in V(P_{k,\rho_3})$ ,  $Q_{\rho_3} - v_0$  is a subpath of  $P_{k,\rho_3}$ . According to **HFinding Algorithm**,  $v_0 \in \{\hat{h}_{k,\rho_1}^{-1}, \hat{h}_{k,\rho_2}^{-1}\}$ . Since both  $Q_{\rho_1}$  and  $Q_{\rho_2}$  are  $o$ -paths and neither  $Q_{\rho_1}$  nor  $Q_{\rho_2}$  contains  $v_0$  for every ordered pair  $(v_{\rho_1}, v_{\rho_2})$ ,  $Q_{\rho_3} \parallel Q_{\rho_1}$  and  $Q_{\rho_3} \parallel Q_{\rho_2}$ .

Q. E. D.

**Theorem 3.1.** *Graphs  $T_1^0, T_2^0$  and  $T_3^0$  are 3-ISTs of  $\mathbb{P}_n$  rooted at apex.*

**Proof.** Let  $\rho_1, \rho_2$  and  $\rho_3$  be color numbers of the vertices  $v, v^+, v^-$  and  $v^{++}$  in  $S_1$ . Since  $E(\hat{H}_{1,\rho_1}) = E(\langle v^{++}, v^+, v, v^- \rangle) \cup \{(r, v^+)\}$ ,  $E(\hat{H}_{1,\rho_2}) = E(\langle v^{++}, v^-, v, v^+ \rangle) \cup \{(r, v^-)\}$  and  $E(\hat{H}_{1,\rho_3}) = E(\langle v^+, v^{++}, v^- \rangle) \cup \{(r, v), (r, v^{++})\}$ , every  $\hat{H}_{1,\rho_i}, i = 1, 2, 3$ , is a tree. According to **HFinding Algorithm**, every  $\hat{H}_{k,i}$  is acyclic, where  $2 \leq k \leq n$ . For connectivity, each  $\hat{h}_{k,i}$  has a layer edge  $(\hat{h}_{k,i}, p(\hat{h}_{k,i})) \in E(\hat{H}_{k,i})$  for connecting subgraphs  $\hat{H}_{k,i}$  and  $\hat{H}_{k-1,i}$ . It follows that  $T_i^0$  is a tree. Actually,  $T_i^0$  is a spanning tree of  $\mathbb{P}_n$  due to  $V(T_i^0) = V(\mathbb{P}_n)$ .

For any vertex  $u_k = (k; x, y) \in V(\mathbb{P}_n)$ , we now want to find three pairwise internally disjoint paths from  $u_k$  to the apex  $r$ . The case  $u_k = (0; 0, 0)$  is trivial. Consider  $k = 1$ . If  $u_k = \hat{h}_{1,0}$ , we get three pairwise internally disjoint paths  $\langle u_k, \hat{h}_{1,\rho_1}, r \rangle$ ,  $\langle u_k, \hat{h}_{1,\rho_2}, r \rangle$  and  $\langle u_k, r \rangle$  from  $u_k$  to  $r$  in  $T_{\rho_1}^0$ ,  $T_{\rho_2}^0$  and  $T_{\rho_3}^0$ , respectively. Similarly, we get three pairwise internally disjoint paths  $\langle u_k, r \rangle$ ,  $\langle u_k, \hat{h}_{1,0}, \hat{h}_{1,\rho_1}, r \rangle$  and  $\langle u_k, r \rangle$  for the case  $u_k = \hat{h}_{1,\rho_1}$ ; paths  $\langle u_k, \hat{h}_{1,\rho_0}, \hat{h}_{1,\rho_1}, r \rangle$ ,  $\langle u_k, r \rangle$  and  $\langle u_k, \hat{h}_{1,\rho_3}, r \rangle$  for  $u_k = \hat{h}_{1,\rho_2}$ ; and paths  $\langle u_k, \hat{h}_{1,\rho_1}, r \rangle$ ,  $\langle u_k, \hat{h}_{1,\rho_2}, r \rangle$  and  $\langle u_k, r \rangle$  for  $u_k = \hat{h}_{1,\rho_3}$ .

For  $k \geq 2$ , let  $u'_j$  be the dummy vertex with dummy edges incident to every vertex of  $\mathcal{P}_{\mathcal{B}}(S_{j+1})$ , where  $1 \leq j \leq k-1$ . From Lemma 3.1, we get pairwise internally disjoint paths  $X_{k,\rho_i}$  from  $u_k$  to  $u'_{k-1}$  in  $T_{\rho_i}^0$  for each  $i = 1, 2, 3$ . By applying Lemma 3.2, we get pairwise internally disjoint paths  $X_{j,\rho_i}$  from  $u'_j$  to  $u'_{j-1}$  in  $T_{\rho_i}^0$  for each  $i$ , where  $2 \leq j \leq k-1$ . Moreover, since  $(\hat{h}_{1,\rho_i}, r) \in E(\hat{H}_{1,\rho_i})$ ,

we get three pairwise internally disjoint paths from  $u_k$  to  $r$ .

Q. E. D.

**Theorem 3.2.** *The construction of 3-ISTs of  $\mathbb{P}_n$  rooted at the apex can be solved in  $O(|V|)$  time, where  $|V|$  is the number of vertices of  $\mathbb{P}_n$ .*

**Proof.** At first, we assign color numbers to the four vertices of  $S_1$  and build three subgraphs  $\hat{H}_{1,i}$ ,  $i = 1, 2, 3$ , on  $S_1$  in constant time. By using **HFinding Algorithm**, the coloring of every cluster is determined by the coloring of its parent block and costs constant time. So, the coloring of  $\mathbb{P}_n$  costs  $O(|V|)$  time. According to  $\pi_0$  coloring,  $\hat{h}_{k,1}$ ,  $\hat{h}_{k,2}$  and  $\hat{h}_{k,3}$ ,  $k = 2, 3, \dots, n$ , of  $S_k$  can be decided in constant time and then every  $\hat{H}_{k,i}$  is established in constant time. Since each edge set  $E(T_i^0)$ ,  $i = 1, 2, 3$ , has  $|V| - 1$  edges and each edge of  $E(T_i^0)$  can be checked in constant time, the finding of  $E(T_i^0)$  costs  $O(|V|)$  time. Therefore, the construction of 3-ISTs of  $\mathbb{P}_n$  rooted at the apex is solved in  $O(|V|)$  time.

Q. E. D.

### 3.2 Rooted at a Vertex on Layer 1

In  $S_1$ , we let  $r$  as the corner vertex and assign colors  $0, \rho_1, \rho_2$  and  $\rho_3$  to vertices  $r, r^+, r^-$  and  $r^{++}$ , respectively. Due to  $r \in V(S_1)$ ,  $r^+, r^-$  and  $r^{++}$  are actually  $\hat{h}_{1,\rho_1}, \hat{h}_{1,\rho_2}$  and  $\hat{h}_{1,\rho_3}$ , respectively. To color  $V(\mathbb{P}_n)$ , we also apply **HFinding Algorithm** iteratively on every block of  $\mathbb{P}_n$ . Since the coloring of a cluster can be decided by its parent block, the  $\pi_0$  coloring of  $S_2$  is determined by the coloring of  $S_1$ . In this way, the coloring and subgraphs of every cluster of  $\mathbb{P}_n$  can be established.

Let  $\hat{H}_{1,\rho_i}^1$  have the vertex set  $V(S_1) \cup \{\Delta\}$ . Let  $E(\hat{H}_{1,\rho_1}^1) = E(\langle r^-, r^{++}, r^+, r \rangle) \cup \{(r^+, \Delta)\}$ ,  $E(\hat{H}_{1,\rho_2}^1) = E(\langle r^+, r^{++}, r^-, r \rangle) \cup \{(r^-, \Delta)\}$  and  $E(\hat{H}_{1,\rho_3}^1) = \{(v, \Delta) | v \in V(S_1)\}$ . Notations  $S_k$ ,  $\hat{h}_{k,i}$  and  $\hat{H}_{k,i}$ , where  $2 \leq k \leq n$  and  $1 \leq i \leq 3$ , refer to Section 3.1. Let  $V(T_i^1) = V(\mathbb{P}_n)$  and  $E(T_i^1) = E(\hat{H}_{1,i}^1) \bigcup_{k=2}^n \bigcup_{a,b=0}^{2^{k-2}-1} E(\hat{H}_i(k; a, b))$  for each  $i$ .

**Theorem 3.3.** *Graphs  $T_1^1, T_2^1$  and  $T_3^1$  are 3-ISTs of  $\mathbb{P}_n$  rooted at a vertex of  $S_1$ .*

**Proof.** Let  $0, \rho_1, \rho_2$  and  $\rho_3$  be color numbers of the vertices  $r, r^+, r^-$  and  $r^{++}$  in  $S_1$ . Since  $E(\hat{H}_{1,\rho_1}^1) = E(\langle r^-, r^{++}, r^+, r \rangle) \cup \{(r^+, \Delta)\}$ ,  $E(\hat{H}_{1,\rho_2}^1) = E(\langle r^+, r^{++}, r^-, r \rangle) \cup \{(r^-, \Delta)\}$  and  $E(\hat{H}_{1,\rho_3}^1) = \{(v, \Delta) | v \in V(S_1)\}$ , every  $\hat{H}_{1,i}^1$ ,  $i = 1, 2, 3$ , is a tree. According to **HFinding Algorithm**, every  $\hat{H}_{k,i}$

is acyclic, where  $2 \leq k \leq n$ . For connectivity, each  $\hat{h}_{k,i}$  has a layer edge  $(\hat{h}_{k,i}, p(\hat{h}_{k,i})) \in E(\hat{H}_{k,i})$  for connecting subgraphs  $\hat{H}_{k,i}$  and  $\hat{H}_{k-1,i}$ , where  $3 \leq k \leq n$ . In addition, each  $\hat{h}_{2,i}$  has a layer edge  $(\hat{h}_{2,i}, p(\hat{h}_{2,i})) \in E(\hat{H}_{2,i})$  for connecting subgraphs  $\hat{H}_{2,i}$  and  $\hat{H}_{1,i}^1$ . It follows that  $T_i^1$  is a tree. Actually,  $T_i^1$  is a spanning tree of  $\mathbb{P}_n$  due to  $V(T_i^1) = V(\mathbb{P}_n)$ .

For any vertex  $u_k = (k; x, y) \in V(\mathbb{P}_n)$ , we now want to find three pairwise internally disjoint paths from  $u_k$  to  $r$ . If  $u_k$  is  $\Delta$ , by the definition of each  $\hat{H}_{1,\rho_i}^1$ , we have pairwise internally paths  $\langle u_k, r^+, r \rangle$ ,  $\langle u_k, r^-, r \rangle$  and  $\langle u_k, r \rangle$  in  $T_{\rho_1}^1$ ,  $T_{\rho_2}^1$  and  $T_{\rho_3}^1$ , respectively, from  $u_k$  to  $r$ . The case  $u_k = r$  is trivial. Consider the case  $k = 1$  and  $u_k \neq r$ . If  $u_k = r^{++}$ , then paths  $\langle u_k, r^+, r \rangle$ ,  $\langle u_k, r^-, r \rangle$  and  $\langle u_k, p(r), r \rangle$  are in  $T_{\rho_1}^1$ ,  $T_{\rho_2}^1$  and  $T_{\rho_3}^1$ , respectively, and are pairwise internally disjoint. Similarly we have three pairwise internally disjoint paths for the case  $u_k$  is a neighbor of  $r$  and leave the proof to the reader.

For  $k \geq 2$ , let  $u'_j$  be the dummy vertex with dummy edges incident to every vertex of  $\mathcal{P}_{\mathcal{B}}(S_{j+1})$ , where  $1 \leq j \leq k-1$ . From Lemma 3.1, we get pairwise internally disjoint paths  $X_{k,\rho_i}$  from  $u_k$  to  $u'_{k-1}$  in  $T_{\rho_i}^1$  for each  $i = 1, 2, 3$ . By applying Lemma 3.2, we get pairwise internally disjoint paths  $X_{j,\rho_i}$  from  $u'_j$  to  $u'_{j-1}$  in  $T_{\rho_i}^1$  for each  $i$ , where  $2 \leq j \leq k-1$ . Let  $X_{1,\rho_i}$  be the path from  $u'_1$  to  $r$  in  $T_{\rho_i}^1$  for each  $i$ . For each  $u'_j$ ,  $1 \leq j \leq k-1$ , if  $v$  is a neighbor of  $u'_j$  and  $\pi_0(v) = 0$ , then, by the definition of each  $E(\hat{H}_{j,\rho_i})$ , either  $v$  or  $v^{++}$  can be on one of the paths in  $\{X_{j,\rho_1}, X_{j,\rho_2}, X_{j,\rho_3}\}$ . Let  $v_i$  be the neighbor of  $u'_1$  in  $X_{1,\rho_i}$  for each  $i$ . Since  $\pi_0(r) = 0$ , by the definition of each  $E(\hat{H}_{2,\rho_i})$ , either  $r$  or  $r^{++}$  is a vertex in  $\{v_1, v_2, v_3\}$ . If  $r \in \{v_1, v_2, v_3\}$ , then we get three pairwise internally disjoint paths  $\langle u'_1, r \rangle$ ,  $\langle u'_1, r^+, r \rangle$  and  $\langle u'_1, r^-, r \rangle$ . Otherwise, we get three pairwise internally disjoint paths  $\langle u'_1, r^{++}, \Delta, r \rangle$ ,  $\langle u'_1, r^+, r \rangle$  and  $\langle u'_1, r^-, r \rangle$ . Therefore, we get three pairwise internally disjoint paths from  $u_k$  to  $r$ .

Q. E. D.

**Theorem 3.4.** *The construction of 3-ISTs of  $\mathbb{P}_n$  rooted at a vertex on layer 1 can be solved in  $O(|V|)$  time, where  $|V|$  is the number of vertices of  $\mathbb{P}_n$ .*

**Proof.** Similar to the proof of Theorem 3.2.

Q. E. D.

### 3.3 Rooted at Vertices on Higher Layers

The construction of 3-ISTs rooted at  $r$ ,  $2 \leq l(r) \leq n$ , has two stages: *Backbone Establishment Stage* and *Branching Stage*. In Backbone Establishment Stage, we shall first define subgraphs on every cluster  $S_k$  that contains an ancestor of  $r$ . Then dedicated layer edges of  $\mathbb{P}_n$  are selected for connecting subgraphs of  $S_k$  and  $S_{k-1}$ , for each  $k = 1, 2, \dots, l(r)$ , to establish tree backbones. We further do an expansion on tree backbones to build 3-ISTs of  $\mathbb{P}_n$  rooted at  $r$  in Branching Stage. The detailed description of the two stages are given in the following two subsections.

#### 3.3.1 Backbone Establishment Stage

A block is an *a-block* if it contains an ancestor of  $r$  (including  $r$ ). Let  $A_k$  denote the *a-block* on layer  $k$  of  $\mathbb{P}_n$ , where  $1 \leq k \leq l(r)$ . Let  $S_k$  be the cluster containing an *a-block* on layer  $k$  and  $\mathcal{BB}_r$  be the subgraph induced by the vertex set  $\bigcup_{k=0}^{l(r)} V(S_k)$ . A block  $B$  of  $\mathcal{BB}_r$  is said to be a *b-block* if  $B$  is not an *a-block*. Every vertex of an *a-block* (respectively, a *b-block*) is called an *a-vertex* (respectively, a *b-vertex*). The corner vertex and inner vertex of  $A_k$  are denoted as  $a_{k,0}$  and  $a_{k,3}$ , respectively, and  $a_{k,0}^+$  and  $a_{k,0}^-$  are also denoted by  $a_{k,1}$  and  $a_{k,2}$ , respectively. Every  $V(S_k)$  can be partitioned into four vertex subsets  $U_{k,0}$ ,  $U_{k,1}$ ,  $U_{k,2}$  and  $U_{k,3}$ . Let  $U_{k,0} = \{a_{k,0}, a_{k,0}^{+3}, a_{k,0}^{+6}, a_{k,0}^{+9}\}$ ,  $U_{k,1} = \{a_{k,0}^{+1}, a_{k,0}^{+2}, a_{k,0}^{+5}, a_{k,0}^{+8}\}$ ,  $U_{k,2} = \{a_{k,0}^{+11}, a_{k,0}^{+10}, a_{k,0}^{+7}, a_{k,0}^{+4}\}$  and  $U_{k,3} = \{a_{k,3}, a_{k,3}^{-1}, a_{k,3}^{-2}, a_{k,3}^{-3}\}$ . Notice that every  $U_{k,i}$  contains exactly one vertex of each block of  $\mathcal{BB}_r$ .

Let  $\pi_1 : V \rightarrow \{0, 1, 2, 3\}$  be a coloring on  $\mathcal{BB}_r$ , where for each vertex  $v \in V(\mathcal{BB}_r)$ ,  $\pi_1(v) \in \{0, 1, 2, 3\}$ . For an  $S_k$ ,  $2 \leq k \leq l(r) - 1$ , if  $A_k$  is the top-left block or top-right block of  $S_k$ , we assign color  $i$  to every vertex of  $U_{k,i}$ . Otherwise, we assign colors 0, 2, 1 and 3 to every vertex of  $U_{k,0}$ ,  $U_{k,1}$ ,  $U_{k,2}$  and  $U_{k,3}$ , respectively. According to the above coloring rules for  $\pi_1$ , we get colorings  $\Pi_0$ ,  $\Pi_1$ ,  $\Pi_2$  and  $\Pi_3$  for a cluster with respect to an  $A_k$  locating at the top-left, top-right, bottom-left and bottom-right block of  $S_k$ , respectively. Figure 3 depicts the four possible  $\pi_1$  colorings  $\Pi_0$ ,  $\Pi_1$ ,  $\Pi_2$  and  $\Pi_3$  of a cluster  $S_k$ . The block having grey vertices along with bold edges indicates an *a-block*.

Let  $h_{k,i}$ ,  $i = 0, 1, 2, 3$ , denote the *a-vertex* of color  $i$  in  $A_k$  and  $h_k = \{h_{k,1}, h_{k,2}, h_{k,3}\}$ . The coloring of  $S_1$  is the same as the coloring of  $A_2$ . That is, if  $v$  is the corner vertex of  $A_2$ , then we assign colors  $\pi_1(v)$ ,  $\pi_1(v^+)$ ,  $\pi_1(v^-)$  and  $\pi_1(v^{++})$  to  $p(v)$ ,  $p(v)^+$ ,  $p(v)^-$  and  $p(v)^{++}$ , respectively. The  $\pi_1$  coloring of  $S_{l(r)}$  is treated as a special case of  $\pi_1$  and determined by the  $\pi_1$  coloring of  $A_{l(r)-1}$  in

RootColoring Algorithm.

---



---

**Input:**  $\pi_1$  coloring of  $A_{l(r)-1}$ .

**Output:**  $\pi_1$  coloring of  $S_{l(r)}$ .

**begin**

Assign color 0 to all vertices of  $U_{l(r),0}$ ;

Case  $\pi_1(p(r)) \in \{0, 3\}$ .

Assign color 3 to all vertices of  $U_{l(r),3}$ ;

If  $\pi_1(p(r)^+) = 1$ , then assign color 1 to all vertices of  $U_{l(r),1}$  and color 2 to  $U_{l(r),2}$   
else assign color 2 to every vertex of  $U_{l(r),1}$  and color 1 to  $U_{l(r),2}$ ;

Case  $\pi_1(p(r)) = 1$ .

Assign color 1 to all vertices of  $U_{l(r),3}$ ;

If  $\pi_1(p(r)^+) = 0$ , then assign color 2 to all vertices of  $U_{l(r),1}$  and color 3 to  $U_{l(r),2}$   
else assign color 3 to every vertex of  $U_{l(r),1}$  and color 2 to  $U_{l(r),2}$ ;

Case  $\pi_1(p(r)) = 2$ .

Assign color 2 to all vertices of  $U_{l(r),3}$ ;

If  $\pi_1(p(r)^+) = 0$ , then assign color 1 to all vertices of  $U_{l(r),1}$  and color 3 to  $U_{l(r),2}$   
else assign color 3 to every vertex of  $U_{l(r),1}$  and color 1 to  $U_{l(r),2}$ ;

Adjust the coloring of  $A_{l(r)}$ :

Assign color  $\pi_1(r^{++})$  to  $a_{l(r),0}$ , and then assign color  $\pi_1(r)$  to  $r^{++}$ ;

Assign color 0 to  $r$ ;

**end**

RootColoring Algorithm

---

We are able to establish three tree backbones  $TB_{r,i}$  on  $\mathcal{BB}_r$  for each  $i = 1, 2, 3$ . Each  $TB_{r,i}$  is regarded as an  $l(r)$ -layered graph. For a  $v \in V(\mathcal{BB}_r)$ , let  $c_i(v)$  denote the child of  $v$  of color  $i$ . Let  $H_{k,i}$  denote a subgraph of an  $S_k$ , where  $1 \leq k \leq l(r)$  and  $1 \leq i \leq 3$ . Let  $V(H_{1,i}) = \{h_{1,0}, h_{1,1}, h_{1,2}, h_{1,3}, \Delta\}$  and  $E(H_{1,i}) = \{(h_{1,i}, \Delta)\}$ . We define two  $o$ -paths

$P_{k,\rho_1} = \langle a_{k,1}, a_{k,1}^{+1}, a_{k,1}^{+2}, \dots, a_{k,1}^{+9} \rangle$ ,  $P_{k,\rho_2} = \langle a_{k,2}, a_{k,2}^{-1}, a_{k,2}^{-2}, \dots, a_{k,2}^{-9} \rangle$  and an  $i$ -path  $P_{k,\rho_3} = \langle a_{k,3}, a_{k,3}^{-1}, a_{k,3}^{-2}, a_{k,3}^{-3} \rangle$

for every  $S_k$ . For convenience, let  $a'_1, a'_2$  and  $a'_3$  denote as  $a_{l(r),1}^{+1}, a_{l(r),2}^{-1}$  and  $a_{l(r),3}^{-1}$ , respectively, and let  $\rho_i = \pi_1(a'_i)$  for each  $i$ . The three subgraphs  $H_{k,\rho_1}$ ,  $H_{k,\rho_2}$  and  $H_{k,\rho_3}$  of  $S_k$  are defined in the following:

$$E(H_{k,\rho_j}) = E(P_{k,\rho_j}) \cup \{(u, v) \in E(S_k) | u \notin V(A_k), \pi_1(u) = \rho_3 \text{ and } \pi_1(v) = \rho_j\} \cup \{(v, p(v)) | v \notin V(A_{l(r)}) \text{ and } \pi_1(v) = \rho_j\}$$
 for  $j = 1, 2$ , and

$$E(H_{k,\rho_3}) = E(P_{k,\rho_3}) \cup \{(u, v) \in E(S_k) | u \notin V(A_k), \pi_1(u) = \rho_3 \text{ and } \pi_1(v) = \rho_1 \text{ or } \rho_2\} \\ \cup \{(v, p(v)) | v \notin V(A_{l(r)}) \text{ and } \pi_1(v) = \rho_3\} \cup \{(v, p(v)) | v \notin V(A_k) \text{ and } \pi_1(v) = 0\}.$$

Among the four  $a$ -vertices of an  $A_k$ ,  $2 \leq k \leq l(r) - 1$ ,  $h_{k,\rho_i}$  is adjacent to a  $b$ -vertex of  $H_{k,\rho_i}$  and its parent, while the other three  $a$ -vertices are isolated in  $H_{k,\rho_i}$ . For connectivity, every  $a$ -vertex of  $H_{k,\rho_i}$  is incident to a child of color  $\rho_i$  in  $H_{k+1,\rho_i}$ . Figure 4 illustrates subgraphs  $H_{k,\rho_1}, H_{k,\rho_2}$  and

$H_{k,\rho_3}$  of  $S_k$ . Bold edges are edges of subgraphs  $H_{k,\rho_i}, 1 \leq i \leq 3$ , and grey vertices are  $a$ -vertices.

Let  $\mathcal{E}_1 = E(\langle r^+, r^{++}, r^-, r \rangle) \cup \{(r^-, p(r))\}$ ,  $\mathcal{E}_2 = E(\langle r^-, r^{++}, r^+, r \rangle) \cup \{(r^+, p(r))\}$  and  $\mathcal{E}_3 = \{(v, p(r)) | v \in V(A_{l(r)})\}$ . For each  $TB_{r,i}$ , let  $R_i$  denote the subgraph with the vertex set  $V(A_{l(r)}) \cup \{p(r)\}$  and an edge set in  $\{\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3\}$ , where  $i = 1, 2, 3$ . The formal definition of each  $TB_{r,i}$  is in the following:

**Definition 3.1.**

(1)  $E(R_{\pi_1(r^-)}) = \mathcal{E}_1$ ,  $E(R_{\pi_1(r^+)}) = \mathcal{E}_2$  and  $E(R_{\pi_1(r^{++})}) = \mathcal{E}_3$ .

(2)  $V(TB_{r,\rho_i}) = V(\mathcal{BB}_r)$ ,

$$E(TB_{r,\rho_i}) = E(H_{1,\rho_i}) \cup E(R_{\rho_i}) \bigcup_{k=2}^{l(r)} \bigcup_{a,b=0}^{2^{k-2}-1} E(H_{\rho_i}(k; a, b)), \text{ where } \rho_i = \pi_1(a'_i) \text{ and } i = 1, 2, 3.$$

**Lemma 3.3.** *The graphs  $TB_1, TB_2$  and  $TB_3$  are spanning trees of  $\mathcal{BB}_r$  rooted at  $r$ .*

**Proof.** Since  $E(H_{1,i}) = \{(h_{1,i}, \Delta)\}, 1 \leq i \leq 3$ , every  $H_{1,i}$  is clearly acyclic. Since each  $H_{k,i}, k = 2, 3, \dots, l(r) - 1$ , is constructed as a tree along with three isolated  $a$ -vertices on  $S_k$ , every  $H_{k,i}$  is acyclic. In the  $S_{l(r)}$ ,  $A_{l(r)}$  and  $p(r)$  form three tree subgraphs  $R_1, R_2$  and  $R_3$ , and the remaining three  $b$ -blocks form tree subgraphs  $H_{l(r),1}, H_{l(r),2}$  and  $H_{l(r),3}$ . In addition,  $H_{l(r),i}$  and  $R_i$  share a common  $a$ -vertex  $u \in \{a_{l(r),1}, a_{l(r),2}, a_{l(r),3}\}$ . For connectivity, each vertex  $v \notin V(A_{l(r)})$  of color  $i$  in  $H_{k,i}$  has an edge  $(v, p(v)) \in E(H_{k,i})$  for connecting subgraphs  $H_{k,i}$  and  $H_{k-1,i}$ , where  $k = 2, 3, \dots, l(r)$  and each  $p(v)$  is in fact an  $a$ -vertex of  $H_{k-1,i}$ . Thus,  $TB_{r,i}$  is a tree. Actually, each  $TB_{r,i}$  is a spanning tree because  $V(TB_{r,i}) = V(\mathcal{BB}_r)$ .

Q. E. D.

For convenience, let  $u'_j$  be the dummy vertex with dummy edges incident to every vertex of  $h_j$  for  $j = 1, 2, \dots, l(r) - 1$  and  $u'_{l(r)}$  be the dummy vertex with dummy edges incident to  $a'_1, a'_2$  and  $a'_3$  in the remaining text of this section.

**Lemma 3.4.** *The paths from  $u'_{k-1}$  to  $u'_k$  in each  $TB_{r,i}$  are pairwise internally disjoint, where  $1 \leq i \leq 3$  and  $2 \leq k \leq l(r) - 1$ .*

**Proof.** Let  $a_{k,0}^{+1}$  and  $a_{k,0}^{+11}$  be  $h_{k,\rho_1}$  and  $h_{k,\rho_2}$ , respectively. Let  $P_{\rho_i}$  be the path from  $c_{\rho_i}(h_{k-1,\rho_i})$  to  $h_{k,\rho_i}$  in  $H_{k,\rho_i}$  for  $i = 1, 2, 3$ . Since  $c_{\rho_i}(h_{k-1,\rho_i}) \in V(P_{k,\rho_i})$ ,  $P_{\rho_i}$  is a subpath of  $P_{k,\rho_i}$ . Then  $P_{\rho_3} \parallel P_{\rho_1}$  and  $P_{\rho_3} \parallel P_{\rho_2}$  because  $P_{k,\rho_3} \parallel P_{k,\rho_1}$  and  $P_{k,\rho_3} \parallel P_{k,\rho_2}$ . It remains to prove  $P_{\rho_1} \parallel P_{\rho_2}$ .

We first consider the case when  $k = 2$ . By the definition of the  $\pi_1$  coloring, the coloring of  $S_1$  is the same as the coloring of  $A_2$ . So,  $a_{k,0}^{+2} = c_{\rho_1}(h_{1,\rho_1})$  and  $a_{k,0}^{+10} = c_{\rho_2}(h_{1,\rho_2})$ . Thus, we get disjoint paths  $P_{\rho_1} = \langle a_{k,0}^{+2}, a_{k,0}^{+1} \rangle$  and  $P_{\rho_2} = \langle a_{k,0}^{+10}, a_{k,0}^{+11} \rangle$ . For the case  $3 \leq k \leq l(r) - 1$ , if  $c_{\rho_1}(h_{k-1,\rho_1})$  is an  $a$ -vertex,  $P_{\rho_1} = \langle h_{k,\rho_1} \rangle$ . By the definition of  $H_{k,\rho_2}$ ,  $h_{k,\rho_1} \notin V(H_{k,\rho_2})$ . That is,  $h_{k,\rho_1} \notin V(P_{\rho_2})$ . Thus,  $P_{\rho_1} \parallel P_{\rho_2}$ . Due to the same reason,  $P_{\rho_1}$  and  $P_{\rho_2}$  are disjoint for the case that  $c_{\rho_2}(h_{k-1,\rho_2})$  is an  $a$ -vertex. Consider the case that both  $c_{\rho_1}(h_{k-1,\rho_1})$  and  $c_{\rho_2}(h_{k-1,\rho_2})$  are  $b$ -vertices. Since  $h_{k-1,\rho_1}$  and  $h_{k-1,\rho_2}$  are not adjacent,  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1})$  and  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_2})$  must be two non-adjacent  $b$ -blocks of  $S_k$ . So,  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1})$  is either  $A_k^-$  or  $A_k^+$ .

*Case 1.*  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1}) = A_k^-$ .

We first assume that  $A_k$  is the top-left block of  $S_k$ . Then  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1})$  is the bottom-left block of  $S_k$  and hence  $h_{k-1,\rho_1}$  is the bottom-left vertex of  $A_{k-1}$ . Based on the coloring rules of  $\pi_1$ , every vertex of  $U_{k,i}$  is assigned the color number  $i$  when  $A_k$  is the top-left block of  $S_k$ . So,  $\pi_1(h_{k,\rho_1}) = \pi_1(a_{k,0}^{+1}) = 1$ . Then  $\rho_1 = 1$  and  $\pi_1(h_{k-1,\rho_1}) = 1$ . However, every bottom-left vertex of an  $a$ -block must be of color 0 (in  $\Pi_2$ ), 3 (in  $\Pi_1$ ) or 2 (in  $\Pi_0$  and  $\Pi_3$ ) which contradicts the fact that  $h_{k-1,\rho_1}$  is the bottom-left vertex of  $A_{k-1}$ . This implies that  $A_k$  can not be the top-left block of  $S_k$ . Due to the same reason, contradiction occurs on the condition that  $A_k$  is the top-right block, bottom-left block or bottom-right block of  $S_k$ .

*Case 2.*  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1}) = A_k^+$ .

Since, by the definition of  $U_{k,1}$ ,  $\pi_1(a_{k,0}^{+1}) = \pi_1(a_{k,0}^{+2})$ , it follows that  $a_{k,0}^{+2}$  is the unique vertex of color  $\rho_1$  in  $\mathcal{C}_{\mathcal{B}}(h_{k-1,\rho_1})$  and hence  $c_{\rho_1}(h_{k-1,\rho_1})$  must be  $a_{k,0}^{+2}$ . So,  $P_{\rho_1} = \langle a_{k,0}^{+2}, a_{k,0}^{+1} \rangle$ . We also get  $P_{\rho_2} = \langle a_{k,0}^{+10}, a_{k,0}^{+11} \rangle$  because both  $a_{k,0}^{+10}$  and  $a_{k,0}^{+11}$  are in  $U_{k,2}$  and are of the same color. Therefore,  $P_{\rho_1} \parallel P_{\rho_2}$ .

Q. E. D.

For any vertex  $v$  of  $\mathcal{BB}_r$ , we want to show the paths from  $v$  to  $r$  in  $TB_1, TB_2$  and  $TB_3$  are pairwise internally disjoint. Each path from  $v$  to  $r$  in a  $TB_{r,i}$ ,  $1 \leq i \leq 3$ , can be regarded as a concatenation of three subpaths, including the subpath from  $v$  to its nearest  $a$ -vertex  $h_{k,i}$ ,  $l(v) \leq l(k)$  (see Lemma 3.5), the subpath from  $h_{k,i}$  to  $h_{l(r)-1,i}$  (see Lemma 3.3), the subpath from  $h_{l(r)-1,i}$  to a vertex  $a$  of  $A_{l(r)}$  (see Corollaries 3.1 and 3.2) and the subpath from  $a$  to  $r$  (see Lemma 3.7). We are ready to give proofs for the above subpaths.



**Lemma 3.5.** *If  $u_k$  is a  $b$ -vertex of  $S_k$ ,  $2 \leq k \leq l(r) - 1$ , then the paths from  $u_k$  to  $u'_k$  in each  $TB_{r,i}$ ,  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Proof.** Let  $a_{k,0}^{+1}$  and  $a_{k,0}^{+11}$  be  $h_{k,\rho_1}$  and  $h_{k,\rho_2}$ , respectively. Since each  $TB_{r,\rho_i}$ ,  $i = 1, 2, 3$ , by Lemma 3.3, is a spanning tree of  $\mathcal{BB}_r$ , we have the path  $P_{\rho_i}$  from  $u_k$  to  $h_{k,\rho_i}$  in  $TB_{r,\rho_i}$  for each  $i$ . We first prove  $V(P_{\rho_3}) \cap V(P_{\rho_j}) = \{u_k\}$ , where  $j = 1, 2$ . If  $u_k$  is a corner vertex, then  $\pi_1(u_k) = 0$  and  $u_k^{++}$  is an inner vertex of color  $\rho_3$ . By the definition of  $H_{k,\rho_3}$ ,  $(u_k, p(u_k)), (u_k^{++}, p(u_k)) \in E(H_{k,\rho_3})$ . Thus,  $\langle u_k, p(u_k), u_k^{++} \rangle$  is a subpath of  $P_{\rho_3}$ . Since  $u_k^{++} \in V(P_{k,\rho_3})$ ,  $P_{\rho_3} - u_k - p(u_k)$  is a subpath of  $P_{k,\rho_3}$ . In contrast,  $P_{\rho_j}$  is a subpath of  $P_{k,\rho_j}$  because  $u_k \in V(P_{k,\rho_j})$ . Moreover, since  $p(u_k) \notin V(P_{k,\rho_j})$  and  $P_{k,\rho_3} \parallel P_{k,\rho_j}$ ,  $V(P_{\rho_3}) \cap V(P_{\rho_j}) = \{u_k\}$ . Otherwise,  $u_k$  is either an inner vertex or adjacent to an inner vertex. Then  $P_{\rho_j} - u_k$  and  $P_{\rho_3} - u_k$  are subpaths of  $P_{k,\rho_j}$  and  $P_{k,\rho_3}$ , respectively. Therefore,  $V(P_{\rho_3}) \cap V(P_{\rho_j}) = \{u_k\}$ . It remains to prove  $V(P_{\rho_1}) \cap V(P_{\rho_2}) = \{u_k\}$ .

*Case 1.*  $u_k$  is an outer vertex.

Let  $u_k = a_{k,0}^{+d}$ , where  $2 \leq d \leq 10$ . Since  $h_{k,\rho_1} = a_{k,0}^{+1}$  and  $h_{k,\rho_2} = a_{k,0}^{+11}$ ,  $P_{\rho_1} = \langle a_{k,0}^{+d}, a_{k,0}^{+(d-1)}, \dots, a_{k,0}^{+1} \rangle$  and  $P_{\rho_2} = \langle a_{k,0}^{+d}, a_{k,0}^{+(d+1)}, \dots, a_{k,0}^{+11} \rangle$ . Clearly,  $V(P_{\rho_1}) \cap V(P_{\rho_2}) = \{u_k\}$ .

*Case 2.*  $u_k$  is an inner vertex.

Every inner vertex is of color  $\rho_3$  and has neighbors  $a_{k,0}^{+d}$  ( $\in U_{k,1}$ ) and  $a_{k,0}^{+(d+2)}$  ( $\in U_{k,2}$ ) of colors  $\rho_1$  and  $\rho_2$ , respectively, where  $d \in \{2, 5, 8\}$ . By the definitions of  $H_{k,\rho_1}$  and  $H_{k,\rho_2}$ ,  $(u_k, a_{k,0}^{+d}) \in E(H_{k,\rho_1})$  and  $(u_k, a_{k,0}^{+(d+2)}) \in E(H_{k,\rho_2})$ . Then paths  $P_{\rho_1} = \langle u_k, a_{k,0}^{+d}, a_{k,0}^{+(d-1)}, \dots, a_{k,0}^{+1} \rangle$  and  $P_{\rho_2} = \langle u_k, a_{k,0}^{+(d+2)}, a_{k,0}^{+(d+3)}, \dots, a_{k,0}^{+11} \rangle$  share the common vertex  $u_k$ .

Q. E. D.

The next result follows immediately.

**Corollary 3.1.** *If  $u_{l(r)}$  is a  $b$ -vertex of  $S_{l(r)}$ , then the paths from  $u_{l(r)}$  to  $u'_{l(r)}$  in each  $TB_{r,i}$ ,  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Lemma 3.6.** *If  $u_k$  is an  $a$ -vertex of  $S_k$ ,  $1 \leq k \leq l(r) - 2$ , then the paths from  $u_k$  to  $u'_{k+1}$  in each  $TB_{r,i}$ ,  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Proof.** Let  $a_{k+1,0}^{+1}$  and  $a_{k+1,0}^{+11}$  be  $h_{k+1,\rho_1}$  and  $h_{k+1,\rho_2}$ , respectively. Let  $B$  be the block containing  $c_{\rho_i}(u_k)$  for  $i = 1, 2, 3$ . By the definition of  $E(H_{k+1,\rho_i})$ , the edge  $(u_k, c_{\rho_i}(u_k)) \in E(H_{k+1,\rho_i})$  for each  $i$ . If  $B$  is an  $a$ -block, paths  $\langle u_k, c_{\rho_i}(u_k), u'_{k+1} \rangle$  in  $TB_{r,\rho_i}$ , for  $i = 1, 2, 3$ , are pairwise internally disjoint.

For the case  $B$  is a  $b$ -block, let  $P_{\rho_i}$  be the path from  $c_{\rho_i}(u_k)$  to  $h_{k+1,\rho_i}$  in  $H_{k+1,\rho_i}$ . Since  $c_{\rho_i}(u_k) \in V(P_{k+1,\rho_i})$ ,  $P_{\rho_i}$  is a subpath of  $P_{k+1,\rho_i}$ . Recall that  $P_{k+1,\rho_3}$  is an  $i$ -path, whereas both  $P_{k+1,\rho_1}$  and  $P_{k+1,\rho_2}$  are  $o$ -paths. Thus,  $P_{\rho_3} \parallel P_{\rho_1}$  and  $P_{\rho_3} \parallel P_{\rho_2}$ . It remains to prove  $P_{\rho_1} \parallel P_{\rho_2}$ . Since  $c_{\rho_1}(u_k) \in U_{k+1,1}$  and  $c_{\rho_2}(u_k) \in U_{k+1,2}$ , we let  $a_{k+1,0}^{+d}$  and  $a_{k+1,0}^{+(d+2)}$  be  $c_{\rho_1}(u_k)$  and  $c_{\rho_2}(u_k)$ , respectively, where  $d \in \{2, 5, 8\}$ . Then paths  $P_{\rho_1} = \langle a_{k+1,0}^{+d}, a_{k+1,0}^{+(d-1)}, \dots, a_{k+1,0}^{+1} \rangle$  and  $P_{\rho_2} = \langle a_{k+1,0}^{+(d+2)}, a_{k+1,0}^{+(d+3)}, \dots, a_{k+1,0}^{+11} \rangle$  are disjoint.

Q. E. D.

**Corollary 3.2.** *If  $u_{l(r)-1}$  is an  $a$ -vertex of  $S_{l(r)-1}$ , then the paths from  $u_{l(r)-1}$  to  $u'_{l(r)}$  in each  $TB_{r,i}$ ,  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Lemma 3.7.** *The paths from  $u'_{l(r)}$  to  $r$  in each  $TB_{r,i}$ , for  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Proof.** Let  $\pi_1(a'_i) = \rho_i$  for  $i = 1, 2, 3$ . Recall that  $(a_{l(r),i}, a'_i) \in E(H_{l(r),\rho_i})$ . Let  $P_{\rho_i} \subseteq R_{\rho_i}$  be the path from  $a_{l(r),i}$  to  $r$  in  $TB_{r,\rho_i}$  for each  $i$ .

*Case 1.*  $r$  is  $a_{l(r),1}$ .

Intuitively,  $P_{\rho_1} = \langle r \rangle$ . Vertices  $r^+$ ,  $r^-$  and  $r^{++}$  are  $a_{l(r),3}$ ,  $a_{l(r),0}$  and  $a_{l(r),2}$ , respectively. Since  $\pi_1(r^+) = \rho_3$  and  $E(R_{\pi_1(r^+)}) = \mathcal{E}_2$ ,  $E(R_{\rho_3}) = \mathcal{E}_2$ . Then  $P_{\rho_3} = \langle r^+, r \rangle$  reveals the fact that  $P_{\rho_3} \subseteq R_{\rho_3}$ . Recall that we assign color  $\pi_1(r^{++})$  to  $a_{l(r),0}$  in RootColoring Algorithm. So  $\pi_1(r^-) = \rho_2$ . Furthermore, since  $E(R_{\pi_1(r^-)}) = \mathcal{E}_1$ , it follows that  $E(R_{\rho_2}) = \mathcal{E}_1$ . Then,  $P_{\rho_2} = \langle r^{++}, r^-, r \rangle$  due to  $P_{\rho_2} \subseteq R_{\rho_2}$ . Clearly,  $P_{\rho_1}, P_{\rho_2}$  and  $P_{\rho_3}$  are pairwise internally disjoint.

*Case 2.*  $r \in \{a_{l(r),0}, a_{l(r),2}, a_{l(r),3}\}$ .

Similar to the proof of *Case 1*.

Q. E. D.

**Lemma 3.8.** *The paths from  $u'_k$  to  $r$  in each  $TB_{r,i}$  are pairwise internally disjoint, where  $1 \leq k \leq l(r) - 1$  and  $1 \leq i \leq 3$ .*

**Proof.** Let  $\rho_i = \pi_1(a'_i)$  for  $i = 1, 2, 3$ . By Lemma 3.4, we have pairwise internally disjoint paths  $X_{j-1,\rho_i}$  from  $u'_{j-1}$  to  $u'_j$ , where  $1 \leq i \leq 3$ ,  $k+1 \leq j \leq l(r) - 1$  and  $h_{j-1,\rho_i}, h_{j,\rho_i} \in V(X_{j-1,\rho_i})$ . For each  $i$ , let  $X_{l(r)-1,\rho_i}$  be the path from  $u'_{l(r)-1}$  to  $u'_{l(r)}$  and let  $X_{l(r),\rho_i}$  be the path from  $u'_{l(r)}$  to  $r$  in  $TB_{r,\rho_i}$ . Let  $Y_{l(r)-1,\rho_i}$  be the subpath of  $X_{l(r)-1,\rho_i}$  from  $h_{l(r)-1,\rho_i}$  to  $a'_i$  in  $TB_{r,\rho_i}$  for each  $i$ . Notably,

$h_{l(r)-1, \rho_{i_1}}, a'_{i_1} \notin V(X_{l(r)-1, \rho_{i_2}})$  if  $i_1 \neq i_2$ . We shall list pairwise internally disjoint paths from  $u'_{l(r)-1}$  to  $r$  in each  $TB_{r, \rho_i}$  to complete the proof.

*Case 1.*  $\pi_1(p(r)) = 0$ .

From **RootColoring Algorithm**, every vertex of  $U_{l(r),1}, U_{l(r),2}$  and  $U_{l(r),3}$  is assigned the color  $\pi_1(p(r)^+), \pi_1(p(r)^-)$  and  $\pi_1(p(r)^{++})$ , respectively. So,  $\pi_1(a'_1) = \pi_1(p(r)^+)$ ,  $\pi_1(a'_2) = \pi_1(p(r)^-)$  and  $\pi_1(a'_3) = \pi_1(p(r)^{++})$ , implying  $p(r)^+, p(r)^-$  and  $p(r)^{++}$  are  $h_{l(r)-1, \rho_1}, h_{l(r)-1, \rho_2}$  and  $h_{l(r)-1, \rho_3}$ , respectively. Since  $c_{\rho_1}(h_{l(r)-1, \rho_1}) = a'_1$  and  $c_{\rho_2}(h_{l(r)-1, \rho_2}) = a'_2$ , it follows that  $Y_{l(r)-1, \rho_1} = \langle h_{l(r)-1, \rho_1}, a'_1 \rangle$  and  $Y_{l(r)-1, \rho_2} = \langle h_{l(r)-1, \rho_2}, a'_2 \rangle$ . Since  $h_{l(r)-1, \rho_1}, h_{l(r)-1, \rho_2}, a'_1, a'_2 \notin V(Y_{l(r)-1, \rho_3})$ ,  $Y_{l(r)-1, \rho_1}, Y_{l(r)-1, \rho_2}$  and  $Y_{l(r)-1, \rho_3}$  are pairwise disjoint. Moreover, by Lemma 3.7, we have pairwise internally disjoint paths  $X_{l(r), \rho_i}$  from  $u'_{l(r)}$  to  $r$ , where  $1 \leq i \leq 3$  and  $a'_i \in V(X_{l(r), \rho_i})$ .

*Case 2.*  $\pi_1(p(r)) = 1$ .

Vertex  $p(r)$  is  $h_{l(r)-1,1}$ . By the definition of  $R_1$ , the path from  $p(r)$  to  $r$  in  $TB_{r,1}$ , denoted  $P_1$ , is a subgraph of  $R_1$ . From **RootColoring Algorithm**, every vertex of  $U_{l(r),3}$  is assigned color 1. If  $\pi_1(p(r)^+) = 0$ , every vertex of  $U_{l(r),1}$  and  $U_{l(r),2}$  is assigned color 2 and 3, respectively. So,  $p(r)^-$  and  $p(r)^{++}$  are  $h_{l(r)-1,3}$  and  $h_{l(r)-1,2}$ , respectively. Then  $Y_{l(r)-1,3} = \langle p(r)^-, a'_2 \rangle$ . Since  $p(r)^-, a'_2 \notin V(Y_{l(r)-1, \rho_3})$ ,  $Y_{l(r)-1,2}$  and  $Y_{l(r)-1,3}$  are disjoint. By Lemma 3.7, we have internally disjoint paths  $X_{l(r), i_2}$  from  $u'_{l(r)}$  to  $r$ , where  $i_2 = 2, 3$  and  $a'_{i_2} \in V(X_{l(r), i_2})$ . Notably, each  $X_{l(r), i_2}$  is a subgraph of  $R_{i_2}$  and shares a unique common vertex  $r$  with  $P_1$ . For the case  $\pi_1(p(r)^+) = 3$ , due to the same reason, we get pairwise internally disjoint paths from  $u'_{l(r)-1}$  to  $r$  in each  $TB_{r,i}$ .

*Case 3.*  $\pi_1(p(r)) \in \{2, 3\}$ .

Similar to the proof of *Case 2*.

Q. E. D.

**Theorem 3.5.** *For a given root vertex  $r$ ,  $2 \leq l(r) \leq n$ , in  $\mathbb{P}_n$ ,  $\mathcal{BB}_r$  has 3-ISTs.*

**Proof.** Let  $\rho_i = \pi_1(a'_i)$  for  $i = 1, 2, 3$ . Let  $v \in V(\mathcal{BB}_r)$ . By Lemma 3.3, we have paths  $P_{\rho_i}$  from  $v$  to  $r$  in  $TB_{r, \rho_i}$  for each  $i$ . We want to show  $P_{\rho_1}, P_{\rho_2}$  and  $P_{\rho_3}$  are pairwise internally disjoint.

*Case 1.*  $l(v) = 0$ .

Vertex  $v$  is  $\Delta$  and adjacent to every vertex of  $h_1$ . By Lemma 3.4, we have pairwise internally disjoint paths  $X_{1, \rho_i}$  from  $v$  to  $u'_2$ , where  $1 \leq i \leq 3$  and  $h_{1, \rho_i}, h_{2, \rho_i} \in V(X_{1, \rho_i})$ . Furthermore, by Lemma 3.8, we

have pairwise internally disjoint paths  $X_{2,\rho_i}$  from  $u'_2$  to  $r$ , where  $1 \leq i \leq 3$  and  $h_{2,\rho_i}, a'_i \in V(X_{2,\rho_i})$ . Let  $Y_{1,\rho_i} = X_{1,\rho_i} - u'_2$  and  $Y_{2,\rho_i} = X_{2,\rho_i} - u'_2$  for each  $i$ . Clearly, paths  $Y_{1,\rho_i}Y_{2,\rho_i}$  from  $v$  to  $r$  are pairwise internally disjoint for  $i = 1, 2, 3$ .

*Case 2.*  $v$  is an  $a$ -vertex and  $1 \leq l(v) \leq l(r) - 2$ .

By Lemma 3.6, we have pairwise internally disjoint paths  $X_{1,\rho_i}$  from  $v$  to  $u'_{l(v)+1}$  in each  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $h_{l(v)+1,\rho_i} \in V(X_{1,\rho_i})$ . Furthermore, by Lemma 3.8, we have pairwise internally disjoint paths  $X_{2,\rho_i}$  from  $u'_{l(v)+1}$  to  $r$ , where  $1 \leq i \leq 3$  and  $h_{l(v)+1,\rho_i} \in V(X_{2,\rho_i})$ . Let  $Y_{1,\rho_i} = X_{1,\rho_i} - u'_{l(v)+1}$  and  $Y_{2,\rho_i} = X_{2,\rho_i} - u'_{l(v)+1}$  for each  $i$ . Clearly, paths  $Y_{1,\rho_i}Y_{2,\rho_i}$  from  $v$  to  $r$  are pairwise internally disjoint for  $i = 1, 2, 3$ .

*Case 3.*  $v$  is an  $a$ -vertex and  $l(v) = l(r) - 1$ .

If  $v = p(r)$ , then, by the definition of each  $R_{\rho_i}$ , we have three pairwise internally disjoint paths  $\langle v, r^-, r \rangle$ ,  $\langle v, r^+, r \rangle$  and  $\langle v, r \rangle$ . Otherwise,  $v \neq p(r)$ . Since  $c_{\rho_i}(v) \in V(P_{k+1,\rho_i})$ ,  $P_{\rho_i}$  is a subpath of  $P_{k+1,\rho_i}$ . So,  $P_{\rho_3} \parallel P_{\rho_1}$  and  $P_{\rho_3} \parallel P_{\rho_2}$  because  $P_{k+1,\rho_3} \parallel P_{k+1,\rho_1}$  and  $P_{k+1,\rho_3} \parallel P_{k+1,\rho_2}$ . Moreover, since  $c_{\rho_1}(v) \in U_{k+1,1}$  and  $c_{\rho_2}(v) \in U_{k+1,2}$  belong to the same block, it follows that  $P_{\rho_1} \parallel P_{\rho_2}$ .

*Case 4.*  $v$  is a  $b$ -vertex and  $2 \leq l(v) \leq l(r) - 1$ .

By Lemma 3.5, we have pairwise internally disjoint paths  $X_{1,\rho_i}$  from  $v$  to  $u'_{l(v)}$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $h_{l(v),\rho_i} \in V(X_{1,\rho_i})$ . Furthermore, by Lemma 3.8, we have pairwise internally disjoint paths  $X_{2,\rho_i}$  from  $u'_{l(v)}$  to  $r$ , where  $1 \leq i \leq 3$  and  $h_{l(v),\rho_i}, a'_i \in V(X_{2,\rho_i})$ . Let  $Y_{1,\rho_i} = X_{1,\rho_i} - u'_{l(v)}$  and  $Y_{2,\rho_i} = X_{2,\rho_i} - u'_{l(v)}$  for each  $i$ . Clearly, paths  $Y_{1,\rho_i}Y_{2,\rho_i}$  from  $v$  to  $r$  are pairwise internally disjoint for  $i = 1, 2, 3$ .

*Case 5.*  $v \in V(A_{l(r)})$ .

The case  $v = r$  is trivial. For cases  $v \in \{r^+, r^-, r^{++}\}$ , we shall list three pairwise internally disjoint paths for each case, where each path is a subpath of  $R_{\rho_i}$ . If  $v = r^+$ , we have paths  $\langle r^+, r^{++}, r^-, r \rangle$ ,  $\langle r^+, r \rangle$  and  $\langle r^+, p(r^+), r \rangle$ . For the case  $v = r^-$ , we have  $\langle r^-, r \rangle$ ,  $\langle r^-, r^{++}, r^+, r \rangle$  and  $\langle r^-, p(r^-), r \rangle$ . And we have  $\langle r^{++}, r^-, r \rangle$ ,  $\langle r^{++}, r^+, r \rangle$  and  $\langle r^{++}, p(r^{++}), r \rangle$  for the case  $v = r^{++}$ .

*Case 6.*  $v \in V(S_{l(r)}) \setminus V(A_{l(r)})$ .

By Corollary 3.1, we get pairwise internally disjoint paths  $X_{1,\rho_i}$  from  $v$  to  $u'_{l(r)}$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $a'_i \in V(X_{1,\rho_i})$ . Furthermore, by Lemma 3.7, we have pairwise internally disjoint

paths  $X_{2,\rho_i}$  from  $u'_{l(r)}$  to  $r$ , where  $1 \leq i \leq 3$  and  $a'_i \in V(X_{2,\rho_i})$ . Let  $Y_{1,\rho_i} = X_{1,\rho_i} - u'_{l(r)}$  and  $Y_{2,\rho_i} = X_{2,\rho_i} - u'_{l(r)}$  for each  $i$ . Clearly, paths  $Y_{1,\rho_i}Y_{2,\rho_i}$  from  $v$  to  $r$  are pairwise internally disjoint for  $i = 1, 2, 3$ .

Q. E. D.

### 3.3.2 Branching Stage

A *subpyramid*  $\mathbb{P}_n(u)$  is a subgraph of  $\mathbb{P}_n$  by letting a vertex  $u \in V(\mathbb{P}_n)$  be the apex of  $\mathbb{P}_n(u)$ . A *flat subpyramid*  $\overline{\mathbb{P}}_n(u)$  is a subpyramid without an apex  $u$  and a layer 1. Let  $V_b$  denote the set of  $b$ -vertices of  $\mathcal{BB}_r$ . The subgraph  $\mathbb{P}_n - \mathcal{BB}_r$  is regarded as an union of flat subpyramids, and every flat subpyramid is defined as  $\overline{\mathbb{P}}_n(p(v))$ , where  $v \in V(A_{l(r)}) \cup V_b$ . In particular,  $\overline{\mathbb{P}}_n(p(v))$  is defined to be empty if  $v \in S_{l(r)}$  and  $l(r) = n$ . For easily describing notations in  $\overline{\mathbb{P}}_n(p(v))$ , we let  $\overline{S}_k$  denote the cluster on layer  $k$  of  $\overline{\mathbb{P}}_n(p(v))$ , where  $2 \leq k \leq n - l(v) + 1$ . That is,  $\overline{S}_k$  is actually a cluster on layer  $l(v) + k - 1$  of  $\mathbb{P}_n$ . For a  $\overline{\mathbb{P}}_n(p(v))$ ,  $\mathcal{P}_B(\overline{S}_2)$  is either  $A_{l(r)}$  or a  $b$ -block of  $\mathcal{BB}_r$ , and  $\mathcal{P}_B(\overline{S}_2)$  is also denoted by  $\overline{S}_1$ . Since  $\overline{S}_1$  is in fact a block of  $\mathcal{BB}_r$ ,  $\overline{S}_1$  has a  $\pi_1$  coloring in Backbone Establishment Stage. Henceforth, every cluster of  $\overline{\mathbb{P}}_n(p(v))$  can have a  $\pi_0$  coloring by applying **HFinding Algorithm** on its parent block. Notations  $\hat{h}_{k,i}$  and  $\hat{H}_{k,i}$ ,  $1 \leq i \leq 3$ , refer to Section 3.1. Let  $T_i(v)$ ,  $1 \leq i \leq 3$ , denote the spanning subgraph of  $\overline{\mathbb{P}}_n(p(v))$  and have the edge set  $\bigcup_{k=2}^{n-l(v)+1} \bigcup_{a,b=0}^{2^{k-2}-1} E(\hat{H}_i(k; a, b))$ . Let  $T_{r,i}^2$ ,  $1 \leq i \leq 3$ , denote the spanning subgraph of  $\mathbb{P}_n$  with the edge set  $E(TB_{r,i}) \cup_{v \in V(A_{l(r)}) \cup V_b} E(T_i(v))$ .

**Lemma 3.9.** *Let  $v \in V(A_{l(r)}) \cup V_b$ . If  $u_k$  is a vertex of  $\overline{S}_k$  in  $\overline{\mathbb{P}}_n(p(v))$ ,  $2 \leq k \leq n - l(v) + 1$ , then the paths from  $u_k$  to  $r$  in each  $T_{r,i}^2$ , for  $i = 1, 2, 3$ , are pairwise internally disjoint.*

**Proof.**  $\overline{S}_1$  is either  $A_{l(r)}$  or a  $b$ -block of  $\mathcal{BB}_r$ . If  $\overline{S}_1 = A_{l(r)}$ , then, by Theorem 3.3, we have pairwise internally disjoint paths from  $u_k$  to  $r$  in each  $T_{r,i}^2$ . Otherwise,  $\overline{S}_1$  is a  $b$ -block of  $\mathcal{BB}_r$ . Let  $\overline{u'_j}$  be the dummy vertex with dummy edges incident to every vertex of  $\mathcal{P}_B(\overline{S_{j+1}})$  for  $j = 1, 2, \dots, k-1$ . Let  $u'_{l(r)}$  be the dummy vertex with dummy edges incident to every vertex of  $\{a'_1, a'_2, a'_3\}$ . Let  $\rho_i = \pi_1(a'_i)$  for each  $i$ . From Lemma 3.1, we get pairwise internally disjoint paths  $X_{k,\rho_i}$  from  $u_k$  to  $\overline{u'_{k-1}}$  in  $T_{\rho_i}(v)$  for each  $i = 1, 2, 3$ . By applying Lemma 3.2, we get pairwise internally disjoint paths  $X_{j,\rho_i}$  from  $\overline{u'_j}$  to  $\overline{u'_{j-1}}$  in  $T_{\rho_i}(v)$  for each  $i$ , where  $2 \leq j \leq k-1$ . Let  $Y_{k,\rho_i} = X_{k,\rho_i} - \overline{u'_{k-1}}$  and  $Y_{j,\rho_i} = X_{j,\rho_i} - \overline{u'_j} - \overline{u'_{j-1}}$  for each  $i, j$ . We now list the three pairwise internally disjoint paths  $Y_{1,\rho_i}$  from  $\overline{u'_1}$  to  $r$  in each  $TB_{r,\rho_i}$  to complete the proof.

Let  $V(\overline{S_1}) = \{v_0, v_{\rho_1}, v_{\rho_2}, v_{\rho_3}\}$  such that  $\pi_1(v_0) = 0$  and  $\pi_1(v_{\rho_i}) = \rho_i$  for  $i = 1, 2, 3$ . Recall that  $v_{\rho_3} = v_0^{++}$ . By the definition of each  $E(\hat{H}_{2,\rho_i})$ , either  $v_{\rho_3}$  or  $v_0$  can be on the path  $X_{2,\rho_3}$ .

*Case 1.*  $\overline{S_1}$  is a  $b$ -block of  $S_{l(r)}$ .

By Lemma 3.7, we have pairwise internally disjoint paths  $Z_{1,\rho_i}$  from  $u'_{l(r)}$  to  $r$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $a'_i \in V(Z_{1,\rho_i})$ . Furthermore, from Corollary 3.2, we get pairwise internally disjoint paths  $P_{\rho_i}$  from  $p(v)$  to  $u'_{l(r)}$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $a'_i \in V(P_{\rho_i})$ . Since  $(v_{\rho_i}, p(v)) \in E(TB_{r,\rho_i})$ ,  $P_{\rho_i} - p(r)$  is the path from  $v_{\rho_i}$  to  $u'_{l(r)}$  in  $TB_{r,\rho_i}$ . If  $v_{\rho_3} \in V(X_{2,\rho_3})$ , we get paths  $Y_{1,\rho_i} = Z_{2,\rho_i}Z_{1,\rho_i} - u'_{l(r)}$  for each  $i$ . Otherwise  $v_0 \in V(X_{2,\rho_3})$ . Since  $\langle v_0, p(v_0), v_{\rho_3} \rangle$  is a subgraph of  $E(H_{l(r),\rho_3})$ , we get the path  $Y_{1,\rho_3} = \langle v_0, p(v_0), v_{\rho_3} \rangle Z_{2,\rho_3}Z_{1,\rho_3} - u'_{l(r)}$ . In addition,  $Y_{1,\rho_1} = Z_{2,\rho_1}Z_{1,\rho_1} - u'_{l(r)}$  and  $Y_{1,\rho_2} = Z_{2,\rho_2}Z_{1,\rho_2} - u'_{l(r)}$ .

*Case 2.*  $\overline{S_1}$  is not a block of  $S_{l(r)}$ .

Notice that  $\overline{S_1}$  is a  $b$ -block and  $l(v) < l(r)$ . By Lemma 3.8, we have pairwise internally disjoint paths  $Z_{1,\rho_i}$  from  $u'_{l(v)}$  to  $r$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $h_{l(v),\rho_i} \in V(Z_{1,\rho_i})$ . Furthermore, from Lemma 3.6, we get pairwise internally disjoint paths  $P_{\rho_i}$  from  $p(v)$  to  $u'_{l(v)}$  in  $TB_{r,\rho_i}$ , where  $1 \leq i \leq 3$  and  $h_{l(v),\rho_i} \in V(P_{\rho_i})$ . Since  $(v_{\rho_i}, p(v)) \in E(TB_{r,\rho_i})$ ,  $P_{\rho_i} - p(r)$  is the path from  $v_{\rho_i}$  to  $u'_{l(v)}$  in  $TB_{r,\rho_i}$ . If  $v_{\rho_3} \in V(X_{2,\rho_3})$ , we get paths  $Y_{1,\rho_i} = Z_{2,\rho_i}Z_{1,\rho_i} - u'_{l(v)}$  for each  $i$ . Otherwise  $v_0 \in V(X_{2,\rho_3})$ . Since  $\langle v_0, p(v_0), v_{\rho_3} \rangle$  is a subgraph of  $E(H_{l(v),\rho_3})$ , we get the path  $Y_{1,\rho_3} = \langle v_0, p(v_0), v_{\rho_3} \rangle Z_{2,\rho_3}Z_{1,\rho_3} - u'_{l(v)}$ . In addition,  $Y_{1,\rho_1} = Z_{2,\rho_1}Z_{1,\rho_1} - u'_{l(v)}$  and  $Y_{1,\rho_2} = Z_{2,\rho_2}Z_{1,\rho_2} - u'_{l(v)}$ .

It concludes that paths  $Y_{k,\rho_i}Y_{k-1,\rho_i} \dots Y_{1,\rho_i}$  from  $u_k$  to  $r$  in  $T_{r,\rho_i}^2$ , for  $i = 1, 2, 3$ , are pairwise internally disjoint.

Q. E. D.

**Theorem 3.6.** *The construction of 3-ISTs of  $\mathbb{P}_n$  rooted at  $r$ ,  $2 \leq l(r) \leq n$ , can be solved in  $O(|V|)$  time, where  $|V|$  is the number of vertices of  $\mathbb{P}_n$ .*

**Proof.** The construction of 3-ISTs rooted at  $r$ ,  $2 \leq l(r) \leq n$ , contains the Backbone Establishment Stage and the Branching Stage. Let  $r = (k; x, y)$ . In the Backbone Establishment Stage, every vertex  $(k - j; \lfloor \frac{x}{2^j} \rfloor, \lfloor \frac{y}{2^j} \rfloor)$ , for  $j = 0, 1, \dots, k - 1$ , is an ancestor of  $r$  and can be decided in  $O(1)$  time, implying every  $a$ -block can be found in  $O(1)$  time. Since every cluster of  $\mathcal{BB}_r$  can be colored by constant time,  $\pi_1$  coloring costs  $O(|V(\mathcal{BB}_r)|)$  time. We then need additional constant

time to have an `RootColoring Algorithm`. Vertices of  $h_k$  of a cluster are decided in constant time and hence every  $H_{k,i}$  is established in constant time. Since each edge set  $E(H_{x,i}), i = 1, 2, 3$  has  $|V(\mathcal{BB}_r)| - 1$  edges and each edge of  $E(H_{x,i})$  can be checked in constant time, the finding of  $E(H_{x,i})$  costs  $O(|V(\mathcal{BB}_r)|)$  time. Therefore, Backbone Establishment Stage costs  $O(|V(\mathcal{BB}_r)|)$  time.

In Branching Stage, every flat subpyramid  $\overline{\mathbb{P}}_n(p(v))$  needs  $O(|V(\overline{\mathbb{P}}_n(p(v)))|)$  time  $\pi_0$  coloring. By Theorem 3.2, the construction of 3-ISTs of every  $\overline{\mathbb{P}}_n(p(v))$  rooted at  $p(v)$  can be solved in  $O(|V(\overline{\mathbb{P}}_n(p(v)))|)$  time, where  $v \in V(A_{l(r)}) \cup V_b$ . The overall time complexity of Branching Stage is  $O(|V(\mathbb{P}_n - \mathcal{BB}_r)|)$ . That is, the construction of 3-ISTs of  $\mathbb{P}_n$  rooted at  $r$ , can be solved in  $O(|V(\mathbb{P}_n)|)$  time.

Q. E. D.

## 4 Conclusion Remarks

In this paper, we present linear time algorithms to construct 3-ISTs in an  $n$ -dimensional pyramid  $\mathbb{P}_n$ . To the best of our knowledge, this is the first result for constructing ISTs in  $\mathbb{P}_n$ . The tree height and the average distance of an IST are considered as two performance metrics in our algorithms. The *height* of a tree  $T$  is the the number of vertices in a longest path from the root to a leaf. The *average distance* of  $T$  is the average of all distances from any vertex to the root. The resulting values are shown in Table 1. In Table 1, the column “height” provides an upper bound to the values of the average distance of a tree, where  $n$  is the dimension of the given network. Columns 4-10 demonstrate average distances for different network scales. The data are calculated on different scenarios to see the tree heights and the average distances for different scales of pyramid networks. Although the values of average distances are associated with a proportional increase in the dimension of the networks, the resulting values of  $T_{\rho_i}^0, T_{\rho_i}^1$  and  $T_{r,\rho_i}^2$  are still bounded by  $2.81n, 2.81n$  and  $6.54n$ , respectively, even when the given network is of order over 1 million vertices.

## References

- [1] F. Bao, Y. Funyu, Y. Hamada, Y. Igarashi, Reliable broadcasting and secure distributing in channel networks, in: ISPAN, 1997, p. 472.
- [2] J. Cheriyan, S. N. Maheshwari, Finding nonseparating induced cycles and independent spanning trees in 3-connected graphs, *Journal of Algorithms* 9 (1988) 507–537.

- [3] D. E. Culler, J. P. Singh, A. Gupta, *Parallel Computer Architecture: A Hardware/Software Approach*, Morgan Kaufmann, San Francisco, CA, 1999.
- [4] Y. Iwasaki, Y. Kajiwara, K. Obokata, and Y. Igarashi, Independent spanning trees of chordal rings, *Information Processing Letters* 69 (1999) 155–160.
- [5] J. S. Kim, H. O. Lee, E. Cheng, L. Liptak, Optimal independent spanning trees on odd graphs, *The Journal of Supercomputing* 56 (2011) 212–225.
- [6] K. Obokata, Y. Iwasaki, F. Bao, and Y. Igarashi, Independent spanning trees of product graphs and their construction, *IEICE Trans. Fundamentals of Electronics, Communications and Computer Sciences*, E79–A (1996) 1894–1903.
- [7] H. S. Shahhoseini, E. S. Kandzi, M. Mollajafari, Nonflat surface level pyramid: a high connectivity multidimensional interconnection network, *The Journal of Supercomputing* 67 (1) (2014) 31–46.
- [8] S. M. Tang, Y. L. Wang, Y. H. Leu, Optimal independent spanning trees on hypercubes, *Journal of Information Science and Engineering* 20 (1) (2004) 143–155.
- [9] S. M. Tang, J. S. Yang, Y. L. Wang, J. M. Chang, Independent spanning trees on multidimensional torus networks, *IEEE Transactions on Computers* 59 (1) (2010) 93–102.
- [10] Y. Wang, J. Fan, X. Jia, H. Huang, An algorithm to construct independent spanning trees on parity cubes, *Theoretical Computer Science* 465 (2012) 61–72.
- [11] J. S. Yang, J. M. Chang, S. M. Tang, Y. L. Wang, On the independent spanning trees of recursive circulant graphs  $G(cd^m; d)$  with  $d > 2$ , *Theoretical Computer Science* 410 (2009) 2001–2010.






Figure 1: A top view of the pyramid  $\mathbb{P}_2$ .




Figure 2: The bold cluster edges establish three subgraphs (a)  $\hat{H}_{k, \rho_1}$ , (b)  $\hat{H}_{k, \rho_2}$ , (c)  $\hat{H}_{k, \rho_3}$  of a cluster  $S_k$ , where the parent vertex of the top-left block is of color 0. The dash lines indicate layer edges.




Figure 3: The four possible  $\pi_1$  colorings  $\Pi_0$ ,  $\Pi_1$ ,  $\Pi_2$  and  $\Pi_3$  of a cluster  $S_k$  when  $A_k$  is (a) the top-left block; (b) the top-right block; (c) the bottom-left block; (d) the bottom-right block of  $S_k$ .




Figure 4: The three subgraphs (a)  $H_{k,\rho_1}$ , (b)  $H_{k,\rho_2}$ , (c)  $H_{k,\rho_3}$  of a cluster  $S_k$ , where the top-left block of  $S_k$  is an  $a$ -block.

Table 1: The table of tree heights and average distances

	$l(r)$	height	$n$ (dimension)						
			2	3	4	5	6	7	10
$ V(RP_n) $			21	85	341	1,365	5,461	21,845	> 1 million
$T_{\rho_1}^0, T_{\rho_2}^0$	0	$5n - 2$	$2.14n$	$2.39n$	$2.53n$	$2.62n$	$2.69n$	$2.73n$	$2.81n$
$T_{\rho_3}^0$	0	$5n - 3$	$1.71n$	$2.06n$	$2.28n$	$2.42n$	$2.52n$	$2.59n$	$2.71n$
$T_{\rho_1}^1, T_{\rho_2}^1$	1	$5n - 2$	$2.14n$	$2.39n$	$2.53n$	$2.62n$	$2.69n$	$2.73n$	$2.81n$
$T_{\rho_3}^1$	1	$5n - 2$	$2.10n$	$2.37n$	$2.53n$	$2.62n$	$2.68n$	$2.73n$	$2.81n$
$T_{r,\rho_1}^2, T_{r,\rho_2}^2$	$\geq 2$	$\leq 5n - 11$ $+5l(r) + \varepsilon$	$\leq 2.14n$	$\leq 3.45n$	$\leq 4.43n$	$\leq 5.11n$	$\leq 5.58n$	$\leq 5.92n$	$\leq 6.54n$
$T_{r,\rho_3}^2$	$\geq 2$	$\leq 9n - 19$ $+4l(r) + \varepsilon$	$\leq 2.50n$	$\leq 2.60n$	$\leq 3.35n$	$\leq 3.99n$	$\leq 4.47n$	$\leq 4.83n$	$\leq 5.48n$
$\varepsilon$ : the distance from an $a$ -vertex of $S_{l(r)}$ to $r$ .									