



HAL
open science

Coxeter groups, symmetries, and rooted representations

Olivier Geneste, Luis Paris

► **To cite this version:**

Olivier Geneste, Luis Paris. Coxeter groups, symmetries, and rooted representations. *Communications in Algebra*, 2018, 46 (5), pp.1996-2002. 10.1080/00927872.2017.1363221 . hal-01404050

HAL Id: hal-01404050

<https://hal.science/hal-01404050>

Submitted on 28 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Coxeter groups, symmetries, and rooted representations

OLIVIER GENESTE
LUIS PARIS

20F55

1 Introduction

Let Γ be a Coxeter graph, and let (W, S) be the Coxeter system of Γ . A *symmetry* of Γ is a permutation g of S such that $m_{g(s),g(t)} = m_{s,t}$ for all $s, t \in S$, where $(m_{s,t})_{s,t \in S}$ is the Coxeter matrix of Γ . Let G be a group of symmetries of Γ . Then G is necessarily finite, and it can be viewed as a group of automorphisms of W . We denote by W^G the subgroup of W fixed under the action of G . Mühlherr [7] and Hée [4], independently of one another, proved that W^G is a Coxeter group. None of them gave explicitly the Coxeter graph $\tilde{\Gamma}$ which defines W^G . However, a third proof, different from the other two, with an explicit description of $\tilde{\Gamma}$, is given in Crisp [2, 3].

Let $\Pi = \{\epsilon_s \mid s \in S\}$ be a set in one-to-one correspondence with S , let V be the real vector space having Π as a basis, and let $\langle \cdot, \cdot \rangle$ be the symmetric bilinear form on V defined by $\langle \epsilon_s, \epsilon_t \rangle = -\cos(\pi/m_{s,t})$ if $m_{s,t} \neq \infty$ and $\langle \epsilon_s, \epsilon_t \rangle = -1$ if $m_{s,t} = \infty$. For every $s \in S$ we define the linear transformation $f_s : V \rightarrow V$ by $f_s(x) = x - 2\langle x, \epsilon_s \rangle \epsilon_s$. Then the map $S \rightarrow \text{GL}(V)$, $s \mapsto f_s$, induces a celebrated faithful linear representation $f : W \rightarrow \text{GL}(V)$, called the *canonical representation* of W (see Bourbaki [1]). In our context, the triple $(V, \langle \cdot, \cdot \rangle, \Pi)$ will be called the *canonical root basis* of Γ .

Let G be a group of symmetries of Γ . Then G acts on V sending ϵ_s to $\epsilon_{g(s)}$ for all $s \in S$, and this action leaves invariant the canonical form $\langle \cdot, \cdot \rangle$. Hence, the canonical representation $f : W \rightarrow \text{GL}(V)$ is equivariant, in the sense that $f(g(w)) = g \circ f(w) \circ g^{-1}$ for all $g \in G$ and all $w \in W$, and therefore f induces a linear representation $f^G : W^G \rightarrow \text{GL}(V^G)$, where $V^G = \{x \in V \mid g(x) = x \text{ for all } g \in G\}$.

A naive question would be: Is $f^G : W^G \rightarrow \text{GL}(V^G)$ the canonical representation of W^G ? A positive answer would provide a way to (re)prove that W^G is a Coxeter group and to determine the Coxeter graph of W^G . Unfortunately, simple calculations show that f^G is not the canonical representation in general. Nevertheless, one can transpose

this question to a larger family of linear representations, the rooted representations introduced by Krammer [5, 6], and, in this context, the answer is yes. Our purpose is to show that.

A *root basis* of Γ is a triple $(V, \langle \cdot, \cdot \rangle, \Pi)$, where V is a finite dimensional real vector space, $\langle \cdot, \cdot \rangle$ is a symmetric bilinear form on V , and $\Pi = \{\epsilon_s \mid s \in S\}$ is a collection of vectors in V in one-to-one correspondence with S , that satisfies the following properties:

- (a) $\langle \epsilon_s, \epsilon_s \rangle = 1$ for all $s \in S$;
- (b) for all $s, t \in S$, $s \neq t$, we have

$$\begin{aligned} \langle \epsilon_s, \epsilon_t \rangle &= -\cos(\pi/m_{s,t}) && \text{if } m_{s,t} \neq \infty, \\ \langle \epsilon_s, \epsilon_t \rangle &\in (-\infty, -1] && \text{if } m_{s,t} = \infty; \end{aligned}$$
- (c) there exists $\chi \in V^*$ such that $\chi(\epsilon_s) > 0$ for all $s \in S$.

As mentioned above, if Π is a basis of V and $\langle \epsilon_s, \epsilon_t \rangle = -1$ whenever $m_{s,t} = \infty$, then $(V, \langle \cdot, \cdot \rangle, \Pi)$ is called the *canonical root basis* of Γ .

This definition is taken from Krammer's thesis [5, 6]. It is both, a generalization of the canonical spaces and canonical forms defined by Bourbaki [1], and a new point of view on the theory of reflection groups developed by Vinberg [8]. Note also that Condition (c) in the above definition often follows from Conditions (a) and (b), but not always (see Krammer [6, Proposition 6.1.2]).

Let $(V, \langle \cdot, \cdot \rangle, \Pi)$ be a root basis of Γ . For every $s \in S$ we define the linear transformation $f_s : V \rightarrow V$ by $f_s(x) = x - 2\langle x, \epsilon_s \rangle \epsilon_s$. The following theorem can be proved for any root basis in the same way as it is proved in Bourbaki [1] for the canonical root basis.

Theorem 1.1 (Krammer [5, 6]) *The map $S \rightarrow \text{GL}(V)$, $s \mapsto f_s$, induces a faithful linear representation $f : W \rightarrow \text{GL}(V)$.*

The representation $f : W \rightarrow \text{GL}(V)$ of Theorem 1.1 is called the *rooted representation* of W associated with $(V, \langle \cdot, \cdot \rangle, \Pi)$.

Let G be a group of symmetries of Γ , and let $(V, \langle \cdot, \cdot \rangle, \Pi)$ be a root basis. As for the canonical root basis, we assume that G embeds in $\text{GL}(V)$, satisfies $g(\epsilon_s) = \epsilon_{g(s)}$ for all $g \in G$ and all $s \in S$, and leaves invariant the form $\langle \cdot, \cdot \rangle$. Then the representation $f : W \rightarrow \text{GL}(V)$ is equivariant in the sense that $f(g(w)) = g \circ f(w) \circ g^{-1}$ for all $g \in G$ and all $w \in W$, and therefore f induces a linear representation $f^G : W^G \rightarrow \text{GL}(V^G)$, where $V^G = \{x \in V \mid g(x) = x \text{ for all } g \in G\}$. The goal of this paper is to prove the following.

Theorem 1.2 (1) *The group W^G is a Coxeter group.*

(2) *Let $\tilde{\Gamma}$ denote the Coxeter graph of W^G . There exists a subset $\tilde{\Pi}$ of V^G such that $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$ is a root basis of $\tilde{\Gamma}$, and the induced representation $f^G : W^G \rightarrow \text{GL}(V^G)$ is the rooted representation associated with $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$. In particular, f^G is faithful.*

A similar approach is adopted in Hée [4, Section 3], with different definitions. One can easily show that the root system obtained from a root basis is a root system in Hée sense [4], and that part of the results of the paper, such as the fact that $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$ is a root basis, can be deduced from Hée [4]. However, to get the explicit expression of the Coxeter graph $\tilde{\Gamma}$ of W^G , one would need extra arguments that can be either a rewrite of Lemma 3.3, or some arguments similar to that given in Crisp [2, 3]. More generally, the whole theorem is more or less in the literature. In particular, as mentioned before, Part (1) is explicit in Mühlherr [7], Hée [4] and Crisp [2, 3]. But, our aim is to provide a new point of view on the question with unified, short and self-contained proofs.

A more precise statement of Theorem 1.2 is given in Section 2. In particular, the Coxeter graph $\tilde{\Gamma}$ and the set $\tilde{\Pi}$ are explicitly described. Section 3 is dedicated to the proofs.

2 Statement

The *length* of an element $w \in W$, denoted by $\text{lg}(w)$, is the shortest length of an expression of w over the elements of S . An expression $w = s_1 \cdots s_\ell$ is called *reduced* if $\ell = \text{lg}(w)$. It is known that, if W is finite, then W has a unique *longest element*, that is, an element $w_0 \in W$ such that $\text{lg}(w) \leq \text{lg}(w_0)$ for all $w \in W$, and this element is an involution (see Bourbaki [1]).

For $X \subset S$, we denote by Γ_X the full subgraph of Γ spanned by X , and by W_X the subgroup of W generated by X . The subgroup W_X is called a *standard parabolic subgroup* of W . By Bourbaki [1], (W_X, X) is a Coxeter system of Γ_X . If W_X is finite, then we denote by w_X the longest element of W_X .

Let G be a group of symmetries of Γ . Now, we define a Coxeter matrix $\tilde{M} = \tilde{M}^G = (\tilde{m}_{X,Y})_{X,Y \in \mathcal{S}}$ (and its associated Coxeter graph, $\tilde{\Gamma}$). This will be the Coxeter matrix (and the Coxeter graph) of W^G (see Theorem 1.2 and Theorem 2.2).

We denote by \mathcal{O} the set of orbits of G in S , and we set $\mathcal{S} = \{X \in \mathcal{O} \mid W_X \text{ is finite}\}$. Then \mathcal{S} is the set of indices of \tilde{M} (which is the set of vertices of $\tilde{\Gamma}$). Let $X, Y \in \mathcal{S}$.

- If $m_{s,t} = 2$ for all $s \in X$ and all $t \in Y$, then we set $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = 2$.
- Let $k \in \{1, 2, 3, 4, 5\}$. If $\Gamma_{X \cup Y}$ is a disjoint union of copies of the Coxeter graph depicted in Figure 2.1 (k), where the vertices corresponding to x_1, x_2, \dots belong to X and the vertices corresponding to y_1, y_2, \dots belong to Y , then we set $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = m$ if $k = 1$, $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = 4$ if $k \in \{2, 3\}$, $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = 8$ if $k = 4$, and $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = 6$ if $k = 5$. In this case we say that (X, Y) is a *bi-orbit of type k* .
- We set $\tilde{m}_{X,Y} = \tilde{m}_{Y,X} = \infty$ in the remaining cases.




Figure 2.1: Bi-orbits.

The next lemma will be used in the definition of the set $\tilde{\Pi}$. It is well-known and can be easily proved using [1, Chapter V, Section 4, Subsection 8].

Lemma 2.1 *Let $(V, \langle \cdot, \cdot \rangle, \Pi)$ be a root basis of Γ . Suppose that W is finite and that Π spans V . Then $\langle \cdot, \cdot \rangle$ is a scalar product, and $(V, \langle \cdot, \cdot \rangle, \Pi)$ is the canonical root basis of Γ . In particular, Π is a basis of V .*

We turn back to the hypothesis of Theorem 1.2, that is, Γ is any Coxeter graph, G is a group of symmetries of Γ , and $(V, \langle \cdot, \cdot \rangle, \Pi)$ is a root basis of Γ . We assume that G embeds in $\text{GL}(V)$ so that the form $\langle \cdot, \cdot \rangle$ is invariant under the action of G , and $g(\epsilon_s) = \epsilon_{g(s)}$ for all $s \in S$ and all $g \in G$.

Let X be an element of \mathcal{S} , that is, an orbit of G in S such that W_X is finite. Set $\Pi_X = \{\epsilon_s \mid s \in X\}$, and denote by V_X the linear subspace of V spanned by Π_X , and by $\langle \cdot, \cdot \rangle_X$ the restriction of $\langle \cdot, \cdot \rangle$ to $V_X \times V_X$. By Lemma 2.1, Π_X is a basis of V_X and $\langle \cdot, \cdot \rangle_X$ is a scalar product. Let $a_X = \sum_{s \in X} \epsilon_s$. Note that $a_X \in V^G$, hence, by the above, $a_X \neq 0$ and $\|a_X\| > 0$. We set $\tilde{\epsilon}_X = \frac{a_X}{\|a_X\|}$ for all $X \in \mathcal{S}$, and $\tilde{\Pi} = \tilde{\Pi}^G = \{\tilde{\epsilon}_X \mid X \in \mathcal{S}\}$. The main result of the paper, with a precise statement, is the following.

Theorem 2.2 (1) *The set $\mathcal{S}_W = \{w_X \mid X \in \mathcal{S}\}$ generates W^G , and (W^G, \mathcal{S}_W) is a Coxeter system of $\tilde{\Gamma}$.*

- (2) The triple $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$ is a root basis of $\tilde{\Gamma}$, and the induced representation $f^G : W^G \rightarrow \text{GL}(V^G)$ is the rooted representation associated with $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$. In particular, f^G is faithful.

Remark The proof of Part (1) of Theorem 2.2 uses the induced representation $f^G : W^G \rightarrow \text{GL}(V^G)$. Nevertheless, the conclusion of Part (1) is always true because there is always a root basis which satisfies the hypothesis of the theorem: the canonical root basis.

3 Proof

We assume given a Coxeter graph Γ , a root basis $(V, \langle \cdot, \cdot \rangle, \Pi)$ of Γ , and a group G of symmetries of Γ . We assume that G embeds in $\text{GL}(V)$, satisfies $g(\epsilon_s) = \epsilon_{g(s)}$ for all $g \in G$ and all $s \in S$, and leaves invariant the form $\langle \cdot, \cdot \rangle$.

Let $f : W \rightarrow \text{GL}(V)$ be the rooted representation of W associated with $(V, \langle \cdot, \cdot \rangle, \Pi)$. From now on, in order to simplify the notations, we will assume that W acts on V via f , and we will write $w(x)$ in place of $f(w)(x)$ for $w \in W$ and $x \in V$. Lemmas 3.1 to 3.4 are preliminaries to the proof of Theorem 2.2. Lemma 3.1 (1) is well-known. It is a direct consequence of Mühlherr [7, Lemma 2.8], and its proof can be found in the beginning of the proof of Mühlherr [7, Theorem 1.3]. Lemma 3.1 (2) is also known. Its proof is implicit in Crisp [2], but, as far as we know, it is not explicitly given anywhere else.

Lemma 3.1 (1) The group W^G is generated by S_W .

- (2) We have $(w_X w_Y)^{\tilde{m}_{X,Y}} = 1$ for all $X, Y \in \mathcal{S}$ such that $\tilde{m}_{X,Y} \neq \infty$.

Proof As mentioned above, the proof of Part (1) can be found in Mühlherr [7]. So, we only need to prove Part (2). Let $X \subset S$ be such that Γ_X is a disjoint union of vertices (i.e. Γ_X has no edge). Then W_X is finite and $w_X = \prod_{s \in X} s$. Let $X = \{s, t\}$ be a pair included in S such that $m_{s,t} = m < \infty$. Then W_X is finite, $w_X = (st)^{\frac{m}{2}}$ if m is even, and $w_X = (st)^{\frac{m-1}{2}} s$ if m is odd. Now, let $X, Y \in \mathcal{S}$. If $m_{s,t} = 2$ for all $s \in X$ and $t \in Y$, then w_X and w_Y commute, hence $(w_X w_Y)^2 = 1$, as w_X and w_Y are both involutions. Suppose that (X, Y) is a bi-orbit of type j , where $j \in \{1, 2, 3, 4, 5\}$. Let $\Gamma_1, \dots, \Gamma_\ell$ be the connected components of $\Gamma_{X \cup Y}$. For $i \in \{1, \dots, \ell\}$, we denote by Z_i the set of vertices of Γ_i , and we set $X_i = X \cap Z_i$ and $Y_i = Y \cap Z_i$. We have $w_X = \prod_{i=1}^{\ell} w_{X_i}$ and $w_Y = \prod_{i=1}^{\ell} w_{Y_i}$. Moreover, using the above observation together

with Theorem 1.1, it is easily checked that $(w_{X_i} w_{Y_i})^{\tilde{m}_{X,Y}} = 1$ for all i . It follows that $(w_X w_Y)^{\tilde{m}_{X,Y}} = \prod_{i=1}^{\ell} (w_{X_i} w_{Y_i})^{\tilde{m}_{X,Y}} = 1$. \square

Lemma 3.2 *Let $X \in \mathcal{S}$. Then one of the following two alternatives holds.*

- (I) Γ_X is a disjoint union of vertices (i.e. Γ_X has no edge).
- (II) There exists $m \in \mathbb{N}$, $m \geq 3$, such that Γ_X is a disjoint union of copies of the Coxeter graph depicted in Figure 2.1 (1).

Proof For $s \in X$ we set $v_s(X) = |\{t \in X \mid m_{s,t} \geq 3\}|$. Since W_X is finite, the connected components of Γ_X are trees (see Bourbaki [1]), hence there exists $s \in X$ such that $v_s(X) \leq 1$. On the other hand, since G acts transitively on X , we have $v_s(X) = v_t(X)$ for all $s, t \in X$. So, either $v_s(X) = 0$ for all $s \in X$, or $v_s(X) = 1$ for all $s \in X$. If $v_s(X) = 0$ for all $s \in X$, then we are in Alternative (I). If $v_s(X) = 1$ for all $s \in X$, then we are in Alternative (II). \square

Let $X \in \mathcal{S}$. We say that X is of *type I* if Γ_X satisfies Condition (I) of Lemma 3.2, and that X is of *type II_m* if Γ_X satisfies Condition (II).

Lemma 3.3 *Let $X, Y \in \mathcal{S}$, $X \neq Y$. Then*

$$\begin{aligned} \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle &= -\cos(\pi/\tilde{m}_{X,Y}) && \text{if } \tilde{m}_{X,Y} \neq \infty, \\ \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle &\in (-\infty, -1] && \text{if } \tilde{m}_{X,Y} = \infty. \end{aligned}$$

Proof Observe that, if $m_{s,t} = 2$ for all $s \in X$ and all $t \in Y$, then $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = 0$ and $\tilde{m}_{X,Y} = 2$. Hence, we can assume that there exist $s \in X$ and $t \in Y$ such that $m_{s,t} \geq 3$. Since G acts transitively on X and leaves invariant Y , it follows that, for all $s \in X$, there exists $t \in Y$ such that $m_{s,t} \geq 3$. Similarly, for all $t \in Y$, there exists $s \in X$ such that $m_{s,t} \geq 3$.

Recall that $a_X = \sum_{s \in X} \epsilon_s$, $a_Y = \sum_{t \in Y} \epsilon_t$, $\tilde{\epsilon}_X = \frac{a_X}{\|a_X\|}$, $\tilde{\epsilon}_Y = \frac{a_Y}{\|a_Y\|}$. Choose $s \in X$ and set $v_X = |\{t \in Y \mid m_{s,t} \geq 3\}|$ and $p_X = \sum_{t \in Y} \langle \epsilon_s, \epsilon_t \rangle = \langle \epsilon_s, a_Y \rangle$. Since G acts transitively on X and leaves invariant Y , these definitions do not depend on the choice of s . Similarly, choose $t \in Y$ and set $v_Y = |\{s \in X \mid m_{s,t} \geq 3\}|$ and $p_Y = \sum_{s \in X} \langle \epsilon_t, \epsilon_s \rangle = \langle \epsilon_t, a_X \rangle$. The hypothesis that there exist $s \in X$ and $t \in Y$ such that $m_{s,t} \geq 3$ implies that $v_X \geq 1$ and $v_Y \geq 1$.

Let $s \in X$ and $t \in Y$. If $m_{s,t} \geq 3$, then $\langle \epsilon_s, \epsilon_t \rangle \leq -\frac{1}{2}$, and if $m_{s,t} = 2$, then $\langle \epsilon_s, \epsilon_t \rangle = 0$. It follows that

$$(3-1) \quad p_X \leq -\frac{v_X}{2}.$$

On the other hand, we have

$$(3-2) \quad |X| v_X = |Y| v_Y.$$

This is the number of edges in Γ connecting an element of X with an element of Y . A direct calculation shows that

$$(3-3) \quad \|a_X\| = \begin{cases} \sqrt{|X|} & \text{if } X \text{ is of type } I, \\ \sqrt{|X|(1 - \cos(\pi/m))} & \text{if } X \text{ is of type } II_m. \end{cases}$$

Finally, by definition of p_X ,

$$(3-4) \quad \langle a_X, a_Y \rangle = |X| p_X.$$

Case 1: X and Y are of type I . Applying Equations (3-2), (3-3), and (3-4) we get

$$(3-5) \quad \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = \frac{p_X \sqrt{v_Y}}{\sqrt{v_X}}.$$

Applying Equation (3-1) to this equality we get $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\frac{\sqrt{v_X v_Y}}{2}$. It follows that, if either $v_X \geq 4$, or $v_Y \geq 4$, or $v_X, v_Y \geq 2$, then $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -1$. If $v_X = 1$, $v_Y \geq 2$ and $p_X \leq -\cos(\pi/4) = -\frac{1}{\sqrt{2}}$, then, by Equation (3-5), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -1$. If $v_X = 1$, $v_Y = 3$ and $p_X = -\cos(\pi/3) = -\frac{1}{2}$, then, by Equation (3-5), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = -\frac{\sqrt{3}}{2} = -\cos(\pi/6)$. In this case (Y, X) is a bi-orbit of type 5 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = 6$. If $v_X = 1$, $v_Y = 2$ and $p_X = -\cos(\pi/3) = -\frac{1}{2}$, then, by Equation (3-5), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = -\frac{\sqrt{2}}{2} = -\cos(\pi/4)$. In this case (Y, X) is a bi-orbit of type 2 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = 4$. If $v_X = 1$, $v_Y = 1$ and $p_X = -\cos(\pi/m)$ with $m \neq \infty$, then, by Equation (3-5), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = -\cos(\pi/m)$. In this case (Y, X) is a bi-orbit of type 1 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = m$. Finally, if $v_X = v_Y = 1$ and $p_X \leq -1$, then, by Equation (3-5), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = p_X \leq -1$. In this case (Y, X) is a bi-orbit of type 1 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = \infty$.

Case 2: X is of type II_m and Y is of type I . Applying Equations (3-2), (3-3), and (3-4) we get

$$(3-6) \quad \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = \frac{p_X \sqrt{v_Y}}{\sqrt{v_X(1 - \cos(\pi/m))}}.$$

Applying Equation (3-1) to this equality, we get

$$(3-7) \quad \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\frac{\sqrt{v_X v_Y}}{2\sqrt{(1 - \cos(\pi/m))}}.$$

If $m \geq 5$, then $\sqrt{1 - \cos(\pi/m)} < \frac{1}{2}$, hence, by Equation (3-7), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\sqrt{v_X v_Y} \leq -1$. So, we can assume that $m \in \{3, 4\}$. Then we have $\sqrt{1 - \cos(\pi/m)} \leq \frac{1}{\sqrt{2}}$ and, by Equation (3-7), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\frac{\sqrt{v_X v_Y}}{\sqrt{2}}$. It follows that, if either $v_X \geq 2$, or $v_Y \geq 2$, then

$\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -1$. If $v_X = 1, v_Y = 1$ and $p_X \leq -\cos(\pi/4) = -\frac{1}{\sqrt{2}}$, then, by Equation (3-6), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -1$. If $v_X = 1, v_Y = 1, p_X = -\cos(\pi/3) = -\frac{1}{2}$ and $m = 4$, then, by Equation (3-6), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = -\frac{\sqrt{2+\sqrt{2}}}{2} = -\cos(\pi/8)$. In this case (Y, X) is a bi-orbit of type 4 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = 8$. If $v_X = 1, v_Y = 1, p_X = -\cos(\pi/3) = -\frac{1}{2}$ and $m = 3$, then, by Equation (3-6), $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = -\frac{1}{\sqrt{2}} = -\cos(\pi/4)$. In this case (Y, X) is a bi-orbit of type 3 and $\tilde{m}_{Y,X} = \tilde{m}_{X,Y} = 4$.

Case 3: X is of type II_m and Y is of type $II_{m'}$. Applying Equations (3-2), (3-3) and (3-4) we get

$$\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle = \frac{p_X \sqrt{v_Y}}{\sqrt{v_X(1 - \cos(\pi/m))(1 - \cos(\pi/m'))}}.$$

Applying Equation (3-1) to this equality we get

$$\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\frac{\sqrt{v_X v_Y}}{2\sqrt{(1 - \cos(\pi/m))(1 - \cos(\pi/m'))}}.$$

Since $\sqrt{(1 - \cos(\pi/m))} \leq \frac{1}{\sqrt{2}}$ and $\sqrt{(1 - \cos(\pi/m'))} \leq \frac{1}{\sqrt{2}}$, it follows that $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle \leq -\sqrt{v_X v_Y} \leq -1$. \square

Lemma 3.4 *Let $X \in \mathcal{S}$, and let $x \in V^G$. Then $w_X(x) = x - 2\langle x, \tilde{\epsilon}_X \rangle \tilde{\epsilon}_X$.*

Proof Let Γ' be a Coxeter graph, and let (W', S') be its associated Coxeter system, such that W' is finite. Let w'_0 be the longest element of W' , and let $(V', \langle \cdot, \cdot \rangle', \Pi')$ be the canonical root basis of Γ' . Then, by Bourbaki [1], $w'_0(\Pi') = -\Pi'$.

Let $X \in \mathcal{S}$. Recall that $\Pi_X = \{\epsilon_s \mid s \in X\}$. By Lemma 2.1 and the above, we have $w_X(\Pi_X) = -\Pi_X$, hence $w_X(a_X) = -a_X$, therefore $w_X(\tilde{\epsilon}_X) = -\tilde{\epsilon}_X$.

Recall that V_X denotes the linear subspace of V spanned by Π_X . For all $x \in V$ and all $u \in W_X$ there exists $y \in V_X$ such that $u(x) = x + y$. This is true by definition for all $s \in X$, hence it is true for all $u \in W_X$. Let $x \in V^G$. Let $y \in V_X$ be such that $w_X(x) = x + y$. Let $y = \sum_{s \in X} \lambda_s \epsilon_s$ be the expression of y in the basis Π_X . For $g \in G$ we have

$$\begin{aligned} x + \sum_{s \in X} \lambda_s \epsilon_s &= w_X(x) = g(w_X)(g(x)) = g(w_X(x)) \\ &= g(x) + \sum_{s \in X} \lambda_s g(\epsilon_s) = x + \sum_{s \in X} \lambda_s \epsilon_{g(s)}, \end{aligned}$$

hence $\lambda_s = \lambda_{g^{-1}(s)}$ for all $s \in X$. Since G acts transitively on X , it follows that $\lambda_s = \lambda_t$ for all $s, t \in X$. So, there exists $\lambda \in \mathbb{R}$ such that $w_X(x) = x + \lambda a_X = x + \lambda \|a_X\| \tilde{\epsilon}_X$.

We have

$$\langle x, \tilde{\epsilon}_X \rangle = \langle w_X(x), w_X(\tilde{\epsilon}_X) \rangle = \langle x + \lambda \|a_X\| \tilde{\epsilon}_X, -\tilde{\epsilon}_X \rangle = -\langle x, \tilde{\epsilon}_X \rangle - \lambda \|a_X\|,$$

hence $\lambda \|a_X\| = -2\langle x, \tilde{\epsilon}_X \rangle$. So, $w_X(x) = x - 2\langle x, \tilde{\epsilon}_X \rangle \tilde{\epsilon}_X$. \square

Proof of Theorem 2.2 We have $\langle \tilde{\epsilon}_X, \tilde{\epsilon}_X \rangle = 1$ for all $X \in \mathcal{S}$ by definition. We have

$$\begin{aligned} \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle &= -\cos(\pi/\tilde{m}_{X,Y}) && \text{if } \tilde{m}_{X,Y} \neq \infty, \\ \langle \tilde{\epsilon}_X, \tilde{\epsilon}_Y \rangle &\in (-\infty, -1] && \text{if } \tilde{m}_{X,Y} = \infty, \end{aligned}$$

by Lemma 3.3. Let $\chi \in V^*$ be such that $\chi(\epsilon_s) > 0$ for all $s \in \mathcal{S}$. Let $\tilde{\chi} : V^G \rightarrow \mathbb{R}$ be the restriction of χ to V^G . Then, for $X \in \mathcal{S}$, $\tilde{\chi}(\tilde{\epsilon}_X) = \frac{1}{\|a_X\|} \sum_{s \in X} \chi(\epsilon_s) > 0$. So, $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi}^G)$ is a root basis of $\tilde{\Gamma}$.

Let $(\tilde{W}, \tilde{\mathcal{S}})$ be a Coxeter system of $\tilde{\Gamma}$, where $\tilde{\mathcal{S}} = \{\tilde{s}_X \mid X \in \mathcal{S}\}$ is a set in one-to-one correspondence with \mathcal{S} . By Lemma 3.1, the map $\tilde{\mathcal{S}} \rightarrow \mathcal{S}_W$, $\tilde{s}_X \mapsto w_X$, induces a surjective homomorphism $\gamma : \tilde{W} \rightarrow W^G$. By Lemma 3.4, the composition $f^G \circ \gamma : \tilde{W} \rightarrow \text{GL}(V^G)$ is the rooted representation associated with $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$. By Theorem 1.1, it follows that $f^G \circ \gamma$ is injective, hence γ is an isomorphism. So, (W, \mathcal{S}_W) is a Coxeter system of $\tilde{\Gamma}$, and $f^G : W^G \rightarrow \text{GL}(V^G)$ is the rooted representation associated with $(V^G, \langle \cdot, \cdot \rangle, \tilde{\Pi})$. \square

References

- [1] **N Bourbaki**, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*, Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968.
- [2] **J Crisp**, *Symmetrical subgroups of Artin groups*, Adv. Math. 152 (2000), no. 1, 159–177.
- [3] **J Crisp**, *Erratum to: “Symmetrical subgroups of Artin groups”*, Adv. Math. 179 (2003), no. 2, 318–320.
- [4] **J-Y Hée**, *Système de racines sur un anneau commutatif totalement ordonné*, Geom. Dedicata 37 (1991), no. 1, 65–102.
- [5] **D Kramer**, *The conjugacy problem for Coxeter groups*, Ph. D. Thesis, Utrecht, 1994.
- [6] **D Kramer**, *The conjugacy problem for Coxeter groups*, Groups Geom. Dyn. 3 (2009), no. 1, 71–171.

- [7] **B Mühlherr**, *Coxeter groups in Coxeter groups*, Finite geometry and combinatorics (Deinze, 1992), 277–287, London Math. Soc. Lecture Note Ser., 191, Cambridge Univ. Press, Cambridge, 1993.
- [8] **È B Vinberg**, *Discrete linear groups that are generated by reflections*, Izv. Akad. Nauk SSSR Ser. Mat. 35 (1971), 1072–1112.

IMB, UMR 5584, CNRS, Univ. Bourgogne Franche-Comté, 21000 Dijon, France

IMB, UMR 5584, CNRS, Univ. Bourgogne Franche-Comté, 21000 Dijon, France

olivier.geneste@u-bourgogne.fr, lparis@u-bourgogne.fr