



HAL
open science

PATIENT-SPECIFIC REGIONALIZATION FOR FOOT PRESSURE ULCER PREVENTION

Vincent Luboz, Antoine Perrier, Marek Bucki, F Cannard, E Champion, N
Vuillerme, Yohan Payan

► **To cite this version:**

Vincent Luboz, Antoine Perrier, Marek Bucki, F Cannard, E Champion, et al.. PATIENT-SPECIFIC REGIONALIZATION FOR FOOT PRESSURE ULCER PREVENTION. 5th World Congress of the World Union of Wound Healing Societies, Sep 2016, Florence, Italy. hal-01404074

HAL Id: hal-01404074

<https://hal.science/hal-01404074>

Submitted on 28 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PATIENT-SPECIFIC REGIONALIZATION FOR FOOT PRESSURE ULCER PREVENTION

V. Luboz³, A. Perrier^{1,2,3}, M. Bucki³, F. Cannard³, E. Champion³, N. Vuillerme^{2,4}, Y. Payan¹

¹TIMC-IMAG; ²AGEIS, Grenoble; ³TexiSense, Montceau-les-Mines; ⁴IUF, France.

Background and Aims

Foot pressure ulcers (PU) mainly result from excessive pressure intensity (internal strains above 50% for about 10 minutes) or prolonged compression (internal strains above 20% for about two hours) [1]. It is therefore crucial to monitor the internal strains using biomechanical models combined with skin surface pressure measurements to improve daily examination. This paper focuses on defining the regions with highest strains and analyzes the variations between patient morphologies.

Methods

Our biomechanical Finite Element (FE) foot atlas [2] is divided into eight anatomical regions: Achilles tendon, ankle, heel, medial foot, first metatarsus, four other metatarsi, hallux, and four other toes. This model is deformed using rigid and elastic transformations [3] to fit three patients' morphologies obtained from Computed Tomography scanners. These four models are used to simulate bipedal standing.

Results

The simulations report high Von Mises (VM) strains variations from one patient to another between regions, cf. figure. The maximum "strain cluster volume" (defined as the volume of the largest group of adjacent FE nodes exhibiting VM strains over the thresholds mentioned above) is always located in the heel region, and ranges between 51 and 74cm³ for VM strains above 20% and between 0 and 21cm³ for VM strains above 50%.

Conclusions

Given the morphological differences and the large variations observed in cluster volumes, an accurate representation of the external and internal structures is necessary to precisely estimate the regions with highest internal strains. This confirms that patient-specific modelling is required in PU prevention.

References

- [1] Loerakker S. et al. *J. Appl. Phys.* 111(4):1168-77, 2011.
- [2] Perrier A. et al. *CMBBE*. 18(1):2024-2025, 2015.
- [3] Bucki M. et al. *MEDIA*, 14:303-317, 2010.

