



**HAL**  
open science

# Iterative inversion of weighted Radon transforms in 3D

F O Goncharov

► **To cite this version:**

| F O Goncharov. Iterative inversion of weighted Radon transforms in 3D. 2016. hal-01405387v1

**HAL Id: hal-01405387**

**<https://hal.science/hal-01405387v1>**

Preprint submitted on 29 Nov 2016 (v1), last revised 14 Sep 2017 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Iterative inversion of weighted Radon transforms in 3D

F.O. Goncharov\*

November 29, 2016

## Abstract

We propose iterative inversion algorithms for weighted Radon transforms in 3D. Our results can be considered as extension to 3D of two-dimensional results of Kunyansky (1992), Novikov (2014), Guillement, Novikov (2014). In addition, we generalize our results to the case of dimension  $n > 3$ .

## 1 Introduction

Generalized Radon transforms arise in many domains of pure and applied mathematics; see e.g. [BQ87], [Dea07], [DB84], [GGV14], [GN16], [Gra91], [Kun92].

In the present article we consider the weighted Radon transforms  $R_W$  defined by the formula

$$R_W f(s, \theta) = \int_{x\theta=s} W(x, \theta) f(x) dx, \quad (s, \theta) \in \mathbb{R} \times \mathbb{S}^2, \quad x \in \mathbb{R}^3, \quad (1.1)$$

where  $W = W(x, \theta)$  is the weight,  $f = f(x)$  is a test function.

In this work we assume that

$$W \in C(\mathbb{R}^3 \times \mathbb{S}^2) \cap L^\infty(\mathbb{R}^3 \times \mathbb{S}^2), \quad (1.2)$$

$$w_{0,0}(x) \stackrel{\text{def}}{=} \frac{1}{4\pi} \int_{\mathbb{S}^2} W(x, \theta) d\theta, \quad w_{0,0}(x) \neq 0, \quad x \in \mathbb{R}^3, \quad (1.3)$$

where  $W$  is complex-valued weight,  $d\theta$  is element of standard measure on  $\mathbb{S}^2$ . If  $W \equiv 1$ , then  $R_W$  is reduced to the classical Radon transform  $\mathbb{R}^3$ , see e.g. [Rad17], [Nat86].

For the test function we assume that

$$f \in L^\infty(\mathbb{R}^3), \quad \text{supp } f \subset D, \quad (1.4)$$

where  $D$  is an open bounded domain (which is fixed a priori).

Let

$$c \stackrel{\text{def}}{=} \inf_{x \in D} |w_{0,0}(x)| > 0, \quad (1.5)$$

where  $D$  is the domain in (1.4) (inequality (1.5) holds by assumptions (1.2)-(1.4)).

We recall that inversion methods for  $R_W$  admit tomographical applications in the framework of the scheme described as follows.

It is well-known that in many tomographies measured data are modeled by weighted ray transforms  $P_w f$  defined by the formula

$$P_w f(x, \alpha) = \int_{\mathbb{R}} w(x + \alpha t, \alpha) f(x + \alpha t) dt, \quad (x, \alpha) \in TS^2, \quad (1.6)$$
$$TS^2 = \{(x, \alpha) \in \mathbb{R}^3 \times \mathbb{S}^2 : x\alpha = 0\},$$

where  $f$  is an object function defined on  $\mathbb{R}^3$ ,  $w$  is the weight function defined on  $\mathbb{R}^3 \times \mathbb{S}^2$ , and  $TS^2$  can be considered as the set of all rays (oriented straight lines) in  $\mathbb{R}^3$ , see e.g. [Cha78], [Nat86], [Kun92].

In addition, in [GN16] (Section 3) it was shown that if  $P_w f$  are given for all rays parallel to some fixed plane  $\Sigma$  in  $\mathbb{R}^3$  then  $R_W f$  with appropriate  $W$  can be obtained by the explicit formulas from  $P_w f$  and  $w$  (in a similar way with the case  $w \equiv 1$ , see [Nat86], Chapter 2, Formula (1.1); see also [Gra91]). Therefore, reconstruction of  $f$  from data modeled by  $P_w f$ , defined by (1.6) and restricted to all rays parallel to  $\Sigma$ , can be reduced to

---

\*CMAP, Ecole Polytechnique, CNRS, Université Paris-Saclay, 91128, Palaiseau, France; email: fedor.goncharov.ol@gmail.com

reconstruction of  $f$  from  $R_W f$ , defined by (1.1). In [GN16] it was also indicated that the reduction from  $P_w f$  to  $R_W f$  with subsequent reconstruction of  $f$  from  $R_W f$  can drastically reduce the impact of the noise in the initial data.

In the present article we use the expansion for weight  $W$ :

$$W(x, \theta(\gamma, \phi)) = \sum_{k=0}^{\infty} \sum_{n=-k}^k w_{k,n}(x) Y_k^n(\gamma, \phi), \quad x \in \mathbb{R}^3, \quad (1.7)$$

$$Y_k^n(\gamma, \phi) \stackrel{\text{def}}{=} p_k^{|n|}(\cos \gamma) e^{in\phi}, \quad (1.8)$$

$$\theta(\gamma, \phi) = (\cos \gamma, \sin \gamma \cos \phi, \sin \gamma \sin \phi) \in \mathbb{S}^2 \subset \mathbb{R}^3, \quad \gamma \in [0, \pi], \quad \phi \in [0, 2\pi], \quad (1.9)$$

where  $p_k^n(x)$ ,  $x \in [-1, 1]$ ,  $k \in \mathbb{N} \cup \{0\}$ ,  $n = \overline{-k, k}$  are the associated Legendre polynomials. Polynomials  $p_k^n(x)$  are well-known in literature (see e.g. [SW16]) and are defined using ordinary Legendre polynomials  $p_k(x)$  by the formulas:

$$p_k^n(x) = (-1)^n (1-x^2)^{n/2} \frac{d^n}{dx^n} (p_k(x)), \quad n, k \in \mathbb{N} \cup \{0\}, \quad (1.10)$$

$$p_k^{-n}(x) = (-1)^n \frac{(k-n)!}{(k+n)!} p_k^n(x), \quad p_k^n(x) \equiv 0 \text{ if } |n| > k, \quad (1.11)$$

$$p_k(x) = \frac{1}{2^k k!} \frac{d^k}{dx^k} [(x^2-1)^k], \quad x \in [-1, 1], \quad k \in \mathbb{N} \cup \{0\}, \quad (1.12)$$

see also [SW16], [ZT79] for other properties of associated Legendre polynomials. In addition, coefficients  $w_{k,n}(x)$  in (1.7) are defined by the formulas:

$$w_{k,n}(x) = c(k, n) \int_0^{2\pi} d\phi e^{-in\phi} \int_0^\pi W(x, \theta(\gamma, \phi)) p_k^{|n|}(\cos \gamma) \sin \gamma d\gamma, \quad (1.13)$$

$$c(k, n) = \frac{(2k+1)(k-|n|)!}{4\pi(k+|n|)!}, \quad k \in \mathbb{N} \cup \{0\}, \quad n = 0, \pm 1, \dots, \pm k,$$

where function  $\theta(\gamma, \phi)$  is defined in (1.9).

Under assumption (1.2), for each fixed  $x$ , series (1.7) converge in  $L^2(\mathbb{S}^2)$ ; see e.g. [SW16] (Chapter 4), [Mor98] (Chapter 2), [ZT79].

We consider also

$$\sigma_{\widetilde{W}, D, m} = \frac{1}{2\pi} \sum_{k=1}^m \sum_{n=-2k}^{2k} \sup_{x \in D} \left| \frac{w_{2k,n}(x)}{w_{0,0}(x)} \right| \quad \text{for } m \in \mathbb{N}, \quad (1.14)$$

$$\sigma_{\widetilde{W}, D, m} = 0 \text{ for } m = 0,$$

$$\sigma_{\widetilde{W}, D, \infty} = \lim_{m \rightarrow \infty} \sigma_{W, D, m}, \quad (1.15)$$

$$W_N(x, \theta(\gamma, \phi)) = \sum_{k=0}^N \sum_{n=-k}^k w_{k,n}(x) Y_k^n(\gamma, \phi), \quad (1.16)$$

$$\widetilde{W}_N(x, \theta(\gamma, \phi)) = \sum_{k=0}^{[N/2]} \sum_{n=-2k}^{2k} w_{2k,n}(x) Y_{2k}^n(\gamma, \phi), \quad (1.17)$$

$$x \in \mathbb{R}^3, \quad \gamma \in [0, \pi], \quad \phi \in [0, 2\pi], \quad (1.18)$$

where coefficients  $w_{k,n}(x)$  are defined in (1.13),  $Y_k^n(\gamma, \phi)$  are defined by (1.8),  $[N/2]$  denotes the integer part of  $N/2$ .

In the present article we obtained, in particular, the following results under assumptions (1.2), (1.3), (1.4):

1. If  $\sigma_{W, D, \infty} < 1$ , then  $R_W$  is injective and, in addition, the inversion of  $R_W f$  is given via formulas (3.2), (3.3); see Section 3 for details.
2. If  $\sigma_{W, D, \infty} \geq 1$ , then  $f$  can be approximately reconstructed from  $R_W f$  as  $f \approx (R_{\widetilde{W}_N})^{-1} R_W f$ , where  $R_{\widetilde{W}_N}$  is defined according to (1.1) for  $\widetilde{W}_N$  defined by (1.17) for  $N = 2m$ , where  $m$  is chosen as the largest while condition  $\sigma_{W, D, m} < 1$  holds. More precisely, approximate inversion of  $R_W f$  is given via the formulas (3.7), (3.8).

In addition, if  $W = W_N$  defined by (1.16) and  $\sigma_{W, D, m} < 1$ ,  $m = [N/2]$ , where  $[N/2]$  denotes the integer part of  $N/2$ , then  $R_{W_N}$  is injective and invertible by formula (3.17); see Section 3 for details.

Prototypes of these results for the weighted Radon transforms in 2D were obtained in [Kun92], [Nov14], [GuNo14].

The present work also continues the studies of [GN16], where approximate inversion of  $R_W$  was realized as  $(R_{W_N})^{-1}$  for  $N = 0$  or by other words as an approximate Chang-type inversion formula.

In Section 2 we give some notations and preliminary results.

Main results of the present work are presented in detail in Section 3.

In Section 4 we generalize results of Sections 2, 3 for the case of dimension  $n \geq 3$ .

Proofs of results of Sections 2, 3, 4 are presented in Sections 5, 6.

## 2 Some preliminary results

### 2.1 Some formulas for $R$ and $R^{-1}$

We recall that for the classical Radon transform  $R$  (formula (1.1) for the case of  $W \equiv 1$ ) the following identity holds (see [Nat86], Theorem 1.2, p.13):

$$R(f *_{\mathbb{R}^3} g) = Rf *_{\mathbb{R}} Rg, \quad (2.1)$$

where  $*_{\mathbb{R}^3}, *_{\mathbb{R}}$  denote the 3D and 1D convolutions (respectively),  $f, g$  are test functions.

The classical Radon inversion formula is defined as follows (see e.g. [Nat86]):

$$R^{-1}q(x) = -\frac{1}{8\pi^2} \int_{\mathbb{S}^{n-1}} q^{(2)}(x\theta, \theta) d\theta, \quad x \in \mathbb{R}^3, \quad (2.2)$$

$$q^{(2)}(s, \theta) = \frac{d^2}{ds^2} q(s, \theta), \quad s \in \mathbb{R}, \theta \in \mathbb{S}^2,$$

where  $q$  is a test function on  $\mathbb{R} \times \mathbb{S}^2$ .

In addition, from the Projection Theorem (see [Nat86], Theorem 1.1, p.11) it follows that:

$$R^{-1}q(x) \stackrel{def}{=} \frac{1}{(2\pi)^{5/2}} \int_{\mathbb{R}} \frac{\rho^2}{2} d\rho \int_{\mathbb{S}^2} \hat{q}(\rho, \omega) e^{i\rho(x\omega)} d\omega, \quad x \in \mathbb{R}^3, \quad (2.3)$$

$$\hat{q}(s, \theta) \stackrel{def}{=} \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} q(t, \theta) e^{-its} dt, \quad (s, \theta) \in \mathbb{R} \times \mathbb{S}^2, \quad (2.4)$$

where  $q(t, \theta)$  is a test function on  $\mathbb{R} \times \mathbb{S}^2$ .

For the case of  $\hat{q}$  even (i.e.  $\hat{q}(s, \theta) = \hat{q}(-s, -\theta)$ ),  $(s, \theta) \in \mathbb{R} \times \mathbb{S}^2$ , where  $\hat{q}$  is defined in (2.4)), formulas (2.3), (2.4) can be rewritten as follows:

$$R^{-1}q = \frac{1}{2\pi} \mathcal{F}[\hat{q}] = \frac{1}{2\pi} \mathcal{F}^{-1}[\hat{q}], \quad (2.5)$$

where  $\mathcal{F}[\cdot], \mathcal{F}^{-1}[\cdot]$  denote the Fourier transform and its inverse in 3D, respectively, and are defined by the formulas (in spherical coordinates):

$$\mathcal{F}[q](\xi) \stackrel{def}{=} \frac{1}{(2\pi)^{3/2}} \int_0^{+\infty} \rho^2 d\rho \int_{\mathbb{S}^2} q(\rho, \omega) e^{-i\rho(\xi\omega)} d\omega, \quad (2.6)$$

$$\mathcal{F}^{-1}[q](\xi) \stackrel{def}{=} \frac{1}{(2\pi)^{3/2}} \int_0^{+\infty} \rho^2 d\rho \int_{\mathbb{S}^2} q(\rho, \omega) e^{i\rho(\xi\omega)} d\omega, \quad \xi \in \mathbb{R}^3, \quad (2.7)$$

where  $q(\rho, \omega)$  is a test-function on  $[0, +\infty) \times \mathbb{S}^2$  (identified with  $\mathbb{R}^3$ ).

### 2.2 Symmetrization of $W$

Let

$$A_W f = R^{-1} R_W f, \quad (2.8)$$

where  $R_W$  is defined in (1.1),  $f$  is a test function, satisfying assumptions (1.4).

Let

$$\widetilde{W}(x, \theta) \stackrel{def}{=} \frac{1}{2} (W(x, \theta) + W(x, -\theta)), \quad x \in \mathbb{R}^3, \theta \in \mathbb{S}^2. \quad (2.9)$$

The following formulas hold:

$$\frac{1}{2}(R_W f(s, \theta) + R_W f(-s, -\theta)) = R_{\widetilde{W}} f(s, \theta), \quad (s, \theta) \in \mathbb{R} \times \mathbb{S}^2, \quad (2.10)$$

$$A_W f = R^{-1} R_{\widetilde{W}} f, \quad (2.11)$$

$$\widetilde{W}(x, \theta(\gamma, \phi)) = \sum_{k=0}^{\infty} \sum_{n=-2k}^{2k} w_{2k,n}(x) Y_{2k}^n(\gamma, \phi), \quad (2.12)$$

$$x \in \mathbb{R}^3, \quad \gamma \in [0, \pi], \quad \phi \in [0, 2\pi],$$

where  $w_{k,n}$  are defined in (1.13),  $\theta(\gamma, \phi)$  is defined in (1.9),  $Y_k^n(\gamma, \phi)$  are defined in (1.8). Identity (2.11) is proved in [GN16] in 3D, where  $\widetilde{W}$  is denoted as  $W_{sym}$ .

Identity (2.12) follows from (1.8) and the following identities:

$$p_k^n(-x) = (-1)^{n+k} p_k^n(x), \quad x \in [-1, 1], \quad (2.13)$$

$$e^{in(\phi+\pi)} = (-1)^n e^{in\phi} \quad (2.14)$$

where  $p_k^n(x)$ ,  $k \in \mathbb{N} \cup \{0\}$ ,  $n = \overline{-k, k}$  are the associated Legendre polynomials, defined by identities (1.10), (1.11), (1.12) (see e.g. [SW16]).

Note also that  $\widetilde{W}_N$  defined by (1.17) is the approximation of  $\widetilde{W}$  defined by (2.9) and

$$\widetilde{W}_N(x, \cdot) \xrightarrow[N \rightarrow \infty]{L^2(\mathbb{S}^2)} \widetilde{W}(x, \cdot) \text{ for each fixed } x \in \mathbb{R}^3. \quad (2.15)$$

Using formulas (2.10)-(2.12) we reduce inversion of  $R_W$  to the inversion of  $R_{\widetilde{W}}$  defined by (1.1) for  $W = \widetilde{W}$ . Prototypes of (2.8), (2.11), (2.12) for the two-dimensional case can be found in [Kun92], [Nov11].

### 2.3 Operators $Q_{W,D,m}$ and numbers $\sigma_{W,D,m}$

Let  $D$  be the domain of (1.4), and  $\chi_D$  denote the characteristic function of  $D$ , i.e.

$$\chi_D \equiv 1 \text{ on } D, \quad \chi_D \equiv 0 \text{ on } \mathbb{R}^3 \setminus D. \quad (2.16)$$

Let

$$Q_{\widetilde{W},D,m} u(x) \stackrel{def}{=} R^{-1}(R_{\widetilde{W},D,m} u)(x), \quad m \in \mathbb{N} \quad (2.17)$$

$$Q_{\widetilde{W},D,m} u(x) = 0 \text{ for } m = 0,$$

$$Q_{\widetilde{W},D,\infty} u(x) \stackrel{def}{=} R^{-1}(R_{\widetilde{W},D,\infty} u)(x), \quad (2.18)$$

where

$$R_{\widetilde{W},D,m} u(s, \theta(\gamma, \phi)) \stackrel{def}{=} \int_{x\theta=s} \left( \sum_{k=1}^m \sum_{n=-2k}^{2k} \frac{w_{2k,n}(x)}{w_{0,0}(x)} Y_{2k}^n(\gamma, \phi) \right) \chi_D(x) u(x) dx, \quad (2.19)$$

$$\begin{aligned} R_{\widetilde{W},D,\infty} u(s, \theta(\gamma, \phi)) &\stackrel{def}{=} \lim_{m \rightarrow \infty} R_{\widetilde{W},D,m} u(s, \theta(\gamma, \phi)) \\ &= \int_{x\theta=s} \left( \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \frac{w_{2k,n}(x)}{w_{0,0}(x)} Y_{2k}^n(\gamma, \phi) \right) \chi_D(x) u(x) dx, \end{aligned} \quad (2.20)$$

$$x \in \mathbb{R}^3, \quad s \in \mathbb{R}, \quad \theta(\gamma, \phi) \in \mathbb{S}^2,$$

where  $R^{-1}$  is defined by (2.2) or (2.3),  $u$  is a test function,  $Y_k^n(\gamma, \phi)$  are defined in (1.8),  $w_{0,0}, w_{2k,n}$  are the Fourier-Laplace coefficients defined by (1.13) and  $w_{2k,n}/w_{0,0}, \chi_D$  are considered as multiplication operators on  $\mathbb{R}^3$ . Note that  $R_{\widetilde{W},D,\infty} f = R_{\widetilde{W}} f$  under assumption (1.4).

Let

$$d_{2k,n}(x) \stackrel{def}{=} R^{-1}(\delta(\cdot) Y_k^n)(x), \quad x \in \mathbb{R}^3, \quad k \in \mathbb{N}, \quad n = \overline{-2k, 2k}, \quad (2.21)$$

where  $R^{-1}$  is defined by (2.3),  $\delta = \delta(s)$  denotes the 1D Dirac delta function,  $Y_k^n$  is defined in (1.8).

**Lemma 1.** *Let  $d_{2k,n}$  be defined by (2.21). Then*

$$d_{2k,n}(x(r, \gamma, \phi)) = \frac{(-1)^k \sqrt{2} \Gamma(\frac{3}{2} + k)}{\pi \Gamma(k)} \frac{Y_{2k}^n(\gamma, \phi)}{r^3}, \quad r > 0, \quad (2.22)$$

where  $\Gamma(\cdot)$  is the Gamma-function,  $x(r, \gamma, \phi)$  is defined by the identity:

$$x(r, \gamma, \phi) = (r \cos \gamma, r \sin \gamma \cos \phi, r \sin \gamma \sin \phi) \in \mathbb{R}^3, \quad \gamma \in [0, \pi], \quad \phi \in [0, 2\pi], \quad r \geq 0. \quad (2.23)$$

In addition, the following inequality holds:

$$|\mathcal{F}[d_{2k,n}](\xi)| \leq 1/2\pi, \quad \xi \in \mathbb{R}^3, \quad (2.24)$$

where  $\mathcal{F}[\cdot]$  is the Fourier transform defined in (2.6).

The following lemma gives some useful expressions for operators  $Q_{\widetilde{W},D,m}$ ,  $Q_{\widetilde{W},D,\infty}$  defined in (2.17), (2.18).

**Lemma 2.** Let operators  $Q_{\widetilde{W},D,m}$ ,  $Q_{\widetilde{W},D,\infty}$  be defined by (2.17), (2.18), respectively, and  $u$  be a test function satisfying (1.4). Then

$$Q_{\widetilde{W},D,m}u = \sum_{k=1}^m \sum_{n=-2k}^{2k} d_{2k,n} *_{\mathbb{R}^3} \frac{w_{2k,n}}{w_{0,0}} u, \quad (2.25)$$

$$Q_{\widetilde{W},D,\infty}u = \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} d_{2k,n} *_{\mathbb{R}^3} \frac{w_{2k,n}}{w_{0,0}} u, \quad (2.26)$$

where coefficients  $w_{k,n}$  are defined in (1.13),  $d_{2k,n}$  is defined by (2.21),  $*_{\mathbb{R}^3}$  denotes the convolution in  $3D$ .

*Remark 1.* Convolution terms in the right hand-side of (2.25), (2.26) are well defined functions in  $L^2(\mathbb{R}^3)$ . This follows from identity (2.22) and Calderón-Zygmund Theorem for convolution-type operators with singular kernels (see [Kna05], p.83, Theorem 3.26).

The following lemma shows that the operators  $Q_{\widetilde{W},D,m}$ ,  $Q_{\widetilde{W},D,\infty}$  are well-defined in  $L^2(\mathbb{R}^3)$ .

**Lemma 3.** Operator  $Q_{\widetilde{W},D,m}$ , defined by (2.17), is a linear bounded operator in  $L^2(\mathbb{R}^3)$  and the following estimate holds:

$$\|Q_{\widetilde{W},D,m}\|_{L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)} \leq \sigma_{\widetilde{W},D,m}, \quad (2.27)$$

where  $\sigma_{\widetilde{W},D,m}$  is defined by (1.14).

If  $\sigma_{\widetilde{W},D,\infty} < +\infty$ , where  $\sigma_{\widetilde{W},D,\infty}$  is defined by (1.15), then  $Q_{\widetilde{W},D,\infty}$ , defined by (2.18), is a linear bounded operator in  $L^2(\mathbb{R}^3)$  and the following estimate holds:

$$\|Q_{\widetilde{W},D,\infty}\|_{L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)} \leq \sigma_{\widetilde{W},D,\infty}. \quad (2.28)$$

**Lemma 4.** Let

$$\sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \left\| \frac{w_{2k,n}}{w_{0,0}} \right\|_{L^2(D)} < +\infty, \quad (2.29)$$

where  $w_{k,n}$  are defined in (1.13). Then

$$R^{-1}R_W f \in L^2(\mathbb{R}^3). \quad (2.30)$$

In addition, the following formula holds:

$$R^{-1}R_W f = w_{0,0}f + \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} d_{2k,n} *_{\mathbb{R}^3} w_{2k,n}f, \quad (2.31)$$

where  $R_W$  is defined by (1.1),  $W$  satisfies (1.2), (1.3),  $f$  satisfies (1.4), operator  $R^{-1}$  is defined by (2.3),  $d_{2k,n}$  is defined by (2.21).

## 3 Main results

### 3.1 Case of $\sigma_{\widetilde{W},D,\infty} < 1$

Let

$$\sigma_{\widetilde{W},D,\infty} < 1, \quad (3.1)$$

where  $\sigma_{\widetilde{W},D,\infty}$  is defined by (1.15).

Inequality (3.1) and upper bound (2.28) in Lemma 3 imply that operator  $I + Q_{\widetilde{W},D,\infty}$  is continuously invertible in  $L^2(\mathbb{R}^3)$  and the following identity holds (in the sense of the operator norm in  $L^2(\mathbb{R}^3)$ ):

$$(I + Q_{\widetilde{W},D,\infty})^{-1} = I + \sum_{j=1}^{\infty} (-Q_{\widetilde{W},D,\infty})^j, \quad (3.2)$$

where  $I$  is the identity operator in  $L^2(\mathbb{R}^3)$ , operator  $Q_{\widetilde{W},D,\infty}$  is defined in (2.18).

**Theorem 1.** *Let conditions (1.2)-(1.4), (3.1) be fulfilled. Then  $R_W$ , defined by (1.1), is injective and the following exact inversion formula holds:*

$$f = (w_{0,0})^{-1}(I + Q_{\widetilde{W},D,\infty})^{-1}R^{-1}R_W f, \quad (3.3)$$

where  $w_{0,0}$  is defined in (1.3), operator  $(I + Q_{\widetilde{W},D,\infty})^{-1}$  is defined by (3.2),  $R^{-1}$  is defined in (2.2).

*Remark 2.* Formula (3.3) can be considered as the following Fredholm's second kind linear integral equation for the  $w_{0,0}f$ :

$$w_{0,0}f + Q_{\widetilde{W},D,\infty}(w_{0,0}f) = R^{-1}R_W f. \quad (3.4)$$

Inequality (3.1) and identity (3.2) imply that equation (3.4) can be solved by the method of successive approximations.

### 3.2 Case of $1 \leq \sigma_{\widetilde{W},D,\infty} < +\infty$

Let

$$\sigma_{\widetilde{W},D,m} < 1, \text{ for some } m \in \mathbb{N} \cup \{0\}, \quad (3.5)$$

$$\sigma_{\widetilde{W},D,\infty} < +\infty. \quad (3.6)$$

Inequality (3.5) and upper bound (2.27) in Lemma 3 imply that  $I + Q_{\widetilde{W},D,m}$  is continuously invertible in  $L^2(\mathbb{R}^3)$  and the following identity holds (in the sense of the operator norm in  $L^2(\mathbb{R}^3)$ ):

$$(I + Q_{\widetilde{W},D,m})^{-1} = I + \sum_{j=1}^{\infty} (-Q_{\widetilde{W},D,m})^j, \quad (3.7)$$

where  $I$  is the identity operator in  $L^2(\mathbb{R}^3)$ , operator  $Q_{\widetilde{W},D,m}$  is defined in (2.17).

**Theorem 2.** *Let conditions (1.2)-(1.4), (3.5), (3.6) be fulfilled. Then*

$$f \approx f_m \stackrel{def}{=} (w_{0,0})^{-1}(I + Q_{\widetilde{W},D,m})^{-1}R^{-1}R_W f, \quad (3.8)$$

$$f = f_m - (w_{0,0})^{-1}(I + Q_{\widetilde{W},D,m})^{-1}R^{-1}R_{\delta W_m} f, \quad (3.9)$$

$$\|f - f_m\|_{L^2(D)} \leq \frac{\|f\|_{\infty}}{2\pi c(1 - \sigma_{\widetilde{W},D,m})} \sum_{k=m+1}^{\infty} \sum_{n=-2k}^{2k} \|w_{2k,n}\|_{L^2(D)} < +\infty, \quad (3.10)$$

where

$$\delta W_m(x, \theta(\gamma, \phi)) \stackrel{def}{=} W(x, \theta(\gamma, \phi)) - \sum_{k=0}^{2m+1} \sum_{n=-k}^k w_{k,n}(x) Y_k^n(\gamma, \phi), \quad (3.11)$$

$$x \in \mathbb{R}^3, \gamma \in [0, \pi], \phi \in [0, 2\pi], m \in \mathbb{N} \cup \{0\}, \quad (3.12)$$

function  $\theta(\gamma, \phi)$  is defined in (1.9), coefficients  $w_{k,n}$  are defined in (1.13),  $Y_k^n$  are defined in (1.8),  $\sigma_{\widetilde{W},D,m}$  is defined in (1.14), operator  $(I + Q_{\widetilde{W},D,m})^{-1}$  is defined in (3.7), constant  $c$  is defined in (1.5).

*Remark 3.* Formula (3.8) can be considered as the following Fredholm's second kind linear integral equation for  $w_{0,0}f$ :

$$w_{0,0}f + Q_{\widetilde{W},D,m}(w_{0,0}f) = R^{-1}R_W f. \quad (3.13)$$

Inequality (3.5) and identity (3.7) imply that equation (3.13) is solvable by the method of successive approximations.

Note also that condition (3.6) can be relaxed to the following one:

$$\sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \left\| \frac{w_{2k,n}}{w_{0,0}} \right\|_{L^2(D)} < +\infty, \quad (3.14)$$

where  $w_{2k,n}$  are defined in (1.13).

Formula (3.8) is an extension to 3D of the Chang-type two-dimensional inversion formulas in [Cha78], [Nov14], [GuNo14]. In addition, formula (3.8) is an extension of approximate inversion formula in [GN16], where this formula was given for  $m = 0$ .

If (3.5) is fulfilled for some  $m \geq 1$ , then  $f_m$  is a refinement of the Chang-type approximation  $f_0$  and, more generally,  $f_j$  is a refinement of  $f_i$  for  $0 \leq i < j \leq m$ . In addition,  $f_j = f_i$  if  $w_{2k,n} \equiv 0$  for  $i < k \leq j$ ,  $n = -2k, 2k$ . Thus, we propose the following approximate reconstruction of  $f$  from  $R_W f$ :

- (i) find maximal  $m$  such that (3.5) is still efficiently fulfilled,
- (ii) approximately reconstruct  $f$  by  $f_m$  using (3.8).

### 3.3 Exact inversion for finite Fourier series weights

Let

$$W = W_N, N \in \mathbb{N} \cup \{0\}, \quad (3.15)$$

where  $W_N$  is defined by (1.16).

Suppose that

$$\sigma_{\widetilde{W}, D, m} < 1 \text{ for } m = [N/2], \quad (3.16)$$

where  $\sigma_{\widetilde{W}, D, m}$  is defined by (1.14),  $[N/2]$  denotes the integer part of  $N/2$ .

**Theorem 3.** *Let conditions (1.2)-(1.4), (3.15), (3.16) be fulfilled. Then  $R_W$  defined by (1.1) is injective and the following exact inversion formula holds:*

$$f = (w_{0,0})^{-1} (I + Q_{\widetilde{W}, D, m})^{-1} R^{-1} R_W f, \quad (3.17)$$

where  $w_{0,0}$  is defined in (1.2),  $(I + Q_{\widetilde{W}, D, m})^{-1}$  is defined by (3.7),  $R^{-1}$  is defined by (2.2).

*Remark 4.* Formula (3.17) can be considered as the following Fredholm's second type linear integral equation for  $w_{0,0} f$ :

$$w_{0,0} f + Q_{\widetilde{W}, D, m} (w_{0,0} f) = R^{-1} R_W f. \quad (3.18)$$

Identity (3.16) imply that (3.18) can be solved by the method of successive approximations.

## 4 Generalization for multidimensions

Formulas (1.1)-(1.4) admit straightforward extensions to the case of dimension  $n > 3$ . In particular, in this case  $R_W f$  is defined as corresponding integrals along  $(n - 1)$ -dimensional hyperplanes in  $\mathbb{R}^n$ ; see e.g. [GN16], [BQ87].

For weight  $W$  we consider Fourier-Laplace expansion:

$$W(x, \theta) = \sum_{k=0}^{\infty} \sum_{i=1}^{a_{k,n}} w_k^i(x) Y_k^i(\theta), \quad x \in \mathbb{R}^n, \theta \in \mathbb{S}^{n-1}, \quad (4.1)$$

where

$$w_k^i(x) = \|Y_k^i\|_{L^2(\mathbb{S}^{n-1})}^{-2} \int_{\mathbb{S}^{n-1}} W(x, \theta) \overline{Y_k^i(\theta)} d\theta, \quad (4.2)$$

$$a_{k,n+1} = \frac{(n+k)!}{k!n!} - \frac{(n+k-2)!}{(k-2)!n!}, \quad n, k \geq 2; a_{0,n} = 1, a_{1,n} = n, \quad (4.3)$$

where  $\{Y_k^i \mid k = \overline{0, \infty}, i = \overline{1, a_{k,n}}\}$  is the Fourier-Laplace basis of harmonics on  $\mathbb{S}^{n-1}$ ,  $\overline{Y_k^i}$  denotes the complex conjugate of  $Y_k^i$ ; see [SW16], [Mor98]. In this work we choose basis  $Y_k^i$  as in [Hig87] without normalizing constants  ${}_n c_L^i$  (i.e.  $\{Y_k^i\}$  are the products of Legendre polynomials with one complex exponent and without any additional constants).

Formula (1.3) can be rewritten now as follows:

$$w_0^1(x) \stackrel{def}{=} \frac{1}{\text{vol}(\mathbb{S}^{n-1})} \int_{\mathbb{S}^{n-1}} W(x, \theta) d\theta, \quad x \in \mathbb{R}^n, w_0^1(x) \neq 0, x \in \mathbb{R}^n, \quad (4.4)$$

where  $\text{vol}(\mathbb{S}^{n-1})$  denotes the standard Euclidean volume of  $\mathbb{S}^{n-1}$ .



Under assumptions (1.2), (1.4) for dimension  $n > 3$ , series (4.1) converge in  $L^2(\mathbb{S}^{n-1})$ ; see e.g. [SW16] (Chapter 4), [Mor98] (Chapter 2), [ZT79].

Formula (2.3) is extended as follows:

$$R^{-1}q(x) = (2\pi)^{1/2-n} \int_{\mathbb{R}} \frac{|\rho|^{n-1}}{2} \int_{\mathbb{S}^{n-1}} \hat{q}(\rho, \theta) e^{i\rho(x\theta)} d\theta, \quad x \in \mathbb{R}^n, \quad (4.5)$$

where  $q(s, \theta)$  is a test function on  $\mathbb{R} \times \mathbb{S}^{n-1}$ ,  $\hat{q}(s, \theta)$  is defined as in (2.4) (with  $\mathbb{S}^2$  replaced by  $\mathbb{S}^{n-1}$ ).

Fourier transforms defined in (2.6), (2.7) are rewritten now as follows:

$$\mathcal{F}[q](\xi) \stackrel{def}{=} (2\pi)^{-n/2} \int_0^{+\infty} \rho^{n-1} d\rho \int_{\mathbb{S}^{n-1}} q(\rho, \omega) e^{-i\rho(\xi\omega)} d\omega, \quad (4.6)$$

$$\mathcal{F}^{-1}[q](\xi) \stackrel{def}{=} (2\pi)^{-n/2} \int_0^{+\infty} \rho^{n-1} d\rho \int_{\mathbb{S}^{n-1}} q(\rho, \omega) e^{i\rho(\xi\omega)} d\omega, \quad \xi \in \mathbb{R}^n, \quad (4.7)$$

where  $q(\rho, \omega)$  is a test function on  $[0, +\infty) \times \mathbb{S}^{n-1}$  (identified with  $\mathbb{R}^n$ ).

In dimension  $n > 3$ , formulas (2.8)-(2.12) remain valid with  $Y_k^m$  defined in (1.8) replaced by general basis of spherical harmonics  $\{Y_k^i\}$  on  $\mathbb{S}^{n-1}$ . In particular, multidimensional analog of formula (2.12) (with  $x \in \mathbb{R}^n$ ,  $\theta \in \mathbb{S}^{n-1}$ ) holds by the following property of harmonic  $Y_k^i(\theta)$ :

$$Y_k^i(-\theta) = (-1)^k Y_k^i(\theta), \quad \theta \in \mathbb{S}^{n-1}, \quad k \in \mathbb{N} \cup \{0\}, \quad i = \overline{1, a_{k,n}}, \quad (4.8)$$

where  $a_{k,n}$  is defined by (4.3). Identity (4.8) reflects the fact that  $Y_k^i(\theta) = Y_k^i(\theta_1, \theta_2, \dots, \theta_n)$ ,  $\theta = (\theta_1, \theta_2, \dots, \theta_n) \in \mathbb{S}^{n-1}$ ,  $i = \overline{1, a_{k,n}}$  is a homogenous polynomial of degree  $k$ , see e.g. [SW16], [Mor98].

The extensions of (1.14), (1.15), (2.17), (2.18), to the case of dimension  $n > 3$ , are defined as follows:

$$\sigma_{\widetilde{W}, D, m} \stackrel{def}{=} (2\pi)^{(1-n)/2} \sum_{k=1}^m \sum_{i=1}^{a_{2k,n}} \sup_{x \in D} \left| \frac{w_{2k}^i(x)}{w_0^1(x)} \right|, \quad (4.9)$$

$$\sigma_{\widetilde{W}, D, \infty} \stackrel{def}{=} \lim_{m \rightarrow +\infty} \sigma_{\widetilde{W}, D, m} \quad (4.10)$$

$$Q_{\widetilde{W}, D, m} u(x) \stackrel{def}{=} R^{-1}(R_{\widetilde{W}, D, m} u)(x), \quad m \in \mathbb{N} \quad (4.11)$$

$$Q_{\widetilde{W}, D, m} u(x) = 0 \quad \text{for } m = 0,$$

$$Q_{\widetilde{W}, D, \infty} u(x) \stackrel{def}{=} R^{-1}(R_{\widetilde{W}, D, \infty} u)(x), \quad (4.12)$$

where

$$R_{\widetilde{W}, D, m} u(s, \theta) \stackrel{def}{=} \int_{x\theta=s} \left( \sum_{k=1}^m \sum_{i=1}^{a_{2k,n}} \frac{w_{2k}^i(x)}{w_0^1(x)} Y_{2k}^i(\theta) \right) \chi_D(x) u(x) dx, \quad (4.13)$$

$$R_{\widetilde{W}, D, \infty} u(s, \theta) \stackrel{def}{=} \lim_{m \rightarrow \infty} R_{\widetilde{W}, D, m} u(s, \theta), \quad (4.14)$$

$$x \in \mathbb{R}^n, \quad s \in \mathbb{R}, \quad \theta \in \mathbb{S}^{n-1},$$

where  $R^{-1}$  is defined in (4.5), coefficients  $w_k^i$  are defined in (4.2).

Formula (2.21) is now rewritten as follows:

$$d_{2k,i}(x) \stackrel{def}{=} R^{-1}(\delta(\cdot) Y_{2k}^i)(x), \quad x \in \mathbb{R}^n, \quad i = \overline{1, a_{k,n}}, \quad (4.15)$$

where  $R^{-1}$  is defined in (4.5),  $Y_{2k}^i$  is the spherical harmonic on  $\mathbb{S}^{n-1}$ .

Results of Lemma 1 remain valid with formula (2.22) replaced by the following one:

$$d_{2k,i}(r, \theta) = c(k, n) \frac{(-1)^k Y_{2k}^i(\theta)}{r^n}, \quad r > 0, \quad \theta \in \mathbb{S}^{n-1}, \quad (4.16)$$

where

$$c(k, n) = \frac{\sqrt{2}\pi^{(1-n)/2} \Gamma(k + \frac{1}{2}) \Gamma(k + \frac{n}{2})}{\Gamma(k) \Gamma(k + \frac{n-1}{2})} \cdot \left( \frac{\Gamma(k+1)}{\Gamma(k + \frac{1}{2})} \right)^{n-2}, \quad (4.17)$$

$\Gamma(\cdot)$  is the Gamma function, and inequality (2.24) is rewritten as follows:

$$|\mathcal{F}[d_{2k,i}](\xi)| \leq (2\pi)^{(1-n)/2}, \quad \xi \in \mathbb{R}^n, \quad (4.18)$$

where  $\mathcal{F}[\cdot]$  is the Fourier transform defined in (4.6),  $d_{2k,i}$  is defined in (4.15). Constant  $c(k, n)$  in (4.17) is obtained using formulas (4.5), (4.15) and Theorems 1, 2 in [Gon16].

Results of Lemma 2 remain valid, for the case of dimension  $n > 3$ , with formulas (2.25), (2.26) rewritten as follows:

$$Q_{\widetilde{W}, D, m} u = \sum_{k=1}^m \sum_{i=1}^{a_{k,n}} d_{2k,i} *_{\mathbb{R}^n} \frac{w_{2k}^i}{w_0^1} u, \quad (4.19)$$

$$Q_{\widetilde{W}, D, \infty} u = \sum_{k=1}^{\infty} \sum_{i=1}^{a_{k,n}} d_{2k,i} *_{\mathbb{R}^n} \frac{w_{2k}^i}{w_0^1} u, \quad (4.20)$$

where coefficients  $w_{2k}^i(x)$ ,  $w_0^1(x)$  are defined in (4.2),  $a_{k,n}$  is defined in (4.3),  $d_{2k,i}$  is defined in (4.16),  $*_{\mathbb{R}^n}$  denotes the convolution in  $\mathbb{R}^n$ .

Results of Lemma 3 remain valid with  $\mathbb{R}^3$  replaced by  $\mathbb{R}^n$ ,  $n > 3$ , where we use definitions (4.9), (4.11), (4.12).

Assumption (2.29) in Lemma 4 is rewritten now as follows:

$$\sum_{k=1}^{\infty} \sum_{i=1}^{a_{k,n}} \left\| \frac{w_{2k}^i}{w_0^1} \right\|_{L^2(D)} < +\infty, \quad (4.21)$$

where coefficients  $w_k^i$  are defined in (4.2),  $a_{k,n}$  is defined in (4.3),  $D$  is the open bounded domain defined according to (1.4) in  $\mathbb{R}^n$ . Under assumption (4.21), result (2.30) of Lemma 4 remains valid, with  $\mathbb{R}^3$  replaced by  $\mathbb{R}^n$  and with the following analog of formula (2.31), for the case of dimension  $n > 3$ :

$$R^{-1} R_W f = w_0^1 f + \sum_{k=1}^{\infty} \sum_{i=1}^{a_{k,n}} d_{2k,i} *_{\mathbb{R}^n} w_{2k}^i f, \quad (4.22)$$

where  $R^{-1}$  is defined in (4.5),  $f$  is a test function satisfying (1.4) (with  $\mathbb{R}^3$  replaced by  $\mathbb{R}^n$ ),  $a_{k,n}$  is defined in (4.3),  $d_{2k,i}$  is defined in (4.16).

Under assumption (3.1) (with  $\sigma_{\widetilde{W}, D, \infty}$  defined in (4.10)), results of Theorem 1 remain valid in dimension  $n > 3$ .

Under assumptions (3.5), (3.6) (with  $\sigma_{\widetilde{W}, D, m}$ ,  $\sigma_{\widetilde{W}, D, \infty}$  defined in (4.9), (4.10); operators  $Q_{\widetilde{W}, D, m}$ ,  $Q_{\widetilde{W}, D, \infty}$  defined in (4.11), (4.12); operator  $R^{-1}$  defined in (4.5); coefficients  $w_{k,n}$  replaced by  $w_k^i$ , defined in (4.2)) results of Theorem 2 remain valid in the case of dimension  $n > 3$ .

Under assumption (3.16) (with  $\sigma_{\widetilde{W}, D, m}$  defined in (4.9)) results of Theorem 3 also remain valid.

## 5 Proofs of Lemma 1, 2, 3, 4

### 5.1 Proof of Lemma 1

We consider  $x(r, \gamma, \phi)$  defined by (2.23) and  $\omega(\gamma, \phi) = x(1, \gamma, \phi)$  (i.e.  $\omega \in \mathbb{S}^2$ ).

Identity (2.23) implies the following expression of the scalar product  $(x\omega)$  in spherical coordinates in  $\mathbb{R}^3$ :

$$(x\omega) = (x(r, \tilde{\gamma}, \tilde{\phi}), \omega(\gamma, \phi)) = r(\cos \gamma \cos \tilde{\gamma} + \sin \gamma \sin \tilde{\gamma} \cos(\phi - \tilde{\phi})), \quad (5.1)$$

where  $\gamma, \tilde{\gamma} \in [0, \pi]$ ,  $\phi, \tilde{\phi} \in [0, 2\pi]$ ,  $r \geq 0$ .

From formulas (1.8), (2.3), (2.21), (5.1) it follows that

$$\begin{aligned} (2\pi)^{5/2} d_{2k,n}(x(r, \tilde{\gamma}, \tilde{\phi})) &= \int_{\mathbb{R}} \frac{\rho^2}{2} d\rho \int_{\mathbb{S}^2} e^{i\rho(x\omega(\gamma, \phi))} Y_{2k}^n(\gamma, \phi) d\omega(\gamma, \phi) \\ &= \int_{\mathbb{R}} \frac{\rho^2}{2} d\rho \int_0^\pi \sin(\gamma) p_k^{n|}(\cos \gamma) d\gamma \int_0^{2\pi} e^{i\rho(x, \omega(\gamma, \phi)) + in\phi} d\phi \\ &= e^{in(\tilde{\phi} - \pi/2)} \int_{\mathbb{R}} \frac{\rho^2}{2} d\rho \int_0^\pi \sin(\gamma) p_{2k}^{n|}(\cos \gamma) e^{i\rho r \cos \gamma \cos \tilde{\gamma}} \int_0^{2\pi} e^{in\phi + \rho r \sin \gamma \sin \tilde{\gamma} \sin \phi} d\phi \\ &= 2\pi e^{in(\tilde{\phi} - \pi/2)} (-1)^n \int_{\mathbb{R}} \frac{\rho^2}{2} d\rho \int_0^\pi \sin(\gamma) p_{2k}^{n|}(\cos \gamma) e^{i\rho r \cos \gamma \cos \tilde{\gamma}} J_n(\rho r \sin \gamma \sin \tilde{\gamma}) d\gamma, \end{aligned} \quad (5.2)$$

where  $J_n(\cdot)$  is the  $n$ -th standard Bessel function of the first kind; see e.g. [Tem11].

Integrand in  $d\gamma$  in the right hand-side of (5.3) was considered in [NPF+06], where the following exact analytic solution was given:

$$\int_0^\pi \sin(\gamma) p_{2k}^{[n]}(\cos \gamma) e^{i\rho r \cos \gamma \cos \tilde{\gamma}} J_n(\rho r \sin \gamma \sin \tilde{\gamma}) d\gamma = 2i^{2k-n} p_{2k}^{[n]}(\cos \tilde{\gamma}) j_{2k}(\rho r), \quad (5.4)$$

where  $j_{2k}$  is the standard spherical Bessel function of order  $2k$ ; see e.g. [Tem11].

From identities (5.3), (5.4) it follows that:

$$\begin{aligned} d_{2k,n}(x(r, \tilde{\gamma}, \tilde{\phi})) &= \frac{(-1)^k}{(2\pi)^{3/2}} p_{2k}^{[n]}(\cos \tilde{\gamma}) e^{in\tilde{\phi}} \int_{\mathbb{R}} \rho^2 j_{2k}(\rho r) d\rho \\ &= \frac{4\sqrt{\pi}(-1)^k \Gamma(\frac{3}{2} + k)}{(2\pi)^{3/2} \Gamma(k)} \frac{p_{2k}^{[n]}(\cos \tilde{\gamma}) e^{in\tilde{\phi}}}{r^3}, \quad r > 0. \end{aligned} \quad (5.5)$$

where  $\Gamma(\cdot)$  is the Gamma function.

Defenition (1.8) and identity (5.5) imply formula (2.22).

Formulas (2.5), (2.13), (2.14), (2.21) imply that

$$(2\pi) d_{2k,n}(x(r, \tilde{\gamma}, \tilde{\phi})) = \mathcal{F}^{-1}[Y_{2k}^n](x(r, \tilde{\gamma}, \tilde{\phi})), \quad r > 0, \tilde{\gamma} \in [0, \pi], \tilde{\phi} \in [0, 2\pi], \quad (5.6)$$

where  $\mathcal{F}^{-1}[\cdot]$  is defined in (2.7),  $Y_k^n$  is defined in (1.8).

From the invertibility of the Fourier transform defined in (2.6), (2.7) and identity (5.6) the following identity holds:

$$2\pi \mathcal{F}[d_{2k,n}] = \mathcal{F} \mathcal{F}^{-1}[Y_{2k}^n] = Y_{2k}^n. \quad (5.7)$$

For  $Y_k^n$  defined in (1.8) the following inequality holds (see e.g. [SW16]):

$$|Y_k^n(\gamma, \theta)| \leq 1, \quad \gamma \in [0, \pi], \theta \in [0, 2\pi]. \quad (5.8)$$

Identities (5.6) and inequality (5.8) imply (2.24).

Note that  $|\mathcal{F}[d_{2k,n}](\xi)|$ ,  $\xi \in \mathbb{R}^3$ , is uniformly bounded by  $1/2\pi$  except only one point  $\xi = 0$ , where direction  $\xi/|\xi| \in \mathbb{S}^2$  is not defined. However, point  $\xi = 0$  is of Lebesgue measure zero and  $\mathcal{F}[d_{2k,n}]$  can be defined with any value at the origin in  $\mathbb{R}^3$ .

Lemma 1 is proved.

## 5.2 Proof of Lemma 2

From identity (2.17) it follows that

$$\begin{aligned} Q_{\tilde{W}, D, m} u &= R^{-1} \left( \sum_{k=1}^m \sum_{n=-2k}^{2k} Y_{2k}^n R \left( \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right) \right) \\ &= R^{-1} \left( \sum_{k=1}^m \sum_{n=-2k}^{2k} (\delta(\cdot) Y_{2k}^n) *_{\mathbb{R}} R \left( \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right) \right) \\ &= R^{-1} \left( \sum_{k=1}^m \sum_{n=-2k}^{2k} R(d_{2k,n}) *_{\mathbb{R}} R \left( \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right) \right) \end{aligned} \quad (5.9)$$

where  $*_{\mathbb{R}}$  denotes the 1D convolution,  $\delta = \delta(s)$  is the 1D-Dirac delta function,  $d_{2k,n}$  is defined by (2.21).

Identities (2.1), (5.9) imply (2.25).

For operator  $Q_{\tilde{W}, D, \infty}$  defined by (2.18) we proceed according to identity (5.9) with  $m \rightarrow +\infty$ . Linearity of operator  $R^{-1}$  defined by (2.2) and identities (2.1), (5.9) imply (2.26).

Lemma 2 is proved.

## 5.3 Proof of Lemma 3

From formula (2.25) and the fact that Fourier transform defined in (2.6) does not change the  $L^2$ -norm we obtain:

$$\begin{aligned} \|Q_{\tilde{W}, D, m} u\|_{L^2(\mathbb{R}^3)} &\leq \sum_{k=1}^m \sum_{n=-2k}^{2k} \left\| d_{2k,n} *_{\mathbb{R}^3} \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right\|_{L^2(\mathbb{R}^3)} \\ &= \sum_{k=1}^m \sum_{n=-2k}^{2k} \left\| \mathcal{F}[d_{2k,n}] \mathcal{F} \left[ \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right] \right\|_{L^2(\mathbb{R}^3)}. \end{aligned} \quad (5.10)$$

From inequalities (2.24), (5.10) we obtain:

$$\begin{aligned}
\|Q_{\widetilde{W},D,m} u\|_{L^2(\mathbb{R}^3)} &\leq \frac{1}{2\pi} \sum_{k=1}^m \sum_{n=-2k}^{2k} \left\| \mathcal{F} \left( \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right) \right\|_{L^2(\mathbb{R}^3)} \\
&= \frac{1}{2\pi} \sum_{k=1}^m \sum_{n=-2k}^{2k} \left\| \frac{w_{2k,n}}{w_{0,0}} \chi_D u \right\|_{L^2(\mathbb{R}^3)} \\
&\leq \sigma_{\widetilde{W},D,m} \|u\|_{L^2(D)},
\end{aligned} \tag{5.11}$$

where  $\sigma_{\widetilde{W},D,m}$  is defined by (1.14).

Inequality (5.11) implies (2.27).

Estimate (2.28) follows from definition (2.18), formula (2.26), linearity of operator  $R^{-1}$  defined by (2.2), inequalities (5.10), (5.11) for  $m \rightarrow +\infty$ .

Lemma 3 is proved.

## 5.4 Proof of Lemma 4

From formulas (1.1), (1.7) it follows that

$$R_W f(s, \theta(\gamma, \phi)) = \sum_{k=0}^{\infty} \sum_{n=-k}^k Y_k^n(\gamma, \phi) R[w_{k,n} f](s, \theta(\gamma, \phi)), \tag{5.12}$$

$$s \in \mathbb{R}, \gamma \in [0, \pi], \phi \in [0, 2\pi], \tag{5.13}$$

where  $\theta(\gamma, \phi)$  is defined in (1.9),  $Y_k^n(\gamma, \phi)$  are defined by (1.8),  $w_{k,n}$  are defined in (1.13).

Formula (2.31) follows from formulas (2.8), (2.11), (2.12), (2.18) and formula (2.26) in Lemma 2, where test function  $u$  is replaced by  $w_{0,0}u$ .

From inequality (2.24), formulas (2.1), (2.31) and the fact that Fourier transform defined in (2.6) does not change the  $L^2$ -norm we obtain:

$$\begin{aligned}
\|R^{-1} R_W f\|_{L^2(\mathbb{R}^3)} &\leq \|w_{0,0} f\|_{L^2(\mathbb{R}^3)} + \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \|d_{2k,n} *_{\mathbb{R}^3} w_{2k,n} f\|_{L^2(\mathbb{R}^3)} \\
&= \|w_{0,0} f\|_{L^2(\mathbb{R}^3)} + \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \|\mathcal{F}[d_{2k,n}] \mathcal{F}[w_{2k,n} f]\|_{L^2(\mathbb{R}^3)} \\
&\leq \|w_{0,0} f\|_{L^2(\mathbb{R}^3)} + \frac{\|f\|_{\infty}}{2\pi} \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \|w_{2k,n}\|_{L^2(D)},
\end{aligned} \tag{5.14}$$

where  $\|\cdot\|_{\infty}$  denotes the  $L^{\infty}$ -norm,  $\mathcal{F}[\cdot]$  is defined in (2.6).

From assumption (2.29) and formula (1.5) it follows that

$$\sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \|w_{2k,n}\|_{L^2(D)} < +\infty. \tag{5.15}$$

Inequalities (5.14), (5.15) and formulas (1.2), (1.3), (1.4) imply that

$$\|R^{-1} R_W f\|_{L^2(\mathbb{R}^3)} \leq \|w_{0,0} f\|_{L^2(\mathbb{R}^3)} + \frac{\|f\|_{\infty}}{2\pi} \sum_{k=1}^{\infty} \sum_{n=-2k}^{2k} \|w_{2k,n}\|_{L^2(D)} < +\infty. \tag{5.16}$$

Lemma 2 is proved.

## 6 Proofs of Theorems 1, 2, 3

### 6.1 Proof of Theorem 1

From formulas (2.18), (2.26), (2.31) we obtain:

$$R^{-1} R_W f = (I + Q_{\widetilde{W},D,\infty})(w_{0,0} f), \tag{6.1}$$

where  $I$  is the identity operator in  $L^2(\mathbb{R}^3)$ ,  $w_{0,0}$  is defined in (1.3).

Assumption (3.1) implies that identity (2.30) holds (by Lemma 4).

Formula (3.3) follows from assumption (3.1), inequality (2.28) in Lemma 3 and formula (6.1).

Theorem 1 is proved.

## 6.2 Proof of Theorem 3

From formulas (2.17), (2.25), (2.31) we obtain:

$$R^{-1}R_W f = (I + Q_{\widetilde{W},D,m})(w_{0,0}f), \quad (6.2)$$

where  $I$  is the identity operator in  $L^2(\mathbb{R}^3)$ ,  $w_{0,0}$  is defined in (1.3).

Assumption (3.16) implies that identity (2.30) holds (by Lemma 4).

Formula (3.17) follows from assumption (3.16), inequality (2.27) in Lemma 3 and formula (6.2).

Theorem 3 is proved.

## 6.3 Proof of Theorem 2

By assumption (3.6) inequality (2.29) holds. Hence, formulas (2.30), (2.31) (in Lemma 4) hold.

Formulas (1.2)-(1.4), assumption (3.5) and inequality (2.27) from Lemma 3 imply that  $f_m \in L^2(\mathbb{R}^3)$ , where  $f_m$  is defined in (3.8).

We split expansion (1.7) of weight  $W$  defined by (1.2) in the following way:

$$W(x, \theta) = W_{N+1}(x, \theta) + \delta W_m(x, \theta), \quad \theta \in \mathbb{S}^2, x \in \mathbb{R}^3, m = [N/2], \quad (6.3)$$

where  $W_{N+1}$  is defined by (1.16),  $[N/2]$  denotes the integer part of  $N/2$ ,  $\delta W_m$  is defined by (3.11).

From formulas (1.2)-(1.4) it follows that

$$R_W f = R_{W_{N+1}} f + R_{\delta W_m} f, \quad (6.4)$$

where  $R_W f, R_{W_{N+1}} f, R_{\delta W_m} f$  are defined by (1.1) for the case of weights  $W, W_{N+1}, \delta W_m$  defined in (1.2), (1.13), (3.11), respectively.

Identities (2.8), (2.11), (6.3), (6.3) imply that

$$R^{-1}R_W f = R^{-1}R_{W_N} f + R^{-1}R_{\delta W_m} f, \quad (6.5)$$

where  $R^{-1}$  is defined by (2.2). By assumption (3.6) inequality (2.29) holds for the cases of weights  $W, W_N, \delta W_m$ , respectively. Therefore, Lemma 4 holds for weights  $W, W_N, \delta W_m$  and, in particular, from identity (2.30) we obtain:

$$R^{-1}R_{W_N} f \in L^2(\mathbb{R}^3), R^{-1}R_{\delta W_m} f \in L^2(\mathbb{R}^3). \quad (6.6)$$

By assumption (3.5) Theorem 3 holds for  $W = W_N, N = 2m$ . Therefore, from formula (3.17) we obtain:

$$f = (w_{0,0})^{-1}(I + Q_{\widetilde{W},D,m})^{-1}R^{-1}R_{W_N} f, \quad (6.7)$$

where operator  $Q_{\widetilde{W},D,m}$  is defined in (2.17) for  $m$  in (3.5).

Formula (3.9) follows from formula (3.8), identities (6.5), (6.6), (6.7).

Identities (3.7), (3.9) and inequality (3.5) imply the following sequence of inequalities:

$$\|f - f_m\|_{L^2(\mathbb{R}^3)} \leq \frac{1}{c} \|(I + Q_{\widetilde{W},D,m})^{-1}\| \cdot \|R^{-1}R_{\delta W_m} f\|_{L^2(\mathbb{R}^3)} \leq \frac{1}{c(1 - \sigma_{\widetilde{W},D,m})} \|R^{-1}R_{\delta W_m} f\|_{L^2(\mathbb{R}^3)} \quad (6.8)$$

where  $c$  is defined in (1.5),  $Q_{\widetilde{W},D,m}$  is defined by (2.17) for  $m$  in (3.5).

From formulas (2.31), (3.11) and according to (5.14) it follows that

$$\|R^{-1}R_{\delta W_m} f\|_{L^2(\mathbb{R}^3)} \leq \frac{\|f\|_{\infty}}{2\pi} \sum_{k=m+1}^{\infty} \sum_{n=-2k}^{2k} \|w_{2k,n}\|_{L^2(D)}, \quad (6.9)$$

where  $R^{-1}$  is defined by (2.2),  $w_{2k,n}$  are defined by (1.13).

Identities (6.8), (6.9) imply (3.10).

Theorem 2 is proved.

## 7 Acknowledgments

The present work was fulfilled in the framework of research conducted under the direction of Prof. R.G. Novikov.

## References

- [BQ87] J. Boman, E. T. Quinto. Support theorems for real-analytic Radon transforms. *Duke Mathematical J.*, 55(4):943–948, 1987.
- [Cha78] L. Chang. A method for attenuation correction in radionuclide computed tomography. *IEEE Transactions on Nuclear Science*, 25(1):638–643, 1978.
- [Dea07] S. R. Deans. *The Radon transform and some of its applications*. Courier Corporation, 2007.
- [DB84] T. Durrani, D. Bisset. The Radon transform and its properties. *Geophysics*, 49(8):1180–1187, 1984.
- [GGV14] I. M. Gel’fand, M. I. Graev, N. Ya. Vilenkin. *Integral geometry and representation theory*, vol. 5. Academic Press, 2014.
- [GN16] F. O. Goncharov, R. G. Novikov. An analog of Chang inversion formula for weighted Radon transforms in multidimensions. *Eurasian Journal of Mathematical and Computer Applications*, 4(2):23–32, 2016.
- [Gon16] F. O. Goncharov, Integrals of spherical harmonics with Fourier exponents in multidimensions. *Preprint*, 2016.
- [Gra91] P. Grangeat. Mathematical framework of cone beam 3D reconstruction via the first derivative of the Radon transform. In *Mathematical methods in tomography*, p. 66–97. Springer, 1991.
- [GuNo14] J-P. Guillemin, R. G. Novikov. Inversion of weighted Radon transforms via finite Fourier series weight approximations. *Inverse Problems in Science and Engineering*, 22(5):787–802, 2014.
- [Hig87] Atsushi Higuchi. Symmetric tensor spherical harmonics on the n-sphere and their application to the de Sitter group  $SO(N, 1)$ . *Journal of mathematical physics*, 28(7):1553–1566, 1987.
- [Kna05] A. W. Knap. *Advanced real analysis*. Springer, 2005.
- [Kun92] L. A. Kunyansky. Generalized and attenuated Radon transforms: restorative approach to the numerical inversion. *Inverse Problems*, 8(5):809, 1992.
- [Mor98] M. Morimoto. *Analytic functionals on the sphere*. American Mathematical Soc., 1998.
- [Nat86] F. Natterer. *The mathematics of computerized tomography*, vol. 32. SIAM, 1986.
- [NPF+06] A. A. Neves, L.A. Padilha, A. F., E. Rodriguez, C. Cruz, L. Barbosa, C. Cesar. Analytical results for a Bessel function times Legendre polynomials class integrals. *Journal of Physics A: Mathematical and General*, 39(18):293, 2006.
- [Nov11] R. G. Novikov. Weighted Radon transforms for which Chang’s approximate inversion formula is exact. *Russian Mathematical Surveys*, 66(2):442–443, 2011.
- [Nov14] R. G. Novikov. Weighted Radon transforms and first order differential systems on the plane. *Moscow mathematical journal*, 14(4):807–823, 2014.
- [Rad17] J. Radon. Über die Bestimmung von Funktionen durch ihre Integralwerte längs gewisser Mannigfaltigkeiten. *Ber. Saechs Akad. Wiss. Leipzig, Math-Phys*, 69:262–267, 1917.
- [SW16] E. M. Stein, G. Weiss. *Introduction to Fourier analysis on Euclidean spaces (PMS-32)*, vol. 32. Princeton University Press, 2016.
- [Tem11] N. M. Temme. *Special functions: An introduction to the classical functions of mathematical physics*. John Wiley & Sons, 2011.
- [ZT79] S. B. Zhizhiashvili, L. V. Topuriya. Fourier-Laplace series on a sphere. *Journal of Soviet Mathematics*, 12(6):682–714, 1979.