



HAL
open science

Spatial patterns of farm land use in fragmented semi-arid landscapes: The case of Lebna catchment (Cap Bon, Tunisia)

Insaf Mekki, Anne Biarnès, Jean-Stéphane Bailly, Fabrice Vinatier, Hichem Chebbi, Tarek Ajmi, Yves Blanca, Frédéric Jacob, Abdelaziz Zairi

► To cite this version:

Insaf Mekki, Anne Biarnès, Jean-Stéphane Bailly, Fabrice Vinatier, Hichem Chebbi, et al.. Spatial patterns of farm land use in fragmented semi-arid landscapes: The case of Lebna catchment (Cap Bon, Tunisia). Agromed International Conference 2016, INRA, Dec 2016, Avignon, France. hal-01408479

HAL Id: hal-01408479

<https://hal.science/hal-01408479>

Submitted on 5 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open licence - etalab

Spatial patterns of farm land use in fragmented semi-arid landscapes: The case of Lebna catchment (Cap Bon, Tunisia)

Insaf Mekki^{±1}, Anne Biarnès², Jean Stéphane Bailly³, Fabrice Vinatier⁴, Hichem Chebbi¹, Tarek Ajmi¹, Yves Blanca², Frédéric Jacob^{1,2}, Abdelaziz Zairi¹

¹ INRGREF, University of Carthage, Tunisia

² IRD, UMR LISAH, France

³ AgroParisTech, UMR LISAH et TETIS, France

⁴ Inra, UMR LISAH, France

* Speaker

± Corresponding author: (insaf.mekki@laposte.net)

Introduction

Mediterranean agricultural landscapes provide ecosystem services driven by land use patterns such as crop spatial distribution (Millennium Ecosystem Assessment, 2005). Assessing the determinants of crop location is therefore a prerequisite for exploring spatially explicit land use scenarios and evaluating the subsequent ecosystem services (Benoît et al., 2012). Although remote sensing techniques are now able to provide time series of land use patterns, they do not permit to identify the driving factors of crop location that are related to farm characteristics. Intra-farm land fragmentation is a typical characteristic of rural landscapes (Demetriou, 2014), and we hypothesized in the current study that it may strongly influence crop spatial patterns. Therefore, the current paper aimed to study the impact of farmland fragmentation in crop location within a semi-arid rural zone of Tunisia, with a focus on spatiotemporal dependencies.

Materials and Methods

The study was conducted over the Lebna catchment (220 km² - Tunisian Cap Bon Peninsula), which is typified by a hilly topography, scattered small farm territories, pasture vegetation, rainfed mixed farming (wheat, fodder, condiment and legume) and livestock. We characterized spatial pattern of crop location at the catchment scale and we analyzed the role of farmland fragmentation in the observed pattern by using Spot imagery based classification and georeferenced database of field observations. The database included 340 fields belonging to 30 farms selected according to their land structure (scattered or aggregated fields) and 1220 neighboring fields. The collected variables were (i) the crop type in all fields for the 2016 cycle, to account for spatial dependencies between neighboring fields, (ii) the crop type in the 340 aforementioned fields for the 2015 cycle, to account for crop rotation driver and (iii) the field context (isolated or non-isolated field relative to the other fields and to the farm facilities). We first mapped crop distribution by applying a Random Forest model on the field database and three Spot images collected between mid-March and late April 2016. Then, univariate analysis permitted to characterize the variability of both crop distribution in neighboring fields and previous crop in the same field, in accordance to crop and field context. Finally, linear discriminant analysis permitted to quantify the ability to predict crop type from the variables describing neighborhood and previous year crop.

Results and Discussion

The distribution of annual crops in the Lebna catchment was mainly characterized by aggregates of several neighboring fields sown with the same crop type. Rules of crop location

depended upon field context. For isolated fields, crop distribution in neighboring fields significantly depended upon the crop sown in the field (Fig. 1, for the case of wheat). Neighboring crops and crop rotation had the same weight on crop choice (Tab. 1). Overall, it seems that farmer strategy rely on agreements with their neighbors to sow the same crop type and therefore to follow the same crop rotation. These agreements are likely to permit management of constraints related to field accessibility and free grazing after crop harvest. In non-isolated fields, such agreements are unnecessary. Distribution of neighboring crops appears to be more random, and crop choice significantly depends more upon crop rotation than upon neighboring crops.




Figure 1. Distribution of neighboring wheat plots according to crop type and field context
W: wheat, CL: condiment or legumes, F: fodder (mainly other cereals).

Table 1. Results of the multi linear regression obtained from 1000 cross-validations and comparison with random distributions.

Tested variables	Average rate of good prediction of the type of crop (standard deviation)	
	Isolated fields	Non-isolated fields
Random distribution	0.34 (0.04)	0.37 (0.04)
Neighboring crops (percentages of neighboring fields sown in each crops type)	0.56 (0.05)	0.49 (0.07)
Main previous year crop	0.56 (0.05)	0.63 (0.07)

Conclusion

Our study revealed the significant effect of farmland fragmentation on the spatial pattern of land use at the catchment scale. For a complete overview of the driver complexity, other potential determinants should also be explored, including productive strategies of farmers and physical characteristics of their fields. Furthermore, our findings underscore the need to deepen our knowledge of existing agreements between farmers to refine our understanding of crops distribution.

Acknowledgements

Financial support for this study was provided by the ANR TRANSMED ALMIRA project (contract ANR-12-TMED-0003-01). The fieldwork was performed within the memorandum of understanding between CRDA Nabeul and INRGREF. We gratefully acknowledge I. Belouafi for participation in the fieldwork. The cooperation of the farmers at Lebna region in providing access to their land and data related to their farms was highly appreciated.

References

- Millennium Ecosystem Assessment, 2005/ <http://www.maweb.org/en/index.aspx>.
- Benoît M., Rizzo D. Marraccini E., Moonen A. C., Galli M., Lardon S., Rapey H., Thenail C., Bonari E., 2012. Landscape agronomy: a new field for addressing agricultural landscape dynamics, *Landscape Ecol.*, 27:1385–1394
- Demetriou D. 2014. The Development of an Integrated Planning and Decision Support System (IPDSS) for Land Consolidation, Springer Theses, DOI: 10.1007/978-3-319-02347-2_2