



HAL
open science

Existence of guided waves due to a lineic perturbation of a 3D periodic medium

Bérangère Delourme, Patrick Joly, Elizaveta Vasilevskaya

► To cite this version:

Bérangère Delourme, Patrick Joly, Elizaveta Vasilevskaya. Existence of guided waves due to a lineic perturbation of a 3D periodic medium. Applied Mathematics Letters, 2016, 10.1016/j.aml.2016.11.017 . hal-01412396

HAL Id: hal-01412396

<https://hal.science/hal-01412396>

Submitted on 8 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence of guided waves due to a lineic perturbation of a 3D periodic medium

Bérangère Delourme^{a,b}, Patrick Joly^b, Elizaveta Vasilevskaya^{a,b}

a: Université Paris 13, Sorbonne Paris Cité, LAGA, UMR 7539, 93430 Villetaneuse, France

b: POEMS, UMR 7231 CNRS/ENSTA/Inria, ENSTA ParisTech, 828 boulevard des Maréchaux, 91762 Palaiseau Cedex, France

Abstract

In this note, we exhibit a three dimensional structure that permits to guide waves. This structure is obtained by a geometrical perturbation of a 3D periodic domain that consists of a three dimensional grating of equi-spaced thin pipes oriented along three orthogonal directions. Homogeneous Neumann boundary conditions are imposed on the boundary of the domain. The diameter of the section of the pipes, of order $\varepsilon > 0$, is supposed to be small. We prove that, for ε small enough, shrinking the section of one line of the grating by a factor of $\sqrt{\mu}$ ($0 < \mu < 1$) creates guided modes that propagate along the perturbed line. Our result relies on the asymptotic analysis (with respect to ε) of the spectrum of the Laplace-Neumann operator in this structure. Indeed, as ε tends to 0, the domain tends to a periodic graph, and the spectrum of the associated limit operator can be computed explicitly.

Keywords : guided waves, periodic media, spectral theory.

AMS codes : 78M35, 35J05, 58C40

1 Statement of the problem

Let ω_1 , ω_2 and ω_3 be three Lipschitz bounded domains of \mathbb{R}^2 of same area ($|\omega_1| = |\omega_2| = |\omega_3|$) containing the origin $(0,0)$, let $\varepsilon > 0$ be a parameter (that is going to be small), and let a_1 , a_2 and a_3 be three positive real numbers. We denote by $(\mathbf{e}_i)_{i \in \{1,2,3\}}$, the standard basis of \mathbb{R}^3 . For any $(k, \ell) \in \mathbb{Z}^2$, we consider the three dimensional domain $D_{k,\ell,3}^\varepsilon$ defined by

$$D_{k,\ell,3}^\varepsilon = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \text{ such that } ((x_1 - a_1 k)/\varepsilon, (x_2 - a_2 \ell)/\varepsilon) \in \omega_3\},$$

which is an unbounded cylinder of constant cross section $\varepsilon\omega_3$. It is infinite along the \mathbf{e}_3 direction (invariant with respect to x_3) and contains the point $(a_1 k, a_2 \ell, 0)$. Similarly, for any $(k, \ell) \in \mathbb{Z}^2$, we define the domains

$$D_{k,\ell,1}^\varepsilon = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \text{ such that } ((x_2 - a_2 k)/\varepsilon, (x_3 - a_3 \ell)/\varepsilon) \in \omega_1\},$$

$$D_{k,\ell,2}^\varepsilon = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \text{ such that } ((x_1 - a_1 k)/\varepsilon, (x_3 - a_3 \ell)/\varepsilon) \in \omega_2\},$$

and we consider the periodic domain Ω_ε given by

$$\Omega_\varepsilon = \bigcup_{i \in \{1,2,3\}} \bigcup_{(k,\ell) \in \mathbb{Z}^2} D_{k,\ell,i}^\varepsilon. \quad (1)$$

The domain Ω_ε is a three dimensional grating of equi-spaced parallel pipes (of constant cross section) oriented along the three orthogonal directions \mathbf{e}_1 , \mathbf{e}_2 and \mathbf{e}_3 . It is a_j -periodic with respect to x_j , $j = 1, 2, 3$. Moreover, the points $(ka_1, \ell a_2, ma_3)$, $(k, \ell, m) \in \mathbb{Z}^3$, belong to Ω_ε .

In order to create guided modes, we introduce a linear defect (see [1]-[5]-[2]) in the periodic structure by modifying the section size of one pipe of the grating (it is conjectured that guided modes cannot appear in the purely periodic structure, see [4] for the proof in the case of a symmetric medium). More precisely, we assume that the domain $D_{0,0,3}^\varepsilon$ is replaced with the domain

$$D_{0,0,3}^{\varepsilon,\mu} = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \text{ such that } (x_1/(\sqrt{\mu}\varepsilon), x_2/(\sqrt{\mu}\varepsilon)) \in \omega_3\},$$




Figure 1: Illustration of the perturbed periodic domain Ω_ε^μ and the limit graph \mathcal{G} .

where μ is a positive parameter. In other words, we enlarge ($\mu > 1$) or shrink ($0 < \mu < 1$) the section of one pipe of the domain by a factor μ (see Fig 1a). The corresponding perturbed domain is denoted by Ω_ε^μ . Its precise definition is given by

$$\Omega_\varepsilon^\mu = \left(\bigcup_{i \in \{1,2\}} \bigcup_{(k,\ell) \in \mathbb{Z}^2} D_{k,\ell,i}^\varepsilon \right) \cup \left(\bigcup_{(k,\ell) \in \mathbb{Z}^2 \setminus \{(0,0)\}} D_{k,\ell,3}^\varepsilon \right) \cup D_{0,0,3}^{\varepsilon,\mu}. \quad (2)$$

Ω_ε^μ is still a_3 -periodic with respect to x_3 . However, the presence of the perturbed pipe $D_{0,0,3}^{\varepsilon,\mu}$ breaks the periodicity with respect to x_1 and x_2 . We emphasize that the domain Ω_ε^μ (as well as Ω_ε) tends to a 3D periodic graph as ε tends to 0.

We look for guided modes, i.e. solutions to the wave equation $\partial_t^2 u - \Delta u = 0$ in Ω_ε^μ , satisfying homogeneous Neumann boundary conditions on $\partial\Omega_\varepsilon^\mu$ (see [7] for the investigation of the Dirichlet case), that propagate along the defect pipe $D_{0,0,3}^{\varepsilon,\mu}$ (i.e. in the \mathbf{e}_3 direction) but stay confined in the transversal directions. More precisely, denoting by B_ε^μ the restriction of the domain Ω_ε^μ to the band $|x_3| < a_3/2$,

$$B_\varepsilon^\mu = \{(x_1, x_2, x_3) \in \Omega_\varepsilon^\mu \text{ such that } |x_3| < a_3/2\}, \quad (3)$$

we search solutions of the form $u(x_1, x_2, x_3, t) = v(x_1, x_2, x_3)e^{i\omega t - \beta x_3}$, where β is a real parameter and $v(x_1, x_2, x_3) \in L_2(B_\varepsilon^\mu)$ is an a_3 -periodic function in x_3 . In fact, it is easily seen that the β -quasi-periodic function $v(x_1, x_2, x_3)e^{-i\beta x_3}$ is an eigenfunction of the operator

$$A_\varepsilon^\mu(\beta) : D(A_\varepsilon^\mu(\beta)) \subset L_2(B_\varepsilon^\mu) \rightarrow L_2(B_\varepsilon^\mu), \quad A_\varepsilon^\mu(\beta) = -\Delta u \text{ in } B_\varepsilon^\mu, \quad (4)$$

with $D(A_\varepsilon^\mu(\beta)) = \{u \in H_\Delta^1(B_\varepsilon^\mu), u|_{\Sigma^+} = e^{-i\beta} u|_{\Sigma^-}, \partial_{x_3} u|_{\Sigma^+} = e^{-i\beta} \partial_{x_3} u|_{\Sigma^-}, \partial_n u|_{\partial B_\varepsilon^\mu \setminus \Sigma^\pm} = 0\}$, where

$$H_\Delta^1(B_\varepsilon^\mu) = \{u \in H^1(B_\varepsilon^\mu), \text{ s.t. } \Delta u \in L_2(B_\varepsilon^\mu)\} \text{ and } \Sigma^\pm = \{(x_1, x_2, x_3) \in \partial B_\varepsilon^\mu, x_3 = \pm a_3/2\}.$$

To study the spectral properties of $A_\varepsilon^\mu(\beta)$, we investigate its (formal) limit $\mathcal{A}^\mu(\beta)$ as ε tends to 0. The operator $\mathcal{A}^\mu(\beta)$ is defined on the limit graph \mathcal{G} (see Fig. 1b) and its spectrum can be explicitly computed. In particular, its spectrum has infinitely many gaps (Lemma 2.1), i.e. open intervals $(a, b) \subset \mathbb{R}$ such that the intersection of $[a, b]$ with the spectrum is reduced to $\{a, b\}$. Moreover, for $\mu < 1$, there is at least one eigenvalue in each gap (Lemma 2.5). Since, in addition, for $\varepsilon > 0$ sufficiently small, the spectrum of $A_\varepsilon^\mu(\beta)$ is close to the spectrum of $\mathcal{A}^\mu(\beta)$, the existence of guided modes is guaranteed (Theorem 3.1).

2 The spectrum of the limit operator $\mathcal{A}^\mu(\beta)$

2.1 Definition of the limit operator $\mathcal{A}^\mu(\beta)$

The limit operator $\mathcal{A}^\mu(\beta)$ is defined on the infinite periodic graph $\mathcal{G} = \bigcap_{\varepsilon>0} B_\varepsilon^\mu$ obtained as the limit of B_ε^μ as ε tends to 0: \mathcal{G} is made of the vertices $\{v_{k,\ell} = (ka_1, \ell a_2, 0), v_{k,\ell}^\pm = (ka_1, \ell a_2, \pm a_3/2), (k, \ell) \in \mathbb{Z}^2\}$ connected by the edges

$$\{e_{k+1/2,\ell} = (v_{k,\ell}, v_{k+1,\ell}), e_{k,\ell+1/2} = (v_{k,\ell}, v_{k,\ell+1}), e_{k,\ell}^\pm = (v_{k,\ell}, v_{k,\ell}^\pm), (k, \ell) \in \mathbb{Z}^2\}.$$

It is a_1 -periodic with respect to x_1 and a_2 -periodic with respect to x_2 (see Fig. 1b).

For any function u defined on \mathcal{G} , we denote by $\mathbf{u}_{k,\ell}$ (resp. $\mathbf{u}_{k,\ell}^\pm$) its value at the vertex $v_{k,\ell}$ (resp. $v_{k,\ell}^\pm$). The restriction of u to the edge $e_{k+1/2,\ell}$ (resp. $e_{k,\ell+1/2}$ and $e_{k,\ell}^\pm$) is denoted by $u_{k+1/2,\ell}(x_1)$ (resp. $u_{k,\ell+1/2}(x_2)$ and $u_{k,\ell}^\pm(x_3)$).

The definition of $\mathcal{A}^\mu(\beta)$ also requires the introduction of the function spaces $L_2^\mu(\mathcal{G})$ and $H^2(\mathcal{G})$ defined as

$$L_2^\mu(\mathcal{G}) = \left\{ u : \|u\|_{L_2^\mu(\mathcal{G})} < +\infty \right\}, \quad H^2(\mathcal{G}) = \left\{ u \in C(\mathcal{G}) : \|u\|_{H^2(\mathcal{G})} < +\infty \right\}, \quad (5)$$

where,

$$\|u\|_{L_2^\mu(\mathcal{G})}^2 = \sum_{(k,\ell) \in \mathbb{Z}^2} \left(w_{k,\ell}^\mu \sum_{\pm} \|u_{k,\ell}^\pm\|_{L_2(e_{k,\ell}^\pm)}^2 + \|u_{k+\frac{1}{2},\ell}\|_{L_2(e_{k+\frac{1}{2},\ell})}^2 + \|u_{k,\ell+\frac{1}{2}}\|_{L_2(e_{k,\ell+\frac{1}{2}})}^2 \right), \quad (6)$$

$$\|u\|_{H^2(\mathcal{G})}^2 = \sum_{(k,\ell) \in \mathbb{Z}^2} \left(\sum_{\pm} \|u_{k,\ell}^\pm\|_{H^2(e_{k,\ell}^\pm)}^2 + \|u_{k+\frac{1}{2},\ell}\|_{H^2(e_{k+\frac{1}{2},\ell})}^2 + \|u_{k,\ell+\frac{1}{2}}\|_{H^2(e_{k,\ell+\frac{1}{2}})}^2 \right), \quad (7)$$

and, for any $(k, \ell) \in \mathbb{Z}^2$, $w_{k,\ell}^\mu$ is the weight coefficient equal to μ for $k = \ell = 0$ and 1 otherwise.

The unbounded limit operator in $L_2^\mu(\mathcal{G})$ has domain

$$D(\mathcal{A}^\mu(\beta)) = \left\{ u \in H^2(\mathcal{G}) : \forall (k, \ell) \in \mathbb{Z}^2, \quad \mathbf{u}_{k,\ell}^+ = e^{-i\beta} \mathbf{u}_{k,\ell}^-, \quad (u_{k,\ell}^+)'(a_3/2) = e^{-i\beta} (u_{k,\ell}^-)'(-a_3/2), \right. \\ \left. u'_{k+\frac{1}{2},\ell}(ka_1) - u'_{k-\frac{1}{2},\ell}(ka_1) + u'_{k,\ell+\frac{1}{2}}(\ell a_2) - u'_{k,\ell-\frac{1}{2}}(\ell a_2) + w_{k,\ell}^\mu \left((u_{k,\ell}^+)'(0) - (u_{k,\ell}^-)'(0) \right) = 0 \right\}, \quad (8)$$

and is defined by

$$\forall u \in D(\mathcal{A}^\mu(\beta)), \quad \mathcal{A}^\mu(\beta)u = -u'' \quad \text{on any edge of the graph } \mathcal{G}. \quad (9)$$

The functions of $D(\mathcal{A}^\mu(\beta))$ are continuous on \mathcal{G} and β quasi-periodic. Moreover, they satisfy the Kirchhoff conditions (8) that enforce the weighted sum of the outward derivatives of u to vanish at each vertex $v_{k,\ell}$ ($(k, \ell) \in \mathbb{Z}^2$). We point out that the perturbation, which results from a geometrical modification of the domain for the problem (4), is taken into account at the limit by means of the Kirchhoff condition (8) at the vertex $v_{0,0}$ ($w_{0,0}^\mu = \mu$). The formal derivation of the limit model can be found in [6]. It is easily verified that the operator $\mathcal{A}^\mu(\beta)$ is self-adjoint (for the weighted scalar product associated with (6)), see also [3]. The objective of the following two sections is to study the spectrum of $\mathcal{A}^\mu(\beta)$.

2.2 Characterization and properties of the essential spectrum of $\mathcal{A}^\mu(\beta)$

By a compact perturbation argument, one can prove that $\sigma_{ess}(\mathcal{A}^\mu(\beta)) = \sigma(\mathcal{A}(\beta))$, where $\mathcal{A}(\beta) = \mathcal{A}^1(\beta)$ is the purely periodic operator corresponding to $\mathcal{A}^\mu(\beta)$ for $\mu = 1$. The computation of its spectrum relies on the Floquet-Bloch theory (see [9]). More precisely, we can prove that $\lambda = \omega^2 \in \sigma(\mathcal{A}(\beta))$ if and only if either $\omega = 0$ and $\beta = 0$ or $\omega \neq 0$ and there exists $(k_1, k_2) \in [0, \pi]^2$ such that

$$\sin(\omega a_2) \sin(\omega a_3) (\cos(\omega a_1) - \cos k_1) + \sin(\omega a_3) \sin(\omega a_1) (\cos(\omega a_2) - \cos k_2) \\ + \sin(\omega a_1) \sin(\omega a_2) (\cos(\omega a_3) - \cos \beta) = 0. \quad (10)$$

Based on the previous characterization, we prove that the operator $\mathcal{A}(\beta)$ has a countable infinity of gaps that can be separated into three categories (see [10] for the proof):

Lemma 2.1 *The following properties hold :*

1- $\sigma_1 \cup \sigma_2 \cup \sigma_3 \subset \sigma(\mathcal{A}(\beta))$, where

$$\sigma_i = \left\{ (\pi n/a_i)^2, n \in \mathbb{Z} \right\} \text{ for } i \in \{1, 2\}, \text{ and } \sigma_3 = \left\{ ((\pm\beta + 2\pi n)/a_3)^2, n \in \mathbb{Z} \right\}.$$

2- For any $\beta \in [0, \pi]$, the operator $\mathcal{A}(\beta)$ has infinitely many gaps whose ends tend to infinity.

3- Let $\mathcal{W}(\beta) = \{\pi n/a_3, n \in \mathbb{N}^*\}$ if $\beta \notin \{0, \pi\}$ and $\mathcal{W}(\beta) = \{\beta/a_3 + (2n+1)\pi/a_3, n \in \mathbb{N}^*\}$ if $\beta \in \{0, \pi\}$,

If an interval (ω_b^2, ω_t^2) is a spectral gap of $\mathcal{A}(\beta)$, then, one of the following possibilities holds:

- (i) $\omega_b^2 \notin \sigma_1 \cup \sigma_2$, $\omega_t^2 \notin \sigma_1 \cup \sigma_2$, and there is a unique $\omega_0 \in (\omega_b, \omega_t) \cap \mathcal{W}(\beta)$.
- (ii) $\omega_b^2 \in \sigma_1 \cup \sigma_2$, $\omega_t^2 \notin \sigma_1 \cup \sigma_2$ and $\mathcal{W}(\beta) \cap (\omega_b, \omega_t) = \emptyset$.
- (iii) $\omega_b^2 \notin \sigma_1 \cup \sigma_2$, $\omega_t^2 \in \sigma_1 \cup \sigma_2$ and $\mathcal{W}(\beta) \cap (\omega_b, \omega_t) = \emptyset$.

2.3 Computation of the discrete spectrum

Let us now determine the discrete spectrum of $\mathcal{A}^\mu(\beta)$. If $\lambda = \omega^2$ is an eigenvalue of $\mathcal{A}^\mu(\beta)$, then the corresponding eigenfunction $u \in D(\mathcal{A}^\mu(\beta))$ satisfies the linear differential equation $u'' + \omega^2 u = 0$ on each edge of the graph \mathcal{G} . Solving explicitly this equation (on each edge), taking into account the quasi-periodicity of u and the Kirchhoff conditions (8), we can replace the eigenvalue problem $\mathcal{A}^\mu(\beta)u = \lambda u$ with a set of finite differences equations for $(\mathbf{u}_{\ell,k})_{(\ell,k) \in \mathbb{Z}^2}$:

Lemma 2.2 *Assume that $\omega \notin \{\pi\mathbb{Z}/a_1\} \cup \{\pi\mathbb{Z}/a_2\} \cup \{\pi\mathbb{Z}/a_3\}$. $u \in D(\mathcal{A}^\mu(\beta))$ is an eigenfunction of $\mathcal{A}^\mu(\beta)$ if and only if $(\mathbf{u}_{k,\ell})_{(k,\ell) \in \mathbb{Z}^2}$ belongs to $\ell_2(\mathbb{Z}^2)$ and satisfies*

$$\begin{cases} \frac{\mathbf{u}_{k+1,\ell}}{\sin(\omega a_1)} + \frac{\mathbf{u}_{k-1,\ell}}{\sin(\omega a_1)} + \frac{\mathbf{u}_{k,\ell+1}}{\sin(\omega a_2)} + \frac{\mathbf{u}_{k,\ell-1}}{\sin(\omega a_2)} - 2g_\beta(\omega)\mathbf{u}_{k,\ell} = 0, & \forall (k,\ell) \in \mathbb{Z}^2 \setminus \{(0,0)\}, \\ \frac{\mathbf{u}_{1,0}}{\sin(\omega a_1)} + \frac{\mathbf{u}_{-1,0}}{\sin(\omega a_1)} + \frac{\mathbf{u}_{0,1}}{\sin(\omega a_2)} + \frac{\mathbf{u}_{0,-1}}{\sin(\omega a_2)} - 2g_\beta(\omega)\mathbf{u}_{0,0} = 2(\mu-1)\frac{\cos(\omega a_3) - \cos\beta}{\sin(\omega a_3)}\mathbf{u}_{0,0}, \end{cases} \quad (11)$$

where we have defined $g_\beta(\omega) = \frac{1}{\tan(\omega a_1)} + \frac{1}{\tan(\omega a_2)} + \frac{\cos(\omega a_3) - \cos\beta}{\sin(\omega a_3)}$.

As well-known, finite difference schemes may be investigated using the discrete Fourier transform

$$\mathcal{F} : \mathbf{v} = (\mathbf{v}_{k,\ell})_{(k,\ell) \in \mathbb{Z}^2} \mapsto \mathcal{F}(\mathbf{v}) = \widehat{\mathbf{v}}, \quad \widehat{\mathbf{v}}(\xi, \eta) = \sum_{k,\ell \in \mathbb{Z}} e^{i(k\xi + \ell\eta)} \mathbf{v}_{k,\ell}, \quad (\xi, \eta) \in [0, 2\pi]^2, \quad (12)$$

where \mathcal{F} is an isometry between $\ell_2(\mathbb{Z}^2)$ and $L_2((0, 2\pi)^2)$. This, together with Lemma 2.2, provides the following characterization for the discrete spectrum of $\mathcal{A}^\mu(\beta)$:

Lemma 2.3 *Assume that $\omega \notin \{\pi\mathbb{Z}/a_1\} \cup \{\pi\mathbb{Z}/a_2\} \cup \{\pi\mathbb{Z}/a_3\}$. $u \in D(\mathcal{A}^\mu(\beta))$ is an eigenfunction of $\mathcal{A}^\mu(\beta)$ if and only if the discrete Fourier transform $\widehat{\mathbf{u}}$ of $(\mathbf{u}_{\ell,k})_{(\ell,k) \in \mathbb{Z}^2}$ belongs to $L_2((0, 2\pi)^2)$ and satisfies*

$$(f(\xi, \eta, \omega) - \phi_\beta(\omega)) \widehat{\mathbf{u}}(\xi, \eta) = (\mu-1)\phi_\beta(\omega)\mathbf{u}_{0,0}. \quad (13)$$

where $\phi_\beta(\omega) = \frac{\cos(\omega a_3) - \cos\beta}{\sin(\omega a_3)}$ and $f(\xi, \eta, \omega) = \frac{\cos\xi - \cos(\omega a_1)}{\sin(\omega a_1)} + \frac{\cos\eta - \cos(\omega a_2)}{\sin(\omega a_2)}$.

Under the assumption of Lemma 2.3, (10) can be written as $f(\xi, \eta, \omega) - \phi_\beta(\omega) = 0$. It follows that $\lambda = \omega^2$ does not belong to $\sigma_{ess}(\mathcal{A}^\mu(\beta))$ if and only if, for any $(\xi, \eta) \in [0, 2\pi]^2$, $\phi_\beta(\omega) - f(\xi, \eta, \omega)$ does not vanish. As a consequence, as soon as $\lambda = \omega^2 \notin \sigma_{ess}(\mathcal{A}^\mu(\beta))$, the function $(\xi, \eta) \mapsto \phi_\beta(\omega)/(\phi_\beta(\omega) - f(\xi, \eta, \omega))$ is continuous and bounded. Then, the inverse discrete Fourier transform can be applied to (13) to obtain

$$\mathbf{u}_{k,\ell} = \frac{(1-\mu)\mathbf{u}_{0,0}}{4\pi^2} \int_{(0,2\pi)^2} \frac{\phi_\beta(\omega)}{\phi_\beta(\omega) - f(\xi, \eta, \omega)} e^{-i(k\xi + \ell\eta)} d\xi d\eta, \quad \forall (k, \ell) \in \mathbb{Z}^2.$$

Writing the previous relation for $k = \ell = 0$ yields the following criterion of existence of an eigenvalue:

Lemma 2.4 Assume that $\omega \notin \{\pi\mathbb{Z}/a_1\} \cup \{\pi\mathbb{Z}/a_2\} \cup \{\pi\mathbb{Z}/a_3\}$ and that $\lambda = \omega^2 \notin \sigma_{\text{ess}}(\mathcal{A}^\mu(\beta))$. Then, λ is an eigenvalue of $\mathcal{A}^\mu(\beta)$ if and only if

$$\mu = 1 - F_\beta(\omega) \quad \text{where} \quad F_\beta(\omega) = \left(\frac{1}{4\pi^2} \int_{(0,2\pi)^2} \frac{\phi_\beta(\omega)}{\phi_\beta(\omega) - f(\xi, \eta, \omega)} d\xi d\eta \right)^{-1}. \quad (14)$$

The study of the behavior of the function F_β leads to the existence of at least one eigenvalue in each gap of $\mathcal{A}^\mu(\beta)$ as soon as $\mu < 1$, the minimal number of eigenvalues in each gap depending on the type of gaps (cf. Lemma. 2.1-3 for the classification):

Lemma 2.5 For $\mu > 1$, the operator $\mathcal{A}^\mu(\beta)$ has no eigenvalue. For $0 < \mu < 1$, let (ω_b^2, ω_t^2) be a spectral gap of the operator $\mathcal{A}^\mu(\beta)$:

- (a) If (ω_b^2, ω_t^2) is a gap of type (i), then $\mathcal{A}^\mu(\beta)$ has at least two eigenvalues $\lambda_1 = \omega_1^2$ and $\lambda_2 = \omega_2^2$ that satisfy $\omega_b < \omega_1 < \omega_0 < \omega_2 < \omega_t$ (see Lemma. 2.1-3 for the definition of ω_0).
- (b) If (ω_b^2, ω_t^2) is a gap of type (ii) or (iii), then $\mathcal{A}^\mu(\beta)$ has at least one eigenvalue $\lambda_1 = \omega_1^2$ such that $\omega_b < \omega_1 < \omega_t$.

The sketch of the proof of the previous lemma is the following, a complete proof being available in [10] (Theorem 5.2.1): First, one can verify that $F_\beta(\omega) \geq 0$ in any gap, which, together with (14) proves that $\mathcal{A}^\mu(\beta)$ has no eigenvalue for $\mu > 1$. Then, if (ω_b^2, ω_t^2) is a gap of type (i), one can show that

$$\lim_{\omega \rightarrow \omega_b^+} (1 - F_\beta(\omega)) = \lim_{\omega \rightarrow \omega_t^-} (1 - F_\beta(\omega)) = 1 \quad \text{and that} \quad \lim_{\omega \rightarrow \omega_0} (1 - F_\beta(\omega)) = 0.$$

By continuity of F_β inside the gap, (a) directly results from the intermediate value theorem and (14). If (ω_b^2, ω_t^2) is a gap of type (ii), the intermediate value theorem also permits us to conclude since

$$\lim_{\omega \rightarrow \omega_b} (1 - F_\beta(\omega)) \leq 0 \quad \text{and} \quad \lim_{\omega \rightarrow \omega_t^-} (1 - F_\beta(\omega)) = 1.$$

A similar argument works for a gap of type (iii).

3 Guided modes for the operator $A_\varepsilon^\mu(\beta)$: an asymptotic result

Finally, thanks to the general result [8] (Theorem 2.13 convergence of the spectrum of $A_\varepsilon^\mu(\beta)$ toward the spectrum of $\mathcal{A}^\mu(\beta)$), we can prove the following result of existence of eigenvalue for the operator $A_\varepsilon^\mu(\beta)$:

Theorem 3.1 Let $\mu \in (0, 1)$, (λ_b, λ_t) be a spectral gap of the operator $\mathcal{A}^\mu(\beta)$ and $\lambda_0 \in (\lambda_b, \lambda_t)$ be a (simple) eigenvalue of this operator. Then, there exists $\varepsilon_0 > 0$ such that if $\varepsilon < \varepsilon_0$ the operator $A_\varepsilon^\mu(\beta)$ has an eigenvalue λ_ε inside a spectral gap $(\lambda_b^\varepsilon, \lambda_t^\varepsilon)$. Moreover, $\lambda_\varepsilon = \lambda_0 + O(\sqrt{\varepsilon})$.

References

- [1] Ammari, H. and Santosa, F., Guided waves in a photonic bandgap structure with a line defect, SIAM Journal on Applied Mathematics, Vol. 64, Num. 6, p 2018-2033, 2004
- [2] Fliss, Sonia, A Dirichlet-to-Neumann approach for the exact computation of guided modes in photonic crystal waveguides, SIAM Journal on Scientific Computing, Vol. 35, Num. 2, p B438–B461, 2013
- [3] Kuchment P., Quantum graphs: an introduction and a brief survey, Analysis on graphs and its applications, Ser. Proc. Sympos. Pure Math., Vol. 77, p. 291–312, Amer. Math. Soc., Providence, RI, 2008
- [4] Friedlander, L., On the spectrum of a class of second order periodic elliptic differential operators, Communications in Mathematical Physics, Vol. 229, p. 49–55, 2002

- [5] Kuchment, P. and Ong, B. S. , On guided waves in photonic crystal waveguides, *Waves in periodic and random media*, Vol. 339, p. 105–115, 2003
- [6] Kuchment, P. and Zeng, H., Convergence of spectra of mesoscopic systems collapsing onto a graph, *Journal of Mathematical Analysis and Applications*, Vol. 258, Num. 2, p. 671-700, 2001
- [7] Nazarov, S.A., On the spectrum of the Laplace operator on the infinite Dirichlet ladder, *St. Petersburg Mathematical Journal*, Vol. 23, p. 1023–1045, 2012
- [8] Post O. , Spectral convergence of quasi-one-dimensional spaces, *Ann. Henri Poincaré*, Vol. 7, Num. 5, p. 933-977, 2006
- [9] Reed, M. and Simon, B., *Methods of modern mathematical physics v. I-IV*, Academic Press,1972-1978
- [10] Vasilevskaya, E., *Open periodic waveguides. Theory and application*, PhD thesis, 2016