



HAL
open science

Time inhomogeneous mutation models with birth-date dependence

Adrien Mazoyer

► **To cite this version:**

Adrien Mazoyer. Time inhomogeneous mutation models with birth-date dependence. 2016. hal-01415995v1

HAL Id: hal-01415995

<https://hal.science/hal-01415995v1>

Preprint submitted on 13 Dec 2016 (v1), last revised 23 Oct 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Time inhomogeneous mutation models with birth-date dependence

Adrien Mazoyer

November 9, 2016

Abstract

The classic Luria-Delbrück model for fluctuation analysis is extended to the case where the split instant distributions of cells are not i.i.d.: the lifetime of each cell is assumed to depend on its birth date. This model takes also into account cell deaths and non exponentially distributed lifetimes. In particular, it is possible to consider subprobability distributions, and model non exponential growth. It leads to a family of probability distributions which depend on the expected number of mutations, the death probability of mutant cells, and the split instant distributions of normal and mutant cells. This is deduced from the Bellman-Harris integral equation, written for the birth-date inhomogeneous case. A new theorem of convergence for the final mutant counts is proved, using an analytic method. Particular examples like the Haldane model, or the case where hazard functions of the split instant distributions are proportional are studied. The Luria-Delbrück distribution with cell deaths is recovered. A computation algorithm for the probabilities is provided.

1 Introduction

Mutation models are probabilistic descriptions of the growth of a population of cells, in which scarce mutations randomly occur. The first objective of these models is to study the distribution of the number of mutant cells at the end of the growth process. The classic mutation models can be interpreted as the result of the three following ingredients (Hamon and Ycart, 2012):

1. a random number of mutations occurring with small probability among a large number of cell divisions. Due to the law of small numbers, the number of mutations approximately follows a Poisson distribution. The expectation of that distribution is the product of the mutation probability by the total number of divisions;
2. from each mutation, a clone of mutant cells growing during a random time. Due to exponential growth, most mutations occur close to the end of the process, and the developing time of a random clone has exponential distribution. The rate of that distribution is the *relative fitness*, i.e. the ratio of the growth rate of normal cells to that of mutants;
3. the number of mutant cells that any clone developing for a given time will produce. The distribution of this number depends on the modelling assumptions, in particular the lifetimes of mutants.

One of the most used mutation models is the well known Luria-Delbrück model (Luria and Delbrück, 1943). Mathematical descriptions were introduced by Lea and Coulson (1949), followed by Armitage (1952) and Bartlett (1955). In that model, division times of mutant cells were supposed to be exponentially distributed. Thus a clone develops according to a Yule process (see Yule (1925, p. 35); Athreya and Ney (1972, p. 109)), and its size at any given time follows a geometric distribution. The distribution of final mutant counts is also explicit when lifetimes of mutant cells are supposed to be constant. This latter

model is called Haldane model by Sarkar (1991); an explicit form of the asymptotic distribution is given by Ycart (2013). General lifetimes have also been studied by Ycart (2013), but no explicit distribution is available apart from the exponential and constant lifetimes. Other extensions of the Luria-Delbrück model take into account the case where cells have a certain probability to die rather than divide (Angerer (2001, Sec. 3.1); Dewanji et al (2005); Komarova et al (2007); Ycart (2014)), where final number of cells are random (Angerer, 2001; Komarova et al, 2007; Ycart and Veziris, 2014), or where the cell divisions are asymmetric (Montgomery-Smith and Oveys, 2016).

In the mutation models cited above, the lifetimes of the cells are supposed to be i.i.d.. This is quite unrealistic. Indeed, during an experiment, a colony of cell grows in an environment which contains a finite amount of resources. Consider two instants s_1 and s_2 such that $s_1 \ll s_2$, then a cell born at time s_1 will complete its lifetime faster than a cell born at time s_2 . The Verhulst model (Verhulst, 1838) is one of the most known deterministic growth model which takes into account this limitation. Logistic-type stochastic models have been described by Allen (2010, Sec. 9.4.2), and mathematically studied by several authors among which Tan (1986); Tan and Piantadosi (1991); Lambert (2005). The independence of lifetimes for single type branching processes was questioned quite early (Kendall, 1952). Experiments have evidenced correlation between a cell and its descendants, and between two sisters conditioning on their mother (Wang et al, 2010). The effects of these correlations and many models have been discussed since then: see Louhichi and Ycart (2015) and references cited therein.

When the lifetimes are i.i.d., using the theory of branching processes (Bellman and Harris, 1952; Athreya and Ney, 1972), the distribution of the total number of mutant cells converges as the initial number of cells tends to infinity to a heavy-tailed distribution. This convergence has also been proved using an analytic way by Bartlett (1955, Sec. 4.31), but for a restrictive case where the fitness is set to 1. (equal growth rates for normal and mutant cells). Stewart et al (1990) proposed an approach to take into account the decreasing rate of division as the cells run out of resources. However, no results for the non i.i.d. lifetimes case have been stated until now. In particular, there is no convergence result for the distribution of final mutant counts.

The main objective of this paper is to extend classic mutation models to the case where the split instant of a cell depends on its birth date. Cells deaths and non exponential lifetimes are also taken into account with this approach. General modelling assumptions are described in section 2. The main tool used in this paper is an extension of the Bellman-Harris integral equation (Bellman and Harris, 1952), which is discussed in section 3. General solutions are also provided. They are used in the asymptotic context of mutation models in section 4. Some examples among which Haldane model and a more general case (non-exponential growth) are provided. The convergence results are finally applied in section 5 to the case where the hazard functions associated to the split instant distribution of normal and mutant cells are proportional. In particular the Luria-Delbrück distribution with cell deaths, denoted here by *LDD* distribution (Ycart, 2014), is recovered. A computation algorithm is proposed in section 6.

2 Hypotheses and models

In this section, the probabilistic model is defined as a tree-indexed process (see Pemantle (1995); Benjamini and Peres (1994) for general references). Denote by \mathbb{T} the infinite complete binary tree and 0 its root. The vertices of \mathbb{T} are interpreted as cells, and each vertex has exactly 2 descendants. If v is a vertex of \mathbb{T} , the number of edges between the root 0 and v is denoted by $|v|$. The binary tree with root v is denoted by \mathbb{T}_v . If v and u are two vertices of \mathbb{T} , $u \preceq v$ is the order relation that holds if u is in the path from 0 to v ; $u \prec v$ holds if u is strictly in the path from 0 to v ; $u \wedge v$ is the most recent common ancestor of u and v . Observe that if $u \preceq v$, then $u \wedge v = u$. Moreover, if v and w are two vertices of \mathbb{T}_u , then $v \wedge w = u$. The mother of the cell v is denoted by \tilde{v} : it is the cell such that $\tilde{v} \preceq v$ and $|\tilde{v}| = |v| - 1$. The length of the edge connecting a cell v to its mother \tilde{v} is interpreted as the lifetime of the cell \tilde{v} . Each cell produces two cells upon completion of its lifetime. Then a cell v_0 which is not the root 0 has a sister

cell v_1 . A vertex v_0 and its sister v_1 satisfy $\tilde{v}_0 = \tilde{v}_1 = v_0 \wedge v_1$.

The evolution of a clone stemming from a single cell at instant 0 will be modeled by a stochastic process $(C_v)_{v \in \mathbb{T}}$ indexed by the binary tree \mathbb{T} . For any $v \in \mathbb{T}$, C_v is a triplet (B_v, D_v, T_v) where B_v describes the nature (mutant or normal) of v , D_v whether v dies or divides, and T_v the instant at which this occurs.

- B_v is a Boolean such that $B_v = 1$ if the cell v is a mutant;
- D_v is a Boolean such that $D_v = 1$ if, upon completion of its lifetime, the cell v dies instead of producing 2 descendants;
- T_v is the instant at which the cell v completes its lifetime, i.e. at which v
 - either dies if $D_v = 1$ (then T_v is called the *death instant* of v);
 - or produces two cells if $D_v = 0$ (then T_v is called the *split instant* of v).

The instant T_v will be called *final instant* if the variable D_v is not precised.

Denote by $\overline{\mathbb{R}}_+ = \mathbb{R}_+ \cup \{+\infty\}$ the extended real line, and by $\mathcal{B}(\overline{\mathbb{R}}_+)$ its Borel σ -field. From the above settings, the stochastic process $(C_v)_{v \in \mathbb{T}}$ is defined on the measurable space (Ω, \mathcal{A}) , where $\Omega = \{0, 1\} \times \{0, 1\} \times \overline{\mathbb{R}}_+$ and its σ -algebra $\mathcal{A} = \mathcal{P}(\{0, 1\}) \times \mathcal{P}(\{0, 1\}) \times \mathcal{B}(\overline{\mathbb{R}}_+)$. The fact that T_v can be infinite will be discussed below. There remains to define a probability distribution on that space. Recall that the birth date of the root is set to 0. Assume that its nature B_0 is known. Note that for any cell $v \neq 0$, the definition of C_v makes sense only if $D_{\tilde{v}} = 0$. Thereafter, if $v \neq 0$, the distributions of the variables B_v , D_v , and T_v are defined conditionally to $D_{\tilde{v}} = 0$. Let π , γ , δ be reals in $(0; 1)$, respectively interpreted as probability of mutation, of dying for a normal cell, of dying for a mutant cell.

Consider a cell $v_0 \neq 0$ and its sister v_1 . Their nature B_{v_0} and B_{v_1} depend only on the nature $B_{\tilde{v}_0}$ of the mother cell:

- if $B_{\tilde{v}_0} = 0$, then:
 - $B_{v_0} = 1$ and $B_{v_1} = 0$ with probability $\pi/2$;
 - $B_{v_0} = 0$ and $B_{v_1} = 1$ with probability $\pi/2$;
 - $B_{v_0} = B_{v_1} = 0$ with probability $1 - \pi$;
- if $B_{\tilde{v}_0} = 1$, then $B_{v_0} = B_{v_1} = 1$ with probability 1.

For any cell v , D_v depends on the nature B_v : if $B_v = 0$, then $D_v = 1$ with probability γ ; if $B_v = 1$, then $D_v = 1$ with probability δ . In other words, upon completion of its lifetime, any normal cell produces one normal and one mutant cell with probability π (this event is called a *mutation*), dies with probability γ or produces two normal cells with probability $1 - \pi - \gamma$. Upon completion of its lifetime, any mutant cell dies with probability δ or produces two mutant cells with probability $1 - \delta$. Moreover, the events of death or mutation do not depend on the final instant of the cell.

For any cell v , its final instant T_v depends on its nature B_v and its birth date, i.e. on the split instant of its mother $T_{\tilde{v}}$: if $B_v = 0$ and $T_{\tilde{v}} = s$, the cumulative distribution function (cdf) of T_v is denoted by $F_\nu(s, \cdot)$; if $B_v = 1$ and $T_{\tilde{v}} = s$, the cdf of T_v is denoted by $F_\mu(s, \cdot)$. These cdfs satisfy $F_\nu(s, t) = 0$ and $F_\mu(s, t) = 0$ for $t \leq s$. Moreover, $F_\nu(s, t)$ and $F_\mu(s, t)$ may be strictly smaller than 1 for any t in $\overline{\mathbb{R}}_+$. This assumption is justified by the practical limitation of the total number of cells by a maximum size. This bound corresponds to the carrying capacity, i.e. the maximum sustainable population: the closer to this bound the number of cells, the slower the growth of the population. In other words, some cells do not produce descendants before the end of the growth process, which is mathematically interpreted as an infinite final instant. This hypothesis requires the notion of subprobability measure on a measurable space (Ω, \mathcal{A}) , i.e. a measure η^* on (Ω, \mathcal{A}) such that $\eta^*(\Omega) \leq 1$. For more details about subprobability

measures, see for example (Nguyen, 2006, p. 170). Consider a subprobability measure η^* on $(\mathbb{R}_+, \mathcal{B}(\mathbb{R}_+))$ such that $\eta^*(\mathbb{R}_+) < 1$. Then its cdf F^* has a limit less than 1 as x tends to infinity. From the measure η^* , the following probability measure η can be defined for any $A \in \mathcal{B}(\overline{\mathbb{R}}_+)$:

$$\eta(A) = \eta^*(A) + (1 - \eta^*(\mathbb{R}_+)) \mathbb{1}_{A \in \mathcal{B}(\overline{\mathbb{R}}_+) \setminus \mathcal{B}(\mathbb{R}_+)}$$

with the associated cdf:

$$F(x) = F^*(x) \mathbb{1}_{x \in [0; +\infty)} + \mathbb{1}_{x = +\infty}. \quad (2.1)$$

The stochastic process $(C_v)_{v \in \mathbb{T}}$ defined above models the evolution of a clone stemming from a single cell at instant 0. Actually, the evolution of a clone stemming from a single cell u at instant $s > 0$ can be deduced from the above process. It consists of a stochastic process $(C_v)_{v \in \mathbb{T}_u}$ indexed by the binary tree \mathbb{T}_u with the same modelling assumptions as above, conditionally to C_u .

There remains to define the dependences between the cells. Consider a cell $v_0 \neq 0$ and its sister v_1 . The final instants T_{v_0} and T_{v_1} are assumed to be independent conditionally to $C_{\tilde{v}_0}$. By extension, the clones $(C_u)_{u \in \mathbb{T}_{v_0}}$ and $(C_u)_{u \in \mathbb{T}_{v_1}}$ are independent conditionally to $C_{\tilde{v}_0}$. Consider now two cells $v \neq 0$ and $w \neq 0$ and their common ancestor $v \wedge w$. Assume that their common ancestor is neither v nor w . Then only one of the daughter cells of $v \wedge w$ is in the path from 0 to v , and its sister is in the path from 0 to w . Thus, according to the previous dependence assumption, the final instants T_v and T_w are independent conditionally to $C_{v \wedge w}$. Therefore, the clones $(C_u)_{u \in \mathbb{T}_v}$ and $(C_u)_{u \in \mathbb{T}_w}$ are independent conditionally to $C_{v \wedge w}$.

Thereafter, the root 0 is assumed to be a normal cell, i.e. $B_0 = 0$. The modelling assumptions can be summarized as follows:

- at time 0, a single normal cell is present;
- the final instant of any cell depends on its nature and its birth date;
- the final instant of a normal cell born at time s is a random variable with cdf $F_\nu(s, \cdot)$ defined on $\overline{\mathbb{R}}_+$;
- upon completion of the lifetime of a normal cell:
 - with probability π one normal and one mutant cells are produced;
 - with probability γ the cell dies out;
 - with probability $1 - \gamma - \pi$ two normal cells are produced;
- the final instant of a mutant cell born at time s is a random variable with cdf $F_\mu(s, \cdot)$ defined on $\overline{\mathbb{R}}_+$;
- upon completion of the lifetime of a mutant cell:
 - with probability δ the cell dies out;
 - with probability $1 - \delta$ two mutant cells are produced;
- for any cell, the events of death or mutation do not depend on its final instant;
- two cells, whatever their nature, are independent conditionally on their common ancestor;
- two clones are independent conditionally on the common ancestor of the two cells which started those clones.

All the results will be given in terms of the bivariate probability generating functions (pgf) of the numbers of normal and mutant cells, and the pgf of the number of mutant cells. The next section is dedicated to the calculation of these functions.

3 Integral equations for probability generating functions

Denote by $\mathcal{N}(s, t)$ the couple of numbers at time t of normal and mutant cells in the clone stemming from a single normal cell born at time s . Its bivariate pgf is defined by:

$$\varphi(y, z, s, t) = \sum_{n, m \geq 0} y^n z^m \mathbb{P}[\mathcal{N}(s, t) = (n, m)].$$

Note that $\varphi(1, z, s, t)$ is the pgf of the number of mutant cells in the clone stemming from a normal cell born at time s . Denote by $M(s, t)$ the number at time t of mutant cells in the clone stemming from a single mutant cell born at time s . Its pgf is defined by:

$$\psi(z, s, t) = \sum_{m \geq 0} z^m \mathbb{P}[M(s, t) = m].$$

One way to study these pgfs is to apply the well known Bellman-Harris integral equation (Bellman and Harris, 1952). However, this equation has been justified only for the case where the lifetimes are i.i.d.. This section extends it to the case where the final instant of a cell depends on its birth date. These equations have already been stated for the case of i.i.d. lifetimes (see for example Kimmel and Axelrod (2002, Chap. 5)). The case of non i.i.d. lifetimes is studied here using similar arguments. The integral equation will be stated first for the bivariate pgf φ . The equation can be written as ordinary differential equation and solved under specific conditions. These results are then applied to obtain results about the distribution of mutant counts at a given time t (Proposition 3.3 and Corollary 3.1).

For any normal cell, the bivariate pgf for the number of progenies is given by:

$$\chi(y, z) = \gamma + \pi yz + (1 - \pi - \gamma)z^2,$$

Consider a normal cell born at time s . Assume this cell completes its lifetime at a given time $u > s$. Then, for times $t < u$ the mother cell is still alone in the corresponding clone. For times $t \geq u$, the number of cells is equal to the sum of cells in all the clones stemming from the first-generation progeny, i.e.:

$$\mathcal{N}(s, t) = \left(\sum_{i=1}^X \mathcal{N}^{(i)}(u, t) \right) \mathbf{1}_{t \geq u} + \mathbf{1}_{t < u},$$

where X is the number of cells in the first-generation progeny and $\mathcal{N}^{(i)}(u, t)$ are i.i.d. copies of the process $\mathcal{N}(u, t)$. In terms of pgf:

$$\varphi(y, z, s, t) = \chi[\varphi(y, z, u, t), \psi(z, u, t)] \mathbf{1}_{t \geq u} + y \mathbf{1}_{t < u}.$$

According to the modelling assumptions, integrating with respect to the distribution $F_\nu(s, \cdot)$ removes the conditioning on the final instant of the cell. Then the following integral equation is obtained:

$$\begin{aligned} \varphi(y, z, s, t) &= \int_s^t \gamma + \pi \varphi(y, z, u, t) \psi(z, u, t) + (1 - \pi - \gamma) \varphi(y, z, u, t)^2 dF_\nu(s, u) \\ &\quad + y(1 - F_\nu(s, t)). \end{aligned} \tag{BH}$$

Similarly as for the homogeneous case (Bellman and Harris, 1952), an intuitive interpretation of (BH) can be given. For a given time $t > s$, there are two possibilities: either the transformation of the particle has taken place after t with probability $1 - F_\nu(s, t)$. Then there is still one normal cell, and $\varphi(y, z, s, t) = y$ (second term in (BH)); or the transformation of the cell has taken place in $[u; u + du]$ (where $s < u \leq t$) with probability $dF_\nu(s, u)$. In that case, either it turns into 2 normal cells with probability $1 - \pi - \gamma$, or into one normal and one mutant with probability π , or dies with probability γ . The two new cells

start independent clones of their own, accounted for by $\varphi(y, z, u, t)^2$ if two normal cells are produced, and by $\varphi(y, z, u, t)\psi(z, u, t)$ if one normal and one mutant are produced.

Until now, there were no specific assumptions on F_ν and F_μ , except their definition domain and the fact that $F_\nu(s, t) = F_\mu(s, t) = 0$ if $t \leq s$. Thereafter, in order to solve (BH), some hypotheses on F_ν are precised. For any $s \geq 0$, let $F(s, \cdot)$ be a cdf on $\overline{\mathbb{R}}_+$ such that $F(s, t) = 0$ if $t \leq s$. The cdf F will satisfy (\mathcal{H}) if there exists a cdf of subprobability on \mathbb{R}_+ , denoted by $F^*(s, \cdot)$, such that the following holds:

(\mathcal{H}_1) the cdf F^* is differentiable with respect to s and t , and decreasing in s ;

(\mathcal{H}_2) $\lim_{t \rightarrow +\infty} F^*(s, t) < 1$ for all $(s, t) \in \mathbb{R}_+^2$ and $F^*(s, t) = 0$ if $t \leq s$;

(\mathcal{H}_3) for any $s \geq 0$, $F(s, \cdot)$ is deduced from $F^*(s, \cdot)$ as (2.1);

(\mathcal{H}_4) the function r defined for all $(s, t) \in \mathbb{R}_+^2$ by:

$$r(s, t) = -\log(1 - F^*(s, t)) ,$$

satisfies for any $t \geq s$:

$$r(s, t) = r(0, t) - r(0, s) .$$

Remark that r is by definition positive, differentiable with respect to s and t , increasing in t , decreasing in s and for any $(s, t) \in \mathbb{R}_+^2$:

$$r(s, t) \leq \lim_{t \rightarrow +\infty} r(0, t) .$$

For the moment, no assumptions on F_μ are required. However, some of the particular cases introduced here will assume that F_μ satisfies (\mathcal{H}) too. In that case, its related function defined in (\mathcal{H}_4) will be denoted by μ . From now on, assume that F_ν satisfies (\mathcal{H}) , and denote by ν its related function defined in (\mathcal{H}_4) . Thus there exists a positive, continuous, \mathbb{R}_+ -valued function λ_ν such that:

$$\nu(s, t) = \int_s^t \lambda_\nu(u) du$$

The function λ_ν can be interpreted as the hazard function associated to F_ν on \mathbb{R}_+ . The cdf $F_\nu(s, \cdot)$ is then defined on $\overline{\mathbb{R}}_+$ for any $s \in \mathbb{R}_+$ by:

$$F_\nu(s, t) = \begin{cases} \left(1 - \exp\left(-\int_s^t \lambda_\nu(u) du\right)\right) \mathbb{1}_{s \leq t} & \text{if } t < +\infty, \\ 1 & \text{if } t = +\infty. \end{cases}$$

Since $\nu(s, t) = \nu(0, t) - \nu(0, s)$, replacing F_ν by its expression in (BH) leads to:

$$\begin{aligned} \varphi(y, z, s, t) e^{-\nu(0, s)} &= \int_s^t [\gamma + \pi \varphi(y, z, u, t) \psi(z, u, t) + (1 - \pi - \gamma) \varphi(y, z, u, t)^2] \lambda_\nu(u) e^{-\nu(0, u)} du \\ &+ y e^{-\nu(0, t)} . \end{aligned}$$

Taking the derivative with respect to s and dividing by $e^{-\nu(0, s)}$ leads to the following Riccati equation:

$$\begin{aligned} \partial_s \varphi(y, z, s, t) &= -\lambda_\nu(s) [\gamma - (1 - \pi \psi(z, s, t)) \varphi(y, z, s, t) \\ &+ (1 - \pi - \gamma) \varphi(y, z, s, t)^2] , \end{aligned} \tag{R1}$$

with the condition $\varphi(y, z, t, t) = y$. Riccati equations may have explicit solutions depending on the coefficients (see for example Kucera (1973); Harko et al (2014)). One of the cases where (R1) can be

solved is when $\gamma = 0$. This case makes sense in the context of mutation models: the objective is to obtain an explicit pgf for the mutant counts only. The number of divisions occurring in dying clones remains bounded and can be neglected. Thus, it can be considered that observed mutants only come from divisions in surviving clones.

In order to simplify expressions, define the following function:

$$I(z, s, t) = \int_s^t \lambda_\nu(u) \psi(z, u, t) du. \quad (3.1)$$

If $\gamma = 0$, (R1) reduces to a Bernoulli equation of order 2. Then the change of variable $\tilde{\varphi} = 1/\varphi$ leads to the solution $\varphi(y, z, s, t)$.

Proposition 3.1. *Assume $\gamma = 0$. The general solution of the Riccati equation (R1) is given by:*

$$\varphi(y, z, s, t) = e^{\pi I(z, s, t) - \nu(s, t)} \left\{ \frac{1}{y} - (1 - \pi) \int_s^t \lambda_\nu(u) e^{\pi I(z, u, t) - \nu(u, t)} du \right\}^{-1}. \quad (3.2)$$

Another case where (R1) has an explicit solution is when $\gamma = \delta$ and F_μ satisfies (\mathcal{H}) such that $\mu(s, t) = \nu(s, t)$ for any (s, t) in \mathbb{R}_+^2 . In that case, ψ is a particular solution, and the general solution of (R1) is explicit (Harko et al, 2014). However, it corresponds to the case where mutant and normal cells have the same death probability and the same final instant distribution. Mutant and normal cells are then indistinguishable, which seems to be of less practical relevance. An explicit solution can also be obtained if $1 - \pi - \gamma = 0$. However, γ has to be less than 0.5 (supercritical process), and π is typically very small. Thus this case is not realistic at all, and will not be studied here. As it will be mentioned further on, one of the best compromise between relevance and generality of hypotheses is to assume that $\gamma = 0$ and F_μ satisfies (\mathcal{H}) such that $\nu(s, t)$ and $\mu(s, t)$ are proportional for any (s, t) in \mathbb{R}_+^2 .

As a direct consequence of Proposition 3.1, setting $y = 1$ and $s = 0$ in (3.2) leads to the following result.

Corollary 3.1. *Assume $\gamma = 0$. The mutant counts at time t starting with a single normal cell at time 0 follows the distribution with pgf:*

$$\phi(z, t) = e^{\pi I(z, 0, t) - \nu(0, t)} \left\{ 1 - (1 - \pi) \int_0^t \lambda_\nu(u) e^{\pi I(z, u, t) - \nu(u, t)} du \right\}^{-1}, \quad (3.3)$$

From now on, γ will be set at 0. Observe that no assumptions on the cdf F_μ are required. In particular, F_μ does not necessarily satisfy (\mathcal{H}) . As long as the pgf ψ is known, the distribution of the mutant counts at a given time is explicit. As an example, consider the Haldane model: the final instants of the normal cell are exponentially distributed and the lifetimes of the mutant cells are equal to a constant a . In this case, λ_ν is a positive constant and ν is given by:

$$\nu(s, t) = \lambda_\nu(t - s),$$

and the cdf $F_\mu(s, \cdot)$ is defined for any $t \geq s$ by:

$$F_\mu(s, t) = \begin{cases} 1 & \text{if } t \geq s + a, \\ 0 & \text{else.} \end{cases}$$

Then F_μ does not satisfy (\mathcal{H}) : property (\mathcal{H}_1) is not satisfied. However, the pgf ψ is easily identifiable. Assume that the lifetime of any mutant cells is equal to a . Considering a cell born at time s , let $b_i(z)$ be the pgf of the size of its clone in the interval $[s + ia; s + (i + 1)a)$. Then $b_0(z) = z$, and for all $i > 0$,

$$b_i(z) = \delta + (1 - \delta)(b_{i-1}(z))^2.$$

Therefore the pgf of the size at time t of a clone starting at time s is:

$$\psi(z, s, t) = \sum_{i \geq 0} b_i(z) \mathbb{1}_{t \in [s+ia; s+(i+1)a)}.$$

Functions $I(z, s, t)$ and $g(z, t)$ can be explicited. This example will be continued in the next section where the asymptotic model is considered.

Assume now that F_μ satisfies (\mathcal{H}) . There exists a function λ_μ , with the same properties as λ_ν , such that $F_\mu(s, \cdot)$ is given by:

$$F_\mu(s, t) = \begin{cases} \left(1 - \exp\left(-\int_s^t \lambda_\mu(u) du\right)\right) \mathbb{1}_{s \leq t} & \text{if } t < +\infty, \\ 1 & \text{if } t = +\infty. \end{cases}$$

The pgf of the number of progenies for any mutant cell is given by:

$$\xi(z) = \delta + (1 - \delta)z^2.$$

By the same reasoning as for φ , the following Riccati equation for ψ is obtained:

$$\partial_s \psi(z, s, t) = -\lambda_\mu(s) [\delta - \psi(z, s, t) + (1 - \delta)\psi(z, s, t)^2], \quad (\text{R2})$$

with the condition $\psi(z, t, t) = z$. The general solution of (R2) can be explicited without specific hypotheses:

Proposition 3.2. *The general solution of the Riccati equation (R2) is given by:*

$$\psi(z, s, t) = \frac{\delta(1 - z) + e^{-\mu^*(s,t)}((1 - \delta)z - \delta)}{(1 - \delta)(1 - z) + e^{-\mu^*(s,t)}((1 - \delta)z - \delta)}, \quad (\text{3.4})$$

where:

$$\mu^*(s, t) = (1 - 2\delta)\mu(s, t).$$

The proof of Proposition 3.2 simply consists in observing that 1 is a particular solution of (R2). Then the general solution (R2) is explicit (Harko et al, 2014) and given by (3.4).

Observe that if λ_μ is a constant, then $\mu(s, t) = \lambda_\mu(t - s)$ and Proposition 3.2 reduces to the example by Athreya and Ney (1972, p. 109). Moreover, if λ_μ is a constant and $\delta = 0$, (R2) reduces to a Bernoulli equation. Its solution is the pgf of the geometric distribution with parameter $e^{-\lambda_\mu(t-s)}$, i.e. the pgf of a Yule process with parameter λ_μ . (see [p. 35] Yule (1925) or Athreya and Ney (1972, p. 109)). As a direct consequence of Proposition 3.2, the distribution of the number of mutant cells in a mutant clone started at time s can be explicited.

Proposition 3.3. *Denote by $(p_k(s, t))_{k \in \mathbb{N}}$ the probabilities of the size at time t of a clone starting from a single mutant cell born at time s . Then:*

$$p_0(s, t) = \frac{\delta(1 - e^{-\mu^*(s,t)})}{1 - \delta - \delta e^{-\mu^*(s,t)}},$$

and for $k \geq 1$,

$$p_k(s, t) = (1 - p_0(s, t))P(s, t)(1 - P(s, t))^{k-1},$$

where:

$$P(s, t) = \frac{(1 - 2\delta)e^{-\mu^*(s,t)}}{1 - \delta - \delta e^{-\mu^*(s,t)}},$$

and:

$$\mu^*(s, t) = (1 - 2\delta)\mu(s, t).$$

In other words, a random variable with pgf $\psi(z, s, t)$ is the following random mixture: either 0 with probability $p_0(s, t)$, or a geometric random variable with parameter $P(s, t)$.

Proposition 3.3. Writing ψ as the following rational function:

$$\begin{aligned}\psi(z, s, t) &= \frac{\delta(1 - e^{-\mu^*(s,t)}) - z(\delta - e^{-\mu^*(s,t)}(1 - \delta))}{(1 - \delta - \delta e^{-\mu^*(s,t)}) - z(1 - \delta)(1 - e^{-\mu^*(s,t)})} \\ &= \frac{n_0(s, t) + zn_1(s, t)}{d_0(s, t) + zd_1(s, t)},\end{aligned}$$

the probabilities $p_k(s, t)$ can be recursively identified:

$$p_0(s, t) = \frac{n_0(s, t)}{d_0(s, t)} = \frac{\delta(1 - e^{-\mu^*(s,t)})}{1 - \delta - \delta e^{-\mu^*(s,t)}},$$

and

$$\begin{aligned}p_1(s, t) &= \frac{n_1(s, t)}{d_0(s, t)} - \frac{d_1(s, t)}{d_0(s, t)}p_0(s, t) \\ &= \frac{(1 - 2\delta)^2 e^{-\mu^*(s,t)}}{d_0(s, t)^2} \\ &= (1 - p_0(s, t)) \frac{(1 - 2\delta)e^{-\mu^*(s,t)}}{d_0(s, t)}.\end{aligned}$$

Denote by $P(s, t)$ the second term of the above product. Then for $k \geq 2$:

$$\begin{aligned}p_k(s, t) &= -\frac{d_1(s, t)}{d_0(s, t)}p_{k-1}(s, t) \\ &= \left(-\frac{d_1(s, t)}{d_0(s, t)}\right)^{k-1} (1 - p_0(s, t))P(s, t) \\ &= (1 - p_0(s, t))P(s, t)(1 - P(s, t))^{k-1}.\end{aligned}$$

□

Observe that if λ_μ is a constant, Proposition 3.3 reduces to formula (3.1) of Ycart (2014). Moreover, the expectation of the size at a finite time t of a mutant clone started at time s is $e^{\mu^*(s,t)}$. Assume there exists $\rho > 0$ such that for any $s \geq 0$:

$$\lambda_\nu(s) = \rho(1 - 2\delta)\lambda_\mu(s).$$

The constant ρ can be interpreted as the instantaneous ratio of hazard functions. The assumption of proportional hazard functions is realistic: in survival analysis, it is known as the Cox proportional-hazard regression model. In that case, (3.1) can be explicitated:

$$\begin{aligned}I(z, s, t) &= \frac{\rho(1 - 2\delta)}{1 - \delta} \log \left[\frac{(1 - 2\delta)e^{(1-\delta)\mu(s,t)}}{(1 - \delta)((1 - z)e^{\mu^*(s,t)} + z) - \delta} \right] \\ &= \nu(s, t) + \frac{\rho(1 - 2\delta)}{1 - \delta} \log \left[\frac{(1 - 2\delta)}{(1 - \delta)((1 - z)e^{\mu^*(s,t)} + z) - \delta} \right].\end{aligned}\tag{3.5}$$

In particular, let f be a continuous, non negative, and increasing function on \mathbb{R}_+ . Let μ be defined for (s, t) in \mathbb{R}_+^2 by:

$$\mu(s, t) = \log \left(\frac{f(t)}{f(s)} \right).$$

Then λ_μ is given by:

$$\lambda_\mu(s) = \frac{f'(s)}{f(s)},$$

and the expectation of the size at a finite time t of a mutant clone started at time s is $(f(t)/f(s))^{1-2\delta}$. In other words, it is possible to fit the average trajectory of the development of the clones to any appropriate function of time. Moreover, if $\delta = 0$, plugging (3.5) in (3.3) and applying the change of variable $w = f(u)/f(t)$ leads to:

$$\begin{aligned} \phi(z, t) &= \left(\frac{f(t)}{f(0)} \right)^{-\rho} \left(1 - z + z \frac{f(0)}{f(t)} \right)^{\pi\rho} \\ &\times \left\{ 1 - (1 - \pi) \int_{\frac{f(0)}{f(t)}}^1 \rho w^{\rho-1} (1 - z + zw)^{-\pi\rho} dw \right\}^{-1}. \end{aligned} \quad (3.6)$$

For example, if f is defined for any $t \geq 0$ by $f(t) = e^t$, (3.6) is given by:

$$\begin{aligned} \phi(z, t) &= e^{-\rho t} (1 - z + ze^{-t})^{-\pi\rho} \\ &\times \left\{ 1 - (1 - \pi) \int_0^t \rho e^{-\rho u} (1 - z + ze^{-u})^{-\pi\rho} du \right\}^{-1}, \end{aligned}$$

which is the inverse of formula (10) of Bartlett (1955, p. 116). Functions with a carrying capacity, such that logistic or Gompertz functions, can also be considered and plugged into (3.6).

Corollary 3.1 is used in the next section in a relevant asymptotic context to get the convergence in distribution of the mutant count when n normal cells are initially present.

4 Asymptotic for mutation models

In this section, the previous results are applied to mutation models. A convergence theorem for the final number of mutant cells is proved, generalizing the analytic method initiated by Bartlett (1955, Sec. 4.31). A mutation model consists of n independent copies of the model described in section 2. Denote by $\phi_n(z, t)$ the pgf for the mutant counts at time t starting with n normal cells at time 0. Because of the independence of the n initial cells, $\phi_n(z, t)$ is the n -th power of (3.3). The main objective of this section is to establish the convergence as n tends to infinity of $\phi_n(z, t)$. Before stating the result, define the function h as:

$$h(z, t) = \frac{1}{1 - e^{-\nu(0,t)}} \int_0^t \psi(z, u, t) \lambda_\nu(u) e^{-\nu(u,t)} du. \quad (4.1)$$

This is a notation in order to simplify expressions, however the function h could be intuitively interpreted as the pgf of the size of any mutant clone at a given time t . This interpretation requires to assume that the birth dates of the initial mutants, i.e. the mutation instants, follow the distribution $\lambda_\nu(u) e^{-\nu(u,t)}$ truncated to $[0; t]$, which is not proven here. The main result of this paper is the following convergence theorem:

Theorem 4.1. *Let $\pi = (\pi_n)_{n \in \mathbb{N}}$ and $t = (t_n)_{n \in \mathbb{N}}$ two sequences, and α a positive real such that:*

$$\lim_{n \rightarrow +\infty} \pi_n = 0, \quad \lim_{n \rightarrow +\infty} t_n = +\infty, \quad \lim_{n \rightarrow +\infty} \pi_n n e^{\nu(0,t_n)} = \alpha. \quad (H1)$$

Assume that:

$$\lim_{n \rightarrow +\infty} I(z, 0, t_n) e^{-\nu(0, t_n)} < +\infty. \quad (\text{H2})$$

As n tends to infinity, the pgf of the number of mutants at time t_n , starting with n normal cells at time 0, converges to the pgf

$$\phi(z) = \exp \{-m(1 - h(z))\}, \quad (4.2)$$

where

$$h(z) = \lim_{n \rightarrow +\infty} h(z, t_n).$$

and

$$m = \alpha(1 - e^{-\nu_\infty})$$

Observe that (4.2) is the pgf of a Poisson compound, with parameter m . From that expression, the parameter m could be seen as the mean number of mutations. This conjecture makes sense, assuming that the number of mutation occasions is at least equivalent to $n(e^{\nu(0, t_n)} - n)$ as n tends to infinity. The main tool required to prove Theorem 4.1 is Lemma 4.1 below.

Lemma 4.1. For any $\pi \in [0; 1[$, $z \in [0; 1]$, $t \in \mathbb{R}_+$, and $s \in [0; t]$, the following bound holds:

$$\left| e^{\pm \pi I(z, s, t)} - (1 \pm I(z, s, t)) \right| \leq \pi^2 e^{I(z, 0, t)}.$$

The proof uses a power series expansion of $e^{\pm \pi I(z, s, t)}$:

Lemma 4.1.

$$e^{\pm \pi I(z, s, t)} = \sum_{k \geq 0} \frac{(\pm \pi I(z, s, t))^k}{k!}.$$

Hence:

$$\begin{aligned} \left| e^{\pm \pi I(z, s, t)} - (1 \pm \pi I(z, s, t)) \right| &\leq \sum_{k \geq 2} |\pm \pi|^k \frac{I(z, s, t)^k}{k!} \\ &\leq \pi^2 e^{I(z, s, t)} \leq \pi^2 e^{I(z, 0, t)}. \end{aligned}$$

□

An analytic proof for the case where mutant and normal cells are exponentially i.i.d. with equal rates has been provided by Bartlett (1955, Sec. 4.31). This approach has been adapted to prove Theorem 4.1.

Theorem 4.1. Define the following two functions:

$$f_1(z, u, t_n, \pi_n) = e^{\pi_n I(z, u, t_n)} - (1 + \pi_n I(z, u, t_n)) \quad \text{and} \quad f_2(z, t_n, \pi_n) = f_1(z, 0, t_n, \pi_n).$$

According to Lemma 4.1:

$$|f_1(z, u, t_n, \pi_n)| \leq \pi_n^2 e^{I(z, 0, t_n)} \quad \text{and} \quad |f_2(z, t_n, \pi_n)| \leq \pi_n^2 e^{I(z, 0, t_n)}.$$

Then, the second factor in (3.3)

$$1 - (1 - \pi_n) \int_0^{t_n} \lambda_\nu(u) e^{\pi_n I(z, u, t_n)} e^{-\nu(u, t_n)} du,$$

can be written as:

$$\begin{aligned}
& 1 - (1 - \pi_n) \left[\int_0^{t_n} \lambda_\nu(u) e^{-\nu(u, t_n)} f_2(z, u, t_n, \pi_n) du + \left(1 - e^{-\nu(0, t_n)} \right) \right. \\
& \quad \left. + \pi_n \int_0^{t_n} I(z, u, t_n) \lambda_\nu(u) e^{-\nu(u, t_n)} du \right] \\
&= e^{-\nu(0, t_n)} - \int_0^{t_n} \lambda_\nu(u) e^{-\nu(u, t_n)} f_2(z, u, t_n, \pi_n) du \\
& \quad + \pi_n \left[1 - e^{-\nu(0, t_n)} - \int_0^{t_n} I(z, u, t_n) \lambda_\nu(u) e^{-\nu(u, t_n)} du + \int_0^{t_n} \lambda_\nu(u) e^{-\nu(u, t_n)} f_2(z, u, t_n, \pi_n) du \right] \\
& \quad + \pi_n^2 \int_0^{t_n} I(z, u, t_n) \lambda_\nu(u) e^{-\nu(u, t_n)} du.
\end{aligned}$$

Let:

$$\begin{aligned}
\mathcal{I}(z, t_n) &= \int_0^{t_n} I(z, u, t_n) \lambda_\nu(u) e^{-\nu(u, t_n)} du \\
&= \left(1 - e^{-\nu(0, t_n)} \right) h(z, t_n) - e^{-\nu(0, t_n)} I(z, 0, t_n),
\end{aligned}$$

and

$$\mathcal{J}(z, t_n, \pi_n) = \int_0^{t_n} \lambda_\nu(u) e^{-\nu(u, t_n)} f_1(z, u, t_n, \pi_n) du.$$

From (H2), the limit as n tends to infinity of $\mathcal{I}(z, t_n)$ exists and is finite. Consider now the following term:

$$e^{-\pi_n I(z, 0, t_n)} \left\{ 1 - (1 - \pi_n) \int_0^{t_n} \lambda_\nu(u) e^{\pi_n I(z, u, t_n)} e^{-\nu(u, t_n)} du \right\}.$$

It can be written as:

$$\begin{aligned}
& (f_2(z, t_n, \pi_n) + 1 - \pi_n I(z, 0, t_n)) \left\{ e^{-\nu(0, t_n)} - \mathcal{J}(z, t_n, \pi_n) \right. \\
& \quad \left. + \pi_n \left(1 - e^{-\nu(0, t_n)} - \mathcal{I}(z, t_n) + \mathcal{J}(z, t_n, \pi_n) \right) + \pi_n^2 \mathcal{I}(z, t_n) \right\} \\
&= (f_2(z, t_n, \pi_n) + 1) \left(e^{-\nu(0, t_n)} - \mathcal{J}(z, t_n, \pi_n) \right) \\
& \quad + \pi_n \left\{ (f_2(z, t_n, \pi_n) + 1) \left(1 - e^{-\nu(0, t_n)} - \mathcal{I}(z, t_n) + \mathcal{J}(z, t_n, \pi_n) \right) \right. \\
& \quad \quad \left. - I(z, 0, t_n) \left(e^{-\nu(0, t_n)} - \mathcal{J}(z, t_n, \pi_n) \right) \right\} \\
& \quad + \pi_n^2 \left\{ (f_2(z, t_n, \pi_n) + 1) \mathcal{I}(z, t_n) - I(z, 0, t_n) \left(1 - e^{-\nu(0, t_n)} - \mathcal{I}(z, t_n) + \mathcal{J}(z, t_n, \pi_n) \right) \right\} \\
& \quad - \pi_n^3 I(z, 0, t_n) \mathcal{I}(z, t_n).
\end{aligned}$$

Multiplying by $e^{\nu(0,t_n)}$:

$$\begin{aligned} \frac{1}{\phi(z, t_n)} &= (f_2(z, t_n, \pi_n) + 1) \left(1 - e^{\nu(0,t_n)} \mathcal{J}(z, t_n, \pi_n) \right) \\ &\quad + \pi_n \left\{ (f_2(z, t_n, \pi_n) + 1) \left(e^{\nu(0,t_n)} - 1 - e^{\nu(0,t_n)} \mathcal{I}(z, t_n) + e^{\nu(0,t_n)} \mathcal{J}(z, t_n, \pi_n) \right) \right. \\ &\quad \quad \left. - I(z, 0, t_n) \left(1 - e^{\nu(0,t_n)} \mathcal{J}(z, t_n, \pi_n) \right) \right\} \\ &\quad + \pi_n^2 \left\{ (f_2(z, t_n, \pi_n) + 1) e^{\nu(0,t_n)} \mathcal{I}(z, t_n) \right. \\ &\quad \quad \left. - I(z, 0, t_n) \left(e^{\nu(0,t_n)} - 1 - e^{\nu(0,t_n)} \mathcal{I}(z, t_n) + e^{\nu(0,t_n)} \mathcal{J}(z, t_n, \pi_n) \right) \right\} \\ &\quad - \pi_n^3 I(z, 0, t_n) e^{\nu(0,t_n)} \mathcal{I}(z, t_n). \end{aligned}$$

Remark now that according to (3.1):

$$\begin{aligned} \mathcal{J}(z, t_n, \pi_n) &\leq \pi_n^2 e^{I(z,0,t_n)} \left(1 - e^{-\nu(0,t_n)} \right) \\ &\leq \pi_n^2 e^{\nu(0,t_n)}. \end{aligned}$$

Then, since $n\pi_n e^{\nu(0,t_n)}$ tends to α as n tends to infinity:

$$\lim_{n \rightarrow +\infty} n e^{\nu(0,t_n)} \mathcal{J}(z, t_n, \pi_n) = 0.$$

Denote by $o(\pi_n, t_n)$ any function such that $no(\pi_n, t_n)$ tends to 0 as n tends to infinity. Then:

$$\begin{aligned} \frac{1}{\phi(z, t_n)} &= f_2 + 1 + \pi_n \left\{ (f_2 + 1) \left(e^{\nu(0,t_n)} (1 - \mathcal{I}(z, t_n)) - 1 \right) - I(z, 0, t_n) \right\} \\ &\quad + \pi_n^2 \left\{ (f_2 + 1) e^{\nu(0,t_n)} \mathcal{I}(z, t_n) - I(z, 0, t_n) \left(e^{\nu(0,t_n)} (1 - \mathcal{I}(z, t_n)) - 1 \right) \right\} \\ &\quad - \pi_n^3 I(z, 0, t_n) e^{\nu(0,t_n)} \mathcal{I}(z, t_n) \\ &\quad + o(\pi_n, t_n) \\ &= 1 + \pi_n \left(e^{\nu(0,t_n)} \left(1 - \left(1 - e^{-\nu(0,t_n)} \right) h(z, t_n) \right) - 1 \right) + o(\pi_n, t_n) \end{aligned}$$

Let $(\phi_n)_{n \in \mathbb{N}}$ be the sequence of functions defined by $\phi_n(z, t_n) = \phi(z, t_n)^n$. Then:

$$\phi_n(z, t_n) = \exp \left\{ -n \log \left(1 + \pi_n \left(e^{\nu(0,t_n)} \left(1 - \left(1 - e^{-\nu(0,t_n)} \right) h(z, t_n) \right) - 1 \right) + o(\pi_n, t_n) \right) \right\},$$

and, since $\pi_n n$ is equivalent to $\alpha e^{-\nu_\infty}$:

$$\lim_{n \rightarrow +\infty} \phi_n(z, t_n) = \exp \left\{ -\alpha (1 - e^{-\nu_\infty}) (1 - h(z)) \right\}.$$

□

Observe that Theorem 4.1 holds whether F_μ satisfies (\mathcal{H}) or not. As an example, consider again the

Haldane model exposed in section 3. The function $h(z, t)$ is then given by:

$$\begin{aligned} h(z, t) &= \frac{1}{1 - e^{-\lambda_\nu t}} \int_0^t \psi(z, u, t) \lambda_\nu e^{-\lambda_\nu(t-u)} du \\ &= \frac{1}{1 - e^{-\lambda_\nu t}} \sum_{i \geq 0} b_i(z) \int_0^t \mathbf{1}_{w \in [ia; (i+1)a)} \lambda_\nu e^{-\lambda_\nu w} dw \\ &= \frac{1}{1 - e^{-\lambda_\nu t}} \sum_{i \geq 0} b_i(z) \left[\left(e^{-\lambda_\nu ia} - e^{-\lambda_\nu t} \right) \mathbf{1}_{t \in [ia; (i+1)a)} \right. \\ &\quad \left. + \left(e^{-\lambda_\nu ia} - e^{-\lambda_\nu(i+1)a} \right) \mathbf{1}_{t \in [(i+1)a; +\infty)} \right]. \end{aligned}$$

Hence the limit of $h(z, t)$ as t tends to infinity:

$$h(z) = \sum_{i \geq 0} b_i(z) e^{-\lambda_\nu ia} (1 - e^{-\lambda_\nu a}).$$

Remark that for $\delta = 0$ and $a = \log(2)$, the result obtained by Ycart (2013) is recovered.

5 Proportional hazard functions

From now on, assume that F_μ satisfies (\mathcal{H}) . Moreover, assume there exists $\rho > 0$ such that for any $s \geq 0$:

$$\lambda_\nu(s) = \rho(1 - 2\delta)\lambda_\mu(s). \quad (\text{H3})$$

An interpretation of assumption (H3) was given at the end of Section 3. In this section, a convergence theorem is deduced from Theorem 4.1, and examples are discussed. In particular, the Luria-Delbrück model with cell deaths is recovered.

Consider first the case where:

$$\mu_\infty = \lim_{t \rightarrow +\infty} \mu(0, t) < +\infty.$$

Under (H3), the function I is given by (3.5). Then hypothesis (H2) is satisfied, and Theorem 4.1 can be applied. Consider now the case where $\mu_\infty = +\infty$. Then:

$$I(z, 0, t) \underset{t \rightarrow +\infty}{\sim} \frac{\delta}{1 - \delta} \mu^*(0, t).$$

Thus:

$$\lim_{t \rightarrow +\infty} \left[e^{-\nu(0, t)} I(z, 0, t) \right] = 0,$$

i.e. the assumption (H2) is still satisfied under (H3), whether μ_∞ is finite or not. Then the following result can be deduced from Theorem 4.1:

Theorem 5.1. *Under (H1) and (H3), as n tends to infinity, the pgf of the number of mutants at time t_n , starting with n normal cells at time 0, converges to the pgf:*

$$\phi(z) = \exp \{ -m(1 - h(z)) \}, \quad (5.1)$$

where

$$h(z) = \frac{1}{1 - e^{-\nu_\infty}} \int_{e^{-\mu_\infty}^1} \frac{\delta(1 - z) + w((1 - \delta)z - \delta)}{(1 - \delta)(1 - z) + w((1 - \delta)z - \delta)} \rho w^{\rho-1} dw,$$

with

$$\mu_\infty = \lim_{t \rightarrow +\infty} \mu(0, t), \quad \mu_\infty^* = (1 - 2\delta)\mu_\infty \quad \text{and} \quad \nu_\infty = \rho\mu_\infty^*.$$

As a general application of Theorem 5.1, let f be a non negative and increasing function on \mathbb{R}_+ , with finite limit f_∞ as t tends to infinity. Let μ be defined for (s, t) in \mathbb{R}_+^2 by:

$$\mu(s, t) = \log \left(\frac{f(t)}{f(s)} \right),$$

Assume that hypothesis (H3) is satisfied. Then:

$$h(z) = \frac{1}{1 - \left(\frac{f^*(0)}{f_\infty} \right)^\rho} \int_{\frac{f^*(0)}{f_\infty}}^1 \frac{\delta(1-z) + w((1-\delta)z - \delta)}{(1-\delta)(1-z) + w((1-\delta)z - \delta)} \rho w^{\rho-1} dw,$$

where $f^*(t) = f^{1-2\delta}$ for any positive t , and $f_\infty^* = f_\infty^{1-2\delta}$. Remark that only the ratio of f_∞ over $f(0)$ has an influence on $h(z)$. Another natural example is the classic case where the final instants of normal and mutant cells are both exponentially distributed, i.e.:

$$\mu(s, t) = \lambda_\mu(t - s), \quad \text{and} \quad \nu(s, t) = \rho(1 - 2\delta)\mu(s, t),$$

where λ_μ is a positive constant. Then the *LDD* distribution (Ycart, 2014) is recovered. Actually, if (H3) is satisfied and $\mu_\infty = +\infty$, the *LDD* distribution can be recovered from Theorem 5.1:

Corollary 5.1. *Under (H1) and (H3), assume that $\mu_\infty = +\infty$. As n tends to infinity, the distribution of the number of mutants at time t_n starting with n normal cells at time 0, converges to the distribution with pgf:*

$$\phi(z) = \exp \{ -\alpha (1 - h(z)) \}, \quad (5.2)$$

where

$$h(z) = \int_0^1 \frac{\delta(1-z) + w((1-\delta)z - \delta)}{(1-\delta)(1-z) + w((1-\delta)z - \delta)} \rho w^{\rho-1} dw.$$

In other words, the *LDD* distribution can be extended to the case where $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ are non-exponential distributions, as long as $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ are cdfs of true measures on \mathbb{R}_+ and the associated hazard functions are proportional.

6 Calculation algorithm

A probability computation algorithm for the distribution of the final mutant counts is described here under the hypotheses of Theorem 5.1. By definition, the pgf ψ is given by:

$$\psi(z, s, t) = \sum_{k \geq 0} p_k(s, t) z^k,$$

where the p_k 's are defined in Proposition 3.3. Thus, (4.1) can be written as:

$$h(z, t) = \sum_{k \geq 0} r_k(t) z^k,$$

where r_k is defined for any $k \geq 0$ by:

$$r_k(t) = \frac{1}{1 - e^{-\nu(0,t)}} \int_0^t p_k(u, t) \lambda_\nu(u) e^{-\nu(u,t)} du.$$

Hence:

$$r_0(t) = \frac{1}{1 - e^{-\nu(0,t)}} \int_{e^{-\mu^*(0,t)}}^1 \frac{\delta - \delta w}{1 - \delta - \delta w} \rho w^{\rho-1} dw,$$

and for all $k > 0$:

$$r_k(t) = \frac{(1-\delta)^{k-1}(1-2\delta)^2}{1-e^{-\nu(0,t)}} \int_{e^{-\mu^*(0,t)}}^1 \frac{(1-w)^{k-1}}{(1-\delta-\delta w)^{k+1}} \rho w^\rho dw.$$

Taking the limit as t tends to infinity:

$$\begin{aligned} r_0 &= \lim_{t \rightarrow +\infty} r_0(t) \\ &= \frac{1}{1-e^{-\nu_\infty}} \int_{e^{-\mu_\infty^*}}^1 \frac{\delta-\delta w}{1-\delta-\delta w} \rho w^{\rho-1} dw, \end{aligned}$$

and for all $k > 0$:

$$\begin{aligned} r_k &= \lim_{t \rightarrow +\infty} r_k(t) \\ &= \frac{(1-\delta)^{k-1}(1-2\delta)^2}{1-e^{-\nu_\infty}} \int_{e^{-\mu_\infty^*}}^1 \frac{(1-w)^{k-1}}{(1-\delta-\delta w)^{k+1}} \rho w^\rho dw. \end{aligned}$$

Then $h(z)$ can be given by:

$$h(z) = \sum_{k \geq 0} r_k z^k.$$

Moreover, (5.1) admits a serie expansion for any $z \in [0; 1]$:

$$\phi(z) = \sum_{k \geq 0} q_k z^k,$$

where the q_k 's can be easily expressed as function of the r_k 's using an algorithm exposed by Embrechts and Hawkes (1982). Firstly:

$$q_0 = \phi(0) = e^{-m(1-r_0)}.$$

The derivative of ϕ with respect to z is given by:

$$\begin{aligned} \frac{d\phi}{dz} &= m \frac{dh}{dz} \phi \\ &= m \left(\sum_{i \geq 1} i r_i z^{i-1} \right) \left(\sum_{j \geq 0} q_j z^j \right) \\ &= m \left(\sum_{i \geq 1, j \geq 0} i r_i q_j z^{i+j-1} \right). \end{aligned}$$

On the other hand:

$$\frac{d\phi}{dz} = \sum_{k \geq 1} k q_k z^{k-1}.$$

Hence for any $k > 0$:

$$\begin{aligned} q_k &= \frac{l}{k} \sum_{\substack{i \geq 1, j \geq 0 \\ i+j=k}} i r_i q_j \\ &= \frac{m}{k} \sum_{i=1}^k i r_i q_{k-i}. \end{aligned}$$

Naturally, if $\mu_\infty = +\infty$, the probabilities of the *LDD* distribution (see Ycart (2014)) are recovered.

7 Conclusion and perspectives

An extension for the classic mutation models to the case where the final instant of a cell depends on its birth date has been proposed. The main results are based on the theory of supercritical branching processes. It led to a family of distributions, modelling asymptotic number of mutants. These distributions depend on the expected number of mutations α , the death probability of mutant cells δ , and the final instant distributions $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ for normal and mutant cells born at a given time s . A convergence theorem for the final count of mutants has been proved for both cases where $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ are defined on $\overline{\mathbb{R}}_+$ or \mathbb{R}_+ . The first case provides the possibility that a cell does not split or die before the end of the experiment. It enables to model more realistic growths, such as logistic growth. The particular case where the hazard functions λ_ν and λ_μ associated to $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ are proportional has been studied. Computation algorithm for probabilities has been described. Moreover, the *LDD* distribution is recovered when $F_\nu(s, \cdot)$ and $F_\mu(s, \cdot)$ are defined on \mathbb{R}_+ and the associated hazard functions are proportional. The consequences for statistical inference and simulation must be developed. Since the R package `flan` (available on CRAN: <https://cran.r-project.org/web/packages/flan/index.html>) provides tools for inference of mutation models for the case where final instants are i.i.d., an extension to the model proposed here is planned.

References

- Allen L (2010) An Introduction to Stochastic Processes with Applications to Biology, 2nd edn. Chapman and Hall/CRC
- Angerer W (2001) An explicit representation of the Luria-Delbrück distribution. *J Math Biol* 42(2):145–174
- Armitage P (1952) The statistical theory of bacterial populations subject to mutation. *J R Statist Soc B* 14:1–40
- Athreya K, Ney P (1972) *Branching Processes*. Springer Berlin Heidelberg
- Bartlett M (1955) *An Introduction to Stochastic Processes, with Special Reference to Methods and Applications*. Cambridge University Press
- Bellman R, Harris T (1952) On age-dependent binary branching processes. *Ann Math* 55(2):280–295
- Benjamini I, Peres Y (1994) Markov chains indexed by trees. *Ann Probab* 22(1):219–243
- Dewanji A, Luebeck E, Moolgavkar S (2005) A generalized Luria-Delbrück model. *Math Biosci* 197(2):140–152
- Embrechts P, Hawkes J (1982) A limit theorem for tails of discrete infinitely divisible laws with applications to fluctuation theory. *J Austral Math Soc Series A* 32:412–422
- Hamon A, Ycart B (2012) Statistics for the Luria-Delbrück distribution. *Elect J Statist* 6:1251–1272
- Harko T, Lobo F, Mak M (2014) Analytical solutions of the Riccati equation with coefficients satisfying integral or differential conditions with arbitrary functions. *Univ J Appl Math* 2(2):109–118
- Kendall D (1952) On the choice of a mathematical model to represent normal bacterial growth. *J R Statist Soc B* 14(1):41–44
- Kimmel M, Axelrod D (2002) *Branching Processes in Biology*. Springer-Verlag New York

- Komarova NL, Wu L, Baldi P (2007) The fixed-size Luria-Delbrück model with a nonzero death rate. *Math Biosci* 210(1):253–290
- Kucera V (1973) A review of the matrix Riccati equation. *Kybernetika* 9(1):42–61
- Lambert A (2005) The branching process with logistic growth. *Ann Appl Probab* 15(2):1506–1535
- Lea D, Coulson C (1949) The distribution of the number of mutants in bacterial populations. *J Genet* 49(3):264–285
- Louhichi S, Ycart B (2015) Exponential growth of bifurcating processes with ancestral dependence. *Adv Appl Probab* 47(2)
- Luria S, Delbrück M (1943) Mutations of bacteria from virus sensitivity to virus resistance. *Genetics* 28(6):491–511
- Montgomery-Smith S, Oveys H (2016) Age-dependent branching processes and applications to the Luria-Delbrück experiment. arXiv preprint arXiv:160806314
- Nguyen H (2006) *An Introduction to Random Sets*. Chapman & Hall/CRC
- Pemantle R (1995) Tree-indexed process. *Statist Sci* 10(2):200–213
- Sarkar S (1991) Haldane’s solution of the Luria-Delbrück distribution. *Genetics* 127:257–261
- Stewart F, Gordon D, Levin B (1990) Fluctuation analysis: the probability distribution of the number of mutants under different conditions. *Genetics* 124(1):175–185
- Tan W (1986) A stochastic Gompertz birth-death process. *Statist Probab Lett* 4(4):25–28
- Tan W, Piantadosi S (1991) On stochastic growth processes with application to stochastic logistic growth. *Statist Sinica* 1:527–540
- Verhulst PF (1838) Notice sur la loi que la population suit dans son accroissement. In: Garnier J, Quetelet A (eds) *Correspondance mathématique et physique*, vol 10, Société Belge de Librairie, Bruxelles, pp 113–121
- Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, Jun S (2010) Robust growth of *Escherichia Coli*. *Curr Biol* 20:1099–1103
- Ycart B (2013) Fluctuation analysis: can estimates be trusted? *PLoS One* 8(12)
- Ycart B (2014) Fluctuation analysis with cell deaths. *J Appl Probab Statist* 9(1):13–29
- Ycart B, Veziris N (2014) Unbiased estimates of mutation rates under fluctuating final counts. *PLoS One* 9(7)
- Yule G (1925) A mathematical theory of evolution, based on the conclusions of Dr. J.C. Willis, F.R.S. *Phil Trans Roy Soc London Ser B* 213:21–87