

Transgenerational effects of ER α 36 over-expression on mammary gland development and molecular phenotype: clinical perspective for breast cancer risk and therapy.

Clémence Chamard-Jovenin¹, Amand Chesnel¹, Emmanuel Bresso², Chloé Morel¹, Charlène Thiébaud¹, Malika Smail-Tabbone², El-Hadi Djermoune¹, Marie-Dominique Devignes², Taha Boukhobza¹ and Hélène Dumond¹.

¹CNRS-Université de Lorraine, UMR 7039, Centre de Recherche en Automatique de Nancy, BP70239, Vandœuvre-lès-Nancy, F-54506, France.

²LORIA, CNRS UMR 7503, INRIA, Villers les Nancy, France.

E-mail : clemence.jovenin@univ-lorraine.fr

Growing source of evidence suggests that exposure to estrogen mimicking agents is a risk factor for breast cancer onset and progression. Long chain alkylphenols are man-made compounds still present in household products, industrial and agricultural processes, leading to a global environmental and human contamination. These molecules are known to exert estrogen-like activities through binding to classical estrogen receptors. Recently, we have demonstrated that a realistic mixture of 4-tert-octylphenol and 4-nonylphenol can stimulate proliferation and modulate epigenetic status of testicular cancer germ cells through a rapid, Estrogen Receptor α 36 (ER α 36)- dependent non genomic pathway (Ajj et al, 2013; doi: 10.1371/journal.pone.0061758). In a retrospective study of breast tumor samples, we also validated ER α 36 expression as a reliable prognostic factor for cancer progression from an estrogen dependent proliferative tumor toward an estrogen dispensable metastatic disease (Chamard-Jovenin et al, 2015; doi: 10.1186/s12918-015-0178-7).

Since high ER α 36 expression enhances expression of migration/invasion markers in breast tumors, we addressed the question of its involvement in response to alkylphenol exposure *in vitro* (MCF-10A mammary epithelial cell line and MCF-7 estrogen-sensitive cancer cells) and *in vivo* (C57BL mice).

A male inherited transgenerational model of exposure to environmentally relevant doses of an alkylphenol mix was set up in C57BL/6J mice to determine whether and how it impacts on mammary gland morphogenesis. Human mammary epithelial MCF-10A cells were exposed to similar doses to decipher the molecular mechanisms involved by a combination of transcriptomic study, cell phenotype analyses, functional and signaling network modeling. The relevance of mouse phenotype extrapolation to human risk is discussed.

Mouse mammary gland exposed transgenerationally to the alkylphenol mix displayed a neoplastic-like histology. This phenotype was correlated with the enhanced proliferation, migration ability and apoptosis resistance observed *in vitro* on human mammary epithelial cells and mediated by the estrogen receptor variant ER α 36.

Since cellular phenotypes are similar *in vivo* and *in vitro* and involve the unique ER α 36 human variant, such consequences of alkylphenol exposure could be extrapolated from mouse model to human. Low dose alkylphenol transgenerational exposure could promote abnormal mammary gland development and subsequently increase the risk of breast cancer at ageing.

This work is supported by ANSES (n°2012-2-014) and INSERM/Plan Cancer (2013-2016).