



HAL
open science

**Looking for new antiplasmodial quinazolines:
DMAP-catalyzed synthesis of 4-benzyloxy- and
4-aryloxy-2-trichloromethylquinazolines and their in
vitro evaluation toward Plasmodium falciparum**

Armand Gellis, Nicolas Primas, Sébastien Hutter, Gilles Lanzada, Vincent Remusat, Pierre Verhaeghe, Patrice Vanelle, Nadine Azas

► **To cite this version:**

Armand Gellis, Nicolas Primas, Sébastien Hutter, Gilles Lanzada, Vincent Remusat, et al.. Looking for new antiplasmodial quinazolines: DMAP-catalyzed synthesis of 4-benzyloxy- and 4-aryloxy-2-trichloromethylquinazolines and their in vitro evaluation toward Plasmodium falciparum. European Journal of Medicinal Chemistry, 2016, 119, pp.34-44. 10.1016/j.ejmech.2016.04.059 . hal-01417007

HAL Id: hal-01417007

<https://hal.science/hal-01417007>

Submitted on 15 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Looking for new antiplasmodial quinazolines: DMAP-catalyzed synthesis of 4-benzyloxy- and 4-aryloxy-2-trichloromethylquinazolines and their *in vitro* evaluation toward *Plasmodium falciparum*

Armand Gellis¹, Nicolas Primas¹, Sébastien Hutter², Gilles Lanzada¹, Vincent Remusat¹, Pierre Verhaeghe^{3*}, Patrice Vanelle¹ and Nadine Azas².

¹Aix-Marseille Université, CNRS, ICR UMR 7273, Laboratoire de Pharmaco-Chimie Radicalaire, Faculté de Pharmacie, 27 Boulevard Jean Moulin – CS30064, 13385 Marseille cedex 05, France.

²Aix-Marseille Université, UMR MD 3, Infections Parasitaires, Transmission et Thérapeutique, Faculté de Pharmacie, 27 Boulevard Jean Moulin – CS30064, 13385 Marseille cedex 05, France.

³ Université Paul Sabatier, Faculté des Sciences Pharmaceutiques – CNRS UPR 8241, Laboratoire de Chimie de Coordination, 205 Route de Narbonne, 31077 Toulouse cedex 04, France.

*Corresponding author: Pierre Verhaeghe, Tel.: +33561333236; E-mail: pierre.verhaeghe@univ-tlse3.fr.

Abstract: A DMAP catalyzed synthesis of new 4-benzyloxy- and 4-aryloxy-2-trichloromethylquinazolines was studied, in a view to react 4-chloroquinazolines with poorly nucleophilic alcohols such as benzylic alcohols, *via* a simple and cheap S_NAr reaction approach. A fast (1 h) general operating procedure, affording good reaction yields, was achieved under microwave irradiation. Thus, a series of 35 molecules was obtained and evaluated *in vitro* on the K1 multi-resistant *P. falciparum* strain, in parallel with a cytotoxicity assessment on the human HepG2 cell line. 5 hit-molecules were identified, presenting both promising antiplasmodial activity ($1.5 \mu M < IC_{50} < 2 \mu M$) and low cytotoxicities ($25 \mu M < CC_{50} < 45 \mu M$). Apart for 2 molecules, the global series displayed a satisfying solubility in the aqueous biological media. Structure-activity relationships showed that the molecules presenting a benzyloxy moiety were less cytotoxic than the ones bearing a phenoxy moiety at position 4 of the quinazoline ring. It also appeared that the introduction of a heteroaryl moiety afforded inactive compounds. Finally, the most active and selective molecules (Selectivity index = 22-27) were the ones presenting either an unsubstituted benzyloxy group or a phenoxy group, this last bearing a *p*-bromo or an *o*-acetyl substituent.

Highlights: ► The DMAP-catalyzed S_NAr reaction between 4-chloro-2-trichloromethylquinazoline and various alcohols or phenols was studied. ► A series of 35 new quinazolines was synthesized in good yields. ► *In vitro* evaluation was made on *Plasmodium falciparum*. ► 5 selective antiplasmodial hit molecules were identified.

Keywords: 2-trichloromethylquinazoline; 4-benzyloxyquinazoline; 4-phenoxyquinazoline; DMAP-catalysis, S_NAr reaction, microwave-assisted chemistry, *Plasmodium falciparum*; antiplasmodial activity; HepG2 cytotoxicity, SARs

1. Introduction

Plasmodium falciparum is the protozoa responsible for cerebral malaria, the leading cause of death among parasitic infections worldwide. According to the 2015 World Malaria Report [1], 214 million people were infected by *Plasmodium* in 2014. It was also estimated that 438.000 people died from this parasitic infection, mainly because of the cerebral form of the disease. About 88% of the estimated deaths occurred in Africa, children under five representing a large majority of the victims.

Since the beginning of 2000, a very significant improvement of the situation has been reached, thanks to the WHO intervention and the involvement of several non-governmental organizations (Bill and Melinda Gates Foundation, Roll Back Malaria, Medicines for Malaria Venture...).

Nevertheless, the emergence of drug-resistant strains of the parasite in Africa, South-East Asia and South America remains a serious concern for the medical and scientific community, in a view to keep on controlling the infection and to try to eradicate the disease. For that purpose, the treatment of the patients infected by *P. falciparum* is based on combination therapies including artemisinin derivatives, in order to cure the disease and avoid the selection or resistant strains and their spreading in the exposed population, which would lead to a very worrying sanitary situation. Concerning artemisinin derivatives, which are nowadays key-molecules for the treatment of *P. falciparum* malaria, resistances are emerging in Asia [2], and it has been demonstrated that they are responsible for therapeutic failures in several infected patients [3]. Moreover, it has also recently been highlighted that the African *Anopheles gambiae* mosquito could transmit such Asian resistant parasites [4], indicating a major worldwide spreading risk. Thus, research efforts have to be maintained so as to discover new chemical entities presenting novel antiplasmodial mechanisms of action, to use in combination therapies with the existing antimalarial drugs, to guaranty their durable efficiency.

Considering all the scaffolds which were studied for their antiplasmodial potential, several bioactive molecules are based on a quinazolinering. For example, febrifugine is natural alkaloid including a quinazolinone moiety which is extracted from the Chinese herb *Dichroafebrifuga*, and which was employed by local people as a medicine against fevers caused by malaria. Some synthetic febrifugine derivatives were prepared and displayed good *in vitro* and *in vivo* potential [5]. Many other quinazoline derivatives were synthesized and evaluated toward *P. falciparum*, in particular, quinazolines bearing, an amino group [6], an alkylamine moiety [7,8] or an anilino substituent [9-11] at position 4 of the quinazoline ring.

In continuation with our research activity focusing on the synthesis of new nitrogen-containing heterocycles with anti-infective potential [12-14], our research group previously reported the synthesis and *in vitro* study of numerous antiplasmodial molecules based on a 2-trichloromethylquinazoline scaffold. Thus, several series bearing an anilino [10], aryl [15], thiophenol [16], sulfonamide [17] or alkynyl moiety [18] at position 4 of the quinazoline ring were prepared and revealed several hit-molecules which are presented in Table 1. Among the antiplasmodial hit-molecules which were identified, a 4-phenoxy-2-trichloromethylquinazoline derivative was also discovered [19], presenting a 50% inhibitory concentration (IC_{50}) of 1.1 μ M and a 50% cytotoxicity concentration (CC_{50}) of 50 μ M (Table

1), reaching an encouraging selectivity index of 45, in comparison with chloroquine and doxycycline. Then, searching for novel analogs presenting an oxygen-containing substituent at position 4 of the quinazoline ring, we present herein the synthesis work which was conducted for reacting 4-chloro-2-trichloromethylquinazoline with poorly nucleophilic benzylic alcohols, to afford the target molecules in a fast, simple and cheap way, using the S_NAr reaction. The *in vitro* biological evaluation and the SARs will then be presented and discussed.

Table 1. *In vitro* antiplasmodial profiles of the previously identified hit-molecules in 2-trichloromethylquinazoline series

Antiplasmodial Activity				
<i>P. falciparum</i> W2 or K1 IC_{50} (μM)	2.5	0.4	1.1	0.9
Cytotoxicity				
HepG2 CC_{50} (μM)	> 125	16	50	> 25
Selectivity Index				
(CC_{50}/IC_{50})	50	40	45	> 28

Reference molecules: Chloroquine (IC_{50} K1 *P. falciparum* = 0.6 μM , CC_{50} HepG2 = 30 μM , SI = 50); Doxycycline (IC_{50} K1 *P. falciparum* = 6 μM , CC_{50} HepG2 = 20 μM , SI = 3.3)

2. Results and discussion

2.1. Synthesis

As aliphatic alcohols display poor nucleophilicity, reacting them with chlorinated azaheterocycles, to provide the corresponding S_NAr products, is difficult and usually requires the preliminary *in situ* formation of the corresponding alcoholate anion. Thus, in quinazoline series, various operating procedures were reported in the literature, using methylate sodium in methanol [20], sodium hydride in dry DMF [21], sodium hydride in dry THF [22,23] or sodium in dry THF [24]. These protocols, leading to the expected products in moderate to good yields (32-84%), present several disadvantages as they consist in 2 step-reactions (formation of the anion followed by the S_NAr), require the use of perfectly dry organic solvents and involve highly reactive and flammable reagents. Moreover, we tried to react 4-chloro-2-trichloromethylquinazolinewith benzyl alcohol, using the reaction conditions we previously reported for the synthesis of 4-phenoxyquinazolines [19] (NaH, DMSO, RT, 24 h) and obtained the expected product in a very low 11% yield (Table 2, entry 13).

Among the available alternative strategies, palladium cross-coupling reactions could have been considered. Nevertheless, because of the expensive costs of both palladium catalysts and ligands, in order to set up a cheap and simple operating procedure, we decided to study the one pot S_NAr reaction between 4-chloro-2-trichloromethylquinazoline **1** and benzyl alcohol. To facilitate the S_NAr reaction between **1** and poorly nucleophilic species like amino-

substituted heterocycles, our group recently reported a synthetic procedure using *N,N*-dimethylaminopyridine (DMAP) as an efficient catalyst, involving the formation of a highly reactive quinazoline intermediate which we isolated and characterized by X-ray diffraction [25]. Then, benefiting from our experience in the use of microwave irradiation for introducing substituents at position 4 of the quinazoline ring [26], and considering that a successful microwave-assisted S_NAr reaction between a 4-chloroquinazoline derivative and ethylene glycol had been reported [27], we focused on a S_NAr reaction protocol combining a DMAP catalysis with microwave irradiation. For that purpose, we first prepared substrate **1** (Figure 1), from commercial 2-methylquinazolin-4(3*H*)-one, according to a previously reported microwave synthesis using PCl₅ in POCl₃ [28].

Figure 1. Microwave-assisted synthesis of substrate **1** from commercial 2-methylquinazolin-4(3*H*)-one.

Then, the DMAP-catalyzed S_NAr reaction between **1** and benzyl alcohol was studied (Figure 2) by varying the solvent nature, the amount of catalyst, the amount of benzyl alcohol, the reaction temperature, the microwave irradiation mode and the reaction time. The experimental results are presented in Table 2.

Figure 2. Studied reaction: preparation of 4-benzyloxy-2-trichloromethylquinazoline **2** from substrate **1**

When using DMAP in the presence of toluene, the reaction yield reached 23%, higher than with DMSO, DMF, or THF. Then, by increasing the amount of DMAP to 0.2 equiv., the reaction yield was improved to 42%. Finally, *via* a 1 h monomode microwave irradiation at 130°C, the use of 0.3 equiv. of DMAP, allowed a decrease in the amount of alcohol (1,2 equiv.) needed and a 53% reaction yield was reached. From that point, decreasing reaction temperature, reaction time, or using a multimode microwave irradiation did not provide satisfying reaction yields.

Table 2. Studied parameters for the DMAP-catalyzed reaction of **1** with benzyl alcohol

Entry	DMAP (mol%)	Reaction conditions ^a	Alcohol equiv.	Solvent	Yield (%)
1	10	1 h, 130 °C/monomode MW	2	toluene	23
2	10	1 h, 150 °C/monomode MW	2	DMSO	5 ^b
3	10	1 h, 150 °C/monomode MW	2	DMF	3 ^b
4	10	1 h, 80 °C/monomode MW	2	THF	Traces ^c
5	20	1 h, 130 °C/monomode MW	2	toluene	42
6	30	0.5 h, 130 °C/monomode MW	2	toluene	27
7	40	1 h, 130 °C/monomode MW	2	toluene	51
8	30	1 h, 130 °C/monomode MW	1.2	toluene	53
9	30	1 h, 130 °C/monomode MW	2	toluene	53
10	30	2 h, 110 °C/monomode MW	2	toluene	39 ^b
11	40	0.5 h, 150 °C/monomode MW	2	toluene	18 ^b
12	40	2 h, 800 W, 110 °C/multimode MW	2	toluene	27
13	-	NaH 95% (2 equiv.), RT, 24 h [19]	1	DMSO	11

^aThe monomode microwave oven used was a Biotage Initiator. The multimode microwave oven used was a Milestone SynthLab Station

^bFormation of resins

^cDetected by LC/MS

For comparison purposes, the same reaction was carried out without DMAP and then conducted with other tertiary amine derivatives (DBU, DABCO and TEA). As shown in Table 3, under these reaction conditions, the presence of DMAP appears mandatory for the reaction to proceed. Moreover, among tertiary amines, only DMAP allowed the synthesis of **2**, considering that substrate **1** was found unconsumed after reaction in the conditions of entries 3 and 4 (only 5% conversion with TEA).

Table 3. Tertiary amine catalysts evaluated for the synthesis of **2**

Entry	Catalyst*	Yield (%)
1	-	0
2	DMAP	53
3	DBU	0
4	DABCO	0

5	TEA	5
*0.3 equiv.		

According to this general procedure, summarized in Figure 3, the S_NAr reaction of **1** with aliphatic alcohols (mainly benzyl alcohols) was then extended to the synthesis of a series of 18 derivatives (molecules **2-19**). As presented in Table 4, the reaction yields varied from moderate (35%) to excellent 97%. In addition, the same protocol was applied to the synthesis of a series of 13 derivatives bearing a phenoxy moiety at position 4 of the quinazoline ring (molecules **20-32**). These molecules were obtained in 38-99% yields. Finally, 4 more derivatives were prepared, bearing a heteroaryloxy moiety at position 4 of the quinazoline ring (molecules **33-36**), in 55-98% yields. Thus, this DMAP-catalyzed operating procedure appeared as generally applicable to aliphatic alcohols, phenols and hydroxyheteroaryl reagents.

Figure 3. Optimal conditions for preparing derivatives belonging to the 4-benzyloxy-2-trichloromethylquinazoline series from substrate **1**

Table 4. Reaction yields and structures of synthesized compounds **2-36**

R-	N°	Yield (%)	R-	N°	Yield (%)
	2	53		20	99
	3	72		21	99
	4	65		22	99
	5	97		23	38
	6	96		24	95

7	91	25	99
8	87	26	47
9	83	27	98
10	81	28	78
11	64	29	94
12	35	30	79
13	68	31	92
14	69	32	89
15	84	33	67
16	41	34	98
17	85	35	67
18	45	36	55
19	68		

Among the 35 synthesized molecules, 2 were previously reported. Thus, compound **2** was mentioned in a 2001 publication [29] but no experimental data is available about it up to day. Concerning quinazoline**27**, it is to note that it had previously been synthesized, according to another protocol [19], in 25% yield. As shown in Figure 4, from the same substrate, the presently described DMAP protocol led to the formation of product**27** in 98% yield and only

1 h, indicating its high efficiency for preparing most of 4-phenoxyquinazolines (apart for molecules **23** and **26**).

Figure 4. Comparison of synthetic routes A and B [19] for the synthesis of compound **27**

2.2. Biological evaluation

All synthesized molecules were then evaluated *in vitro* on the multi-resistant K1 *P. falciparum* strain, by determining their 50% inhibitory concentration (IC_{50}), and compared with 2 antimalarial drug-compounds: chloroquine and doxycycline. In parallel, these molecules were assessed *in vitro* on the HepG2 human hepatocyte cell line, by determining their 50% cytotoxic concentrations (CC_{50}) and comparing them to the one of doxorubicin, used as a cytotoxic reference drug-compound, in order to calculate their respective selectivity index ($SI = CC_{50}/IC_{50}$). The results are presented in Table 5 and highlight 5 hit-molecules (**2**, **6**, **16**, **20**, **26**), presenting both IC_{50} values $< 2 \mu M$ and CC_{50} values $> 25 \mu M$, reaching selectivity indexes ranging between 17 and 27.

Table 5. *In vitro* antiplasmodial and cytotoxicity evaluation of the 4-benzyloxy-2-trichloro-methylquinazolines series

Molecule	R-	HepG2 CC_{50} (μM)	PfK1 IC_{50} (μM)	SI ^e
2		43.8	2.0	21.9
3		50.9	4.0	12.7
4		90.3	$> 10^d$	< 9

5	33.4	2.5	13.4
6	34.9	1.8	19.4
7	29.9	2.2	13.6
8	> 10 ^c	2.3	> 4.3
9	27.8	2.8	9.9
10	33.3	> 10 ^d	< 3.3
11	118.4	3.9	30.6
12	39.9	3.3	12.1
13	25.1	3.5	7.2
14	30.2	4.2	7.2
15	24.6	2.4	10.2
16	25.7	1.5	17.1
17	28.7	3.1	9.3
18	31.25	2.6	12.0
19	62.5	3.8	16.4
20	45	1.8	25
21	36.7	2.3	16.0

22	26.2	1.9	13.8
23	> 8 ^c	1.6	> 5
24	49.5	2.3	21.5
25	1.0	1.7	0.6
26	40.0	1.5	26.7
27	0.5	1.9	0.3
28	75.0	10.0	7.5
29	43.7	4.4	9.9
30	57.0	> 10 ^d	< 5.7
31	5.3	0.9	5.9
32	22.1	1.2	18.4
33	62.7	> 10 ^d	< 6.3
34	4.9	> 10 ^d	< 0.5
35	30.3	> 10 ^d	< 3.0
36	59.0	> 10 ^d	< 5.9
Doxorubicine^a	0.2	-	-
Chloroquine^b	30.0	0.6	50
Doxycycline^b	20.0	6.0	3.3

^aDoxorubicine was used as a cytotoxic reference-drug;

^bChloroquine and doxycycline were used as antimalarial reference-drugs;

^cCC₅₀ could not be determined because of a lack of solubility of the tested molecule in the culture medium;

^dIC₅₀ was not reached at the highest tested concentration (10 μM).

^eSelectivity index (SI) was calculated according to the formula: $SI = CC_{50}/IC_{50}$;

In bold: hit-compounds (IC₅₀ ≤ 2 μM, CC₅₀ ≥ 25 μM and SI ≥ 17)

2.3. Structure-activity relationships

Apart compounds **8** and **23**, all tested molecules presented satisfying solubilities in the biological media as the determination of their *in vitro* activities was not hindered by any precipitation. In the benzyloxy series, no cytotoxicity was identified whereas, in the phenoxy series, compounds **25**, **27** and **31**, presenting a substituent at the *meta* position of the benzene ring, displayed low CC₅₀ values (0.5-5 μM). In the heteroaryloxy series, compound **34** also appeared slightly cytotoxic (CC₅₀ = 5 μM), because of the presence of two bromine atoms.

Concerning antiplasmodial activity, the molecules belonging to the heteroaryloxy series were not active. In the benzyloxy series, compounds **2**, **6** and **16** appeared as hit-molecules with IC₅₀ values ranging from 1.5 to 2 μM and selectivity indexes being around 20, in comparison with the values noted for reference drugs. Compound **2**, bearing a benzyloxy group at position 4 of the quinazoline ring, was found more active and selective than its phenoxy analog (CI₅₀ = 3,1 μM / SI = 16) [19]. In most cases (apart for nitrated molecule **6**), the substitution of the benzene ring of the benzyloxy group was globally deleterious toward antiplasmodial activity, nevertheless, considering compound **16**, it can be noted that the introduction of a substituent on the benzylic carbon, for example *via* a nitrile function, is compatible with maintaining biological activity.

In the phenoxy series, compounds **20** and **26** were identified as hit-molecules, displaying IC₅₀ values of 1.8 and 1.5 μM and selectivity indexes above 25. When comparing *para*-brominated molecule **20** to its chlorinated analog (Table 1), in accordance with the SAR conclusions that we previously reported [19], it can be concluded that the substitution of the benzene ring by a chlorine or bromine atom is favorable toward the antiplasmodial activity if located at the *para* position of the benzene ring, chlorine remaining slightly more favorable. Indeed, we also identified that the introduction of an acetyl substituent at the *ortho* position of the benzene ring (compound **26**) allowed antiplasmodial activity. This selective activity is decreased when changing the substitution position or when changing the acetyl group into a formyl analog (compound **28**). To validate the key role played by the 2-trichloromethylquinazoline scaffold of hit-molecule **26**, its dehalogenated analog **37** was prepared in two steps (Figure 5).

Figure 5. Two step synthesis of molecule **37**

As presented in Figure 6, the antiplasmodial activity of compound **26** depends from the presence of a trichloromethyl group at position 2 of the quinazoline ring, in accordance with the results that we previously reported. Thus, compounds **2** (in the original benzyloxy series) and **26** (first active *o*-substituted phenoxy derivative) appear as novel hit-molecules and extend the available SAR data in the 2-trichloromethylquinazoline series.

Figure 6. Comparison of the *in vitro* antiplasmodial activities of molecules **26** and **37**

Conclusion

Five antiplasmodial hit-molecules were identified from an *in vitro* screening of 35 new 2-trichloromethylquinazoline derivatives bearing an oxygen-containing moiety at position 4 of the quinazoline ring. Structure-activity relationships showed that the derivatives presenting a benzyloxy, a *p*-bromophenoxy or an *o*-acetylphenoxy moiety were the most active and selective ones. The global series was prepared from 4-chloro-2-trichloromethylquinazoline *via* a DMAP catalyzed S_NAr reaction with benzylic alcohols, phenols or hydroxyheterocycles, under microwave irradiation. This operating procedure appears as an efficient, simple, fast and cheap operating protocol for introducing a large diversity of alkoxy, phenoxy or heteroaryloxy substituents at position 4 of 4-chloroquinazolines, substrates of wide pharmaceutical interest.

3. Experimental

3.1. Chemistry

3.1.1. General

Melting points were determined on a Köfler melting point apparatus and are uncorrected. Elemental analyses were carried out at the Spectropole, Faculté des Sciences de Saint-Jérôme (Marseille) with a ThermoFinnigan EA1112 analyzer and a BruckerNonius diffractometer. NMR spectra were recorded on a Bruker ARX 200 spectrometer at the Faculté de Pharmacie of Marseille (200 MHz ^1H NMR: reference CHCl_3 $\delta = 7.26$, and 50 MHz ^{13}C : reference CHCl_3 $\delta = 76.9$). The following adsorbent was used for column chromatography: silica gel 60 (Merck, particle size 0.063-0.200 mm, 70-230 mesh ASTM). TLC was performed on 5 cm x 10 cm aluminum plates coated with silica gel 60F-254 (Merck) in an appropriate eluent. Visualization was performed with ultraviolet light (234 nm). Purity of synthesized compounds was checked by LC/MS analyses, which were realized at the Faculté de Pharmacie of Marseille with a Thermo Scientific Accela High Speed LC System[®] coupled using a single quadrupole mass spectrometer Thermo MSQ Plus[®]. The RP-HPLC column is a ThermoHypersil Gold[®] 50x2.1 mm (C18 bounded), with particles of a diameter of 1.9 mm. The volume of sample injected on the column is 1 μL . Chromatographic analysis, total duration of 8 min, is on the gradient of the following solvents: $t=0$ min, methanol/water 50:50; $0 < t < 4$ min, linear increase in the proportion of water to a methanol/water ratio of 95:5; $4 < t < 6$ min, methanol/water 95:5; $6 < t < 7$ min, linear decrease in the proportion of water to return to a methanol/water ratio of 50:50; $6 < t < 7$ min, methanol/water 50:50. The water used was buffered with ammonium acetate 5 mM. The flow rate of the mobile phase was 0.3 mL/min. The retention times (t_R) of the molecules analyzed are indicated in min. The microwave reactions were performed using multimode reactors: ETHOS Synth Lab station and MicroSYNTH[®] Lab terminal 1024 (Ethos start, Milestone Inc.); or monomode reactors: Biotage Initiator[®] classic in sealed vials.

2-Methylquinazolin-4(3*H*)-one was purchased from Sigma Aldrich.

3.1.2. 4-Chloro-2-trichloromethylquinazoline(1)

4-Chloro-2-trichloromethylquinazoline **1** was prepared as described in the literature [28]. Yield 82%. White solid. Mp 127 °C (lit. 127 °C). ^1H NMR (CDCl_3 , 200 MHz) $\delta = 7.82-$

7.90 (m, 1H), 8.03-8.12 (m, 1H), 8.20-8.24 (m, 1H), 8.33-8.38 (m, 1H). C₉H₄Cl₄N₂, MW: 282.18 g/mol.

3.1.3. General procedure for the preparation of compounds 2 to 34

A mixture of 4-chloro-2-trichloromethylquinazoline **1** (0.2 g, 0.71 mmol), DMAP (26 mg, 0.21 mmol, 0.3 equiv) and adequate alcohol derivative (0.85mmol, 1.2 equiv) in toluene (3 mL) was introduced in miniaturized sealed reactor (5 mL). The reaction mixture was irradiated in a monomode microwave oven, for 1 h at 130 °C. After removal of the toluene under reduced pressure, the residue was purified by silica gel column chromatography and recrystallized from appropriate solvent.

3.3.4. 4-(Benzyloxy)-2-(trichloromethyl)quinazoline (2)

Yield 53%. Beige powder. Mp 115 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.23 (dd, *J* = 1 Hz; 8 Hz, 1H), 8.07 (d, *J* = 8 Hz, 1H), 7.88 (td, *J* = 1 Hz; 5 Hz, 1H), 7.66-7.58 (m, 3H), 7.45-7.37 (m, 3H), 5.74 (s, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.7, 160.1, 150.4, 135.6, 134.4, 129.0, 128.7, 128.6, 123.7, 115.5, 97.3, 69.4. LC-MS (ESI+) *t_R* 5.38 min, *m/z* [M+H]⁺ 351.79/354.40/355.98. MW: 353.63 g/mol. Anal. Calcd for C₁₆H₁₁Cl₃N₂O : C, 54.34; H, 3.14; N, 7.92. Found: C, 54.45; H, 3.11; N, 7.98.

3.1.5. 4-(4-Methylbenzyloxy)-2-(trichloromethyl)quinazoline (3)

Yield 72%. Pale beige powder. Mp 130°C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.23 (d, *J* = 8 Hz, 1H), 8.07 (d, *J* = 8 Hz, 1H), 7.90 (t, *J* = 8 Hz, 1H), 7.67-7.53 (m, 1H), 7.53-7.49 (d, *J* = 8 Hz, 3H), 7.23-7.19 (d, *J* = 8 Hz, 1H), 5.70 (s, 2H), 2.37 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.6, 160.1, 150.4, 138.5, 134.4, 132.6, 129.2, 129.3, 128.7, 128.6, 123.7, 115.6, 97.6, 69.5, 27.3. LC-MS (ESI+) *t_R* 5.68 min, *m/z* [M+H]⁺ 367.11/368.98/371.30. MW: 367.66 g/mol. Anal. Calcd for C₁₇H₁₃Cl₃N₂O : C, 55.54; H, 3.56; N, 7.62. Found: C, 55.39; H, 3.53; N, 7.51.

3.1.6. 4-(4-Methoxybenzyloxy)-2-(trichloromethyl)quinazoline (4)

Yield 65%. Pale beige powder. Mp 86 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.21 (d, *J* = 8 Hz, 1H), 8.06 (d, *J* = 8 Hz, 1H), 7.90 (t, *J* = 8 Hz, 1H), 7.67-7.54 (m, 3H), 6.94-6.90 (d, *J* = 8 Hz, 2H), 5.68 (s, 2H), 3.82 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.6, 159.9, 150.4, 134.4, 131.0, 128.7, 128.6, 127.7, 126.9, 123.7, 115.6, 114.1, 114.0, 97.2, 69.4, 55.3.

LC-MS (ESI+) t_R 5.33 min, m/z $[M+H]^+$ 382.94/384.77/386.88. MW: 383.66 g/mol. Anal. Calcd for $C_{17}H_{13}Cl_3N_2O_2$: C, 53.22; H, 3.42; N, 7.30. Found: C, 53.13; H, 3.39; N, 7.51.

3.1.7. 4-(4-Chlorobenzoyloxy)-2-(trichloromethyl)quinazoline (5)

Yield 97%. Beige powder. Mp 95 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.22 (d, J = 8 Hz, 1H), 8.08 (d, J = 8 Hz, 1H), 7.90 (td, J = 2 Hz; 7 Hz, 1H), 7.69-7.61 (m, 1H), 7.57-7.53 (d, J = 8 Hz, 2H), 7.39-7.34 (d, J = 8 Hz, 2H), 5.69 (s, 2H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 167.4, 160.0, 150.4, 134.5, 134.1, 130.5, 128.8, 128.7, 123.6, 115.5, 97.1, 68.7. LC-MS (ESI+) t_R 5.59 min, m/z $[M+H]^+$ 386.61/388.96/390.91. MW: 388.08 g/mol. Anal. Calcd for $C_{16}H_{10}Cl_4N_2O$: C, 49.52; H, 2.60; N, 7.22. Found: C, 49.70; H, 2.58; N, 7.14.

3.1.8. 4-(4-Nitrobenzoyloxy)-2-(trichloromethyl)quinazoline (6)

Yield 96%. Yellow powder. Mp 199 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.34-8.30 (m, 3H), 8.18-8.15 (m, 1H), 8.04-7.98 (m, 1H), 7.85-7.74 (m, 3H), 5.88 (s, 2H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 167.1, 159.8, 150.5, 147.9, 142.8, 134.8, 129.3, 129.1, 129.0, 123.9, 123.4, 115.3, 97.0, 68.0. LC-MS (ESI+) t_R 5.06 min, m/z $[M+H]^+$ 397.93/399.91/402.06. MW: 398.63 g/mol. Anal. Calcd for $C_{16}H_{10}Cl_3N_3O_3$: C, 48.21; H, 2.53; N, 10.54. Found: C, 48.33; H, 2.55; N, 10.41.

3.1.9. 4-(3-Nitrobenzoyloxy)-2-(trichloromethyl)quinazoline (7)

Yield 91%. Yellow powder. Mp 147 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.27-8.24 (m, 1H), 8.19-8.14 (m, 2H), 7.88-7.81 (m, 3H), 7.69-7.54 (m, 2H), 5.95 (s, 2H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 167.1, 159.7, 150.3, 148.0, 134.9, 133.5, 131.7, 130.1, 129.3, 128.9, 128.8, 124.9, 123.2, 115.1, 97.2, 67.8. LC-MS (ESI+) t_R 4.95 min, m/z $[M+H]^+$ 397.94/399.89/401.96. MW: 398.63 g/mol. Anal. Calcd for $C_{16}H_{10}Cl_3N_3O_3$: C, 48.21; H, 2.53; N, 10.54. Found: C, 48.03; H, 2.56; N, 10.66.

3.1.10. 4-(2-Nitrobenzoyloxy)-2-(trichloromethyl)quinazoline (8)

Yield 87%. Yellow powder. Mp 141 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.28-8.23 (m, 1H), 8.14-8.07 (m, 2H), 7.94 (td, J = 2 Hz; 7 Hz, 1H), 7.82-7.48 (m, 4H), 6.13 (s, 2H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 167.0, 159.9, 150.5, 148.2, 134.7, 133.6, 131.7, 130.3, 129.2, 129.0, 128.9, 125.1, 123.5, 115.3, 96.9, 66.1. LC-MS (ESI+) t_R 4.82 min, m/z

[M+H]⁺ 397.91/400.02/401.86. MW: 398.63 g/mol. Anal. Calcd for C₁₆H₁₀Cl₃N₃O₃ : C, 48.21; H, 2.53; N, 10.54. Found: C, 48.31; H, 2.50; N, 10.39.

3.1.11. 2-(Trichloromethyl)-4-(4-(trifluoromethyl)benzyloxy)quinazoline (9)

Yield 83%. Pale yellow powder. Mp 130 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.27-8.23 (m, 1H), 8.18-8.12 (m, 1H), 8.04-7.98 (m, 1H), 7.97-7.89 (m, 1H), 7.75-7.64 (m, 4H), 5.79 (s, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.3, 160.0, 150.4, 139.6, 134.7, 129.0, 128.9, 126.7, 125.6, 125.5, 123.5, 121.3, 115.4, 97.1, 68.5. LC-MS (ESI+) *t*_R 5.52 min, m/z [M+H]⁺ 420.88/422.92/424.93. MW: 421.63 g/mol. Anal. Calcd for C₁₇H₁₀Cl₃F₃N₂O : C, 48.43; H, 2.39; N, 6.64. Found: C, 48.59; H, 2.41; N, 6.58.

3.1.12. 4-(4-Isopropylbenzyloxy)-2-(trichloromethyl)quinazoline (10)

Yield 81%. Pale beige powder. Mp 121 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.23 (d, *J* = 8 Hz, 1H), 8.07 (d, *J* = 8 Hz, 1H), 7.93-7.85 (m, 1H), 7.67-7.53 (m, 3H), 7.28 (d, *J* = 8 Hz, 2H), 5.72 (s, 2H), 2.94 (sept, *J* = 7 Hz, 1H), 1.27 (d, *J* = 7 Hz, 6H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.6, 160.2, 150.4, 149.5, 134.4, 133.0, 129.3, 128.7, 128.6, 126.7, 123.8, 115.6, 97.3, 68.5, 34.0, 24.0. LC-MS (ESI+) *t*_R 5.73 min, m/z [M+H]⁺ 395.10/396.96/399.02. MW: 395.71 g/mol. Anal. Calcd for C₁₉H₁₇Cl₃N₂O : C, 57.67; H, 4.33; N, 7.08. Found: C, 57.76; H, 4.35; N, 7.21.

3.1.13.4-(3,4-Dimethoxybenzyloxy)-2-(trichloromethyl)quinazoline (11)

Yield 64%. Beige powder. Mp 127 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.22 (d, *J* = 8 Hz, 1H), 8.06 (d, *J* = 8 Hz, 1H), 7.90 (t, *J* = 8 Hz, 1H), 7.63 (t, *J* = 8 Hz, 1H), 7.20-7.16 (m, 2H), 6.89 (d, *J* = 8 Hz, 1H), 5.68 (s, 2H), 3.91 (s, 3H), 3.89 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.6, 160.1, 150.4, 149.4, 149.0, 134.4, 129.0, 128.7, 128.6, 128.1, 126.8, 123.7, 122.2, 115.6, 112.6, 111.0, 97.3, 69.7, 56.0, 55.9. LC-MS (ESI+) *t*_R 4.95 min, m/z [M+H]⁺ 413.00/414.75/416.93. MW: 413.68 g/mol. Anal. Calcd for C₁₈H₁₅Cl₃N₂O₃ : C, 52.26; H, 3.65; N, 6.77. Found: C, 52.09; H, 3.64; N, 6.72.

3.1.14.4-(1-Phenylethoxy)-2-(trichloromethyl)quinazoline (12)

Yield 35%. Beige powder. Mp 141 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.28 (dd, *J* = 1 Hz; 8 Hz, 1H), 8.05 (d, *J* = 8 Hz, 1H), 7.89 (t, *J* = 8 Hz, 1H), 7.70-7.56 (m, 3H),

7.41-7.29 (m, 3H), 6.58 (q, $J = 7$ Hz, 1H), 1.84 (d, $J = 7$ Hz, 3H). ^{13}C NMR (50 MHz, CDCl_3) $\delta = 167.1, 160.1, 150.4, 141.2, 135.5, 134.3, 129.2, 129.0, 128.8, 128.5, 128.1, 126.9, 126.8, 123.7, 115.7, 97.2, 76.0, 22.1$. LC-MS (ESI+) t_R 5.83 min, m/z $[\text{M}+\text{H}]^+$ 366.97/368.97/371.04. MW: 367.66 g/mol. Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{Cl}_3\text{N}_2\text{O}$: C, 55.54; H, 3.56; N, 7.62. Found: C, 55.59; H, 3.59; N, 7.71.

3.1.15. 4-(1-(4-Chlorophenyl)ethoxy)-2-(trichloromethyl)quinazoline (13)

Yield 68%. Brown powder. Mp 86 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) $\delta = 8.22$ (d, $J = 8$ Hz, 1H), 8.05 (d, $J = 8$ Hz, 1H), 7.90 (td, $J = 1$ Hz; 7 Hz, 1H), 7.66(td, $J = 1$ Hz; 7 Hz, 1H), 7.52 (d, $J = 8$ Hz, 2H), 7.32 (d, $J = 8$ Hz, 2H), 6.53 (q, $J = 7$ Hz, 1H), 1.82 (d, $J = 7$ Hz, 3H). ^{13}C NMR (50 MHz, CDCl_3) $\delta = 166.9, 160.0, 150.4, 139.7, 134.4, 133.9, 128.8, 128.7, 128.3, 123.5, 115.6, 97.1, 75.2, 22.0$. LC-MS (ESI+) t_R 5.83 min, m/z $[\text{M}+\text{H}]^+$ 400.96/402.81/404.83. MW: 402.10 g/mol. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{Cl}_4\text{N}_2\text{O}$: C, 50.78; H, 3.01; N, 6.97. Found: C, 50.48; H, 3.03; N, 6.90.

3.1.16. 4-(1-(2,4-Dichlorophenyl)ethoxy)-2-(trichloromethyl)quinazoline (14)

Yield 69%. Brown powder. Mp 123 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) $\delta = 8.31$ (d, $J = 8$ Hz, 1H), 8.07 (d, $J = 8$ Hz, 1H), 7.96-7.87 (m, 1H), 7.69 (t, $J = 7$ Hz, 1H), 7.49-7.39 (m, 2H), 7.20 (dd, $J = 2$ Hz; 8 Hz, 1H), 6.78 (q, $J = 7$ Hz, 1H), 1.81 (d, $J = 7$ Hz, 3H). ^{13}C NMR (50 MHz, CDCl_3) $\delta = 165.3, 159.2, 150.8, 135.3, 131.6, 130.8, 129.4, 129.3, 129.0, 128.6, 123.4, 115.4, 96.9, 72.7, 21.0$. LC-MS (ESI+) t_R 6.02 min, m/z $[\text{M}+\text{H}]^+$ 434.74/436.86/438.81. MW: 436.55 g/mol. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{Cl}_5\text{N}_2\text{O}$: C, 46.77; H, 2.54; N, 6.42. Found: C, 46.41; H, 2.52; N, 6.51.

3.1.17. 4-(1-Phenylprop-2-ynyloxy)-2-(trichloromethyl)quinazoline (15)

Yield 84%. Pale yellow powder. Mp 120 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) $\delta = 8.28$ -8.24 (m, 1H), 8.07 (d, $J = 8$ Hz, 1H), 7.95-7.87 (m, 1H), 7.82-7.77 (m, 2H), 7.69-7.61 (m, 1H), 7.47-7.39 (m, 3H), 7.26-7.24 (m, 1H), 2.75 (d, $J = 2$ Hz, 1H). ^{13}C NMR (50 MHz, CDCl_3) $\delta = 166.3, 159.7, 150.6, 136.3, 134.7, 129.4, 128.8, 128.7, 128.4, 123.8, 115.4, 97.0, 80.1, 76.4, 68.6$. LC-MS (ESI+) t_R 5.20 min, m/z $[\text{M}+\text{H}]^+$ 377.11/379.05/381.08. MW: 377.65 g/mol. Anal. Calcd for $\text{C}_{18}\text{H}_{11}\text{Cl}_3\text{N}_2\text{O}$: C, 57.25; H, 2.94; N, 7.42. Found: C, 57.61; H, 2.95; N, 7.35.

3.1.18. 2-Phenyl-2-(2-(trichloromethyl)quinazolin-4-yloxy)acetonitrile (16)

Yield 41%. Pale yellow powder. Mp 141 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.36-8.21 (m, 1H), 8.14 (d, *J* = 8 Hz, 1H), 7.99 (td, *J* = 2 Hz; 8 Hz, 1H), 7.80-7.68 (m, 3H), 7.54-7.49 (m, 3H), 7.14 (s, 1H). ¹³C NMR (50 MHz, CDCl₃) δ = 166.3, 159.7, 150.6, 136.3, 134.7 129.4, 128.8, 128.7, 128.4, 123.8, 116.0, 114.7, 96.5, 65.9. LC-MS (ESI+) *t*_R 4.73 min, *m/z* [M+H]⁺ 377.99/380.01/381.99. MW: 378.64 g/mol. Anal. Calcd for C₁₇H₁₀Cl₃N₃O : C, 53.93; H, 2.66; N, 11.10. Found: C, 53.89; H, 2.70; N, 11.12.

3.1.19.4-((6-Nitrobenzo[d][1,3]dioxol-5-yl)methoxy)-2-(trichloromethyl)quinazoline (17)

Yield 85%. Pale yellow powder. Mp 181 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.25 (d, *J* = 8 Hz, 1H), 8.10 (d, *J* = 8 Hz, 1H), 7.98-7.92(m, 1H), 7.74-7.65 (m, 2H), 7.27-7.24 (m, 1H), 6.13 (s, 2H), 6.08 (s, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.1, 159.9, 152.2, 150.5, 147.8, 142.3, 134.8 129.0, 128.9, 123.5, 115.3, 109.1, 106.0, 103.2, 96.9, 66.2. LC-MS (ESI+) *t*_R 5.00 min, *m/z* [M+H]⁺ 441.87/443.97/445.95. MW: 442.64 g/mol. Anal. Calcd for C₁₇H₁₀Cl₃N₃O₅ : C, 46.13; H, 2.28; N, 9.49. Found: C, 45.91; H, 2.27; N, 9.57.

3.1.20.(E)-4-(3-(4-Nitrophenyl)allyloxy)-2-(trichloromethyl)quinazoline (18)

Yield 45%. Yellow powder. Mp 140 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.29 (m, 4H), 7.98-7.89(m, 1H), 7.73-7.65 (m, 1H), 7.58-7.24 (m, 2H), 6.98 (d, *J* = 16 Hz, 1H), 6.80-6.66 (m, 1H), 5.41 (dd, *J* = 1 Hz; 6 Hz, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.3, 160.0, 150.4, 147.4, 142.6, 134.6, 133.0, 131.9, 129.6, 129.1, 128.9, 128.8 127.6, 127.3, 124.1 123.8, 123.5, 115.4, 96.3, 67.7. LC-MS (ESI+) *t*_R 5.34 min, *m/z* [M+H]⁺ 423.94/425.92/427.96. MW: 424.67 g/mol. Anal. Calcd for C₁₈H₁₂Cl₃N₃O₃ : C, 50.91; H, 2.85; N, 9.89. Found: C, 51.08; H, 2.86; N, 9.75.

3.1.21. 4-(3-(Benzyloxy)benzyloxy)-2-(trichloromethyl)quinazoline (19)

Yield 68%. Yellow powder. Mp 96 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.23(dd, *J* = 1 Hz; 8 Hz, 1H), 8.10(d, *J* = 8 Hz, 1H), 7.91 (td, *J* = 1 Hz; 8 Hz, 1H), 7.69-7.61 (m, 1H), 7.47-7.18 (m, 8H), 5.72 (s, 1H), 5.72 (s, 2H), 5.10 (s, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.6, 160.1, 159.0, 150.4, 137.2, 136.9, 134.5, 130.0, 129.7, 128.9, 128.8, 128.6, 128.0, 127.5, 123.7, 121.4, 115.5, 115.3, 115.0, 97.2, 70.1, 69.3. LC-MS (ESI+) *t*_R 5.87 min, *m/z* [M+H]⁺ 458.74/460.72/462.84. MW: 459.75 g/mol. Anal. Calcd for C₂₃H₁₇Cl₃N₂O₂ : C, 60.09; H, 3.73; N, 6.09. Found: C, 59.87; H, 3.70; N, 5.98.

3.1.22. 4-(4-Bromophenoxy)-2-(trichloromethyl)quinazoline (20)

Yield 99%. White powder. Mp 179 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.42-8.38 (m, 1H), 8.16(d, *J* = 8 Hz, 1H), 8.02 (td, *J* = 1 Hz; 8 Hz, 1H), 7.81-7.73 (m, 1H), 7.59 (d, *J* = 8 Hz, 2H), 7.30 (d, *J* = 8 Hz, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 164.4, 157.5, 148.7, 148.6, 132.6, 130.0, 126.8, 126.6, 121.1, 120.9, 112.8, 96.7. LC-MS (ESI+) *t_R* 5.38 min, *m/z* [M+H]⁺ 418.26/420.82/422.93. MW: 418.5 g/mol. Anal. Calcd for C₁₅H₈BrCl₃N₂O : C, 43.05; H, 1.93; N, 6.69. Found: C, 43.25; H, 1.95; N, 6.58.

3.1.23. 4-(3-Bromophenoxy)-2-(trichloromethyl)quinazoline (21)

Yield 99%. White powder. Mp 114 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.41-8.37 (m, 1H), 8.18-8.14 (m, 1H), 8.01 (td, *J* = 1 Hz; 8 Hz, 1H), 7.81-7.73 (m, 1H), 7.63-7.62 (m, 1H), 7.48-7.30 (m, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 166.8, 160.0, 152.5, 151.2, 135.1, 130.5, 129.3, 129.1, 125.1, 123.6, 122.3, 120.4, 115.2, 96.6. LC-MS (ESI+) *t_R* 5.43 min, *m/z* [M+H]⁺ 418.18/420.78/422.82. MW: 418.5 g/mol. Anal. Calcd for C₁₅H₈BrCl₃N₂O : C, 43.05; H, 1.93; N, 6.69. Found: C, 42.91; H, 1.92; N, 6.76.

3.1.24. 4-(2-Bromophenoxy)-2-(trichloromethyl)quinazoline (22)

Yield 99%. White powder. Mp 138 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.51-8.47 (m, 1H), 8.17 (d, *J* = 8 Hz, 1H), 8.02 (td, *J* = 1 Hz; 8 Hz, 1H), 7.84-7.67 (m, 2H), 7.49-7.39 (m, 2H), 7.28-7.17 (m, 1H). ¹³C NMR (50 MHz, CDCl₃) δ = 166.7, 159.9, 151.1, 149.4, 135.0, 133.4, 129.2, 128.9, 128.4, 127.4, 123.8, 123.7, 116.2, 114.9, 96.5. LC-MS (ESI+) *t_R* 5.05 min, *m/z* [M+H]⁺ 418.07/420.73/422.87. MW: 418.5 g/mol. Anal. Calcd for C₁₅H₈BrCl₃N₂O : C, 43.05; H, 1.93; N, 6.69. Found: C, 42.89; H, 1.91; N, 6.81.

3.1.25. 4-(2-(Trichloromethyl)quinazolin-4-yloxy)benzotrile (23)

Yield 38%. White powder. Mp 194 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.41 (d, *J* = 8 Hz, 1H), 8.19(d, *J* = 8 Hz, 1H), 8.07-8.00 (m, 1H), 7.84-7.78 (m, 3H), 7.57 (d, *J* = 8 Hz, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 166.4, 159.7, 155.4, 151.3, 135.4, 133.8, 129.6, 129.2, 123.5, 122.7, 118.3, 115.0, 109.9, 96.5. LC-MS (ESI+) *t_R* 4.46 min, *m/z* [M+H]⁺ 364.10/365.99/367.99. MW: 364.61 g/mol. Anal. Calcd for C₁₆H₈Cl₃N₃O : C, 52.71; H, 2.21; N, 11.52. Found :C, 52.71; H, 2.21; N, 11.52.

3.1.26.1-(4-((2-(Trichloromethyl)quinazolin-4-yl)oxy)phenyl)ethanone (24)

Yield 95%. White powder. Mp 149 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.41 (d, *J* = 8 Hz, 1H), 8.18-7.97(m, 4H), 7.78 (t, *J* = 8 Hz, 1H), 7.52 (d, *J* = 8 Hz, 2H), 2.65 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 196.9, 166.7, 159.9, 155.8, 151.2, 135.2, 134.7, 130.0,

129.4, 129.1, 123.6, 121.6, 115.2, 96.6, 26.7. LC-MS (ESI+) t_R 4.48 min, m/z $[M+H]^+$ 381.07/383.07/385.02. MW: 381.64 g/mol. Anal. Calcd for $C_{17}H_{11}Cl_3N_2O_2$: C, 53.50; H, 2.91; N, 7.34. Found :C, 53.71; H, 2.89; N, 7.41.

3.1.27.1-(3-(2-(Trichloromethyl)quinazolin-4-yloxy)phenyl)ethanone (25)

Yield 99%. White powder. Mp 106 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.43 (dd, J = 1 Hz; 8 Hz, 1H), 8.17 (d, J = 8 Hz, 1H), 8.05-7.89(m, 3H), 7.83-7.75 (m, 1H), 7.67-7.54 (m, 2H), 2.64 (s, 3H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 197.1, 167.0, 159.9, 151.1, 138.5, 135.1, 129.7, 129.3, 129.0, 126.3, 125.8, 123.6, 121.5, 115.2, 96.7, 26.8. LC-MS (ESI+) t_R 4.47 min, m/z $[M+H]^+$ 381.07/383.05/384.98. MW: 381.64 g/mol. Anal. Calcd for $C_{17}H_{11}Cl_3N_2O_2$: C, 53.50; H, 2.91; N, 7.34. Found :C, 53.21; H, 2.90; N, 7.42.

3.1.28. 1-(2-(2-(Trichloromethyl)quinazolin-4-yloxy)phenyl)ethanone (26)

Yield 47%. White powder. Mp 147 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.46 (d, J = 8 Hz, 1H), 8.12 (d, J = 8 Hz, 1H), 8.05-7.89 (m, 2H), 7.83-7.75 (m, 1H), 7.68-7.59(m, 1H), 7.46-7.33 (m, 2H), 2.50 (s, 3H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 197.4, 167.5, 159.9, 151.2, 150.6, 135.1, 133.6, 131.3, 130.4, 129.4, 129.0, 126.3, 123.7, 115.2, 96.8, 29.7. LC-MS (ESI+) t_R 4.16 min, m/z $[M+H]^+$ 381.03/383.03/384.99. MW: 381.64 g/mol. Anal. Calcd for $C_{17}H_{11}Cl_3N_2O_2$: C, 53.50; H, 2.91; N, 7.34. Found :C, 53.27; H, 2.93; N, 7.29.

3.1.29.4-(3-Methoxyphenoxy)-2-trichloromethylquinazoline (27)

Yield 98%. White powder. Mp 101 °C (Lit. 101 °C [19]), (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.39-8.43 (m, 1H), 8.13-8.17 (m, 1H), 7.95-8.03 (m, 1H), 7.72-7.80 (m, 1H), 7.32-7.40 (m, 1H), 6.96-7.05 (m, 2H), 6.83-6.89 (m, 1H), 3.83 (s, 3H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 167.4, 160.4, 160.1, 153.1, 151.0, 134.8, 129.7, 129.1, 128.9, 123.7, 115.4, 113.5, 112.0, 107.2, 96.8, 55.5. LC-MS (ESI+) t_R 5.10 min, m/z $[M+H]^+$ 369.01/371.15/373.09. MW: 369.63 g/mol. Anal. Calcd for $C_{16}H_{11}Cl_3N_2O_2$: C, 51.99; H, 3.00; N, 7.58. Found :C, 52.16; H, 3.07; N, 7.67.

3.1.30. 2-(2-(Trichloromethyl)quinazolin-4-yloxy)benzaldehyde (28)

Yield 78%. White powder. Mp 111 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 10.19 (s, 1H), 8.47 (dd, J = 1 Hz; 8 Hz, 1H), 8.17 (d, J = 8 Hz, 1H), 8.06-7.98 (m, 2H), 7.87-7.69

(m, 2H), 7.52-7.45 (m, 2H). ^{13}C NMR (50 MHz, CDCl_3) δ = 188.4, 167.5, 163.1, 153.3, 151.2, 146.7, 135.5, 130.1, 129.5, 129.1, 126.7, 123.6, 123.3, 120.9, 114.9, 96.5. LC-MS (ESI+) t_R 4.20 min, m/z $[\text{M}+\text{H}]^+$ 367.03/369.02/371.03. MW: 367.61 g/mol. Anal. Calcd for $\text{C}_{16}\text{H}_9\text{Cl}_3\text{N}_2\text{O}_2$: C, 52.28; H, 2.47; N, 7.62. Found : C, 52.59; H, 2.45; N, 7.68.

3.1.31.4-(Perfluorophenoxy)-2-(trichloromethyl)quinazoline (29)

Yield 94%. White powder. Mp 139 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) δ = 8.43 (dd, J = 1 Hz; 8 Hz, 1H), 8.22 (d, J = 8 Hz, 1H), 8.07 (td, J = 1 Hz; 8 Hz, 1H), 7.88-7.80 (m, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ = 165.4, 159.5, 151.4, 135.7, 129.9, 129.2, 123.4, 114.0, 96.0. LC-MS (ESI+) t_R 5.49 min, m/z $[\text{M}+\text{H}]^+$ 428.71/430.78/432.67. MW: 429.56 g/mol. Anal. Calcd for $\text{C}_{15}\text{H}_4\text{Cl}_3\text{F}_5\text{N}_2\text{O}$: C, 41.94; H, 0.94; N, 6.52. Found : C, 42.12; H, 0.97; N, 6.42.

3.1.32. 3-Methoxy-5-nitro-4-(2-(trichloromethyl)quinazolin-4-yloxy)benzaldehyde (30)

Yield 79%. Yellow powder. Mp 171 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) δ = 10.28 (s, 1H), 8.49-8.46 (m, 2H), 8.24-8.05 (m, 3H), 7.86 (t, J = 6 Hz, 1H), 3.86 (s, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ = 186.2, 166.6, 159.6, 152.8, 151.4, 148.0, 146.5, 135.6, 135.1, 129.8, 129.6, 129.3, 123.5, 115.6, 114.3, 111.5, 96.2, 56.9. LC-MS (ESI+) t_R 4.01 min, m/z $[\text{M}+\text{H}]^+$ 441.87/443.71/445.72. MW: 442.64 g/mol. Anal. Calcd for $\text{C}_{17}\text{H}_{10}\text{Cl}_3\text{N}_3\text{O}_5$: C, 46.13; H, 2.28; N, 9.49. Found : C, 45.95; H, 2.30; N, 9.55.

3.1.33. 4-(4-Chloro-3-methylphenoxy)-2-(trichloromethyl)quinazoline (31)

Yield 92%. White powder. Mp 98 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) δ = 8.37 (dd, J = 1 Hz; 8 Hz, 1H), 8.16-8.12 (m, 1H), 8.02-7.94 (m, 1H), 7.79-7.70 (m, 1H), 7.41 (d, J = 8 Hz, 1H), 7.31-7.30 (m, 1H), 7.20 (dd, J = 1 Hz; 8 Hz, 1H), 2.41 (s, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ = 166.9, 159.9, 151.0, 150.4, 137.3, 134.9, 131.1, 129.6, 129.1, 128.9, 123.7, 123.5, 120.1, 115.2, 96.7, 20.3. MW: 388.08 g/mol. HRMS (ESI): m/z $[\text{M} + \text{Na}]^+$ calcd for $[\text{C}_{16}\text{H}_{10}\text{Cl}_4\text{N}_2\text{O}]^+$: 408.94395; found: 408.94429.

3.1.34. 4-[(2,4-Dichlorophenyl)oxy]-2-(trichloromethyl)quinazoline (32)

Yield 89%. White powder. Mp 110 °C, (isopropanol). ^1H NMR (200 MHz, CDCl_3) δ = 8.43-8.48 (m, 1H), 8.18 (d, J = 8 Hz, 1H), 8.06-7.98 (m, 1H), 7.84-7.75 (m, 1H), 7.54-7.33 (m, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ = 166.5, 159.8, 151.2, 147.0, 135.2, 131.9, 130.2, 129.4, 129.0, 128.0, 127.9, 124.6, 123.6, 114.8, 96.4. LC-MS (ESI+) t_R 4.60 min, m/z $[\text{M}+\text{H}]^+$

408.05/410.09/411.98.MW: 408.49 g/mol. Anal. Calcd for C₁₅H₇Cl₅N₂O : C, 44.10; H, 1.73; N, 6.86. Found : C, 44.50; H, 1.71; N, 6.97.

3.1.35.2-Methyl-3-(2-(trichloromethyl)quinazolin-4-yloxy)-4H-pyran-4-one (33)

Yield 67%. Beige powder. Mp202 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.41 (dd, *J* = 1 Hz; 8 Hz, 1H), 8.15 (d, *J* = 8 Hz, 1H), 7.99 (td, *J* = 1 Hz; 8 Hz, 1H), 7.82-7.72 (m, 2H), 6.49 (d, *J* = 4 Hz, 1H), 2.36 (s, 3H). ¹³C NMR (50 MHz, CDCl₃) δ = 171.9, 165.9, 159.8, 154.4, 151.3, 135.4, 135.1, 129.2, 128.9, 126.8, 123.8, 117.0, 113.0, 96.5, 15.2. LC-MS (ESI+) *t*_R3.29 min, *m/z* [M+H]⁺370.97/372.99/375.00.MW: 371.60 g/mol. Anal. Calcd for C₁₅H₉Cl₃N₂O₃ : C, 48.48; H, 2.44; N, 7.54. Found :C, 48.38; H, 2.41; N, 7.62.

3.1.36. 4-(5,7-Dibromoquinolin-8-yloxy)-2-(trichloromethyl)quinazoline (34)

Yield 98%. Brown powder. Mp186 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.77-8.55 (m, 3H), 8.21-8.12 (m, 2H), 8.04(t, *J* = 8 Hz, 1H), 7.83 (t, *J* = 6 Hz, 1H), 7.56-7.53 (m, 1H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.13, 159.6, 151.4, 151.3, 146.5, 142.2, 136.2, 135.1, 133.3, 129.3, 128.9, 127.8, 124.1, 122.8, 119.5, 116.9, 114.8, 96.4. LC-MS (ESI+) *t*_R5.37 min, *m/z* [M+H]⁺546.08/547.52/549.54.MW: 548.44 g/mol. Anal. Calcd for C₁₈H₈Br₂N₃O : C, 39.42; H, 1.47; N, 7.66. Found: C, 39.49; H, 1.50; N, 7.51.

3.1.37.4-(Quinolin-2-yloxy)-2-(trichloromethyl)quinazoline (35)

Yield 67%. Yellow powder. Mp 163 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.52(d, *J* = 8 Hz, 1H), 8.34 (d, *J* = 8 Hz, 1H), 8.18 (d, *J* = 8 Hz, 1H), 8.07-7.72 (m, 5H), 7.64-7.49 (m, 2H). ¹³C NMR (50 MHz, CDCl₃) δ = 166.9, 159.8, 157.4, 151.3, 146.3, 140.1, 135.3, 130.5, 129.4, 129.0, 128.4, 127.3, 126.8, 124.0, 115.7, 115.2, 96.5. LC-MS (ESI+) *t*_R4.60 min, *m/z* [M+H]⁺389.95/391.95/394.06.MW: 390.65 g/mol. Anal. Calcd for C₁₈H₁₀Cl₃N₃O : C, 55.34; H, 2.58; N, 10.76. Found :C, 55.13; H, 2.56; N, 10.91.

3.1.38.4-(2-(Trichloromethyl)quinazolin-4-yloxy)-2H-chromen-2-one (36)

Yield 55%. Yellow powder. Mp237 °C, (isopropanol). ¹H NMR (200 MHz, CDCl₃) δ = 8.45 (d, *J* = 8 Hz, 1H), 8.26 (d, *J* = 8 Hz, 1H), 8.10 (t, *J* = 8 Hz, 1H), 7.92-7.80 (m, 2H), 7.68-7.60 (m, 1H), 7.47-7.30 (m, 2H), 6.96 (s, 1H). ¹³C NMR (50 MHz, CDCl₃) δ = 165.3, 161.6, 159.7, 153.7,151.7135.9, 133.0, 130.1, 129.6, 124.4, 123.1, 122.9, 117.3, 115.5, 115.1, 104.3, 96.1.

LC-MS (ESI+) t_R 4.60 min, m/z $[M+H]^+$ 406.93/408.93/411.18. MW: 407.63 g/mol. Anal. Calcd for $C_{18}H_9Cl_3N_2O_3$: C, 53.04; H, 2.23; N, 6.87. Found: C, 52.86; H, 2.24; N, 6.92.

3.1.39. Preparation of 1-(2-(2-methylquinazolin-4-yloxy)phenyl)ethanone (37)

Molecule **37** was prepared in two steps from commercial 2-methylquinazolin-4(3*H*)-one. This last was first chlorinated at position 4, by using $POCl_3$, according to a previously microwave-assisted reported procedure [25], leading to 4-chloro-2-methylquinazoline in 73% yield. Then, the general DMAP-catalyzed operating procedure described in § 3.1.3. was applied to this intermediate product, leading to molecule **37** in 45% yield, as a beige powder. Mp 118 °C, (isopropanol). 1H NMR (200 MHz, $CDCl_3$) δ = 8.33 (d, J = 8 Hz, 1H), 7.95-7.83 (m, 3H), 7.65-7.56 (m, 2H), 7.42-7.25 (m, 2H), 2.59 (s, 3H), 2.47 (s, 3H). ^{13}C NMR (50 MHz, $CDCl_3$) δ = 197.9, 166.3, 163.7, 152.0, 150.9, 134.3, 133.3, 132.0, 130.1, 127.1, 127.0, 126.0, 123.8, 123.4, 114.3, 30.1, 26.2. LC-MS (ESI+) t_R 4.60 min, m/z $[M+H]^+$ 279.19. MW: 278.31 g/mol. Anal. Calcd for $C_{17}H_{14}N_2O_2$: C, 73.37; H, 5.07; N, 10.07. Found: C, 73.82; H, 5.11; N, 9.93.

3.2. Biology

3.2.1. *In vitro* Antiplasmodial evaluation

In this study, a K1 culture-adapted *P. falciparum* strain resistant to chloroquine, pyrimethamine and proguanil was used in an *in vitro* culture. Maintenance in continuous culture was done as described previously by Trager and Jensen [30]. Cultures were maintained in fresh A+ human erythrocytes at 2.5% hematocrit in complete medium (RPMI 1640 with 25 mM HEPES, 25 mM $NaHCO_3$, 10% of A+ human serum) at 37 °C under reduced O_2 atmosphere (gas mixture 14% O_2 , 6% CO_2 , and 80% N_2). Parasitaemia was maintained daily between 1% and 6%. The *P. falciparum* drug susceptibility test was carried out by comparing quantities of DNA in treated and control cultures of parasite in human erythrocytes according to a SYBR Green I fluorescence-based method [31] using a 96-well fluorescence plate reader. Compounds, previously dissolved in DMSO (final concentration less than 0.5% v/v) were incubated in a total assay volume of 200 μ L (RPMI, 4% hematocrit and 1% parasitaemia) for 72 h in a humidified atmosphere (14% O_2 and 6% CO_2) at 37 °C, in 96-well flat bottom plates. Duplicate assays were performed for each sample. After incubation, 125 μ L supernatant was discarded and cells were washed twice with 125 μ L 1X PBS. 15 μ L re-suspended cells were transferred to 96-well flat bottom nonsterile black plates

(Greiner Bio-one) already containing 15 μL of the SYBR Green I lysis buffer (2X SYBR Green I, 20 mM Tris base pH 7.5, 20 mM EDTA, 0.008% w/v saponin, 0.08% w/v Triton X-100). Negative control, treated by solvents (DMSO or H_2O) and positive controls (chloroquine and doxycycline) were added to each set of experiments. Plates were incubated for 15 min at 37 °C and then read on a TECAN Infinite F-200 spectrophotometer with excitation and emission wavelengths at 485 and 535 nm, respectively. The concentrations of compounds required to induce a 50% decrease of parasite growth (IC_{50} K1) were calculated from three independent experiments.

3.2.2. *In vitro* Cytotoxicity evaluation

HepG2 cell line was maintained at 37 °C, 6% CO_2 , 14% O_2 , 80% N_2 , with 90% humidity in RPMI supplemented with 10% foetal bovine serum, 1% L-glutamine (200 mM) and penicillin (100 U/mL) / streptomycin (100 $\mu\text{g}/\text{mL}$) (complete RPMI medium). The evaluation of the tested molecules cytotoxicity on the HepG2 (hepatocarcinoma cell line purchased from ATCC, ref HB-8065) cell line was performed according to the method of Mosmann [32] with slight modifications. Briefly, $5 \cdot 10^3$ cells in 100 μL of complete medium were inoculated into each well of 96-well plates and incubated at 37 °C in a humidified 6% CO_2 . After 24 h incubation, 100 μL of medium with various product concentrations dissolved in DMSO (final concentration less than 0.5% v/v) were added and the plates were incubated for 72 h at 37 °C. Triplicate assays were performed for each sample. Each plate-well was then microscope-examined for detecting possible precipitate formation before the medium was aspirated from the wells. 100 μL of MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide) solution (0.5 mg/mL in medium without FCS) were then added to each well. Cells were incubated for 2 h at 37 °C. After this time, the MTT solution was removed and DMSO (100 μL) was added to dissolve the resulting blue formazan crystals. Plates were shaken vigorously (700 rpm) for 10 min. The absorbance was measured at 570 nm with 630 nm as reference wavelength using a BIO-TEK ELx808 Absorbance Microplate Reader. DMSO was used as blank and doxorubicin (purchased from Sigma Aldrich) as positive control. Cell viability was calculated as percentage of control (cells incubated without compound). The 50% cytotoxic concentration (CC_{50}) was determined from the dose–response curve by using the TableCurve software 2D v.5.0. CC_{50} values represent the mean value calculated from three independent experiments.

4. Acknowledgement

This work was supported by the Centre National de la Recherche Scientifique (CNRS) and by Aix-Marseille Université (AMU).

5. References

- [1] World Health Organization (WHO). World Malaria Report 2015. Geneva, Switzerland: World Health Organization (WHO), <http://www.who.int/malaria/publications/world-malaria-report-2015/report/en/>
- [2] A. P. Phyto, S. Nkhoma, K. Stepniewska, E. A. Ashley, S. Nair, R. McGready, C. L. Moo, S. Al-Saai, A. M. Dondorp, K. M. Lwin, P. Singhasivanon, N. P. J. Day, N. J. White, T. J. C. Anderson, F. Nosten, Emergence of artemisinin-resistant malaria on the west border of Thailand: a longitudinal study. *Lancet* 379 (2012) 1960-1966.
- [3] M. D. Spring, J. T. Lin, J. E. Manning, P. Vanachayangkul, S. Somethy, R. Bun, Y. Se, S. Chann, M. Ittiverakul, P. Sia-ngam, W. Kuntawunginn, M. Arsanok, N. Buathong, S. Chaorattanakawee, P. Gosi, W. Ta-aksorn, N. Chanarat, S. Sundrakes, N. Kong, T. K. Heng, S. Nou, P. Teja-isavadharm, S. Pichyangkul, S. T. Phann, S. Balasubramanian, J. J. Juliano, S. R. Meshnick, C. M. Chour, S. Prom, C. A. Lanteri, C. Lon, D. L. Saunders, Dihydroartemisinin-piperaquine failure associated with a triple mutant including kelch13 C580Y in Cambodia: an observational cohort study. *Lancet Inf. Dis.* 15 (2015) 683-691.
- [4] B. St Laurent, B. Miller, T. A. Burton, C. Amaratunga, S. Men, S. Sovannaroth, M. P. Fay, O. Miotto, R. W. Gwadz, J. M. Anderson, R. M. Fairhurst, Artemisinin-resistant *Plasmodium falciparum* clinical isolates can infect diverse mosquito vectors of Southeast Asia and Africa. *Nat. Commun.* 6 (2015) 8614.
- [5] S. Zhu, J. Wang, G. Chandrashekar, E. Smith, X. Liu, Y. Zhang, Synthesis and evaluation of 4-quinazolinone compounds as potential antimalarial agents. *Eur. J. Med. Chem.* 45 (2010) 3864-3869.
- [6] C. Mendoza-Martinez, J. Correa-Basurto, R. Nieto-Meneses, A. Marquez-Navarro, R. Aguilar-Suarez, M. Dinora Montero-Cortes, B. Noguera-Torres, E. Suarez-Contreras, N. Galindo-Sevilla, A. Rojas-Rojas, A. Rodriguez-Lezama, F. Hernandez-Luis, Design,

synthesis and biological evaluation of quinazoline derivatives as anti-trypanosomatid and anti-plasmodial agents, *Eur. J. Med. Chem.* 96 (2015) 296–307.

[7] N. A. Malmquist, T. A. Moss, S. Mecheri, A. Scherf, M. J. Fuchter, Small-molecule histone methyltransferase inhibitors display rapid antimalarial activity against all blood stage forms in *Plasmodium falciparum*, *Proc. Natl. Acad. Sci.* 109 (2012) 16708–16713.

[8] S. Sundriyal, N. A. Malmquist, J. Caron, S. Blundell, F. Liu, X. Chen, N. Srimongkolpithak, J. Jin, S. A. Charman, A. Scherf, M. J. Fuchter, Development of diaminoquinazoline histone lysine methyltransferase inhibitors as potent blood-stage antimalarial compounds, *ChemMedChem*. 9 (2014) 2360–2373.

[9] A. Mishra, K. Srivastava, R. Tripathi, S. K. Puri, S. Batra, Search for new pharmacophores for antimalarial activity. Part III: synthesis and bioevaluation of new 6-thioureido-4-anilinoquinazolines, *Eur. J. Med. Chem.* 44 (2009) 4404–4412.

[10] P. Verhaeghe, N. Azas, S. Hutter, C. Castera-Ducros, M. Laget, A. Dumètre, M. Gasquet, J.-P. Reboul, S. Rault, P. Rathelot, P. Vanelle, Synthesis and *in vitro* antiplasmodial evaluation of 4-anilino-2-trichloromethylquinazolines, *Bioorg. Med. Chem.* 17 (2009) 4313–4322.

[11] Y. Kabri, N. Azas, A. Dumètre, S. Hutter, M. Laget, P. Verhaeghe, A. Gellis, P. Vanelle, Original quinazoline derivatives displaying antiplasmodial properties, *Eur. J. Med. Chem.* 45 (2010) 616–622.

[12] S. Djekou, A. Gellis, J. Maldonado, M. P. Crozet, P. Vanelle, A radical synthetic approach to the new potentially bioactive pyrimidinones, *Heterocycles* 55 (2001) 535–544.

[13] P. Verhaeghe, P. Rathelot, S. Rault, P. Vanelle, Convenient preparation of original vinylic chlorides with antiparasitic potential in quinoline series, *Lett. Org. Chem.* 3 (2006) 891–897.

[14] N. Primas, P. Suzanne, P. Verhaeghe, S. Hutter, C. Kieffer, M. Laget, A. Cohen, J. Broggi, J.-C. Lancelot, A. Lesnard, P. Dallemagne, P. Rathelot, S. Rault, P. Vanelle, N. Azas,

Synthesis and *in vitro* evaluation of 4-trichloromethylpyrrolo[1,2-a]quinoxalines as new antiplasmodial agents, *Eur. J. Med. Chem.* 83 (2014) 26-35.

[15] P. Verhaeghe, N. Azas, M. Gasquet, S. Hutter, C. Ducros, M. Laget, S. Rault, P. Rathelot, P. Vanelle, Synthesis and antiplasmodial activity of new 4-aryl-2-trichloromethylquinazolines, *Bioorg. Med. Chem. Lett.* 18 (2008) 396–401.

[16] P. Verhaeghe, A. Dumètre, C. Castera-Ducros, S. Hutter, M. Laget, C. Fersing, M. Prieri, J. Yzombard, F. Sifredi, S. Rault, P. Rathelot, P. Vanelle, N. Azas, 4-Thiophenoxy-2-trichloromethylquinazolines display *in vitro* selective antiplasmodial activity against the human malaria parasite *Plasmodium falciparum*, *Bioorg. Med. Chem. Lett.* 21 (2011) 6003–6006.

[17] N. Primas, P. Verhaeghe, A. Cohen, C. Kieffer, A. Dumètre, S. Hutter, S. Rault, P. Rathelot, N. Azas, P. Vanelle, A New Synthetic Route to Original Sulfonamide Derivatives in 2-Trichloromethylquinazoline Series: A Structure-Activity Relationship Study of Antiplasmodial Activity, *Molecules.* 17 (2012) 8105–8117.

[18] C. Kieffer, P. Verhaeghe, N. Primas, C. Castera-Ducros, A. Gellis, R. Rosas, S. Rault, P. Rathelot, P. Vanelle, Sonogashira cross-coupling reaction in 4-chloro-2-trichloromethylquinazoline series is possible despite a side dimerization reaction, *Tetrahedron* 69 (2013) 2987-2995.

[19] C. Castera-Ducros, N. Azas, P. Verhaeghe, S. Hutter, P. Garrigue, A. Dumètre, L. Mbatchi, M. Laget, V. Remusat, F. Sifredi, S. Rault, P. Rathelot, P. Vanelle, Targeting the human malaria parasite *Plasmodium falciparum*: *In vitro* identification of a new antiplasmodial hit in 4-phenoxy-2-trichloromethylquinazoline series, *Eur. J. Med. Chem.* 46 (2011) 4184–4191.

[20] A. R. Genady, promising carboranylquinazolines for boron neutron capture therapy: synthesis, characterization and *in vitro* toxicity evaluation, *Eur. J. Med. Chem.* 44 (2009) 409–416.

- [21] R. A. Smits, I. J. P. de Esch, O. P. Zuiderveld, J. Broeker, K. Sansuk, E. Guaita, G. Coruzzi, M. Adami, E. Haaksma, R. Leurs, Discovery of quinazolines as histamine H4 receptor inverse agonists using a scaffold hopping approach, *J. Med. Chem.* 51 (2008) 7855–7865.
- [22] A. Purohit, R. Benetti, M. Hayes, M. Guaraldi, M. Kagan, P. Yalamanchili, F. Su, M. Azure, M. Mistry, M. Yu, S. Robinson, D. D. Dischino, D. Casebier, Quinazoline derivatives as MC-I inhibitors: evaluation of myocardial uptake using positron emission tomography in rat and non-human primate, *Bioorg. Med. Chem. Lett.* 17 (2007) 4882–4885.
- [23] T-A. N. Pham, Z. Yang, Y. Fang, J. Luo, J. Lee, H. Park, Synthesis and biological evaluation of novel 2,4-disubstituted quinazoline analogues as GPR119 agonists, *Bioorg. Med. Chem.* 21 (2013) 1349–1356.
- [24] M. Spulak, J. Pourova, M. Voprsalova, J. Mikusek, J. Kunes, J. Vacek, M. Ghavre, N. Gathergood, M. Pour, Novel bronchodilatoryquinazolines and quinoxalines: synthesis and biological evaluation, *Eur. J. Med. Chem.* 74 (2014) 65–72.
- [25] A. Gellis, C. Kieffer, N. Primas, G. Lanzada, M. Giorgi, P. Verhaeghe, P. Vanelle, A new DMAP-catalyzed and microwave-assisted approach for introducing heteroaryl amino substituents at position 4 of the quinazoline ring, *Tetrahedron* 70 (2014) 8257–8266.
- [26] Y. Kabri, P. Verhaeghe, A. Gellis, P. Vanelle, Regioselective Suzuki-Miyaura Reaction: application the microwave-promoted synthesis of 4,7-diarylquinazolines, *Molecules* 15 (2010) 2949–2961.
- [27] H-Y Li, Y. Wang, W. T. McMillen, A. Chatterjee, J. E. Toth, S. R. Mundla, M. Voss, R. D. Boyer, J. Scott Sawyer, A concise synthesis of quinazoline TGF- β RI inhibitor through one-pot three-component Suzuki-Miyaura/etherification and imidate-amide rearrangement reactions, *Tetrahedron* 63 (2007) 11763–11770.
- [28] P. Verhaeghe, P. Rathelot, A. Gellis, S. Rault, P. Vanelle, Highly efficient microwave assisted α -trichlorination reaction of α -methylated nitrogen containing heterocycles, *Tetrahedron*. 62 (2006) 8173–8176.

- [29] D. B. Jordan, G. S. Basarab, D-I. Liao, W. M. P. Johnson, K. N. Winzenberg, D. A. Winkler, Structure-based design of inhibitors of the rice blast fungal enzyme trihydroxynaphthalene reductase, *J. Mol. Graphics Modell.* 19 (2001) 434–447.
- [30] W. Trager, J.B. Jensen, Human malaria parasites in continuous culture, *Science.* 193 (1976) 673–675.
- [31] W.A. Guiguemde, A.A. Shelat, D. Bouck, S. Duffy, G.J. Crowther, P.H. Davis, D.C. Smithson, M. Connelly, J. Clark, F. Zhu, M.B. Jiménez-Díaz, M.S. Martinez, E.B. Wilson, A.K. Tripathi, J. Gut, E.R. Sharlow, I. Bathurst, F. El Mazouni, J.W. Fowble, I. Forquer, P.L. McGinley, S. Castro, I. Angulo-Barturen, S. Ferrer, P.J. Rosenthal, J.L. Derisi, D.J. Sullivan, J.S. Lazo, D.S. Roos, M.K. Riscoe, M.A. Phillips, P.K. Rathod, W.C. Van Voorhis, V.M. Avery, R.K. Guy, Chemical genetics of *Plasmodium falciparum*, *Nature.* 465 (2010) 311–315.
- [32] T. Mosmann, Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods.* 65 (1983).

Graphical Abstract: