



**HAL**  
open science

## Erratum: Spectral analysis of semigroups and growth-fragmentation equations

Stéphane Mischler

► **To cite this version:**

Stéphane Mischler. Erratum: Spectral analysis of semigroups and growth-fragmentation equations. 2016. hal-01422273

**HAL Id: hal-01422273**

**<https://hal.science/hal-01422273>**

Preprint submitted on 24 Dec 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ERRATUM: SPECTRAL ANALYSIS OF SEMIGROUPS AND GROWTH-FRAGMENTATION EQUATIONS

S. MISCHLER

**Version of December 24, 2016**

ABSTRACT. We correct the article [S. Mischler, J. Scher, Spectral analysis of semigroups and growth-fragmentation equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 33 (2016), no. 3, 849898].

In the article [2], there was an error in Step 1 in the proof of Theorem 2.1 that we were not able to fix, with fatal repercussions on most of the abstract spectral analysis results (Corollary 2.4, Theorem 3.1, Theorem 3.3 and Theorem 3.5 in [2]). However, we claim that Corollary 2.5 is correct and that we may slightly modify the statements of the other above mentioned abstract results in order to make them correct and then to use these variants in order to repair the proof of [2, Theorem 1.1].

We employ the notation and assumptions of [2], except that we replace assumption **(H2)** by the assumptions **(h2)** or **(h2')** below. More precisely, we consider a Banach space  $X$  and the generator  $\Lambda$  of a strongly continuous semigroup  $S_\Lambda$  on  $X$ . We assume that  $\Lambda$  splits as

$$\Lambda = \mathcal{A} + \mathcal{B}$$

where  $\mathcal{B}$  is the generator of a strongly continuous semigroup  $S_\mathcal{B}$  and  $\mathcal{A}$  is  $\mathcal{B}$ -bounded. We also assume that the operators  $\mathcal{A}$  and  $S_\mathcal{B}$  satisfy one of the two following regularizing properties of an iterated enough convolution product:

**(h2)** there exist  $\zeta \in (0, 1]$  and  $\zeta' \in [0, \zeta)$  such that  $\mathcal{A} \in \mathcal{B}(X_{\zeta'}, X)$  and there exists an integer  $n \geq 1$  such that for any  $a > a^*$ , there holds

$$\forall t \geq 0, \quad \|(\mathcal{A}S_\mathcal{B})^{(*n)}(t)\|_{\mathcal{B}(X, X_\zeta)} \leq C_{a,n,\zeta} e^{at}$$

for a constant  $C_{a,n,\zeta} \in (0, \infty)$ ,

or

**(h2')** there exist  $\zeta \in [-1, 0)$  and  $\zeta' \in (\zeta, 0]$  such that  $\mathcal{A} \in \mathcal{B}(X, X_{\zeta'})$  and there exists an integer  $n \geq 1$  such that for any  $a > a^*$ , there holds

$$\forall t \geq 0, \quad \|(S_\mathcal{B}\mathcal{A})^{(*n)}(t)\|_{\mathcal{B}(X_\zeta, X)} \leq C_{a,n,\zeta} e^{at}$$

for a constant  $C_{a,n,\zeta} \in (0, \infty)$ .

It is worth recalling that  $X_\zeta$  stands here for the abstract Sobolev space associated to  $\mathcal{B}$  (and thus also to  $\Lambda$ ) and  $\zeta \in \mathbb{R}$ .

Replacing hypothesis **(H2)** in Theorem 2.1 by hypothesis **(h2)** or **(h2')**, we denote as Theorem E.2.1 the resulting statement. We denote similarly Corollary E.2.4, Theorem E.3.1, Theorem E.3.3 and Theorem E.5.3.

We are going now to briefly explain how to modify the arguments presented in [2] in order to prove the above erratum results. More details and more general versions of these results are available in [1].

*Sketch of the proof of Theorem E.2.1.* We first consider the case when we assume **(h2)** instead of **(H2)**. We fix  $a \in (a^*, a')$ . On the one hand, from **(h2)**, we have

$$\forall z \in \Delta_a, \quad \|(\mathcal{A}\mathcal{R}_{\mathcal{B}}(z))^n\|_{X \rightarrow X_\zeta} \leq C'_{a,n},$$

for a constant  $C'_{a,n} \in (0, \infty)$  only depending on  $a$ ,  $a^*$  and  $C_{a,n,\zeta}$ . On the other hand, using the same interpolation argument as in Step 1 in the proof of [2, Theorem 2.1], we have

$$\forall z \in \Delta_a, \quad \|\mathcal{R}_{\mathcal{B}}(z)\|_{X_\zeta \rightarrow X_{\zeta'}} \leq C_{a,\varepsilon} \langle z \rangle^{-(\zeta - \zeta' - \varepsilon)},$$

for any  $\varepsilon > 0$  and for a constant  $C_{a,\varepsilon} \in (0, \infty)$  only depending on  $a$ ,  $\varepsilon$  and the dissipativity estimate on  $\mathcal{B}$ . Writing

$$(\mathcal{A}\mathcal{R}_{\mathcal{B}}(z))^{n+1} = \mathcal{A}\mathcal{R}_{\mathcal{B}}(z)(\mathcal{A}\mathcal{R}_{\mathcal{B}}(z))^n$$

and using the available estimates on  $(\mathcal{A}\mathcal{R}_{\mathcal{B}}(z))^n$ ,  $\mathcal{R}_{\mathcal{B}}(z)$  and  $\mathcal{A}$ , we immediately obtain that

$$\forall z \in \Delta_a, \quad \|(\mathcal{A}\mathcal{R}_{\mathcal{B}}(z))^{n+1}\|_{\mathcal{B}(X)} \leq C_{a,\alpha} \frac{1}{\langle z \rangle^\alpha},$$

for any  $\alpha \in (0, \zeta - \zeta')$  and a constant  $C_{a,\alpha} \in (0, \infty)$ . That last estimate is nothing but [2, (2.12)] and thus concludes the proof of Step 1 in the proof of [2, Theorem 2.1]. The other steps are unchanged in the case we make the assumption **(h2)**.

We now explain how to adapt the proof in the case we assume **(h2')**. More details are given in [3, 1]. The first step consists in writing

$$(R_{\mathcal{B}}(z)\mathcal{A})^{n+1} = (R_{\mathcal{B}}(z)\mathcal{A})^n R_{\mathcal{B}}(z)\mathcal{A}$$

and to prove in a similar way as above that

$$\forall z \in \Delta_a, \quad \|(R_{\mathcal{B}}(z)\mathcal{A})^{n+1}\|_{\mathcal{B}(X)} \leq C_{a,\alpha} \frac{1}{\langle z \rangle^\alpha},$$

for any  $\alpha \in (0, \zeta' - \zeta)$  and a constant  $C_{a,\alpha} \in (0, \infty)$ . The end of the proof is exactly the same as in the proof of [2, Theorem 2.1], except that we use the left factorization identity

$$\mathcal{R}_\Lambda(z) = \mathcal{R}_{\mathcal{B}}(z) - \mathcal{R}_{\mathcal{B}}(z)\mathcal{A}\mathcal{R}_\Lambda(z)$$

and the left factorization Duhamel formula

$$S_\Lambda = S_\mathcal{B} + S_\mathcal{B}\mathcal{A} * S_\Lambda$$

instead of the right factorization ones.

The other parts of the proof are unchanged.  $\square$

Next, the proofs of Corollary E.2.4, [2, Corollary 2.5], Theorem E.3.1, Theorem E.3.3 and Theorem E.5.3 are exactly the same as the original ones except that we use Theorem E.2.1 instead of [2, Theorem 2.1].

For the fragmentation equation, assumption **(h2)** is fulfilled for the same splitting  $\Lambda = \mathcal{A} + \mathcal{B}$  as introduced in [2, Section 4] because of [2, Lemma 4.8] and [2, Lemma 4.9]. For the fragmentation equation, the proof of [2, Theorem 1.1] is then unchanged (using the erratum versions of the abstract spectral analysis results).

We finally consider the age structured population equation [2, (1.42)-(1.44)] and we recall that it equivalently writes

$$\partial_t f = \Lambda f = \mathcal{A}f + \mathcal{B}f,$$

with  $\mathcal{A}$  and  $\mathcal{B}$  defined on  $X := M^1(\mathbb{R})$  by

$$\begin{aligned} (\mathcal{A}f)(x) &:= \delta_{x=0} \int_0^\infty K(y) f(y) dy \\ (\mathcal{B}f)(x) &:= -\partial_x f(x) - f(x), \end{aligned}$$

with  $K \in C_b^1(\mathbb{R}_+) \cap L^1(\mathbb{R}_+)$ ,  $\|K\|_{L^1} > 1$ . We observe that because of the regularity assumption made on  $K$ , we have  $\mathcal{A} \in \mathcal{B}(X_{-1}, X)$ . On the other hand,  $\mathcal{B}$  is  $-1$ -dissipative in  $X$  from [2, Lemma 4.13]. Both together, we deduce that

$$\forall t \geq 0, \quad \|S_\mathcal{B}(t)\mathcal{A}\|_{\mathcal{B}(X_{-1}, X)} \leq C e^{-t},$$

for a constant  $C \in (0, \infty)$ , and then that condition **(h2')** holds true. For the age structured population equation, the proof of [2, Theorem 1.1] is then a consequence of the new abstract spectral analysis results Theorem E.3.3 and Theorem E.5.3 exactly as explained in [2, Section 4.3] and [2, Section 6.1]. More developments on the spectral analysis of the age structured population model are presented in [3, 1].

## REFERENCES

- [1] MISCHLER, S. Semigroups in Banach spaces. Work in progress.
- [2] MISCHLER, S., AND SCHER, J. Spectral analysis of semigroups and growth-fragmentation equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 33, 3 (2016), 849–898.
- [3] MISCHLER, S., AND WENG, Q. Relaxation in time elapsed neuron network models in the weak connectivity regime. (2015) hal-01148645

STÉPHANE MISCHLER  
UNIVERSITÉ PARIS-DAUPHINE, PSL RESEARCH UNIVERSITY,  
CNRS, UMR [7534], CEREMADE,  
PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY  
75775 PARIS CEDEX 16 FRANCE  
E-MAIL: [mischler@ceremade.dauphine.fr](mailto:mischler@ceremade.dauphine.fr)