



HAL
open science

On the utility of neural perturbation experiments for identifying the neural components of an integrated system.

Laurent Goffart

► To cite this version:

Laurent Goffart. On the utility of neural perturbation experiments for identifying the neural components of an integrated system.. Multiscale analysis of neural systems: taking the challenge SERIOUSLY, Feb 2010, Gif-sur-Yvette, France. ⟨hal-01440282⟩

HAL Id: hal-01440282

<https://hal.science/hal-01440282v1>

Submitted on 29 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

On the utility of neural perturbation experiments for identifying the neural components of an integrated system.



Laurent Goffart, CNRS


Institut des Neurosciences Cognitives de la Méditerranée
UMR 6193 CNRS – Aix Marseille Universités, Marseille, France

Contact : laurent.goffart @incm.cnrs-mrs.fr


Living animals are sensorimotor systems endowed of properties that allow them to **explore** and interact with their environment. The orienting reaction is one of these systems by which animals can establish **equilibrium** between their endogen properties and the constraints imposed by the milieu. Indeed, variations in the environment (like the sudden appearance of an event) or fluctuations in the inner state (e.g., the hypovolemia) often **break** this equilibrium and lead to coordinated movements which are aimed at **restoring** it.

The orienting reaction can be viewed as a **transition**, whose detailed analysis can provide a picture of the dynamic properties of a neurobiological system, at different **time scales** (e.g., from the systematic orientation to its habituation) and **spatial scales** (e.g., from the saccade and its underlying neural network to the bistable properties of recruited neurons). Most often, this reaction consists in orienting the sensory organs toward the **source** of external events.


Many neuronal groups are involved in generating a reaction that **fits** with the spatiotemporal properties of targeted events. These groups are distributed in several brain regions whose interactions generate streams which are more or less **independent** or **cooperative**. Among those regions, the medio-posterior cerebellum (MPC) is known to play a major role in the **adaptability** of the orienting reaction.




Constant error in the cat



Gaze (eye + head) dysmetria



Eye-head coupling



Cerebellar control of foveating saccades










FIG. 10. Scheme by which the saccade-related neurons in the cFN (SRFN) influence the generation of the horizontal component of saccades. See text for additional details. Arrows, excitatory connections; connecting lines ended by a circle, inhibitory connections. The connecting lines have different thickness for clarity purposes.




Subcortical control of foveation



Bilateral and distributed encoding of foveal target position



A small injection of muscimol in the rostral SC also causes a fixation offset



Predictions : inactivation of IBN area should lead to a **fixation offset** because of dysmetric fixational saccades (no shift)

Role of pause neurons : microstimulation of OPN region suppresses the generation of saccades (Keller 1976)
Its inactivation does not alter fixation (Kaneko 1996) → Test of H-V coupling.

On-going projects

- Subcortical control of fixational saccades in the non-human primate (NHP)** : cerebellar control : with Ulrich Büttner (Ludwig-Maximilians-University München, Germany); collicular control : with Rich Krauzlis (Salk Institute for Biological Studies, San Diego, USA).
- Control of head movements and its coupling with the saccade oculomotor control in the NHP** : cerebellar control : with Julie Quinet (Univ. Rochester, USA); head motor primitives : with Longtang Chen (Univ. Texas Medical Branch, Galveston, USA); neuro-anatomy of eye and head motor networks : with Patrice Coulon et Hélène Bras (P3M, France)
- Control of saccades toward a moving target in the NHP** : cerebellar control : with Kikuro Fukushima (Hokkaido Univ. School of Medicine, Sapporo, Japan)
- Development of an invertebrate neurobiological model (praying mantis)** : behavioral study : with Samuel Rossal (Institut für Biologie, Freiburg, Allemagne); with Yoshitomi Yamawaki (Fukuoka University, Fukuoka, Japan).

Selected publications

- Goffart L. & Pélisson D. Cerebellar contribution to the spatial encoding of orienting gaze shifts in the head-free cat. *Journal of Neurophysiology* 72 : 2547-2550, 1994.
- Goffart L. & Pélisson D. Changes in initiation of orienting gaze shifts after muscimol inactivation of the caudal fastigial nucleus in the cat. *Journal of Physiology* 503.3 : 657-671, 1997.
- Goffart L. & Pélisson D. Orienting gaze shifts during muscimol inactivation of the caudal fastigial nucleus in the cat. I. Gaze dysmetria. *Journal of Neurophysiology* 79 : 1942-1958, 1998.
- Goffart L., Pélisson D. & Guillaume A. Orienting gaze shifts during muscimol inactivation of the caudal fastigial nucleus in the cat. II. Dynamics and eye-head coupling. *Journal of Neurophysiology* 79 : 1959-1976, 1998.
- Goffart L., Guillaume A. & Pélisson D. Compensation for gaze perturbations during inactivation of the caudal fastigial nucleus in the monkey. *Annals of the New York Academy of Sciences* 1004 : 220-228, 2003.
- Goffart L., Chen L.L. & Sparks D.L. Deficits in saccades and fixation during muscimol inactivation of the caudal fastigial nucleus in the rhesus monkey. *Journal of Neurophysiology* 92 : 3351-3367, 2004.
- Quinet J. & Goffart L. Saccadic dysmetria in head-unrestrained gaze shifts after muscimol inactivation of the caudal fastigial nucleus in the monkey. *Journal of Neurophysiology* 93 : 2343-2349, 2005.
- Quinet J. & Goffart L. Head unrestrained gaze shifts after muscimol injection in the caudal fastigial nucleus of the monkey. *Journal of Neurophysiology* 98 : 3269-3283, 2007.
- Goffart L. Saccadic eye movements. In : Squire L. (Ed.) *New Encyclopedia of Neuroscience*, Academic Press, Oxford, 2008.
- Hafed Z., Goffart L. & Krauzlis R. Superior colliculus inactivation causes stable offsets in eye position during tracking. *Journal of Neuroscience* 28 : 8124-8137, 2008.
- Hafed Z.M., Goffart L. & Krauzlis R.J. A neural mechanism for microsaccade generation in the primate superior colliculus. *Science* 323 : 940-943, 2009.
- Quinet J. & Goffart L. Electrical microstimulation of the fastigial oculomotor region in the head unrestrained monkey. *Journal of Neurophysiology* 102 : 320-336, 2009.
- Guerrasio L., Quinet J., Büttner U. & Goffart L. The fastigial oculomotor region and the control of foveation during fixation. *Journal of Neurophysiology*, 2010 (in press)