



HAL
open science

DEGREES OF ITERATES OF RATIONAL MAPS ON NORMAL PROJECTIVE VARIETIES

Nguyen-Bac Dang

► **To cite this version:**

Nguyen-Bac Dang. DEGREES OF ITERATES OF RATIONAL MAPS ON NORMAL PROJECTIVE VARIETIES . 2017. hal-01444252v2

HAL Id: hal-01444252

<https://hal.science/hal-01444252v2>

Preprint submitted on 29 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DEGREES OF ITERATES OF RATIONAL MAPS ON NORMAL PROJECTIVE VARIETIES

NGUYEN-BAC DANG

ABSTRACT. Let X be a normal projective variety defined over an algebraically closed field of arbitrary characteristic. We study the sequence of intermediate degrees of the iterates of a dominant rational selfmap of X , recovering former results by Dinh, Sibony [DS05b], and by Truong [Tru16]. Precisely, we give a new proof of the submultiplicativity properties of these degrees and of their birational invariance. Our approach exploits intensively positivity properties in the space of numerical cycles of arbitrary codimension. In particular, we prove an algebraic version of an inequality first obtained by Xiao [Xia15] and Popovici [Pop16], which generalizes Siu's inequality to algebraic cycles of arbitrary codimension. This allows us to show that the degree of a map is controlled up to a uniform constant by the norm of its action by pull-back on the space of numerical classes in X .

CONTENTS

Introduction	2
Organization of the paper	6
Acknowledgements	6
1. Chow group	6
1.1. General facts	6
1.2. Intersection with Cartier divisors	8
1.3. Characteristic classes	8
2. Space of numerical cycles	9
2.1. Definitions	9
2.2. Algebra structure on the space of numerical cycles	12
2.3. Pullback on dual numerical classes	14
2.4. Canonical morphism	14
2.5. Numerical spaces are finite dimensional	16
3. Positivity	17
3.1. Pseudo-effective and numerically effective cones	17
3.2. Pliant classes	17
3.3. Basepoint free cone on normal projective varieties	18
3.4. Siu's inequality in arbitrary codimension	21
3.5. Norms on numerical classes	23
4. Relative numerical classes	25
4.1. Relative classes	25
4.2. Pullback and pushforward	27
4.3. Restriction to a general fiber and relative canonical morphism	28
5. Application to dynamics	30
5.1. Degrees of rational maps	30
5.2. Sub-multiplicativity	31

Date: November 29, 2017.

Research supported by the ERC grant Nonarcomp no. 307856.

5.3. Norms of operators associated to rational maps	32
6. Semi-conjugation by dominant rational maps	34
7. Mixed degree formula	36
7.1. Positivity estimate of the diagonal	36
7.2. Submultiplicativity of mixed degrees	41
7.3. Proof of Theorem 4	42
8. Kähler case	44
Appendix A. Comparison with Fulton's approach	45
References	47

INTRODUCTION

Let $f : X \dashrightarrow X$ be any dominant rational self-map of a normal projective variety X of dimension n defined over an algebraically closed field κ of arbitrary characteristic. Given any big and nef (e.g ample) Cartier divisor ω_X on X , and any integer $0 \leq k \leq n$, one defines the k -th degree of f as the intersection product:

$$\deg_{k,H_X}(f) = (\pi_1^* H_X^{n-k} \cdot \pi_2^* H_X^k),$$

where π_1 and π_2 are the projections from the graph of f in $X \times X$ onto the first and the second factor respectively.

The behaviour of the sequence of degrees $(\deg_{k,H_X}(f^p))_p$ and especially its growth when $p \rightarrow +\infty$ is of crucial importance in the study of the dynamics of f . It controls the topological entropy of f ([Yom87], [Gro87], [DS05b]), and the growth of the number of isolated periodic points when the period goes to infinity ([DNT16]). Its understanding is also the key for the construction of ergodic invariant measures ([BS92], [Sib99],[DS05a], [Gue05], [DTV10], [DS10], [DDG11]). The growth of the degrees appears as an essential tool when exploring algebraic properties of groups of birational transformations of projective varieties (see [Giz80], [DF01], [Can11], [BD15], [BC16] for surfaces, [DS04a] for the study of commutative automorphism groups in dimension ≥ 3 and [CZ12], [Zha14] for some characterizations of positive entropy automorphisms in higher dimension).

Our main theorem can be stated as follows.

Theorem 1. *Let X be a normal projective variety of dimension n and let H_X be a big and nef Cartier divisor on X .*

- (i) *There is a positive constant $C > 0$ such that for any dominant rational self-maps f, g on X , one has:*

$$\deg_{k,H_X}(f \circ g) \leq C \deg_{k,H_X}(f) \deg_{k,H_X}(g).$$

- (ii) *For any big nef Cartier divisor H'_X on X , there exists a constant $C > 0$ such that for any rational self-map f on X , one has:*

$$\frac{1}{C} \leq \frac{\deg_{k,H_X}(f)}{\deg_{k,H'_X}(f)} \leq C.$$

Observe that Theorem 1.(ii) implies that the degree growth of f is a birational invariant, in the sense that there is a positive constant C such that for any birational map $g : X' \dashrightarrow X$ with X' projective, and any big nef Cartier divisor $H_{X'}$ on X' , one has

$$\frac{1}{C} \leq \frac{\deg_{k,H_X}(f^p)}{\deg_{k,H_{X'}}(g^{-1} \circ f^p \circ g)} \leq C,$$

for any $p \in \mathbb{N}$. Fekete's lemma and Theorem 1.(i) also imply the existence of the dynamical degree (first introduced in [RS97] for rational maps of the projective space) as the following quantity:

$$\lambda_k(f) := \lim_{p \rightarrow +\infty} \deg_{k, H_X}(f^p)^{1/p} .$$

The independence of $\lambda_k(f)$ under the choice of H_X , and its birational invariance are the consequence of Theorem 1.(ii) .

When $\kappa = \mathbb{C}$, Theorem 1 was proved by Dinh and Sibony in [DS05b], and further generalized to compact Kähler manifolds in [DS04b]. The core of their argument relied on a procedure of regularization for closed positive currents of any bidegree ([DS04b, Theorem 1.1]) and was therefore transcendental in nature. By the Lefschetz principle Dinh and Sibony's argument also proves that the k -th dynamical degree of any rational dominant map is well-defined when κ is a field of characteristic zero. Recently, Truong [Tru15] managed to get around this problem and proved Theorem 1 for arbitrary smooth varieties using an appropriate Chow-type moving lemma. He went further in [Tru16] and obtained Theorem 1 for any normal variety in all characteristic by applying de Jong's alteration theorem ([Jon96]). Note however that he had to deal with correspondences since a rational self-map can only be lifted as a correspondence through a general alteration map. Our approach avoids this technical difficulty.

To illustrate our method, let us explain the proof of Theorem 1, when X is smooth, $k = 1$ and f, g are regular following the method initiated in [BFJ08, Proposition 3.1]. Recall that a divisor α on X is *pseudo-effective* and one writes $\alpha \geq 0$ if for any ample Cartier divisor H on X , and any rational $\epsilon > 0$, a suitable multiple of the \mathbb{Q} -divisor $\alpha + \epsilon H$ is linearly equivalent to an effective one.

Recall also the fundamental Siu inequality¹ ([Laz04, Theorem 2.2.13], [Cut15]) which states:

$$\alpha \leq n \frac{(\alpha \cdot \beta^{n-1})}{(\beta^n)} \beta, \quad (1)$$

for any nef divisor α , and any big and nef divisor β .

Since the pullback by a dominant morphism of a big nef divisor remains big and nef, we may apply (1) to $\alpha = g^* f^* H_X$ and $\beta = f^* H_X$, and we get

$$g^* f^* H_X \leq n \frac{\deg_{k, H_X}(f)}{(H_X^n)} g^* H_X .$$

Intersecting with the cycle H_X^{n-1} yields the submultiplicativity of the degrees with the constant $C = n/(H_X^n)$.

We observe that the previous inequality (1) can be easily extended to complete intersections by cutting out by suitable ample sections. In particular, we get a positive constant C such that for any big nef divisors α and β , one has:

$$\alpha^k \leq C \frac{(\alpha^k \cdot \beta^{n-k})}{(\beta^n)} \beta^k . \quad (2)$$

Such inequalities have been obtained by Xiao ([Xia15]) and Popovici ([Pop16]) in the case $\kappa = \mathbb{C}$. Their proof uses the resolution of complex Monge-Ampère equations and yields a constant $C = \binom{k}{n}$. On the other hand, our proof applies in arbitrary characteristic and in fact to more general classes than complete intersection ones. We refer to Theorem 3 below and the discussion preceding it for more details. Note however that we only obtain $C = (n - k + 1)^k$, far from the expected optimal constant of Popovici and Xiao. Once (2) is proved, Theorem 1 follows by a similar argument as in the case $k = 1$.

¹this inequality is also referred to as the weak transcendental holomorphic Morse inequality in [LX15]

Going back to the case where X is a complex smooth projective variety, recall that the degree of f is controlled up to a uniform constant by the norm of the linear operator $f^{\bullet,k}$, induced by pullback on the de Rham cohomology space $H_{dR}^{2k}(X)_{\mathbb{R}}$ ([DS05b, Lemma 4]). One way to construct $f^{\bullet,k}$ is to use the Poincaré duality isomorphisms $\psi_X : H_{dR}^{2k}(X, \mathbb{R}) \rightarrow H_{2n-2k}(X, \mathbb{R})$, $\psi_{\Gamma_f} : H_{dR}^{2k}(\Gamma_f, \mathbb{R}) \rightarrow H_{2n-2k}(\Gamma_f, \mathbb{R})$ where $H_k(X, \mathbb{R})$ denotes the k -th simplicial homology group of X . The operator $f^{\bullet,k}$ is then defined following the commutative diagram below:

$$\begin{array}{ccccc} H_{dR}^{2k}(\Gamma_f, \mathbb{R}) & \xrightarrow{\psi_{\Gamma_f}} & H_{2n-2k}(\Gamma_f, \mathbb{R}) & \xrightarrow{\pi_{1*}} & H_{2n-2k}(X, \mathbb{R}) \\ \pi_2^* \uparrow & & & & \downarrow \psi_X^{-1} \\ H_{dR}^{2k}(X, \mathbb{R}) & \xrightarrow{f^{\bullet,k}} & & & H_{dR}^{2k}(X, \mathbb{R}), \end{array}$$

where Γ_f is a desingularization of the graph of f in $X \times X$, and π_1, π_2 are the projections from Γ_f onto the first and second factor respectively.

To generalize this control to our setting we need to find a replacement for the de Rham cohomology group $H_{dR}^{2k}(X)_{\mathbb{R}}$ and define suitable pullback operators. When X is smooth, one natural way to proceed is to consider the spaces $N^k(X)_{\mathbb{R}}$ of algebraic \mathbb{R} -cycles of codimension k modulo numerical equivalence. The operator $f^{\bullet,k}$ is then simply given by the composition $\pi_{1*} \circ \pi_2^* : N^k(X)_{\mathbb{R}} \rightarrow N^k(X)_{\mathbb{R}}$.

When X is singular, then the situation is more subtle because one cannot intersect arbitrary cycle classes in general ². One can consider two natural spaces of numerical cycles $N^k(X)_{\mathbb{R}}$ and $N_k(X)_{\mathbb{R}}$ on which pullback operations and pushforward operations by proper morphisms are defined respectively. More specifically, the *space of numerical k -cycles* $N_k(X)_{\mathbb{R}}$ is defined as the group of \mathbb{R} -cycles of dimension k modulo the relation $z \equiv 0$ if and only if $(p^*z \cdot D_1 \cdot \dots \cdot D_{e+k}) = 0$ for any proper flat surjective map $p : X' \rightarrow X$ of relative dimension e and any Cartier divisors D_i on X' . One can prove that $N_k(X)_{\mathbb{R}}$ is a finite dimensional vector space and one defines $N^k(X)_{\mathbb{R}}$ as its dual $\text{Hom}(N_k(X)_{\mathbb{R}}, \mathbb{R})$.

Note that our presentation differs slightly from Fulton's definition (see Appendix A for a comparison), but we also recover the main properties of the numerical groups.

As in the complex case, we are able to construct Poincaré duality maps $\psi_X : N^k(X)_{\mathbb{R}} \rightarrow N_{n-k}(X)_{\mathbb{R}}$ and $\psi_{\Gamma_f} : N^k(\Gamma_f)_{\mathbb{R}} \rightarrow N_{n-k}(\Gamma_f)_{\mathbb{R}}$, but they are not necessarily isomorphisms due to the presence of singularities. As a consequence, we are only able to define a linear map $f^{\bullet,k}$ as $f^{\bullet,k} := \pi_{1*} \circ \psi_{\Gamma_f} \circ \pi_2^* : N^k(X)_{\mathbb{R}} \rightarrow N_{n-k}(X)_{\mathbb{R}}$ between two distinct vector spaces. Despite this limitation, we prove a result analogous to one of Dinh and Sibony. The next theorem was obtained by Truong for smooth varieties ([Tru16, Theorem 1.1.(5)]).

Theorem 2. *Let X be a normal projective variety of dimension n . Fix any norms on $N^k(X)_{\mathbb{R}}$ and $N_{n-k}(X)_{\mathbb{R}}$, and denote by $\|\cdot\|$ the induced operator norm on linear maps from $N^k(X)_{\mathbb{R}}$ to $N_{n-k}(X)_{\mathbb{R}}$. Then there is a constant $C > 0$ such that for any rational selfmap $f : X \dashrightarrow X$, one has:*

$$\frac{1}{C} \leq \frac{\|(f)^{\bullet,k}\|}{\text{deg}_{k,H_X}(f)} \leq C. \quad (3)$$

Our proof of Theorem 2 exploits a natural notion of positive classes in $N^k(X)_{\mathbb{R}}$ combined with a strengthening of (2) to these classes that we state below (see Theorem 3).

²an arbitrary curve can only be intersected with a Cartier divisor, not with a general Weil divisor.

To simplify our exposition, let us suppose again that X is smooth. As in codimension 1, one can define the *pseudo-effective cone* $\text{Psef}^k(X)$ as the closure in $N^k(X)_{\mathbb{R}}$ of the cone generated by effective cycles of codimension k . Its dual with respect to the intersection product is the *nef cone* $\text{Nef}^{n-k}(X)$, which however does not behave well when $k \geq 2$ (see [DELV11]). Some alternative notions of positive cycles have been introduced by Fulger and Lehmann in [FL14b], among which the notion of basepoint free classes emerges. Basepoint free classes have many good properties such as being both pseudo-effective and nef, being invariant by pull-backs by morphisms and by intersection products, and forming a salient convex cone with non-empty interior. Denote by $\text{BPF}^k(X)$ the cone of basepoint free classes. It is defined as the closure in $N^k(X)_{\mathbb{R}}$ of the cone generated by \mathbb{R} -cycles of the form $p_*(D_1 \cdot \dots \cdot D_{e+k})$ where D_i are ample Cartier \mathbb{R} -divisors and $p : X' \rightarrow X$ is a flat surjective proper morphism of relative dimension e .

For basepoint free classes, we are able to prove the following generalization of (2).

Theorem 3. *Let X be a normal projective variety of dimension n . Then there exists a constant $C > 0$ such that for any basepoint free class $\alpha \in \text{BPF}^k(X)$, for any big nef divisor β , one has in $N^k(X)_{\mathbb{R}}$:*

$$\alpha \leq C \frac{(\alpha \cdot \beta^{n-k})}{(\beta^n)} \times \beta^k. \quad (4)$$

Theorem 2 follows from (4) by observing that $f^{\bullet,k} \text{BPF}^k(X) \subset \text{Psef}^k(X)$, so that the operator norm $\|f^{\bullet,k}\|$ can be computed by evaluating $f^{\bullet,k}$ only on basepoint free classes.

In the singular case, the proof of Theorem 2 is completely similar but the spaces $N^k(X)_{\mathbb{R}}$ and $N_{n-k}(X)_{\mathbb{R}}$ are not necessarily isomorphic in general. As a consequence, several dual notions of positivity appear in $N^k(X)_{\mathbb{R}}$ and $N_k(X)_{\mathbb{R}}$ that make the arguments more technical.

Finally, using the techniques developed in this paper, we give a new proof of the product formula of Dinh, Nguyen, Truong ([DN11, Theorem 1.1],[DNT12, Theorem 1.1]) which they proved when $\kappa = \mathbb{C}$ and which was later generalized by Truong ([Tru16, Theorem 1.1.(4)]) to normal projective varieties over any field.

The setup is as follows. Let $q : X \rightarrow Y$ be any proper surjective morphism between normal projective varieties, and fix two big and nef divisors H_X, H_Y on X and Y respectively. Consider two dominant rational self-maps $f : X \dashrightarrow X, g : Y \dashrightarrow Y$, which are semi-conjugated by q , i.e. which satisfy $q \circ f = g \circ q$. To simplify notation we shall write $X/qY \xrightarrow[f]{g} X/qY$ when these assumptions hold true.

Recall that the k -th relative degree of $X/qY \xrightarrow[f]{g} X/qY$ is given by the intersection product

$$\text{reldeg}_k(f) := (\pi_1^*(H_X^{\dim X - \dim Y - k} \cdot q^* H_Y^{\dim Y}) \cdot \pi_2^* H_X^k),$$

where π_1 and π_2 are the projections from the graph of f in $X \times X$ onto the first and the second component respectively. One can show a relative version of Theorem 1 (see Theorem 5.2.1), and define as in the absolute case, the k -th relative dynamical degree $\lambda_k(f, X/Y)$ as the limit $\lim_{p \rightarrow +\infty} \text{reldeg}_k(f^p)^{1/p}$. It is also a birational invariant, and does not depend on the choices of H_X and H_Y . When $q : X \dashrightarrow Y$ is merely rational and dominant, then we define (see Section 6) the k -th relative degree of f by replacing X with the normalization of graph of q . We prove the following theorem.

Theorem 4. *Let X, Y be normal projective varieties. For any dominant rational self-maps $f : X \dashrightarrow X, g : Y \dashrightarrow Y$ which are semi-conjugated by a dominant rational map $q : X \dashrightarrow Y$, we have*

$$\lambda_k(f) = \max_{\max(0, k-l) \leq j \leq \min(k, e)} (\lambda_{k-j}(g) \lambda_j(f, X/Y)). \quad (5)$$

Observe that we are only able to relate the dynamical degrees of a rational map with the relative dynamical degrees and the dynamical degree on the base. Unfortunately, we do not know how to control the degree growth of f with respect to the degree growth of g on the base Y and on the fibers of $q : X \rightarrow Y$ except when $e = 0$, since in that case Theorem 1.(2) proves that $1/C \leq \deg_{k,H_X}(f)/\deg_{k,H_Y}(g) \leq C$.

Our proof follows closely Dinh and Nguyen's method from [DN11] and relies on a fundamental inequality (see Corollary 7.1.5 below) which follows from Künneth formula at least when $\kappa = \mathbb{C}$. To state it precisely, consider $\pi : X' \rightarrow X$ a surjective generically finite morphism and $q : X \rightarrow Y$ a surjective morphism where X', X and Y are normal projective varieties such that $n = \dim X = \dim X'$ and such that $l = \dim Y$. We prove that for any basepoint free classes $\alpha \in \text{BPF}^k(X')$ and $\beta \in \text{BPF}^{n-k}(X')$, one has:

$$(\beta \cdot \alpha) \leq C \sum_{\max(0, k-l) \leq j \leq \min(k, e)} U_j(\alpha) \times (\beta \cdot \pi^*(q^*H_Y \cdot H_X^j)), \quad (6)$$

where H_Y and H_X are big and nef divisors on Y and X respectively, and $U_j(\alpha)$ is the intersection product given by $U_j(\alpha) = (\pi^*(q^*H_Y^{l-k+j} \cdot H_X^{e-j}) \cdot \alpha)$.

In the singular case, Truong has obtained this inequality using Chow's moving intersection lemma. We replace this argument by a suitable use of Siu's inequality and Theorem 3 in order to prove a positivity property for a class given by the difference between a basepoint free class in $X' \times X'$ and the fundamental class of the diagonal of X' in $X' \times X'$ (see Theorem 7.1.1). Inequality (6) is a weaker version of [DN11, Proposition 2.3] proved by Dinh-Nguyen when Y is a complex projective variety, and was extended to a field arbitrary characteristic by Truong when Y is smooth ([Tru16, Lemma 4.1]).

Organization of the paper. In the first Sections 1 and 2, we review the background on the Chow groups and recall the definitions of the spaces of numerical cycles and provide their basic properties. In §3, we discuss the various notions of positivity of cycles and prove Theorem 3. In §4, we define relative numerical cycles and canonical morphisms which are the analogous to the Poincaré morphisms ψ_X in a relative setting. In §5, we prove Theorem 1, Theorem 2 and Theorem 4. Finally we give an alternate proof of Dinh-Sibony's theorem in the Kähler case ([DS05b, Proposition 6]) in §8 using Popovici [Pop16] and Xiao's inequality [Xia15]. Note that these inequalities allow us to avoid regularization techniques of closed positive currents but rely on a deep theorem of Yau. In Appendix A, we prove that our presentation and Fulton's definition of numerical cycles are equivalent, hence proving that any numerical cycles can be pulled back by a flat morphism.

Acknowledgements. Firstly, I would like to thank my advisor C. Favre for his patience and our countless discussions on this subject. I thank also S. Boucksom for some helpful discussions and for pointing out the right argument for the appendix, T. Truong, L. Fantini, J. Xie, M. Fulger, B. Lehmann and R. Mboro for their precious comments on my previous drafts and for providing me some references.

1. CHOW GROUP

1.1. General facts. Let X be a normal projective variety of dimension n defined over an algebraically closed field κ of arbitrary characteristic.

The space of cycles $Z_k(X)$ is the free abelian group generated by irreducible subvarieties of X of dimension k , and $Z_k(X)_{\mathbb{Q}}, Z_k(X)_{\mathbb{R}}$ will denote the tensor products $Z_k(X) \otimes_{\mathbb{Z}} \mathbb{Q}$ and $Z_k(X) \otimes_{\mathbb{Z}} \mathbb{R}$.

Let $q : X \rightarrow Y$ be a morphism where Y is a normal projective variety. Since X and Y are respectively projective, the map q is proper. Following [Ful98], we define the proper pushforward of the cycle $[V] \in Z_k(X)$ as the element of $Z_k(Y)$ given by:

$$q_*[V] = \begin{cases} 0 & \text{if } \dim(q(V)) < \dim V \\ [\kappa(\eta) : \kappa(q(\eta))] \times [q(V)] & \text{if } \dim V = \dim(q(V)), \end{cases}$$

where V is an irreducible subvariety of X of dimension k , η is the generic point of V and $\kappa(\eta)$, $\kappa(q(\eta))$ are the residue fields of the local rings \mathcal{O}_η and $\mathcal{O}_{q(\eta)}$ respectively. We extend this map by linearity and obtain a morphism of abelian groups $q_* : Z_k(X) \rightarrow Z_k(Y)$.

Let C be any closed subscheme of X of dimension k and denote by C_1, \dots, C_r its k -dimensional irreducible components. Then C defines a fundamental class $[C] \in Z_k(X)$ by the following formula:

$$[C] := \sum_{i=1}^r l_{\mathcal{O}_{C_i, C}}(\mathcal{O}_{C_i, C})[C_i],$$

where $l_A(M)$ denotes the length of an A -module M ([Eis95, section 2.4]).

For any flat morphism $q : X \rightarrow Y$ of relative dimension e between normal projective varieties, we can define a flat pullback of cycles $q^* : Z_k(Y) \rightarrow Z_{k+e}(X)$ (see [Ful98, section 1.7]). If C is any subscheme of Y of dimension k , the cycle $q^*[C]$ is by definition the fundamental class of the scheme-theoretic inverse by q :

$$q^*[C] := [q^{-1}(C)] \in Z_{k+e}(X).$$

Let W be a subvariety of X of dimension $k+1$ and φ be a rational map on W . Then we define a cycle on X by:

$$[\text{div}(\varphi)] := \sum \text{ord}_V(\varphi)[V],$$

where the sum is taken over all irreducible subvarieties V of dimension k of X . A cycle α defined this way is rationally equivalent to 0 and in that case we shall write $\alpha \sim 0$.

The k -th Chow group $A_k(X)$ of X is the quotient of the abelian group $Z_k(X)$ by the free group generated by the cycles that are rationally equivalent to zero. We denote by $A_\bullet(X)$ the abelian group $\bigoplus A_k(X)$.

We recall now the functorial operations on the Chow group, which result from the intersection theory developed in [Ful98].

Theorem 1.1.1. *Let $q : X \rightarrow Y$ be a morphism between normal projective varieties. Then we have:*

- (1) *The morphism of abelian groups $q_* : Z_k(X) \rightarrow Z_k(Y)$ induces a morphism of abelian groups $q_* : A_k(X) \rightarrow A_k(Y)$.*
- (2) *If the morphism q is flat of relative dimension e , then the morphism $q^* : Z_k(Y) \rightarrow Z_{k+e}(X)$ induces a morphism of abelian groups $q^* : A_k(Y) \rightarrow A_{k+e}(X)$.*

Assertion (1) is proved in [Ful98, Theorem 1.4] and assertion (2) is given in [Ful98, Theorem 1.7].

Remark 1.1.2. Let $q : X \rightarrow Y$ is a flat morphism of normal projective varieties. Suppose $\alpha \in A_k(Y)$ is represented by an effective cycle $\alpha \sim \sum n_i [V_i]$ where the n_i are positive integers. Then $q^*\alpha$ is also represented by an effective cycle.

Any cycle $\alpha \in Z_0(X)_{\mathbb{Z}}$ is of the form $\sum n_i [p_i]$ with $p_i \in X(\kappa)$ and $n_i \in \mathbb{Z}$. We define the degree of α to be $\text{deg}(\alpha) := \sum n_i$ and we shall write:

$$(\alpha) := \text{deg}(\alpha) = \sum n_i.$$

The morphism of abelian groups $\text{deg} : Z_0(X)_{\mathbb{Z}} \rightarrow \mathbb{Z}$ induces a morphism of abelian groups $\text{deg} : A_0(X) \rightarrow \mathbb{Z}$.

1.2. Intersection with Cartier divisors. Let X be a normal projective variety and D be a Cartier divisor on X . Let V be a subvariety of of dimension k in X and denote by $j : V \hookrightarrow X$ the inclusion of V in X . We define the intersection of D with $[V]$ as the class:

$$D \cdot [V] := j_*[D'] \in A_{k-1}(X),$$

where D' is a Cartier divisor on V such that the line bundles $j^*\mathcal{O}_X(D)$ and $\mathcal{O}_V(D')$ are isomorphic. We extend this map by linearity into a morphism of abelian groups $D \cdot : Z_k(X) \rightarrow A_{k-1}(X)$.

Theorem 1.2.1. *Let X be a normal projective variety and D be a Cartier divisor on X . The map $D \cdot : Z_k(X) \rightarrow A_{k-1}(X)$ induces a morphism of abelian groups $D \cdot : A_k(X) \rightarrow A_{k-1}(X)$. Moreover, the following properties are satisfied:*

(1) *For all Cartier divisors D and D' on X , for all class $\alpha \in A_k(X)$, we have:*

$$(D' + D) \cdot \alpha = D' \cdot \alpha + D \cdot \alpha.$$

(2) *(Projection formula) Let $q : X \rightarrow Y$ be a morphism between normal projective varieties. Then for all class $\beta \in A_k(X)$ and all Cartier divisor D on Y , we have in $A_{k-1}(Y)$:*

$$q_*(q^*D \cdot \beta) = D \cdot q_*(\beta).$$

Definition 1.2.2. *For all normal projective varieties X , the group $\text{Cl}^k(X)$ is the free group generated by elements of the form $D_1 \cdot \dots \cdot D_k$ where D_1, \dots, D_k are Cartier divisors on X .*

1.3. Characteristic classes.

Definition 1.3.1. *Let X be a normal projective variety of dimension n and L be a line bundle on X . There exists a Cartier divisor D on X such that the line bundles L and $\mathcal{O}_X(D)$ are isomorphic. We define the first Chern class of L as:*

$$c_1(L) := [D] \in A_{n-1}(X).$$

Definition 1.3.2. *Let X be a normal projective variety and E be a vector bundle of rank $e + 1$ on X . Given any vector bundle E on X , we shall denote by $\mathbb{P}(E)$ the projective bundle of hyperplanes in E following the convention of Grothendieck. Let p be the projection from $\mathbb{P}(E^*)$ to X and $\xi = c_1(\mathcal{O}_{\mathbb{P}(E^*)}(1))$. We define the i -th Segre class $s_i(E)$ as the morphism $s_i(E) \lrcorner : A_{\bullet}(X) \rightarrow A_{\bullet-i}(X)$ given by:*

$$s_i(E) \lrcorner \alpha := p_*(\xi^{e+i} \cdot p^*\alpha). \quad (7)$$

Remark 1.3.3. When X is smooth of dimension n , we can define an intersection product on the Chow groups $A_k(X) \times A_l(X) \rightarrow A_{n-k-l}(X)$ (see [Ful98, Definition 8.1.1]) which is compatible with the intersection with Cartier divisors and satisfies the projection formula (see [Ful98, Example 8.1.7]). Applying the projection formula to (7), we get

$$s_i(E) \lrcorner \alpha = p_*(\xi^{e+i}) \cdot \alpha,$$

so that $s_i(E)$ is represented by an element in $A_{n-i}(X)$. To simplify we shall also denote $s_i(E)$ this element.

As Segre classes of vector bundles are operators on the Chow groups $A_{\bullet}(X)$, the composition of such operators defines a product.

Theorem 1.3.4. *(cf [Ful98, Proposition 3.1]) Let $q : X \rightarrow Y$ be a morphism between normal projective varieties. For any vector bundle E and F on Y , the following properties hold.*

(1) *For all $\alpha \in A_k(Y)$ and all $i < 0$, we have $s_i(E) \lrcorner \alpha = 0$.*

- (2) For all $\alpha \in A_k(Y)$, we have $s_0(E) \lrcorner \alpha = \alpha$.
- (3) For all integers i, j , we have $s_i(E) \lrcorner (s_j(F) \lrcorner \alpha) = s_j(F) \lrcorner (s_i(E) \lrcorner \alpha)$.
- (4) (Projection formula) For all $\beta \in A_k(X)$ and any integer i , we have $q_*(s_i(q^*E) \lrcorner \beta) = s_i(E) \lrcorner q_*\beta$.
- (5) If the morphism $q : X \rightarrow Y$ is flat, then for all $\alpha \in A_k(Y)$ and any integer i , we have $s_i(q^*E) \lrcorner q^*\alpha = q^*(s_i(E) \lrcorner \alpha)$.

The i -th Chern class $c_i(E)$ of a vector bundle E on X is an operator $c_i(E) : A_\bullet(X) \rightarrow A_{\bullet-i}$ defined formally as the coefficients in the inverse power series:

$$(1 + s_1(E)t + s_2(E)t^2 + \dots)^{-1} = 1 + c_1(E)t + \dots + c_{r+1}(E)t^{r+1}.$$

A direct computation yields for example $c_1(E) = -s_1(E)$, $c_2(E) = (s_1(E)^2 - s_2(E))$.

Definition 1.3.5. Let X be a normal projective variety. The abelian group $A^k(X)$ is the subgroup of $\text{Hom}(A_\bullet(X), A_{\bullet-k}(X))$ generated by product of Chern classes $c_{i_1}(E_1) \cdots c_{i_p}(E_p)$ where i_1, \dots, i_p are integers satisfying $i_1 + \dots + i_p = k$ and where E_1, \dots, E_p are vector bundles over X . We denote by $A^\bullet(X)$ the group $\bigoplus A^k(X)$.

Observe that by definition, $A^k(X)$ contains the image of $\text{Cl}^k(X)$.

Recall that the Grothendieck group $K^0(X)$ is the free group generated by vector bundles on X quotiented by the subgroup generated by relations of the form $[E_1] + [E_3] - [E_2]$ where there is an exact sequence of vector bundles:

$$0 \longrightarrow E_1 \longrightarrow E_2 \longrightarrow E_3 \longrightarrow 0.$$

Moreover, the group $K^0(X)$ has a structure of rings given by the tensor product of vector bundles.

Recall also that the Chern character is the unique morphism of rings $\text{ch} : (K^0(X), +, \otimes) \rightarrow (A^\bullet(X), +, \cdot)$ satisfying the following properties (see [Ful98, Example 3.2.3]).

- (1) If L is a line bundle on X , then one has:

$$\text{ch}(L) = \sum_{i \geq 0} \frac{c_1(L)^i}{i!}.$$

- (2) For any morphism $q : X' \rightarrow X$ and any vector bundle E on X , we have $q^* \text{ch}(E) = \text{ch}(q^*E)$.

For any vector bundle E on X , we will denote by $\text{ch}_k(E)$ the term in $A^k(X)$ of $\text{ch}(E)$.

We recall Grothendieck-Riemann-Roch's theorem for smooth varieties.

Theorem 1.3.6. (see [Ful98, Corollary 18.3.2]) Let X be a smooth variety. Then the Chern character induces an isomorphism:

$$\text{ch} \lrcorner [X] : E \in K^0(X) \otimes \mathbb{Q} \rightarrow \text{ch}(E) \lrcorner [X] \in A_\bullet(X) \otimes \mathbb{Q}.$$

2. SPACE OF NUMERICAL CYCLES

2.1. Definitions. In all this section, X, Y, X_1, X_2, X_3 and X' are normal projective varieties and X is of dimension n . Two cycles α and β in $Z_k(X)$ are said to be numerically equivalent and we will denote by $\alpha \equiv \beta$ if for all flat morphisms $p_1 : X_1 \rightarrow X$ of relative dimension e and all Cartier divisors D_1, \dots, D_{e+k} in X_1 , we have:

$$(D_1 \cdots D_{e+k} \cdot q^*\alpha) = (D_1 \cdots D_{e+k} \cdot q^*\beta).$$

Definition 2.1.1. The group of numerical classes of dimension k is the quotient $N_k(X) = Z_k(X) / \equiv$.

By construction, the group $N_k(X)$ is torsion free and there is a canonical surjective morphism $A_k(X) \rightarrow N_k(X)$ for any integer k .

Remark 2.1.2. Observe also that for $k = 0$, two cycles are numerically equivalent if and only if they have the same degree. Since smooth points are dense in X (see [Dan94, Theorem II.4.7]) and are of degree 1, this proves that the degree realizes the isomorphism $N_0(X) \simeq \mathbb{Z}$.

We set $N_k(X)_{\mathbb{Q}}$ and $N_k(X)_{\mathbb{R}}$ the two vector spaces obtained by tensoring by \mathbb{Q} and \mathbb{R} respectively.

Remark 2.1.3. This definition allows us to pullback numerical classes by any flat morphism $q : X \rightarrow Y$ of relative dimension e . Our presentation is slightly different from the classical one given in [Ful98, Section 19.1]. We refer to Appendix A for a proof of the equivalence of these two approaches.

Proposition 2.1.4. *Let $q : X \rightarrow Y$ a morphism. Then the morphism of groups $q_* : Z_k(X) \rightarrow Z_k(Y)$ induces a morphism of abelian groups $q_* : N_k(X) \rightarrow N_k(Y)$.*

Proof. Let n be the dimension of X and l be the dimension of Y , and let α be a cycle in $Z_k(X)$ such that α is numerically trivial. We need to prove that $q_*\alpha$ is also numerically trivial.

Take $p_1 : Y_1 \rightarrow Y$ a flat morphism of relative dimension e_1 . Let X_1 be the fibred product $X \times_Y Y_1$ and let p'_1 and q' be the natural projections from X_1 to X and Y_1 respectively.

$$\begin{array}{ccc} X_1 & \xrightarrow{p'_1} & X \\ \downarrow q' & & \downarrow q \\ Y_1 & \xrightarrow{p_1} & Y \end{array}$$

The morphism p'_1 is flat and q' is proper. Pick any cycle γ whose class is in $\text{CI}^{e_1+k}(Y_1)$. We want to prove that $(\gamma \cdot p_1^* q_* \alpha) = 0$. By [Ful98, Proposition 1.7], we have that $p_1^* q_* \alpha = q'_* p_1'^* \alpha$ in $Z_{e_1+k}(Y_1)$. Applying the projection formula, we get:

$$\gamma \cdot p_1^* q_* \alpha = \gamma \cdot q'_* p_1'^* \alpha \sim q'_*(q'^* \gamma \cdot p_1'^* \alpha).$$

Because p'_1 is flat and $q'^* \gamma \in \text{CI}^{e_1+k}(X_1)$, we have $(q'^* \gamma \cdot p_1'^* \alpha) = 0$ so that $(\gamma \cdot p_1^* q_* \alpha) = 0$ as required. \square

The numerical classes defined above are hard to manipulate, we want to define a pullback of numerical classes by any proper morphism. We proceed and define dual classes.

We denote by $Z^k(X) = \text{Hom}_{\mathbb{Z}}(Z_k(X), \mathbb{Z})$ the space of cocycles. If $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 , then any element $\gamma \in \text{CI}^{e_1+k}(X_1)$ induces an element $[\gamma]$ in $Z^k(X)$ by the following formula:

$$[\gamma] : \alpha \in Z_k(X) \rightarrow (\gamma \cdot p_1^* \alpha) \in \mathbb{Z}. \quad (8)$$

Definition 2.1.5. *The abelian group $N^k(X)$ is the subgroup of $Z^k(X)$ generated by elements of the form $[\gamma]$ where $\gamma \in \text{CI}^{e_1+k}(X_1)$ and X_1 is flat over X of relative dimension e_1 .*

Remark 2.1.6. By definition, the map $\text{deg} : Z_0(X) \rightarrow \mathbb{Z}$ is naturally an element of $Z^0(X)$. Moreover, one has using Theorem 1.3.4.(2) that:

$$z \in Z_0(X) \rightarrow (s_0(E) \lrcorner z) = \text{deg}(z) \in \mathbb{Z},$$

for any vector bundle E on X . Hence, deg defines an element of $N^0(X)$ by definition of Segré classes (Definition 7).

Proposition 2.1.7. *By definition of the numerical equivalence relation, any element of $N^k(X)$ induces an element of the dual $\text{Hom}_{\mathbb{Z}}(N_k(X), \mathbb{Z})$. Hence, we can define a natural pairing between $N^k(X)$ and $N_k(X)$. For any normal projective variety, the pairing $N^k(X) \times N_k(X) \rightarrow \mathbb{Z}$ is non degenerate.*

Proof. It follows directly from the definition of $N^k(X)$ and $N_k(X)$. □

A priori, an element of $N^k(X)$ is a combination of elements $[\gamma_1] + [\gamma_2] + \dots + [\gamma_i]$. The following proposition proves one can always take $i = 1$ at least if we tensor all spaces by \mathbb{Q} .

Proposition 2.1.8. *Any element of $N^k(X)$ is induced by $\gamma \in \text{CI}^{e_1+k}(X_1)_{\mathbb{Q}}$ where $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 .*

Proof. By an immediate induction argument, we are reduced to prove the assertion for the sum of two elements $[\gamma_1] + [\gamma_2]$ where $\gamma_i \in \text{CI}^{e_i+k}(X_i)_{\mathbb{Q}}$ and $p_i : X_i \rightarrow X$ are flat morphisms of relative dimension e_1 and e_2 respectively.

Let us consider X' the fibre product $X_1 \times X_2$ over X and p'_i the flat projections from X' to X_i for $i = 1, 2$. By linearity, we only need to show that there exists an element $\gamma'_1 \in \text{CI}^{e_1+e_2+k}(X')$ such that $[\gamma'_1] = [\gamma_1]$ in $N^k(X)$.

$$\begin{array}{ccc}
 & X_1 \times X_2 & \\
 p'_2 \swarrow & & \searrow p'_1 \\
 X_1 & & X_2 \\
 p_1 \searrow & & \swarrow p_2 \\
 & X &
 \end{array}$$

Take an ample Cartier divisor H_{X_2} on X_2 and λ_2 an integer such that $p_{2*}H_{X_2}^{e_2} \sim \lambda_2[X]$. Setting $\gamma'_1 = \frac{1}{\lambda_2} p_1^* H_{X_2}^{e_2} \cdot p_2^* \gamma_1$, we need to prove that for any $\alpha \in Z_k(X)$, one has $(\gamma_1 \cdot p_1^* \alpha) = (\gamma'_1 \cdot p_2^* p_1^* \alpha)$. By [Ful98, Proposition 1.7], we have the equality $p'_{2*} p_1^* H_{X_2}^{e_2} = p_1^* p_{2*} H_{X_2}^{e_2}$ in $Z^{e_2}(X_2)$, hence:

$$p'_{2*} p_1^* H_{X_2}^{e_2} = \lambda_2 p_1^*[X].$$

Since X_1 is reduced and $p_1^*[X]$ is a cycle of codimension 0 in X_1 , we have $p_1^*[X] = [X_1]$. Hence by the projection formula, we have:

$$\begin{aligned}
 \frac{1}{\lambda_2} p'_{2*} (p_2^* (\gamma_1 \cdot p_1^* \alpha) \cdot p_1^* H_{X_2}^{e_2}) &= \frac{1}{\lambda_2} (p_1^* \alpha \cdot \gamma_1) \cdot p_{2*} p_1^* H_{X_2}^{e_2} \\
 &= \frac{1}{\lambda_2} (p_1^* \alpha \cdot \gamma_1) \cdot \lambda_2 [X_1] \\
 &= p_1^* \alpha \cdot \gamma_1.
 \end{aligned}$$

In particular, the degrees are equal and $[\gamma_1] = [\gamma'_1] \in N^k(X)$ as required.

By the same argument, there exists a class $\gamma'_2 \in \text{CI}^{e_1+e_2+k}(X_1 \times X_2)$ such that $[\gamma_2] = [\gamma'_2] \in N^k(X)$, hence $[\gamma_1] + [\gamma_2] = [\gamma'_1] + [\gamma'_2] = [\gamma'_1 + \gamma'_2] \in N^k(X)$ as required. □

Definition 2.1.9. *We define $N_{\bullet}(X)$ (resp. $N^{\bullet}(X)$) by $\bigoplus_k N_k(X)$ (resp. $\bigoplus_k N^k(X)$).*

2.2. Algebra structure on the space of numerical cycles. We now define a structure of algebra on $N^\bullet(X)$, and prove that $N_\bullet(X)$ has a structure of $N^\bullet(X)$ module.

Pick $\gamma \in \text{CI}^{e_1+k}(X_1)_{\mathbb{Q}}$ where $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 . The element γ induces a morphism in the Chow group:

$$\gamma_{\perp} \cdot : \alpha \in A_l(X) \rightarrow p_{1*}(\gamma \cdot p_1^* \alpha) \in A_{l-k}(X). \quad (9)$$

The morphism $\gamma_{\perp} \cdot : A_l(X) \rightarrow A_{l-k}(X)$ induces a morphism of abelian groups from $N_l(X)$ to $N_{l-k}(X)$.

Proposition 2.2.1. *Any element $\alpha \in N^k(X)$ induces a morphism $\alpha_{\perp} \cdot : N_\bullet(X) \rightarrow N_{\bullet-k}(X)$ such that the following conditions are satisfied.*

- (i) *If α is induced by $\gamma \in \text{CI}^{e_1+k}(X_1)_{\mathbb{Q}}$ where $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 , then for any integer l and any $z \in N_l(X)$, one has in $N_{l-k}(X)$:*

$$\alpha_{\perp} z = \gamma_{\perp} z.$$

- (ii) *For any $\alpha, \beta \in N^k(X)$ and any $z \in N_l(X)$, we have:*

$$(\alpha + \beta)_{\perp} z = \alpha_{\perp} z + \beta_{\perp} z.$$

Proof. Let us consider $\alpha \in N^k(X)$ and suppose it is induced by $\gamma_1 \in \text{CI}^{e_1+k}(X_1)_{\mathbb{Q}}$ where $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 . We define the map $\alpha_{\perp} \cdot$ as :

$$\alpha_{\perp} z = \gamma_1_{\perp} z,$$

for any $z \in N_l(X)$. We show that the morphism does not depend on the choice of the class γ_1 and (i) follows from Proposition 2.1.8. Once (i) is satisfied, then (ii) follows directly from the linearity of the intersection product.

Suppose that $[\gamma_1] = [\gamma_2] \in N^k(X)$ where $\gamma_2 \in \text{CI}^{e_2+k}(X_2)_{\mathbb{Q}}$ and $p_2 : X_2 \rightarrow X$ is a flat morphism of relative dimension e_2 , then we need to prove that:

$$p_{1*}(\gamma_1 \cdot p_1^* z) \equiv p_{2*}(\gamma_2 \cdot p_2^* z),$$

for any fixed $z \in Z_l(X)$. Take $\beta \in \text{CI}^{e_3+l-k}(X_3)$ where $p_3 : X_3 \rightarrow X$ is flat morphism of relative dimension e_3 , we only need to show that:

$$(\beta \cdot p_3^* p_{1*}(\gamma_1 \cdot p_1^* z)) = (\beta \cdot p_3^* p_{2*}(\gamma_2 \cdot p_2^* z)).$$

Let X'_1 and X'_2 the fibre products $X_1 \times X_3$ and $X_2 \times X_3$, and $p'_1 : X'_1 \rightarrow X_3$, $p'_3 : X'_1 \rightarrow X_1$, $q_2 : X'_2 \rightarrow X_3$, $q_3 : X'_2 \rightarrow X_2$ be the corresponding flat projection morphisms such that we obtain the following commutative diagrams:

$$\begin{array}{ccc} & X'_1 & \\ & \swarrow p'_3 & \searrow p'_1 \\ X_1 & & X_3 \\ & \searrow p_1 & \swarrow p_3 \\ & X & \end{array} \quad \begin{array}{ccc} & X'_2 & \\ & \swarrow q_3 & \searrow q_2 \\ X_2 & & X_3 \\ & \searrow p_2 & \swarrow p_3 \\ & X & \end{array}$$

As above, we have $p_3^* p_{1*} = p'_1 p'^*_3$, hence:

$$\begin{aligned}
(\beta \cdot p_3^* p_{1*}(\gamma_1 \cdot p_1^* z)) &= (\beta \cdot p_1'^* p_3'^*(\gamma_1 \cdot p_1^* z)) \\
&= (p_1'^* \beta \cdot p_3'^*(\gamma_1 \cdot p_1^* z)) \\
&= (p_3'^* \gamma_1 \cdot p_1'^* p_3'^* z \cdot p_1'^* \beta) \\
&= (\gamma_1 \cdot p_3'^* p_1'^*(p_3^* z \cdot \beta)) \\
&= (\gamma_1 \cdot p_1^* p_{3*}(p_3^* z \cdot \beta)) \\
&= (\gamma_2 \cdot p_2^* p_{3*}(p_3^* z \cdot \beta)).
\end{aligned}$$

By a similar argument, we show that $(\beta \cdot p_3^* p_{2*}(\gamma_2 \cdot p_2^* z)) = (\gamma_2 \cdot p_2^* p_{3*}(p_3^* z \cdot \beta))$ which implies the desired equality:

$$(\beta \cdot p_3^* p_{1*}(\gamma_1 \cdot p_1^* z)) = (\beta \cdot p_3^* p_{2*}(\gamma_2 \cdot p_2^* z)).$$

□

Proposition 2.2.2. *There exists a unique structure of commutative graded ring with unit (\deg) on $N^\bullet(X)$ compatible with the action \lrcorner on $N_\bullet(X)$. Hence, the abelian group $N_\bullet(X)$ has the structure of a graded $N^\bullet(X)$ -module.*

Proof. Take $\alpha_1 \in N^k(X)$ and $\alpha_2 \in N^l(X)$ and define $\varphi \in Z^{k+l}(X)$ by the formula:

$$\varphi : z \in Z_{k+l}(X) \rightarrow (\alpha_1 \lrcorner (\alpha_2 \lrcorner z)).$$

We prove that φ is an element of $N^{k+l}(X)$.

By linearity, we can suppose that α_i is induced by $\gamma_i \in \text{CI}^{k+e_i}(X_i)$ where $p_i : X_i \rightarrow X$ is a flat morphism of relative dimension e_i for $i = 1, 2$. Let $X' = X_1 \times_X X_2$ be the fibre product, let p_1' and p_2' be the projections from X' to X_1 and X_2 respectively such that we have the commutative diagram:

$$\begin{array}{ccc}
& X' & \\
p_2' \swarrow & & \searrow p_1' \\
X_1 & & X_2 \\
p_1 \searrow & & \swarrow p_2 \\
& X &
\end{array}$$

By the projection formula, we obtain for all $z \in Z_{k+l}(X)$:

$$\varphi(z) = (p_1'^* \gamma_2 \cdot p_2'^* \gamma_1 \cdot p_2'^* p_1^* z). \quad (10)$$

In particular, we have shown that φ is induced by $p_1'^* \gamma_2 \cdot p_2'^* \gamma_1 \in \text{CI}^{e_1+e_2+k+l}(X')$, hence φ is an element of $N^{k+l}(X)$. Moreover, the commutativity of the intersection product in (10) proves that $(\alpha_2 \lrcorner (\alpha_1 \lrcorner z)) = (\alpha_1 \lrcorner (\alpha_2 \lrcorner z))$ for any $z \in N_{k+l}(X)$, hence $\alpha_1 \cdot \alpha_2 = \alpha_2 \cdot \alpha_1$.

Pick a vector bundle E on X . As the element $\deg \in N^0(X)$ is equal to $z \rightarrow (s_0(E) \lrcorner z)$ in $N^0(X)$ (see Remark 2.1.6), we get using Theorem 1.3.4.(2) that:

$$(\alpha \lrcorner z) = (\alpha \lrcorner (s_0(E) \lrcorner z)) = (s_0(E) \lrcorner (\alpha \lrcorner z)) = ((\alpha \cdot \deg) \lrcorner z) = (\deg \cdot \alpha) \lrcorner z,$$

for any $z \in N_l(X)$ and any $\alpha \in N^l(X)$. Hence, \deg is a unit of $N^\bullet(X)$.

□

2.3. Pullback on dual numerical classes. Let us consider $q : X \rightarrow Y$ a proper morphism. We define for any integer k the pullback $q^* : N^k(Y) \rightarrow \text{Hom}_{\mathbb{Z}}(N_k(X), \mathbb{Z})$ as the dual of the pushforward operation $q_* : N_k(X) \rightarrow N_k(Y)$ with respect to the pairing $N^k(X) \times N_k(X) \rightarrow \mathbb{Z}$ defined in Proposition 2.1.7.

Proposition 2.3.1. *Let $q : X \rightarrow Y$ be a proper morphism. The morphism q^* induces a morphism of graded rings $q^* : N^\bullet(Y) \rightarrow N^\bullet(X)$ which satisfies the projection formula:*

$$\forall \alpha \in N^k(Y), \forall z \in N_l(X), q_*(q^* \alpha \frown z) = \alpha \frown q_* z.$$

Proof. We only need to prove that the image $q^*(N^k(Y))$ is contained in $N^k(X)$ and that the projection formula is satisfied as it directly implies that $q^* : N^\bullet(Y) \rightarrow N^\bullet(X)$ is a morphism of rings since:

$$(\alpha \cdot \beta) \frown q_* z = q^*(\alpha \cdot \beta) \frown z = \alpha \frown q_*(q^* \beta \frown z) = (q^* \alpha \cdot q^* \beta) \frown z,$$

for any $\alpha \in N^k(Y)$, $\beta \in N^l(Y)$ and any $z \in N_{k+l}(X)$.

Consider a class $\alpha \in N^k(Y)$ which is induced by $\gamma \in \text{CI}^{e_1+k}(Y_1)$ where $p_1 : Y_1 \rightarrow Y$ is a flat proper morphism of relative dimension e_1 . Setting X_1 to be the fibre product $Y_1 \times X$ and p'_1, q' the projections from X_1 to X and X_1 respectively, one remarks using the equality $q'_* p'^*_1 = p^*_1 q_*$ ([Ful98, Proposition 1.7]) that $q^* \alpha$ is induced by $q'^* \gamma$, hence $q^* \alpha \in N^k(X)$ as required. And the projection formula follows easily from the projection formula on divisors (Theorem 1.2.1.(ii)). \square

Let us sum up all the properties of numerical classes proven so far :

Theorem 2.3.2. *Let $q : X \rightarrow Y$ be a proper morphism. For any integer $0 \leq k \leq \dim X$ and $0 \leq l \leq \dim Y$:*

- (i) *The pushforward morphism $q_* : Z_k(X) \rightarrow Z_k(Y)$ induces a morphism of abelian groups $q_* : N_k(X) \rightarrow N_k(Y)$.*
- (ii) *The dual morphism $q^* : Z^l(Y) \rightarrow Z^l(X)$ maps $N^l(Y)$ into $N^l(X)$.*
- (iii) *The induced morphism $q^* : N^\bullet(Y) \rightarrow N^\bullet(X)$ preserves the structure of graded rings.*
- (iv) *(Projection formula) For all $\alpha \in N^l(Y)$ and all $z \in N_k(X)$, we have $q_*(q^* \alpha \frown z) \equiv \alpha \frown q_* z$ in $N_{k-l}(Y)$.*

2.4. Canonical morphism.

Theorem 2.4.1. *There exists a unique canonical morphism $\psi_X : N^k(X) \rightarrow N_{n-k}(X)$ which satisfies the following properties.*

- (i) *The image of the morphism $\text{deg} : Z_0(X) \rightarrow \mathbb{Z}$ seen as an element of $Z^0(X)$ is given by $\psi_X(\text{deg}) = [X]$.*
- (ii) *The morphism ψ_X is $N^k(X)$ -equivariant, i.e for all $\alpha \in N^k(X)$ and all $\beta \in N^l(X)$, we have:*

$$\psi_X(\alpha \cdot \beta) = \alpha \frown \psi_X(\beta).$$

- (iii) *Suppose $q : X \rightarrow Y$ is a generically finite morphism where Y is of dimension n , then we have the following identity:*

$$q_* \circ \psi_X \circ q^* = \text{deg}(q) \times \psi_Y.$$

Proof. We define ψ_X for any $\alpha \in N^k(X)$ by setting:

$$\psi_X(\alpha) = \alpha \frown [X].$$

By construction, we have that $\psi_X(\text{deg}) = [X]$ and (ii) follows directly from the definition and Proposition 2.2.2.

Assertion (iii) is then a consequence of the projection formula (see Theorem 2.3.2.(iv)) and the fact that $q_*[X] = \text{deg}(q)[Y]$.

Let us prove that ψ_X is unique. Suppose that $\varphi : N^k(X) \rightarrow N_{n-k}(X)$ satisfies the hypothesis of the theorem. Since $\varphi(\text{deg}) = [X]$ and since deg is the unit element of the ring $N^\bullet(X)$, we have that for any $\alpha \in N^k(X)$, $\alpha = \alpha \cdot \text{deg}$. By (ii),

$$\varphi(\alpha) = \varphi(\alpha \cdot \text{deg}) = \alpha \lrcorner \varphi(\text{deg}) = \alpha \lrcorner [X] = \psi_X(\alpha),$$

as required. □

Now we prove some properties of ψ_X in some particular cases.

Theorem 2.4.2. *The following properties are satisfied.*

- (i) *If X is smooth, then for all integers $0 \leq k \leq n$, the induced morphism $\psi_X : N^k(X)_\mathbb{Q} \rightarrow N_{n-k}(X)_\mathbb{Q}$ is an isomorphism.*
- (ii) *If X is smooth and $q : X \rightarrow Y$ is a surjective generically finite morphism where Y is a normal projective variety. Then we have for all integer k :*

$$q^*(\psi_Y(N^{n-k}(Y)_\mathbb{Q})^\perp) = q^*(N^k(Y)_\mathbb{Q}) \cap \text{Ker}(q_* \circ \psi_X : N^k(X)_\mathbb{Q} \rightarrow N_{n-k}(Y)_\mathbb{Q}). \quad (11)$$

Proof. (i) Let us show that ψ_X is surjective. By the Grothendieck-Riemann-Roch's theorem (Theorem 1.3.6), the Chern character induces an isomorphism:

$$\text{ch} \lrcorner [X] : E \in K^0(X) \otimes \mathbb{Q} \rightarrow \text{ch}(E) \lrcorner [X] \in A_\bullet(X) \otimes \mathbb{Q}.$$

This implies that the morphism $\psi_X : N^k(X)_\mathbb{Q} \rightarrow N_{n-k}(X)_\mathbb{Q}$ is surjective because any Chern class is the image of a product of Cartier divisors by a flat map (see Remark 1.3.3).

We now prove that $\psi_X : N^k(X)_\mathbb{Q} \rightarrow N_{n-k}(X)_\mathbb{Q}$ is injective. Take $\alpha_1 \in N^k(X)_\mathbb{Q}$ such that $\psi_X(\alpha_1) = 0$. By Proposition 2.1.8, the class α_1 is induced by $\gamma_1 \in \text{CI}^{e_1+k}(X_1)_\mathbb{Q}$ where $p_1 : X_1 \rightarrow X$ is a flat morphism of relative dimension e_1 . The condition $\psi_X(\alpha_1) = 0$ is equivalent to the equality $p_{1*}\gamma_1 = 0 \in N_{n-k}(X)$. We need to show that $(\gamma_1 \cdot p_1^*z) = 0$ for any cycle $z \in Z_k(X)$. As X is smooth, we may compute intersection products inside the Chow group $A_\bullet(X)$ directly by Remark 1.3.3 and we get:

$$(\gamma_1 \cdot p_1^*z) = (p_{1*}(\gamma_1 \cdot p_1^*z)) = (p_{1*}\gamma_1 \cdot z) = 0$$

as the class $z \in N_k(X)$ is the image of an element of $N^{n-k}(X)_\mathbb{Q}$ by surjectivity of ψ_X .

(ii) We have the following series of equivalence:

$$\begin{aligned} \beta \in \psi_Y(N^{n-k}(Y)_\mathbb{Q})^\perp &\Leftrightarrow \forall \alpha \in N^{n-k}(Y)_\mathbb{Q}, (\beta \lrcorner \psi_Y(\alpha)) = 0 \\ &\Leftrightarrow \forall \alpha \in N^{n-k}(Y)_\mathbb{Q}, (\beta \lrcorner (q_*\psi_X q^*\alpha)) = 0 \\ &\Leftrightarrow \forall \alpha \in N^{n-k}(Y)_\mathbb{Q}, (q^*\beta \cdot q^*\alpha) = 0 \\ &\Leftrightarrow \forall \alpha \in N^{n-k}(Y)_\mathbb{Q}, (\alpha \lrcorner q_*\psi_X q^*\beta) = 0 \\ &\Leftrightarrow q^*\beta \in \text{Ker}(q_* \circ \psi_X : N^k(X)_\mathbb{Q} \rightarrow N_{n-k}(Y)_\mathbb{Q}), \end{aligned}$$

where the second equivalence follows from Theorem 2.4.2.(3), the third and the fourth equivalence from the projection formula, and the last equivalence is a consequence of the fact that ψ_X is self-adjoint :

$$(\beta \lrcorner \psi_Y(\alpha)) = (\beta \lrcorner (\alpha \lrcorner [Y])) = (\alpha \lrcorner (\beta \lrcorner [Y])) = (\alpha \lrcorner \psi_Y(\beta)),$$

where $\alpha \in N^k(Y)$ and $\beta \in N^{n-k}(Y)$. □

Remark 2.4.3. The proof of Theorem 2.4.2.(i) shows that when X is smooth, $N_k(X)_{\mathbb{Q}}$ is the quotient of $Z_k(X)_{\mathbb{Q}}$ by cycles $z \in Z_k(X)_{\mathbb{Q}}$ such that for any cycle $z' \in Z_{n-k}(X)_{\mathbb{Q}}$, one has $(z \cdot z') = 0$.

Remark 2.4.4. When X is smooth and when $\kappa = \mathbb{C}$, denote by $\text{Alg}^k(X)$ the subgroup of the de Rham cohomology $H^{2k}(X, \mathbb{C})$ generated by algebraic cycles of dimension k in X . Then there is a surjective morphism $\text{Alg}^k(X) \rightarrow N^k(X)_{\mathbb{Q}}$

2.5. Numerical spaces are finite dimensional.

Theorem 2.5.1. *Both \mathbb{Q} -vector spaces $N_k(X)_{\mathbb{Q}}$ and $N^k(X)_{\mathbb{Q}}$ are finite dimensional.*

Proof. If X is smooth, then using Remark 2.4.3, $N_k(X)_{\mathbb{Q}}$ is the quotient of $Z_k(X)_{\mathbb{Q}}$ by the equivalence relation which identifies cycles α and β in $Z_k(X)_{\mathbb{Q}}$ if for any cycle $z \in Z_{n-k}(X)_{\mathbb{Q}}$, $(z \cdot \alpha) = (z \cdot \beta)$. In particular, the vector-space $N_k(X)_{\mathbb{Q}}$ is finitely generated (see [Mil13, Theorem 23.6] for a reference), and so is $N^k(X)_{\mathbb{Q}}$ using Theorem 2.4.2.(i).

If X is not smooth, by DeJong's alteration theorem (cf [Jon96, Theorem 4.1]), there exists a smooth projective variety X' and a generically finite surjective morphism $q : X' \rightarrow X$. We only need to show that the pushforward $q_* : N_k(X')_{\mathbb{Q}} \rightarrow N_k(X)_{\mathbb{Q}}$ is surjective. Indeed this first implies that $N_k(X)_{\mathbb{Q}}$ is finite dimensional. Since the natural pairing $N^k(X)_{\mathbb{Q}} \times N_k(X)_{\mathbb{Q}} \rightarrow \mathbb{Q}$ is non degenerate we get an injection of $N^k(X)_{\mathbb{Q}}$ onto $\text{Hom}_{\mathbb{Q}}(N_k(X)_{\mathbb{Q}}, \mathbb{Q})$ which is also finite dimensional.

We take V an irreducible subvariety of codimension k in X . If $\dim q^{-1}(V) = \dim V$, then the class $q_*[q^{-1}(V)]$ in $N_{\dim V}(X)_{\mathbb{Q}}$ is represented by a cycle of dimension $\dim V$ which is included in V . As V is irreducible, we have $q_*[q^{-1}(V)] \equiv \lambda[V]$ for some $\lambda \in \mathbb{N}^*$.

If the dimension of $q^{-1}(V)$ is strictly greater than V , we take W an irreducible component of $q^{-1}(V)$ such that its image by $q|_W : W \rightarrow V$ is dominant. We write the dimension of W as $\dim V + r$ where $r > 0$ is an integer. Fix an ample divisor H_X on X . The class $H_X^r \lrcorner [W] \in N_{\dim V}(X')_{\mathbb{Q}}$ is represented by a cycle of dimension $\dim V$ in W . So the image of the class $q_*(H_X^r \lrcorner [W]) \in N_{\dim V}(X)_{\mathbb{Q}}$ is a multiple of $[V]$ which implies the surjectivity of q_* . □

Corollary 2.5.2. *For any integer $0 \leq k \leq n$, the pairing $N^k(X)_{\mathbb{R}} \times N_k(X)_{\mathbb{R}} \rightarrow \mathbb{R}$ is perfect.*

Corollary 2.5.3. *Suppose that the dimension of X is $2n$, then the morphism $\psi_X : N^n(X)_{\mathbb{Q}} \rightarrow N_n(X)_{\mathbb{Q}}$ is an isomorphism.*

Proof. We apply (11) to an alteration X' of X where $q : X' \rightarrow X$ is a proper surjective morphism and where X' is a smooth projective surface. This proves that $\psi_X : N^n(X)_{\mathbb{Q}} \rightarrow N_n(X)_{\mathbb{Q}}$ is surjective. By duality, this gives that $\psi_X : N^n(X)_{\mathbb{Q}} \rightarrow N_n(X)_{\mathbb{Q}}$ is injective. As a consequence, we have that $\psi_X : N^n(X)_{\mathbb{Q}} \rightarrow N_n(X)_{\mathbb{Q}}$ is an isomorphism. □

Corollary 2.5.4. *Let X be a complex normal projective variety with at most rational singularities. We suppose that X is numerically \mathbb{Q} -factorial in the sense of [BdFFU15]. Then the morphisms $\psi_X : N^1(X)_{\mathbb{Q}} \rightarrow N_{n-1}(X)_{\mathbb{Q}}$ and $\psi_X : N^{n-1}(X)_{\mathbb{Q}} \rightarrow N_1(X)_{\mathbb{Q}}$ are isomorphisms.*

Proof. Using [BdFFU15, Theorem 5.11], then any Weil divisor which is numerically \mathbb{Q} -Cartier is \mathbb{Q} -Cartier. In particular, $\psi_X : N^1(X)_{\mathbb{Q}} \rightarrow N_{n-1}(X)_{\mathbb{Q}}$ is surjective. Using (11) to an alteration of X' applied to $k = 1$, we have that $\psi_X : N^1(X)_{\mathbb{Q}} \rightarrow N_{n-1}(X)_{\mathbb{Q}}$ is injective. Hence $N^1(X)_{\mathbb{Q}}$ and $N_{n-1}(X)_{\mathbb{Q}}$ are isomorphic and by duality $N^{n-1}(X)_{\mathbb{Q}}$ and $N_1(X)_{\mathbb{Q}}$ are also isomorphic. □

Example 2.5.5. When $X = X(\Delta)$ be a toric variety associated to a complete fan Δ . The map $\psi_X : N^1(X)_{\mathbb{Q}} \rightarrow N_{n-1}(X)_{\mathbb{Q}}$ is an isomorphism if and only if Δ is a simplicial fan using [Ful93, exercise p65 section 3.4].

3. POSITIVITY

The notion of positivity is relatively well understood for cycles of codimension 1 and of dimension 1. For cycles of intermediate dimension this situation is however more subtle and was only recently seriously considered (see [DELV11], [CC15], [CLO16] and the recent series of papers by Fulger and Lehmann ([FL14a], [FL14b])).

For our purpose, we will first review the notions of pseudo-effectivity and numerically effective classes. Then we generalize the construction of the basepoint free cone introduced by [FL14b] to normal projective varieties. This cone is suitable for stating generalized Siu's inequalities (see Section 3.4).

3.1. Pseudo-effective and numerically effective cones. As in the previous section, X is a normal projective variety of dimension n . To ease notation we shall also write $N^k(X)$ and $N_k(X)$ for the real vector spaces $N^k(X)_{\mathbb{R}}$ and $N_k(X)_{\mathbb{R}}$.

Definition 3.1.1. *A class $\alpha \in N_k(X)$ is pseudo-effective if it is in the closure of the cone generated by effective classes. This cone is denoted $\text{Psef}_k(X)$.*

When $k = 1$, $\text{Psef}_1(X)$ is the Mori cone (see e.g [KM98, Definition 1.17]), and when $k = n - 1$, $\text{Psef}_{n-1}(X)$ is the classical cone of pseudo-effective divisors, its interior being the big cone.

Definition 3.1.2. *A class $\beta \in N^k(X)$ is numerically effective (or nef) if for any class $\alpha \in \text{Psef}_{n-k}(X)$, $(\beta \frown \alpha) \geq 0$. We denote this cone by $\text{Nef}^k(X)$.*

When $k = 1$, the cone $\text{Nef}^1(X)$ is the cone of numerically effective divisors, its interior is the ample cone.

We can define a notion of effectivity in the dual $N^k(X)$.

Definition 3.1.3. *A class $\alpha \in N^k(X)$ is pseudo-effective if $\psi_X(\alpha) \in \text{Psef}_{n-k}(X)$. We will write this cone as $\text{Psef}^k(X)$.*

Definition 3.1.4. *A class $z \in N_k(X)$ is numerically effective if for any class $\alpha \in \text{Psef}^k(X)$, one has $(\alpha \frown z) \geq 0$. This cone is denoted $\text{Nef}_k(X)$.*

By convention, we will write $\alpha \leq \beta$ (resp. $\alpha \leq \beta$) for any $\alpha, \beta \in N_k(X)$ (resp. $\alpha, \beta \in N^k(X)$) if $\beta - \alpha \in \text{Psef}_k(X)$ (resp. $\beta - \alpha \in \text{Psef}^k(X)$).

When X is smooth, the morphism ψ_X induces an isomorphism between $N^k(X)$ and $N_{n-k}(X)$, and we can identify these cones:

$$\begin{aligned} \text{Nef}^k(X) &= \text{Nef}_{n-k}(X), \\ \text{Psef}^k(X) &= \text{Psef}_{n-k}(X). \end{aligned}$$

3.2. Pliant classes. We recall the definition of pliant classes introduced in [FL14b, Definition 3.1] and their main properties.

Definition 3.2.1. *The pliant cone $\text{PL}^\bullet(X)$ is defined as the convex cone generated by product of Schur polynomials of globally generated vector bundle.*

We denote by $\text{PL}^k(X)$ the set of pliant classes of codimension k in X .

Theorem 3.2.2. *(see [FL14b, Theorem 1.3]) The pliant cone $\text{PL}^k(X)$ satisfies the following properties.*

- (i) *The cone $\text{PL}^k(X)$ is a closed convex salient cone with non-empty interior in $N^k(X)_{\mathbb{R}}$.*
- (ii) *The cone $\text{PL}^k(X)$ contains product of ample Cartier divisors in its interior.*
- (iii) *For all integer k, l , we have $\text{PL}^k(X) \cdot \text{PL}^l(X) \subset \text{PL}^{k+l}(X)$.*

(iv) For any (proper) morphism $q : X \rightarrow Y$, one has that $q^* \text{PL}^k(Y) \subset \text{PL}^k(X)$.

We recall another proposition which we will reuse in our proofs.

Proposition 3.2.3. (cf [FL14b, Example 3.13]) *Let \mathbb{G} be a Grassmannian variety. Then $\text{PL}^k(\mathbb{G}) = \text{Psef}^k(\mathbb{G})$.*

3.3. Basepoint free cone on normal projective varieties. In this section, we define a cone $\text{BPF}^k(X)$ and prove in Corollary 3.3.4 that this cone is equal to the basepoint free cone defined by Fulger-Lehmann when X is smooth. This generalizes [FL14b, Theorem 1.7] to normal projective varieties and our proof follows closely Fulger-Lehmann's approach.

Recall that a complete intersection $\gamma \in \text{CI}^{k+e}(X')$ on X' where $p : X' \rightarrow X$ is a flat morphism of relative dimension e and where X' is an equidimensional projective scheme induces naturally (see Definition 2.1.5) an element $[\gamma] \in \text{N}^k(X)_{\mathbb{R}} = \text{Hom}_{\mathbb{R}}(\text{N}_k(X)_{\mathbb{R}}, \mathbb{R})$ by intersecting the class γ with the pullback by p of a k -dimensional cycle in X . We also refer to Proposition 2.2.2 for the definition of the product $\text{N}^k(X)_{\mathbb{R}} \times \text{N}^l(X)_{\mathbb{R}} \rightarrow \text{N}^{k+l}(X)_{\mathbb{R}}$.

Definition 3.3.1. *The cone $\text{BPF}^k(X)$ is the closure of the convex cone in $\text{N}^k(X)_{\mathbb{R}}$ generated by products of the form $[\gamma_1] \cdot \dots \cdot [\gamma_l]$ where each γ_i is a product of $e_i + k_i$ ample Cartier divisors on an equidimensional projective scheme X_i which is flat over X of relative dimension e_i and where k_i are integers satisfying $k_1 + \dots + k_l = k$.*

Remark 3.3.2. By definition, the cone $\text{BPF}^k(X)$ contains the products of ample Cartier divisors and Segre classes of anti-ample vector bundles.

Recall also that if $q : X \rightarrow Y$ is a flat morphism of relative dimension e between projective schemes, then the pushforward is well-defined on numerical cycles $q_* : \text{N}^k(X)_{\mathbb{R}} \rightarrow \text{N}^{k-e}(Y)_{\mathbb{R}}$ (see Corollary A.5).

Theorem 3.3.3. *The cone $\text{BPF}^k(X)$ is the smallest cone which satisfies the following properties.*

- (i) *The cone $\text{BPF}^k(X)$ is a salient, closed, convex cone with non-empty interior in $\text{N}^k(X)_{\mathbb{R}}$.*
- (ii) *The cone $\text{BPF}^k(X)$ contains products of ample Cartier divisors in its interior.*
- (iii) *For all integer k and l , we have $\text{BPF}^k(X) \cdot \text{BPF}^l(X) \subset \text{BPF}^{k+l}(X)$.*
- (iv) *For any (proper) morphism $q : X \rightarrow Y$, we have $q^* \text{BPF}^k(Y) \subset \text{BPF}^k(X)$.*
- (v) *For any integer k , we have $\text{BPF}^k(X) \subset \text{Nef}^k(X) \cap \text{Psef}^k(X)$.*
- (vi) *In codimension 1, one has $\text{BPF}^1(X) = \text{Nef}^1(X)$.*
- (vii) *For any flat morphism $q : X \rightarrow Y$ between equidimensional projective schemes of relative dimension e and any integer $k \geq e$, we have $q_* \text{BPF}^k(X) \subset \text{BPF}^{k-e}(Y)$.*

Proof. We prove successively the items (iii), (vii), (v), (vi), (iv), (ii) and (i).

(iii), (vii) This follows from the definition of $\text{BPF}^k(X)$.

(v) It is sufficient to prove that for any effective cycle $z \in Z_{n-l}(X)$ and any basepoint free class $\alpha \in \text{BPF}^k(X)$, then $\alpha \cdot z \in \text{Psef}_{n-k-l}(X)$. Indeed, apply this successively to $z = [X]$ and $z \in \text{Psef}_k(X)$ give the inclusions $\text{BPF}^k(X) \subset \text{Psef}^k(X)$ and $\text{BPF}^k(X) \subset \text{Nef}^k(X)$. By definition of basepoint free classes and by linearity, we can suppose that α is equal to a product $[\gamma_1] \cdot \dots \cdot [\gamma_p]$ where $\gamma_i \in \text{CI}^{e_i+k_i}(X_i)_{\mathbb{R}}$ are products of ample Cartier divisors on X_i where $p_i : X_i \rightarrow X$ is a flat proper morphism of relative dimension e_i and where k_i are integers such that $k_1 + \dots + k_p = k$. By definition, one has $[\gamma_1] \cdot z = p_{1*}(\gamma_1 \cdot p_1^* z)$. Because the cycle z is pseudo-effective, the cycle $p_1^* z$ remains pseudo-effective as p_1 is a flat morphism. As γ_1 is a positive combination of products of ample Cartier divisors, we deduce that the cycle $\gamma_1 \cdot p_1^* z$ is pseudo-effective. Hence, $[\gamma_1] \cdot z \in \text{Psef}_{n-l_1-l}(X)$. Iterating the same argument, we get that $\alpha \cdot z \in \text{Psef}_{n-k-l}(X)$ as required.

(vi) The interior of $\text{Nef}^1(X)$ is equal to the ample cone of X so by definition:

$$\text{Int}(\text{Nef}^1(X)) \subset \text{BPF}^1(X).$$

As the closure of the ample cone is the nef cone by [Laz04, Theorem 1.4.21.(i)], one gets $\text{Nef}^1(X) \subset \text{BPF}^1(X)$. Conversely, the cone $\text{BPF}^1(X)$ is included in the cone $\text{Nef}^1(X)$, so we get $\text{BPF}^1(X) = \text{Nef}^1(X)$.

(iv) By linearity and stability by products, we are reduced to treat the case of a class $[D]$ induced by an ample Cartier divisor on Y_1 where $p_1 : Y_1 \rightarrow Y$ is a flat proper morphism, and prove that $q^*[D]$ is a limit of ample Cartier divisors on a flat variety over X . Let X_1 be the fibre product of Y_1 and X and let q' be the natural projection from X_1 to Y_1 , observe that $q^*[D]$ is induced by q'^*D which remains nef on X_1 as q' is proper. In particular, it is the limit of ample divisors on $N^1(X_1)$.

(i) Take $\alpha \in \text{BPF}^k(X)$ such that $-\alpha \in \text{BPF}^k(X)$. Then for all $z \in \text{Psef}_k(X)$, one has that $(\alpha \perp z) = 0$ as α is nef by (5). Since effective classes of dimension k generate $Z_k(X)$, it follows that $(\alpha \perp z) = 0$ for any $z \in N_k(X)_{\mathbb{R}}$ which implies by definition that $\alpha = 0$. This shows $\text{BPF}^k(X)$ is salient.

(ii) We show now that $\text{BPF}^k(X)$ contains product of ample divisors in its interior. To do so we prove that $\text{PL}^k(X) \subset \text{BPF}^k(X)$ for any integer $k \geq 1$.

For $k = 1$, $\text{BPF}^1(X) = \text{Nef}^1(X)$, and by definition, the divisor h is ample so it is in the interior of the nef cone and we are done. Take a globally generated line bundle E of rank r on X and consider the induced morphism ϕ given by:

$$\phi : X \rightarrow \mathbb{G} = G(r, \mathbb{P}(H^0(X, E)^*)).$$

Since $\text{PL}^k(X) = \phi^* \text{PL}^k(\mathbb{G})$ and since these cones are preserved by pullbacks, we are then reduced to proving that $\text{PL}^k(\mathbb{G}) \subset \text{BPF}^k(\mathbb{G})$. Denote by $G = \text{PGL}(H^0(X, E)^*)$ and consider a class $\alpha \in N^k(\mathbb{G})_{\mathbb{R}}$. Since \mathbb{G} is smooth, $\psi_{\mathbb{G}} : N^k(\mathbb{G})_{\mathbb{R}} \rightarrow N_{n-k}(\mathbb{G})_{\mathbb{R}}$ is an isomorphism by Theorem 2.4.2 and α is represented by an effective cycle $z \in Z_{n-k}(\mathbb{G})_{\mathbb{R}}$. Consider the closed scheme W in $G \times \mathbb{G}$ given by:

$$W = \{(g, g \cdot x)\}_{g \in G, x \in z} \subset G \times \mathbb{G}.$$

By construction, W is a projective scheme and the projection $p : W \rightarrow \mathbb{G}$ onto \mathbb{G} is a flat morphism. Denote by $q : W \rightarrow G$ the projection onto g . By construction the general fiber of q over g is numerically equivalent to z and we have that:

$$p_* q^* H_G^{\dim G} = \alpha \in N_{n-k}(\mathbb{G})_{\mathbb{R}},$$

where H_G is an ample divisor on G . This proves that $\alpha \in \text{BPF}^k(\mathbb{G})$ since $q^* H_G$ is a nef divisor as required.

Since $\text{PL}^k(X)$ has non-empty interior in $N^k(X)_{\mathbb{R}}$ by Theorem 3.2.2.(ii), we have proved (ii).

Let us prove that the cone $\text{BPF}^k(X)$ is the smallest cone satisfying the seven conditions. Denote by BPF' the minimal cone satisfying these conditions. We have that $\text{BPF}'^k(X) \subset \text{BPF}^k(X)$ by minimality. Take $q : X_1 \rightarrow X$ a flat morphism of relative dimension e where X_1 is an equidimensional projective scheme and consider $\alpha \in \text{Cl}^{k+e}(X_1)$ a product of ample Cartier divisors on X_1 . Since $q_* : N^k(X_1)_{\mathbb{R}} \rightarrow N^{k-e}(X)_{\mathbb{R}}$ and since $\alpha \in \text{BPF}'^{k+e}(X_1)$, we have that $q_* \alpha \in \text{BPF}'^k(X)$ by (vii), hence $\text{BPF}^k(X) \subset \text{BPF}'^k(X)$ as required. \square

We recall Fulger-Lehmann's construction of the basepoint free cone. A class $\alpha \in N_{n-k}(X)_{\mathbb{R}}$ is strongly basepoint free if there is:

- an equidimensional quasi-projective scheme U of finite type over κ ,
- a flat proper morphism $s : U \rightarrow X$
- and a proper morphism $p : U \rightarrow W$ of relative dimension $n - k$ to a quasi-projective scheme W such that each component of U surjects onto W

such that

$$s|_{F_p}([F_p]) = \alpha,$$

where $[F_p]$ is the fundamental class of a general fiber of p . We denote by $\text{BPF}^{k'}(X)$ the closure of the convex cone generated by strongly basepoint free classes in this sense.

Corollary 3.3.4. *Suppose X is smooth, then the cone $\text{BPF}^k(X)$ is equal to the basepoint free cone $\text{BPF}^{k'}(X)$.*

Proof. Since the basepoint free cone satisfies the conditions of Theorem 3.3.3, one has that $\text{BPF}^k(X) \subset \text{BPF}^{k'}(X)$. Let us prove the reverse inclusion $\text{BPF}^{k'}(X) \subset \text{BPF}^k(X)$. Take $p : U \rightarrow W$ a projective morphism onto an equidimensional quasi-projective variety W where U is a quasi-projective scheme and a flat map $s : U \rightarrow X$ such that $s_*[F_p] = \alpha$ where F_p is a general fiber of p . Take H_W an ample divisor on W , then the class α satisfies:

$$\alpha = s_* p^* H_W^{k+e} \in N_{n-k}(X)_{\mathbb{R}}.$$

Choose an ample divisor H on U , since the class $p^* H_W$ is nef, for any $\epsilon > 0$, the divisor $p^* H_W + \epsilon H$ is ample.

Since the morphism $s : U \rightarrow X$ is also quasi-projective and there exists an integer l (which depends on ϵ) such that the following diagram is commutative

$$\begin{array}{ccc} & \mathbb{P}_X^l & \\ i_\epsilon \nearrow & & \searrow \pi \\ U & \xrightarrow{s} & X \end{array}$$

where $i_\epsilon : U \rightarrow \mathbb{P}_X^l$ is an immersion induced by $p^* H_W + \epsilon H$ and $\pi : \mathbb{P}_X^l \rightarrow X$ is the flat projection onto X .

Let ξ be the relative class $c_1(\mathcal{O}_{\mathbb{P}_X^l}(1))$ on \mathbb{P}_X^l , then one has that for any cycle $z \in Z_k(X)_{\mathbb{R}}$:

$$((p^* H_W + \epsilon H)^{k+e} \cdot s^* z) = (\xi^{k+e} \cdot \pi^* z),$$

since $i_\epsilon^* \xi = p^* H_W + \epsilon H$. Hence, we obtain:

$$(s_*(p^* H_W + \epsilon H)^{k+e} \cdot z) = (\pi_* \xi^{k+e} \cdot z).$$

Since the class ξ^{k+e} is nef and since $\pi_*(\xi^{k+e}) \subset \text{BPF}^k(X)$. Taking the limit as $\epsilon \rightarrow 0$, we have that $s_*(p^* H_W + \epsilon H)^{k+e} \rightarrow \alpha = s_* p^* H_W^{k+e}$, hence $\alpha \in \text{BPF}^k(X)$ since each class $s_*(p^* H_W + \epsilon H)^{k+e} \in N^k(X)_{\mathbb{R}}$ belongs to $\text{BPF}^k(X)$. □

We give here a detailed proof of the fact that the pseudo-effective cone is salient (see also [FL14b, Corollary 3.17]). The proof uses a useful proposition that we will use later on.

Proposition 3.3.5. *Let $\alpha \in \text{Psef}_{n-k}(X)$ be a pseudo-effective class on X and $\gamma \in \text{BPF}^{n-k}(X)$ be class lying in the interior of the basepoint free cone. Then we have $(\gamma \perp \alpha) = 0$ if and only if $\alpha = 0$.*

Proof. Let us fix two basepoint free classes β and γ in $N^{n-k}(X)$, and a norm $\|\cdot\|$ on $N^{n-k}(X)_{\mathbb{R}}$. As γ is in the interior of $\text{BPF}^{n-k}(X)$ by Theorem 3.3.3.(2), there exists a positive constant $C > 0$ such that for any $\beta \in \text{BPF}^{n-k}(X)$, one has:

$$C\|\beta\|_{N^{n-k}(X)_{\mathbb{R}}}\gamma - \beta \in \text{BPF}^{n-k}(X).$$

Intersecting with $\alpha \in \text{Psef}_{n-k}(X)$ and using Theorem 3.3.3.(5), we have that $(\beta \cdot \alpha) = 0$. Since the basepoint free cone $\text{BPF}^{n-k}(X)$ generates all $N^{n-k}(X)_{\mathbb{R}}$ by Theorem 3.3.3.(1), we have proved that $(\beta' \cdot \alpha) = 0$ for any $\beta' \in N^{n-k}(X)$, hence $\alpha = 0$ as required. \square

Corollary 3.3.6. *The pseudo-effective cone $\text{Psef}_{n-k}(X)$ is a closed, convex, full dimensional salient cone in $N_{n-k}(X)_{\mathbb{R}}$.*

Proof. We take $u \in \text{Psef}_{n-k}(X)$ such that $-u \in \text{Psef}_{n-k}(X)$, then for any ample Cartier divisor H_X on X , the products $(H_X^{n-k} \cdot u)$ and $(-u \cdot H_X^{n-k})$ are non-negative hence $(u \cdot H_X^{n-k}) = 0$. This implies that $u = 0$ by Proposition 3.3.5. \square

3.4. Siu's inequality in arbitrary codimension. We recall Siu's inequality:

Proposition 3.4.1. ([Laz04, Theorem 2.2.13]) *Let V be a closed subscheme of dimension r in X and let A, B be two \mathbb{Q} -divisors nef on X such that $A|_V$ is big, then we have in $N_{k-1}(X)$,*

$$B \cdot [V] \leq \frac{r((A^{r-1} \cdot B) \cdot [V])}{(A^r \cdot [V])} A \cdot [V].$$

Remark 3.4.2. The case $V = X$ is a consequence of the bigness criterion given in [Laz04, Theorem 2.2.13].

Remark 3.4.3. The proof of the previous proposition implies that $B|_V \leq r(A^{r-1} \cdot B \cdot [V]) / (A^r \cdot [V]) \times A|_V$ in the Chow group $A^1(V)$. However, since we want to work in the numerical group, we compare these classes in X (we look at their pushforward by the inclusion of V in X).

Proof. The proof is the same as in [Laz04, Theorem 2.2.13], that is to find a section of the line bundle $\mathcal{O}_V(m(A - B))$. We choose m general elements E_i of $|B|$ and consider the exact sequence:

$$0 \longrightarrow \mathcal{O}_V(mA - mB) \longrightarrow \mathcal{O}_V(mA) \longrightarrow \mathcal{O}_{\cup E_i}(mA) \longrightarrow 0.$$

Taking long exact sequence associated, one obtains the minoration:

$$h^0(V, \mathcal{O}_V(mA - mB)) \geq h^0(V, \mathcal{O}_V(mA)) - h^0(\cup_{i=1}^m E_i, \mathcal{O}_{\cup_{i=1}^m E_i}(mA)).$$

Observe that $[\cup E_i] = \sum_{i=1}^m [E_i] = mB \cdot [V]$. Applying [GGJ⁺16, Corollary 3.6.3] to the nef divisor A , we get $h^0(V, \mathcal{O}_V(mA)) = m^r / (r!) (A^r \cdot [V]) + o(m^r)$ and

$$h^0(\cup E_i, \mathcal{O}_{\cup_{i=1}^m E_i}(mA)) = \sum_{i=1}^m \frac{m^{r-1}}{(r-1)!} A^{r-1} \cdot B \cdot [V] + o(m^r).$$

Hence,

$$h^0(V, \mathcal{O}_V(mA - mB)) \geq \frac{m^r}{r!} (A^r - rA^{r-1} \cdot B) \cdot [V] + o(m^r).$$

In particular, this implies the required inequality. \square

The next result is a key for our approach to controlling degrees of dominant rational maps.

Theorem 3.4.4. *Let k be an integer and V be a closed subscheme of dimension r in X . For any Cartier divisors $\alpha_1, \dots, \alpha_k$ and β which are big and nef on V , then there exists a constant $C > 0$ depending only on r and k such that:*

$$(\alpha_1 \cdot \dots \cdot \alpha_k)_\perp[V] \leq (r - k + 1)^k \frac{(\alpha_1 \cdot \dots \cdot \alpha_k \cdot \beta^{r-k})_\perp[V]}{(\beta^r)_\perp[V]} \times \beta^k_\perp[V] \in N_{r-k}(X).$$

Remark 3.4.5. Observe that $(\beta^n) > 0$ since β is big.

Proof. By continuity, we can suppose that α_i and b are ample Cartier divisors. We apply successively Siu's inequality by restriction to subschemes representing the classes $\alpha_2 \cdot \dots \cdot \alpha_k_\perp[V]$, $\beta \cdot \alpha_3 \cdot \dots \cdot \alpha_k_\perp[V]$, \dots , $\beta^{k-1} \cdot \alpha_k_\perp[V]$:

$$\begin{aligned} \alpha_1 \cdot \alpha_2 \cdot \dots \cdot \alpha_k_\perp[V] &\leq (r - k + 1) \frac{(\alpha_1 \cdot \dots \cdot \alpha_k \cdot \beta^{r-k})_\perp[V]}{(\beta^{r-k+1} \cdot \alpha_2 \cdot \dots \cdot \alpha_k)_\perp[V]} \times \beta \cdot \alpha_2 \cdot \dots \cdot \alpha_k_\perp[V], \\ \beta \cdot \alpha_2 \cdot \dots \cdot \alpha_k_\perp[V] &\leq (r - k + 1) \frac{(\beta^{r-k+1} \cdot \alpha_2 \cdot \dots \cdot \alpha_k)_\perp[V]}{(\beta^{r-k+2} \cdot \alpha_3 \cdot \dots \cdot \alpha_k)_\perp[V]} \times \beta^2 \cdot \alpha_3 \cdot \dots \cdot \alpha_k_\perp[V], \\ \dots &\dots \\ \beta^{k-1} \cdot \alpha_k_\perp[V] &\leq (r - k + 1) \frac{(\beta^{r-1} \cdot \alpha_k)_\perp[V]}{(\beta^r)_\perp[V]} \times \beta^k_\perp[V]. \end{aligned}$$

This gives the required inequality:

$$\alpha_1 \cdot \dots \cdot \alpha_k_\perp[V] \leq (n - k + 1)^k \frac{(\alpha_1 \cdot \dots \cdot \alpha_k \cdot \beta^{r-k})_\perp[V]}{(\beta^r)_\perp[V]} \times \beta^k_\perp[V].$$

□

Corollary 3.4.6. *Let k be an integer, then for any $a \in \text{BPF}^k(X)$ and any big nef Cartier divisor b on X , one has:*

$$a \leq (n - k + 1)^k \frac{(a \cdot b^{n-k})}{(b^n)} \times b^k.$$

Proof. By linearity and stability by product, we just need to prove the inequality for $a = D_1 \cdot \dots \cdot D_{e_1+k} \in \text{CI}^{e_1+k}(X_1)$, where D_i are ample Cartier divisors X_1 , where $p_1 : X_1 \rightarrow X$ is a flat proper morphism of relative dimension e_1 . We apply Theorem 3.4.4 to $a' = D_{e_1+1} \cdot \dots \cdot D_{e_1+k} \cdot Z$ and $b' = b|_Z$ where $Z = D_1 \cdot \dots \cdot D_{e_1}$. We obtain:

$$a \leq (n - k + 1)^k \frac{(a \cdot b^{n-k})}{(b^n \cdot Z)} \times b^k \cdot Z.$$

As the restriction of p_1 on Z is generically finite, by the projection formula, we get:

$$a \leq (n - k + 1)^k \frac{(a \cdot b^{n-k})}{(b^n)} \times b^k.$$

□

The previous inequality can be applied when we have positivity hypothesis on a birational model as follows.

Corollary 3.4.7. *Let X, Y be two normal projective varieties of dimension n . Let β be a class in $\text{BPF}^k(Y)$, we suppose there exists a birational morphism $q : X \rightarrow Y$ and an ample Cartier divisor A on X such that $A^k \leq q^* \beta$. Then there exists a class $\beta^* \in N_k(X)_\mathbb{R} \cap \text{Psef}_k(X)$ such that for any class $\alpha \in \text{BPF}^k(X)$, we have:*

$$\alpha \leq (\alpha_\perp \beta^*) \times \beta.$$

Proof. We just have to set $\beta^* = \frac{(n-k+1)^k}{(A^n)} q_* \psi_X(A^{n-k})$. □

Remark 3.4.8. We conjecture that for any basepoint free class $a \in \text{BPF}^k(X)$ and any big nef divisor b , one has

$$a \leq \binom{n}{k} \frac{(a \cdot b^{n-k})}{(b^n)} b^k. \quad (12)$$

One can show that this inequality (if true) is optimal since equality can happen when X is an abelian variety.

3.5. Norms on numerical classes. In this section, the positivity properties combined with Siu's inequality allows us to define some norms on $N_k(X)_{\mathbb{R}}$ and on $N^k(X)_{\mathbb{R}}$.

3.5.1. Norms on $N_k(X)_{\mathbb{R}}$. Let $k \leq n$ be an integer and let $\gamma \in \text{BPF}^k(X)$ be a basepoint free class on X . Any cycle $z \in N_k(X)_{\mathbb{R}}$ can be written $z = z^+ - z^-$ where z^+ and z^- are pseudo-effective. We define :

$$F_\gamma(z) := \inf_{\substack{z = z^+ - z^- \\ z^+, z^- \in \text{Psef}_k(X)}} \{(\gamma \cdot z^+) + (\gamma \cdot z^-)\}. \quad (13)$$

Proposition 3.5.1. *For any class $\gamma \in \text{BPF}^k(X)$ lying in the interior of the basepoint free cone, the function F_γ defines a norm on $N_k(X)_{\mathbb{R}}$. In particular, if we fix a norm $\|\cdot\|_{N_k(X)_{\mathbb{R}}}$ on $N_k(X)_{\mathbb{R}}$, there exists a constant $C > 0$ such that for any pseudo-effective class $z \in \text{Psef}_k(X)$, one has:*

$$\frac{1}{C} \|z\|_{N_k(X)_{\mathbb{R}}} \leq (\gamma \cdot z) \leq C \|z\|_{N_k(X)_{\mathbb{R}}}. \quad (14)$$

Proof. The only point to clarify is that $F_\gamma(z) = 0$ implies $z = 0$. Observe that Proposition 3.3.5 implies the result for $z \in \text{Psef}_k(X)$. In general, pick any two sequences $(z_p^+)_{p \in \mathbb{N}}$ and $(z_p^-)_{p \in \mathbb{N}}$ in $\text{Psef}_k(X)$ such that $z = z_p^+ - z_p^-$ and such that $\gamma \cdot z_p^+ + \gamma \cdot z_p^- \rightarrow 0$. Since z_p^+ and z_p^- are pseudo-effective and γ is basepoint free, it follows from Theorem 3.3.3.(5) that

$$\lim_{p \rightarrow +\infty} (\gamma \cdot z_p^+) = \lim_{p \rightarrow +\infty} (\gamma \cdot z_p^-) = 0.$$

As γ lies in the interior of $\text{BPF}^k(X)$, given any β in $\text{BPF}^k(X)$, one has that $C\gamma - \beta$ is still in $\text{BPF}^k(X)$ for some sufficiently large constant $C > 0$. Intersecting with the pseudo-effective classes z_p^+ and z_p^- and using Theorem 3.3.3.(5), we have $\lim_{p \rightarrow \infty} (\beta \cdot z_p^+) = \lim_{p \rightarrow \infty} (\beta \cdot z_p^-) = 0$, thus $(\beta \cdot z) = 0$. Since the basepoint free cone $\text{BPF}^k(X)$ generates all $N^k(X)$ by Theorem 3.3.3.(1), we conclude that $z = 0$ as required. □

3.5.2. Norms on $N^k(X)_{\mathbb{R}}$.

Definition 3.5.2. *We define the subcone $\text{BPF}_0^k(X)$ of $\text{BPF}^k(X)$ as the classes $\alpha \in \text{BPF}^k(X)$ such that for any birational map $q : X' \rightarrow X$, there exists an ample Cartier divisor A on X' such that $q^* \alpha \geq A^k$.*

Proposition 3.5.3. *When X is smooth, the cone $\text{BPF}_0^1(X)$ is equal to the big nef cone. In particular BPF_0^k is neither closed nor open in general.*

Proof. Take $\alpha \in N^1(X)_{\mathbb{R}}$ a big nef divisor. Then for any birational map $q : X' \rightarrow X$ and any ample Cartier divisor A , one has by Theorem 3.4.4 applied to A and $q^* \alpha$:

$$A \leq n \frac{(A \cdot q^* \alpha^{n-1})}{(\alpha^n)} q^* \alpha.$$

Hence, $\alpha \in \text{BPF}_0^1(X)$. Conversely, take a class $\alpha \in \text{BPF}_0^1(X)$, then there exists an ample divisor A on X such that $\alpha \geq A$. Since ample divisors are big, we have that α is big. Moreover, since $\text{BPF}^1(X) = \text{Nef}^1(X) \cap \text{Psef}^1(X)$, we have that α is big and nef as required. \square

Proposition 3.5.4. *The cone $\text{BPF}_0^k(X)$ is a convex open subset of $\text{BPF}^k(X)$ that contains the classes induced by products of big nef divisors.*

Proof. The cone $\text{BPF}_0^k(X)$ contains the products of big and nef Cartier divisors. The fact that $\text{BPF}_0^k(X)$ is convex is a consequence of Siu's inequality. We take two elements α and β in $\text{BPF}_0^k(X)$ and any birational map $q : X' \rightarrow X$. By definition, there exists some ample Cartier divisors A and B on X' such that $q^*\alpha \geq A^k$ and $q^*\beta \geq B^k$. As A and B are ample, there is a constant $C > 0$ such that $A^k \geq CB^k$ using the generalization of Siu's inequality (Theorem 3.4.4). This proves that $q^*(t \times \alpha + (1-t) \times \beta) \geq (tC + (1-t)) \times B^k$ for any $t \in [0, 1]$. Hence $t \times \alpha + (1-t) \times \beta \in \text{BPF}_0^k(X)$ and the cone $\text{BPF}_0^k(X)$ is convex.

We prove that $\text{BPF}_0^k(X)$ is an open subset of $\text{BPF}^k(X)$. We take $\alpha \in \text{BPF}_0^k(X)$. We take any ample Cartier divisor H_X on X such that $\alpha - tH_X^k$ is in $\text{BPF}^k(X)$ for small $t > 0$. We just need to show that $\alpha - tH_X^k$ stays in $\text{BPF}_0^k(X)$ when t is small enough. Let $q : X' \rightarrow X$ be a birational map where X' is projective and normal. By definition of α , there exists an ample Cartier divisor A on X' such that $q^*\alpha \geq A^k$. By Siu's inequality, there exists a constant C such that:

$$q^*H_X^k \leq C \frac{(A^k \cdot q^*H_X^{n-k})}{(H_X^n)} \times A^k.$$

This implies the inequality:

$$q^*\beta - tq^*H_X^k \geq \left(1 - tC \frac{A^k \cdot H_X^{n-k}}{H_X^n}\right) \times A^k. \quad (15)$$

As $A^k \leq q^*\alpha$, we have the following upper bound:

$$(A^k \cdot H_X^{n-k}) \leq (q^*\alpha \cdot q^*H_X^{n-k}).$$

We get the following minoration which depends only on α and H_X :

$$1 - t \frac{C(\alpha \cdot H_X^{n-k})}{(H_X^n)} \leq 1 - t \frac{C(A^k \cdot H_X^{n-k})}{(H_X^n)}. \quad (16)$$

Using (15) and (16), one gets that for $t < \frac{(H_X^n)}{C(\alpha \cdot H_X^{n-k})}$, the class $\alpha - tH_X^k$ is in $\text{BPF}_0^k(X)$. \square

Remark 3.5.5. The cone $\text{BPF}_0^k(X)$ is not always equal to the cone generated by complete intersections. Following [LX15, Example 9.6], there exists a smooth toric threefold such that the cone generated by complete intersections in $\text{N}_1(X)_{\mathbb{R}}$ is not convex, so it cannot be equal to $\text{BPF}_0^2(X)$ using the following proposition.

Let X be a normal projective variety of dimension n . Any class $\alpha \in \text{N}^k(X)_{\mathbb{R}}$ can be decomposed as $\alpha^+ - \alpha^-$ where α^+ and α^- are basepoint free classes. For any $\gamma \in \text{BPF}_0^{n-k}(X)$, we define the function:

$$G_\gamma(\alpha) := \inf_{\substack{\alpha = \alpha^+ - \alpha^- \\ \alpha^+, \alpha^- \in \text{BPF}^k(X)}} \{(\gamma \cdot \alpha^+) + (\gamma \cdot \alpha^-)\}. \quad (17)$$

Proposition 3.5.6. *For any $\gamma \in \text{BPF}_0^{n-k}(X)$, the function G_γ defines a norm on $\mathbb{N}^k(X)_\mathbb{R}$. In particular, for any norm $\|\cdot\|_{\mathbb{N}^k(X)_\mathbb{R}}$ on $\mathbb{N}^k(X)_\mathbb{R}$, there is a constant $C > 0$ such that for any class $\alpha \in \text{BPF}^k(X)$:*

$$\frac{1}{C} \|\alpha\|_{\mathbb{N}^k(X)_\mathbb{R}} \leq (\gamma \cdot \alpha) \leq C \|\alpha\|_{\mathbb{N}^k(X)_\mathbb{R}}.$$

Proof. The only fact which is not immediate is the fact that $G_\gamma(\alpha) = 0$ implies $\alpha = 0$. We are reduced to treat the case where $\alpha \in \text{BPF}^k(X)$.

Suppose first that X is smooth. Since γ belongs to the interior of the basepoint free cone by Proposition 3.5.4, one has that for any basepoint free class $\beta \in \text{BPF}^{n-k}(X)$, there exists a constant $C > 0$ such that:

$$C \|\beta\| \|\gamma - \beta\| \in \text{BPF}^{n-k}(X).$$

In particular, since α is nef, one has:

$$0 = G_\gamma(\alpha) = C \|\beta\| (\gamma \cdot \alpha) \geq (\beta \cdot \alpha) \geq 0.$$

Hence $(\beta \cdot \alpha) = 0$ for any basepoint free class $\beta \in \text{BPF}^{n-k}(X)$ and $\alpha = 0 \in \mathbb{N}^k(X)_\mathbb{R}$ since the basepoint free cone generates all $\mathbb{N}^{n-k}(X)_\mathbb{R}$ by Theorem 3.3.3.(i).

Suppose that X is not smooth. Fix an ample Cartier divisor H_X on X . Take an alteration $\pi : X' \rightarrow X$ of X . Since the morphism $\pi^* : \mathbb{N}^k(X)_\mathbb{R} \rightarrow \mathbb{N}^k(X')_\mathbb{R}$ is injective, we are reduced to prove that $\pi^*\alpha = 0$. Consider $\beta \in \text{BPF}^{n-k}(X)$, we have by the projection formula that:

$$(\pi^*\gamma \cdot \pi^*\alpha) = (\alpha \cdot \gamma).$$

Since γ belongs to the interior of the basepoint free cone, there exists a constant $C > 0$ such that:

$$H_X^{n-k} \leq C\gamma.$$

In particular, this implies that:

$$(\pi^*H_X^{n-k} \cdot \pi^*\alpha) = (H_X^{n-k} \cdot \alpha) = 0.$$

Since π^*H_X is a big nef Cartier divisor, the class $\pi^*H_X^{n-k}$ belongs to $\text{BPF}_0^{n-k}(X')$ by Proposition 3.5.4, hence $\pi^*\alpha = 0$ by the previous argument. \square

Remark 3.5.7. In fact, the above proof gives a stronger statement: for any generically finite morphism $q : X' \rightarrow X$ and any $\gamma \in \text{BPF}_0^{n-k}(X)$, the function $G_{q^*\gamma}$ defines a norm on $\mathbb{N}^k(X')_\mathbb{R}$.

4. RELATIVE NUMERICAL CLASSES

4.1. Relative classes. In this section, we fix $q : X \rightarrow Y$ a surjective proper morphism between normal projective varieties where $\dim X = n$, $\dim Y = l$ and we denote by $e = \dim X - \dim Y$ the relative dimension of q .

Definition 4.1.1. *The abelian group $\mathbb{N}_k(X/Y)$ is the subgroup of $\mathbb{N}_k(X)$ generated by classes of subvarieties V of X such that the image $q(V)$ is a point in Y .*

Observe that by definition, there is a natural injection from $\mathbb{N}_k(X/Y)$ into $\mathbb{N}_k(X)$:

$$0 \longrightarrow \mathbb{N}_k(X/Y) \longrightarrow \mathbb{N}_k(X).$$

Definition 4.1.2. *The abelian group $\mathbb{N}^k(X/Y)$ is the quotient of $Z^k(X)$ by the equivalence relation \equiv_Y where $\alpha \equiv_Y 0$ if for any cycle $z \in Z_k(X)$ whose image by q is a collection of points in Y , we have $(\alpha \cdot z) = 0$.*

Therefore, one has the following exact sequence:

$$N^k(X) \longrightarrow N^k(X/Y) \longrightarrow 0.$$

As before, we write $N_k(X/Y)_{\mathbb{R}} = N_k(X/Y) \otimes_{\mathbb{Z}} \mathbb{R}$, $N^k(X/Y)_{\mathbb{R}} = N^k(X/Y) \otimes \mathbb{R}$, $N_{\bullet}(X/Y) = \bigoplus N_k(X/Y)$ and $N^{\bullet}(X/Y) = \bigoplus N^k(X/Y)$.

Proposition 4.1.3. *The abelian groups $N_k(X/Y)$ and $N^k(X/Y)$ are torsion free and of finite type. Moreover, the pairing $N_k(X/Y)_{\mathbb{Q}} \times N^k(X/Y)_{\mathbb{Q}} \rightarrow \mathbb{Q}$ induced by the pairing $N_k(X)_{\mathbb{Q}} \times N^k(X)_{\mathbb{Q}} \rightarrow \mathbb{Q}$ is perfect.*

Proof. Since $N_k(X/Y)$ is a subgroup of $N_k(X)$, it is torsion free and of finite type. The group $N^k(X/Y)$ is also torsion free. Indeed pick $\alpha \in Z^k(X)$ such that $p\alpha \equiv_Y 0$ for some integer p , then for any cycle z whose image by q is a union of points, we have $(p\alpha \lrcorner z) = p(\alpha \lrcorner z) = 0$ hence $\alpha \equiv_Y 0$. Finally, since there is a surjection from $N^k(X)$ to $N^k(X/Y)$, the group $N^k(X/Y)$ is also of finite type.

Let us show that the pairing is well defined and non degenerate. Take a cycle $z \in Z_k(X)_{\mathbb{Q}}$ such that $q(z)$ is a finite number of points in Y , then if $\alpha \in N^k(X)$ such that its image is 0 in $N^k(X/Y)$, then $(\alpha \lrcorner z) = 0$ and the pairing $N^k(X/Y) \times N_k(X/Y) \rightarrow \mathbb{Z}$ is well-defined. Let us suppose that for any $\alpha \in N^k(X/Y)_{\mathbb{Q}}$, $(\alpha \lrcorner z) = 0$. This implies that for any $\beta \in N^k(X)$, the intersection product $(\beta \lrcorner z) = 0$, thus $z \equiv 0$. Conversely, suppose that $(\alpha \lrcorner z) = 0$ for any $z \in N_k(X/Y)$, then by definition $\alpha \equiv_Y 0$. □

Example 4.1.4. When Y is a point, we have $N_k(X/Y) = N_k(X)$ and $N^k(X/Y) = N^k(X)$.

Example 4.1.5. If the morphism $q : X \rightarrow Y$ is finite, then we have $N^0(X/Y) = N_0(X/Y) = \mathbb{Z}$ and $N^k(X/Y) = N_k(X/Y) = \{0\}$ for $k \geq 1$.

Example 4.1.6. When $k = 1$, the group $N_1(X/Y)$ is generated by curves contracted by q so that $N^1(X/Y)$ is the relative Neron-Severi group and its dimension is the relative Picard number.

Remark 4.1.7. When k is greater than the relative dimension, the relative classes might not be trivial. For example if $q : X \rightarrow Y$ is a birational map, then $e = 0$ but the space $N^1(X/Y)_{\mathbb{R}}$ is generated by classes of exceptional divisors of q .

Proposition 4.1.8. *The intersection product on $N^{\bullet}(X)$ induces a structure of algebra on $N^{\bullet}(X/Y)$. Moreover, the action from $N^{\bullet}(X)$ on $N_{\bullet}(X)$ induces an action from $N^{\bullet}(X/Y)$ on $N_{\bullet}(X/Y)$, so that the vector space $N_{\bullet}(X/Y)_{\mathbb{R}}$ becomes a $N^{\bullet}(X/Y)_{\mathbb{R}}$ -module.*

Proof. Observe that if $z \in Z_k(X)$ such that $q(z)$ is a union of points in Y and $\alpha \in N^l(X)$, then $\alpha \lrcorner z$ lies in $N_{k-l}(X/Y)$. Indeed, by definition, the class $\alpha \lrcorner z$ is represented by a cycle supported in z , so its image by q is a collection of points in Y .

Let us now prove that the product is well-defined in $N^{\bullet}(X/Y)$. Take $\alpha \in N^k(X)$ such that $\alpha = 0$ in $N^k(X/Y)$ and $\beta \in N^l(X)$, we must prove that $\alpha \cdot \beta = 0$ in $N^k(X/Y)$. Pick a cycle $z \in Z_{k+l}(X)$ whose image by q is a collection of points, by the properties of the intersection product, $((\alpha \cdot \beta) \lrcorner z) = (\alpha \lrcorner (\beta \lrcorner z))$. As $\beta \lrcorner z$ is in $N_k(X/Y)$, we get that $((\alpha \cdot \beta) \lrcorner z) = 0$ as required. □

As an illustration, we give an explicit description of these groups in a particular example.

Proposition 4.1.9. *Suppose $q : X = \mathbb{P}(E) \rightarrow Y$ where E is a vector bundle of rank $e + 1$ on Y . Then for any integer $0 \leq k \leq e$, one has:*

$$N_k(X/Y) = \mathbb{Z} \xi^{e-k} \frown q^*[pt],$$

$$N^k(X/Y) = \mathbb{Z} \xi^k,$$

where $\xi = c_1(\mathcal{O}_{\mathbb{P}(E)}(1))$.

Proof. Since the pairing $N^k(X/Y) \times N_k(X/Y) \rightarrow \mathbb{Z}$ is non degenerate and since $(\xi^k \frown (\xi^{e-k} \frown q^*[pt])) = 1$, the second equality is an immediate consequence of the first one. We suppose first that $k > 0$. Pick $\alpha \in Z_k(X)$ which defines a class in $N_k(X/Y)$. Using [Ful98, Theorem 3.3.(b)], α is rationally equivalent to $\sum_{e-k \leq i \leq e} \xi^i \frown q^* \alpha_i$ where α_i is an element of the Chow group $A_{k-e+i}(Y)$. Since the image of α by q is a union of points in Y , we have that $q_* \alpha = 0$ in $A_k(Y)$. Observe that

$$q_*(\xi^e \frown q^* \alpha_e) = \alpha_e,$$

and that for any $i < e$, one has that

$$q_*(\xi^i \frown q^* \alpha_i) = 0$$

since the support of the cycle α_i is of dimension $k - e + i < k$ and $q_*(\xi^e \frown q^* \alpha_i)$ belongs to $A_k(Y)$. Hence the conditions $q_* \alpha = 0$ implies that $\alpha_e = 0$ in $A_k(Y)_{\mathbb{Q}}$. Since $\xi^i \frown \alpha$ defines also a class in $N_{k-i}(X/Y)$, this implies also that $\alpha_{e-i} = 0$ in $A_{k-e+i}(Y)$ for any $i < k$. We have finally that in $N_k(X/Y)$:

$$\alpha = \xi^{e-k} \frown q^* \alpha_{e-k}.$$

Since α_{e-k} belongs to $A_0(Y)$ and $N_0(Y) = \mathbb{Z}[pt]$, the \mathbb{Z} -module $N_k(X/Y)$ is generated by $\xi^{e-k} \frown q^*[pt]$ for $k > 0$.

For $k = 0$, the groups $N_0(X)$ and $N_0(X/Y)$ are isomorphic to \mathbb{Z} , so we get the desired conclusion. \square

4.2. Pullback and pushforward. In this section, we fix any two (proper) surjective morphisms $q_1 : X_1 \rightarrow Y_1$, $q_2 : X_2 \rightarrow Y_2$ between normal projective varieties. To simplify the notation, we write $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ when we have two regular maps $f : X_1 \rightarrow X_2$ and $g : Y_1 \rightarrow Y_2$ such that $q_2 \circ f = g \circ q_1$. When $f : X_1 \dashrightarrow X_2$ and $g : Y_1 \dashrightarrow Y_2$ are merely rational maps, then we write $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$.

Proposition 4.2.1. *Let $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ be a morphism. Then the morphism of abelian groups $f_* : N_k(X_1) \rightarrow N_k(X_2)$ induces a morphism of abelian groups $f_* : N_k(X_1/Y_1) \rightarrow N_k(X_2/Y_2)$.*

Proof. Take a cycle $z \in Z_k(X_1)$ such that $q_1(z)$ is a union of points of Y_1 . Then the image of the cycle z by $q_2 \circ f$ is also a union of points of Y_2 due to the fact that $q_2 \circ f = g \circ q_1$. Hence f_* maps $N_k(X_1/Y_1)$ to $N_k(X_2/Y_2)$. \square

Proposition 4.2.2. *Let $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ be a morphism. Then the morphism of graded rings $f^* : N^\bullet(X_1) \rightarrow N^\bullet(X_2)$ induces a morphism of graded rings $f^* : N^\bullet(X_1/Y_1) \rightarrow N^\bullet(X_2/Y_2)$.*

Proof. This results follows immediately by duality from the previous proposition since the pairing $N^k(X_i/Y_i) \times N_k(X_i/Y_i) \rightarrow \mathbb{Z}$ is non degenerate. \square

4.3. Restriction to a general fiber and relative canonical morphism. Recall that $\dim X = n$, $\dim Y = l$ and that the relative dimension of $q : X \rightarrow Y$ is e .

Proposition 4.3.1. *There exists a unique class $\alpha_{X/Y} \in N^l(X)_{\mathbb{Q}}$ satisfying the following conditions.*

- (1) *The image $\psi_X(\alpha_{X/Y})$ belongs to the subspace $N_e(X/Y)_{\mathbb{Q}}$ of $N_e(X)_{\mathbb{Q}}$.*
- (2) *For any class $\beta \in N_l(X)_{\mathbb{Q}}$, $q_*\beta = (\alpha_{X/Y} \lrcorner \beta) [Y]$.*

Moreover, for any open subset V of Y such that the restriction q to $U = q^{-1}(V)$ is flat, and for all $y \in V$ and any irreducible subvariety F of the scheme-theoretic fiber X_y , we have:

$$\psi_X(\alpha_{X/Y}) = [X_y] = k[F],$$

where k is a rational number which only depends on F and where $[X_y]$ (resp. $[F]$) denotes the fundamental class of X_y (resp. F).

Remark 4.3.2. Recall that by generic flatness (see [FGI⁺05, Theorem 5.12]), one can always find an open subset V of Y such that the restriction of q to $q^{-1}(V)$ is flat over V .

Proof. Fix an ample Cartier divisor H_Y on Y , we set

$$\alpha_{X/Y} := \frac{1}{(H_Y^l)} q^* H_Y^l \in N^l(X)_{\mathbb{Q}}.$$

By definition, it satisfies (1) and (2) follows from the projection formula (Theorem 2.3.2.(4)). Let us show that any class satisfying (1) and (2) is unique. Suppose there is another one $\alpha' \in N^l(X)_{\mathbb{Q}}$. Then for any class $\beta \in N_l(X)_{\mathbb{Q}}$, $((\alpha_{X/Y} - \alpha') \lrcorner \beta) = 0$ so that $\alpha = \alpha'$ since the pairing $N^l(X)_{\mathbb{Q}} \times N_l(X)_{\mathbb{Q}} \rightarrow \mathbb{Q}$ is non degenerate.

Let us prove the last assertion. By generic flatness [FGI⁺05, Theorem 5.12], Let V be an open subset of Y such that the restriction $q|_{q^{-1}(V)} : q^{-1}(V) \rightarrow V$ is flat and such that the dimension of every fiber is e . Since H_Y is ample, we can find some hyperplanes of $H_i \subset Y$ such that $H_1 \cap \dots \cap H_l$ represents the class H_Y^l and such that $H_1 \cap \dots \cap H_l \subset V$. In particular, by [Ful98, Proposition 2.3.(d)], the pullback $q^* H_Y^l$ is represented by a cycle in the fiber of $H_1 \cap \dots \cap H_l$. Denote by $i : V \rightarrow Y$ and $j : U \rightarrow X$ the inclusion maps of V and U into Y and X respectively. The morphisms j and i are open embedding hence are flat. Moreover we have the following commutative diagram.

$$\begin{array}{ccc} U & \xrightarrow{i} & X \\ \downarrow q|_U & & \downarrow q \\ V & \xrightarrow{j} & Y \end{array}$$

Since the class H_Y^l is equal in $A_0(Y)$ to:

$$H_Y^l = \sum a_i [p_i] \tag{18}$$

where $p_i \in V(\kappa)$ are points in V and a_i are positive integers satisfying $\sum a_i = (H_Y^l)$. Using [Ful98, Example 2.4.2], one has that for any $\beta \in A_l(X)$:

$$(q^* H_Y^l \lrcorner \beta) = (q|_U j^*(H_Y^l) \lrcorner i^* \beta).$$

Using (18), one obtains in $A_e(X)$:

$$q|_U j^* H_Y^l = q|_U j^* (\sum a_i [p_i]) = \sum a_i [q^{-1}(p_i)],$$

which is well-defined since the restriction of q on U is flat. By [Ful98, Theorem 10.2], we have that $[X_{p_i}] = [X_y] \in N_e(X)$ for any $p_i, y \in V$. In particular, we have:

$$\psi_X(q^* H_Y^l) = (\sum a_i) [X_y] = (H_Y^l) [X_y] \in N_e(X),$$

where y is a point in V , which proves that $\psi_X(\alpha_{X/Y}) = [X_y]$ in $N_e(X)_{\mathbb{Q}}$ for any point y in V . By the Stein factorization theorem, there exists a morphism $q' : X \rightarrow Y'$ with connected fibres and a finite morphism $f : Y' \rightarrow Y$ such that $q = q' \circ f$. Since $(H_Y^l)[X_y] = q'^* H_{Y'}^l = q'^* f^* H_Y^l$ and since $f^* H_Y^l \in N^l(Y')_{\mathbb{R}}$ which is canonically isomorphic to \mathbb{R} , we have that $f^* H_Y^l = p \cdot [y'] \in N^l(Y')_{\mathbb{R}}$ where p is an integer and where $[y']$ is a general point in $f^{-1}(y)$. We have thus proven that:

$$[X_y] = \frac{p}{(H_Y^l)} [q'^{-1}(y')] \in N_e(X),$$

and $q'^{-1}(y')$ is an irreducible component of X_y as required. \square

The class previously constructed allows us to define a restriction morphism.

Definition 4.3.3. *Suppose that $\dim Y = l$ and that H_Y is an ample Cartier divisor on Y , then we define $\text{Res}_{X/Y} : N_{\bullet}(X)_{\mathbb{Q}} \rightarrow N_{\bullet-l}(X/Y)_{\mathbb{Q}}$ by setting:*

$$\text{Res}_{X/Y}(\beta) := \alpha_{X/Y} \lrcorner \beta.$$

This morphism does not depend on the choice of H_Y .

We shall denote by $\text{Res}_{X/Y}^* : \beta \in N^{\bullet}(X/Y)_{\mathbb{Q}} \rightarrow \alpha_{X/Y} \cdot \beta \in N^{\bullet+l}(X)_{\mathbb{Q}}$ the dual morphism induced by $\text{Res}_{X/Y}$.

Proposition 4.3.4. *Recall that $\dim Y = l$. The following properties are satisfied.*

(1) *For any class $\alpha \in N^{\bullet}(X)_{\mathbb{Q}}$, one has:*

$$\psi_X \circ \text{Res}_{X/Y}^*(\alpha) = \text{Res}_{X/Y} \circ \psi_X(\alpha).$$

(2) *For any morphism $X'/q'Y' \xrightarrow{f} X/qY$ where $\dim X' = \dim X = n$ and $\dim Y' = \dim Y = l$ such that the topological degree of g is d , we have for any $\alpha \in N^{k-l}(X/Y)_{\mathbb{Q}}$:*

$$d \times \text{Res}_{X'/Y'}^* \circ f^* \alpha = f^* \circ \text{Res}_{X/Y}^* \alpha.$$

The definition of the restriction morphism gives a natural way to generalize the definition of the canonical morphism $\psi_X : N^k(X) \rightarrow N_{n-k}(X)$ to the relative case.

Definition 4.3.5. *Recall that the relative dimension of the morphism $q : X \rightarrow Y$ is e . For any integer $k \geq 0$, we define the canonical morphism $\psi_{X/Y}$ by:*

$$\psi_{X/Y} := \psi_X \circ \text{Res}_{X/Y}^* : \beta \in N^k(X/Y)_{\mathbb{Q}} \rightarrow \psi_X(\alpha_{X/Y} \cdot \beta) \in N_{e-k}(X/Y)_{\mathbb{Q}}.$$

Remark 4.3.6. When $k > e$ by convention the map $\psi_{X/Y}$ is zero.

We give here a situation where this map is an isomorphism.

Proposition 4.3.7. *Suppose $q : X \rightarrow Y$ is a smooth morphism of relative dimension e , then for any integer $0 \leq k \leq e$, the map $\psi_{X/Y} : N^k(X/Y)_{\mathbb{Q}} \rightarrow N_{e-k}(X/Y)_{\mathbb{Q}}$ is an isomorphism.*

Proof. Since the pairing $N^k(X/Y)_{\mathbb{Q}} \times N_k(X/Y)_{\mathbb{Q}} \rightarrow \mathbb{Q}$ is perfect by Proposition 4.1.3, we have that the dual morphism $\psi_{X/Y}^* : N^{e-k}(X/Y)_{\mathbb{Q}} \rightarrow N_k(X/Y)_{\mathbb{Q}}$ of $\psi_{X/Y}$ is surjective whenever $\psi_{X/Y} : N^k(X/Y)_{\mathbb{Q}} \rightarrow N_{e-k}(X/Y)_{\mathbb{Q}}$ is injective. We are thus reduced to prove the injectivity of $\psi_{X/Y} : N^k(X/Y)_{\mathbb{Q}} \rightarrow N_{e-k}(X/Y)_{\mathbb{Q}}$. Take $a \in N^k(X/Y)_{\mathbb{Q}}$ such that $\psi_{X/Y}(a) = 0$, and choose a class $\alpha \in N^k(X)_{\mathbb{Q}}$ representing a . We fix a subvariety V of dimension k in a fiber X_y of q where y is a point in Y . We need to prove that $(\alpha \lrcorner [V]) = 0$.

By Proposition 4.3.1, the condition $\psi_{X/Y}(\alpha) = 0$ implies that:

$$\alpha \lrcorner [X_y] = 0 \in N_{e-k}(X)_{\mathbb{Q}}.$$

Since X_y is smooth, by Theorem 2.4.2, there exists a class $\beta \in N^{e-k}(X_y)_{\mathbb{Q}}$ such that:

$$\beta \lrcorner [X_y] = [V].$$

In particular, we get:

$$(\alpha \lrcorner [V]) = (\alpha \lrcorner (\beta \lrcorner [X_y])) = (\beta \lrcorner (\alpha \lrcorner [X_y])) = 0$$

as required. \square

Example 4.3.8. If $X = \mathbb{P}(E)$ where E is a vector bundle on Y , then Proposition 4.1.9 implies that $\psi_{X/Y} : N^k(X/Y)_{\mathbb{Q}} \rightarrow N_{e-k}(X/Y)_{\mathbb{Q}}$ is an isomorphism for any integer $0 \leq k \leq e$.

Example 4.3.9. If X is the blow-up of $\mathbb{P}^1 \times \mathbb{P}^1$ at a point and q is the projection from $\mathbb{P}^1 \times \mathbb{P}^1$ to the first component $Y = \mathbb{P}^1$ composed with the blow-down from X to $\mathbb{P}^1 \times \mathbb{P}^1$. Then the morphism $\psi_{X/Y} : N^0(X/Y)_{\mathbb{Q}} \rightarrow N_1(X/Y)_{\mathbb{Q}}$ is not surjective and $\psi_{X/Y} : N^1(X/Y)_{\mathbb{Q}} \rightarrow N_0(X/Y)_{\mathbb{Q}}$ is not injective.

5. APPLICATION TO DYNAMICS

In this section, we shall consider various normal projective varieties X_i and Y_i respectively of dimension n and l and we write $e = n - l$. Recall from Section 4.2 that the notation $X_i/q_i Y_i$ means that $q_i : X_i \rightarrow Y_i$ is a surjective morphism of relative dimension e and that $X/q \xrightarrow{f} X'/q' Y'$ means that $f : X \dashrightarrow X'$ and $g : Y \dashrightarrow Y'$ are dominant rational maps such that $q' \circ f = g \circ q$. We shall also fix H_{X_i} and H_{Y_i} big and nef Cartier divisors on X_i and Y_i respectively.

In this section we prove Theorem 1 and Theorem 2. They will follow from Theorem 5.2.1 and Theorem 5.3.2 respectively.

5.1. Degrees of rational maps.

Definition 5.1.1. Let us consider a rational map $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ and let Γ_f (resp. Γ_g) be the normalization of the graph of f (resp. g) in $X_1 \times X_2$ (resp. $Y_1 \times Y_2$) such that we have the following diagram.

$$\begin{array}{ccccc}
 & & \Gamma_f & & \\
 & \swarrow \pi_1 & & \searrow \pi_2 & \\
 X_1 & \overset{f}{\dashrightarrow} & & \dashrightarrow & X_2 \\
 \downarrow q_1 & & \varpi & & \downarrow q_2 \\
 Y_1 & \overset{g}{\dashrightarrow} & & \dashrightarrow & Y_2 \\
 & \swarrow \pi'_1 & & \searrow \pi'_2 & \\
 & & \Gamma_g & &
 \end{array}$$

The k -th relative degree of f is defined by the formula:

$$\text{reldeg}_k(f) := (\pi_1^*(H_{X_1}^{e-k} \cdot (q_1^* H_{Y_1})^l) \cdot \pi_2^*(H_{X_2})^k).$$

When Y_1 and Y_2 are reduced to a point, we simply write $\text{deg}_k(f) = \text{reldeg}_k(f)$.

Remark 5.1.2. If $e = 0$, then $\text{reldeg}_k(f) = (q_1^* H_{Y_1}^l)$ if $k = 0$ and $\text{reldeg}_k(f) = 0$ for $k > 0$.

Note that the degrees always depend on the choice of the big nef divisors, but to simplify the notations, we deliberately omit it.

We now explain how to associate to any rational map $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ a pullback operator $(f, g)^{\bullet, k}$.

Definition 5.1.3. Let $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ be a rational map and let π_1 and π_2 be the projections from the graph of f in $X_1 \times X_2$ onto the first and the second factor respectively. We define the linear morphisms $(f, g)^{\bullet, k}$ and $(f, g)_{\bullet, k}$ by the following formula:

$$(f, g)^{\bullet, k} : \alpha \in N^k(X_2/Y_2)_{\mathbb{R}} \longrightarrow (\pi_{1*} \circ \psi_{\Gamma_f/\Gamma_g} \circ \pi_2^*)(\alpha) \in N_{e-k}(X_1/Y_1)_{\mathbb{R}}.$$

$$(f, g)_{\bullet, k} : \beta \in N^k(X_1/Y_1)_{\mathbb{R}} \longrightarrow (\pi_{2*} \circ \psi_{\Gamma_f/\Gamma_g} \circ \pi_1^*)(\beta) \in N_{e-k}(X_2/Y_2)_{\mathbb{R}}.$$

Remark 5.1.4. When Y_1 and Y_2 are reduced to a point, then we simply write $f^{\bullet, k}(\alpha) := (f, \text{Id}_{\{pt\}})^{\bullet, k}(\alpha)$ and $f_{\bullet, k}(\beta) := (f, \text{Id}_{\{pt\}})_{\bullet, k}(\beta)$.

Remark 5.1.5. Since $N^k(X/Y) = 0$ and $N_{e-k}(X) = 0$ when $k > e$, it implies that $(f, g)^{\bullet, k}$ and $(f, g)_{\bullet, k}$ are identically zero for any $k > e$.

5.2. Sub-multiplicativity.

Theorem 5.2.1. Let us consider the composition $X_1/q_1 Y_1 \xrightarrow{f_1} X_2/q_2 Y_2 \xrightarrow{f_2} X_3/q_3 Y_3$ of dominant rational maps. Then for any integer $0 \leq k \leq e$, there exists a constant $C > 0$ which depends only on the choice of H_{X_2} , H_{Y_2} , k , l and e such that:

$$\text{reldeg}_k(f_2 \circ f_1) \leq C \text{reldeg}_k(f_1) \text{reldeg}_k(f_2).$$

More precisely, $C = (e - k + 1)^k / (H_{X_2}^e \cdot q_2^* H_{Y_2}^l)$.

Proof. We denote by Γ_{f_1} (resp. $\Gamma_{f_2}, \Gamma_{g_1}, \Gamma_{g_2}$) the normalization of the graph of f_1 (resp. f_2, g_1, g_2) and π_1, π_2 (resp. $\pi_3, \pi_4, \pi'_1, \pi'_2$ and π'_3, π'_4) the projections onto the first and the second factor respectively. We set Γ as the graph of the rational map $\pi_3^{-1} \circ f_2 \circ \pi_1 : \Gamma_{f_1} \dashrightarrow \Gamma_{f_2}$, u and v the projections from Γ onto Γ_{f_1} and Γ_{f_2} and ϖ_i the restriction on Γ_{f_i} of the projection from $X_i \times X_{i+1}$ to $Y_i \times Y_{i+1}$ for each $i = 1, 2$. We have thus the following diagram.

$$\begin{array}{ccccc}
 & & \Gamma & & \\
 & u \swarrow & & \searrow v & \\
 & \Gamma_{f_1} & & \Gamma_{f_2} & \\
 \pi_1 \swarrow & & & & \swarrow \pi_3 \\
 X_1 & \dashrightarrow & X_2 & \dashrightarrow & X_3 \\
 \downarrow q_1 & & \downarrow q_2 & & \downarrow q_3 \\
 Y_1 & \dashrightarrow & Y_2 & \dashrightarrow & Y_3 \\
 \pi'_1 \swarrow & & \pi'_2 \swarrow & & \pi'_3 \swarrow \\
 & \Gamma_{g_1} & & \Gamma_{g_2} & \\
 \pi_2 \searrow & & \pi_4 \searrow & & \pi_4 \searrow
 \end{array}
 \tag{19}$$

By Proposition 4.3.1 applied to $q_2 \circ \pi_2 \circ u : \Gamma \rightarrow Y_2$, the class $\psi_{\Gamma}(u^* \pi_2^* q_2^* H_{Y_2}^l)$ is represented by the fundamental class $[V]$ where V is a subscheme of dimension e in Γ which is a general fiber of $q_2 \circ \pi_2 \circ u$. We apply Theorem 3.4.4 by restriction to V to the class $a = v^* \pi_4^* H_{X_3}^k \lrcorner [V]$ and $b = u^* \pi_2^* H_{X_2} \lrcorner [V]$. We obtain:

$$v^* \pi_4^* H_{X_3}^k \lrcorner [V] \leq (e - k + 1)^k \frac{(v^* \pi_4^* H_{X_3}^k \cdot u^* \pi_2^* H_{X_2}^{e-k} \lrcorner [V])}{(u^* \pi_2^* H_{X_2}^e \lrcorner [V])} u^* \pi_2^* H_{X_2}^k \lrcorner [V] \in N_{e-k}(\Gamma). \quad (20)$$

Let us simplify the right hand side of inequality (20). Since $\pi_2 \circ u = \pi_3 \circ v$, $\psi_\Gamma(u^* \pi_2^* q_2^* H_{Y_2}^l) = [V] \in N_e(\Gamma)$ and since the morphism v is generically finite, one has that:

$$(v^* \pi_4^* H_{X_3}^k \cdot u^* \pi_2^* H_{X_2}^{e-k} \lrcorner [V]) = (v^*(\pi_4^* H_{X_3}^k \cdot \pi_3^* H_{X_2}^{e-k} \cdot \pi_3^* q_2^* H_{Y_2}^l)) = d \times \text{reldeg}_k(f_2), \quad (21)$$

where d is the topological degree of v . The same argument gives:

$$(u^* \pi_2^* H_{X_2}^e \lrcorner [V]) = d \times (H_{X_2}^e \cdot q_2^* H_{Y_2}^l). \quad (22)$$

Using (21), (22), inequality (20) can be rewritten as:

$$u^* \pi_2^* q_2^* H_{Y_2}^l \cdot v^* \pi_4^* H_{X_3}^k \leq C \text{reldeg}_k(f_2) u^* \pi_2^* H_{X_2}^k \cdot u^* \pi_2^* q_2^* H_{Y_2}^l \in N^{l+k}(\Gamma),$$

where $C = (e - k + 1)^k / (H_{X_2}^e \cdot q_2^* H_{Y_2}^l)$. Since the class $u^* \pi_1^* H_{X_1}^{e-k} \in N^{e-k}(\Gamma)$ is nef, we can intersect this class in the previous inequality to obtain:

$$(u^*(\pi_1^* H_{X_1}^{n-l-k} \cdot \pi_2^* q_2^* H_{Y_2}^l) \cdot v^* \pi_4^* H_{X_3}^k) \leq C' \text{reldeg}_k(f_2) (u^* \pi_2^* H_{X_2}^k \cdot u^* \pi_2^* q_2^* H_{Y_2}^l \cdot u^* \pi_1^* H_{X_1}^{n-l-k}). \quad (23)$$

Let us simplify the expressions in inequality (23). Because $\pi_2^* q_2^* H_{Y_2}^l = \varpi_1^* \pi_2^* H_{Y_2}^l$ and $\text{deg}_l(g_1) = (\pi_2'^* H_{Y_2}^l)$, we deduce that:

$$\pi_2^* q_2^* H_{Y_2}^l = \frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} \varpi_1^* \pi_1^* H_{Y_1}^l = \frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} \pi_1^* q_1^* H_{Y_1}^l. \quad (24)$$

Applying (24), the inequality (23) can be translated as:

$$\frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} (u^* \pi_1^* (H_{X_1}^{n-l-k} \cdot q_1^* H_{Y_1}^l) \cdot v^* \pi_4^* H_{X_3}^k) \leq C \frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} \text{reldeg}_k(f_2) (u^*(\pi_2^* H_{X_2}^k \cdot \pi_1^* q_1^* H_{Y_1}^l \cdot \pi_1^* H_{X_1}^{n-l-k})).$$

We obtain thus:

$$\frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} \text{reldeg}_k(f_2 \circ f_1) \leq C \frac{\text{deg}_l(g_1)}{(H_{Y_1}^l)} \text{reldeg}_k(f_1) \text{reldeg}_k(f_2).$$

This concludes the proof of the inequality after dividing by $\text{deg}_l(g_1)/(H_{Y_1}^l)$. \square

5.3. Norms of operators associated to rational maps. The proof of Theorem 2 relies on an easy but crucial lemma which is as follows.

Lemma 5.3.1. *Let us consider $(V, \|\cdot\|)$ a finite dimensional normed \mathbb{R} -vector space and let \mathcal{C} be a closed convex cone with non-empty interior in V . Then there exists a constant $C > 0$ such that any vector $u \in V$ can be decomposed as $v = v^+ - v^-$ where u^+ and u^- are in \mathcal{C} such that:*

$$\|v^{+/-}\| \leq C \|v\|.$$

Proof. Let us define the map $f : V \rightarrow \mathbb{R}^+$ given by:

$$f(v) = \inf\{\|v'\| + \|v' - v\| \mid v' \in \mathcal{C}, v' - v \in \mathcal{C}\}.$$

We check easily that f defines a norm on V which is similar to the proof of Proposition 3.5.1. Since V is finite dimensional, there exists a constant C such that for any $v \in V$, one has:

$$f(v) \leq C \|v\|,$$

Hence $\|v^+\| \leq C \|v\|$ and $\|v^-\| \leq C \|v\|$. \square

Theorem 5.3.2. *Let $X/qY \xrightarrow[-g]{f} X/qY$ be a rational map. We fix an integer $k \leq e$, some norms on $N^k(X/Y)_{\mathbb{R}}$, on $N_{e-k}(X/Y)_{\mathbb{R}}$. Then there is a constant $C > 0$ such that for any rational map $X/qY \xrightarrow[-g]{f} X/qY$, we have:*

$$\frac{1}{C} \leq \frac{\|(f, g)^{\bullet, k}\|}{\text{reldeg}_k(f)} \leq C.$$

In particular, the k -th relative dynamical degree of f satisfies the following equality:

$$\lambda_k(f, X/Y) = \lim_{p \rightarrow +\infty} \|(f^p, g^p)^{\bullet, k}\|^{1/p}.$$

Moreover, when Y is reduced to a point, we obtain:

$$\lambda_k(f) = \lim_{p \rightarrow +\infty} \|(f^p)^{\bullet, k}\|^{1/p}.$$

Remark 5.3.3. The proof of Theorem 2 follows directly from Theorem 5.3.2 since $N^k(X/Y) = N^k(X)$ and $N_{e-k}(X/Y) = N_{e-k}(X)$ when Y is reduced to a point.

Proof. We denote by π_1 and π_2 the projections from the graph Γ_f of f onto the first and the second component respectively. Since we want to control the norm of $f^{\bullet, k}$ by the k -th relative degree of f , we first find an appropriate norm to relate the norm on $N_{e-k}(X)_{\mathbb{R}}$ with an intersection product. As $N_{e-k}(X/Y)_{\mathbb{R}}$ is a subspace of $N_{e-k}(X)_{\mathbb{R}}$, we can extend the norm $\|\cdot\|_{N_{e-k}(X/Y)_{\mathbb{R}}}$ into a norm on $N_{e-k}(X)_{\mathbb{R}}$. As $N_{e-k}(X)_{\mathbb{R}}$ is a finite dimensional vector space and since H_X^{e-k} is a class in the interior of the basepoint free cone $\text{BPF}^{e-k}(X)$, we can suppose by equivalence of norms that the norm on $N_{e-k}(X)_{\mathbb{R}}$ given by

$$\|z\| = \inf_{\substack{z = z^+ - z^- \\ z^+, z^- \in \text{Psef}_{e-k}(X)}} \{(H_X^{e-k} \frown z^+) + (H_X^{e-k} \frown z^-)\}$$

as in Proposition 3.5.1.

Let us prove that the lower bound of $\|(f, g)^{\bullet, k}\|/\text{reldeg}_k(f)$ is 1. We denote by $\varphi : N^k(X) \rightarrow N^k(X/Y)$ the canonical surjection. Since H_X^k is basepoint free, it implies that the class $(f, g)^{\bullet}(\varphi(H_X^k)) \in N_{e-k}(X/Y)_{\mathbb{R}} \subset N_{e-k}(X)_{\mathbb{R}}$ is pseudo-effective. In particular, this implies that its norm is exactly $\text{reldeg}_k(f)$. We have thus by definition:

$$\frac{\|(f, g)^{\bullet, k}\|}{\text{reldeg}_k(f)} = \left(\frac{\|(f, g)^{\bullet, k}\|}{\|(f, g)^{\bullet, k} \varphi(H_X^k)\|} \right) \geq 1,$$

as required.

Let us find an upper bound for $\|(f, g)^{\bullet, k}\|/\|(f, g)^{\bullet, k} \varphi(H_X^k)\|$. First we fix a morphism $s : N^k(X/Y)_{\mathbb{R}} \rightarrow N^k(X)_{\mathbb{R}}$ such that $\varphi \circ s = \text{Id}$. Take $\alpha \in N^k(X/Y)_{\mathbb{R}}$ of norm 1, then the class $u = s(\alpha) \in N^k(X)_{\mathbb{R}}$ is a representant of α . By construction, the norm of u is bounded by $\|u\|_{N^k(X)_{\mathbb{R}}} \leq C_1 \|\alpha\|_{N^k(X/Y)_{\mathbb{R}}} = C_1$ where C_1 is the norm of the operator s . Since by Proposition 4.3.4.(2), $\text{Res}_{\Gamma_f/\Gamma_g}^* \circ \pi_2^* = (1/\text{deg}_i(g)) \times \pi_2^* \circ \text{Res}_{X/Y}^*$, we have therefore:

$$(f, g)^{\bullet, k} \alpha = \frac{1}{\text{deg}_i(g)} \times \pi_{1*} \circ \psi_{\Gamma_f} \circ \pi_2^* \circ \text{Res}_{X/Y}^*(\alpha) = \text{Res}_{X/Y} f^{\bullet, k} u.$$

By Theorem 3.3.3, the pliant cone $\text{BPF}^k(X)$ has a non-empty interior in $N^k(X)_{\mathbb{R}}$ and we can apply Lemma 5.3.1. There exists a constant $C_2 > 0$ which depends only on $\text{BPF}^k(X)$ and the choice of the norm on $N^k(X)_{\mathbb{R}}$ such that the class u can be decomposed as $u = u_1 - u_2$ where $u_i \in \text{BPF}^k(X)$

such that $\|u_i\|_{N^k(X)_\mathbb{R}} \leq C_2 \|u\|_{N^k(X)_\mathbb{R}}$ for $i = 1, 2$. We set $\alpha_i = \varphi(u_i)$ for all $i \in \{1, 2\}$. By the triangular inequality, we have:

$$\frac{\|(f, g)^{\bullet, k} \alpha\|_{N_{e-k}(X/Y)_\mathbb{R}}}{\|(f, g)^{\bullet, k} \varphi(H_X^k)\|} \leq \frac{\|(f, g)^{\bullet, k} \alpha_1\|_{N_{e-k}(X)_\mathbb{R}}}{\|(f, g)^{\bullet, k} \varphi(H_X^k)\|} + \frac{\|(f, g)^{\bullet, k} \alpha_2\|_{N_{e-k}(X)_\mathbb{R}}}{\|(f, g)^{\bullet, k} \varphi(H_X^k)\|}.$$

We have to find an upper bound of $\|(f, g)^{\bullet, k} \alpha_i\|_{N_{e-k}(X/Y)_\mathbb{R}}$ for each $i = 1, 2$. Applying Corollary 3.4.6 to $a = \pi_2^* u_i$ and $b = \pi_2^* H_X$ and then composing with $\text{Res}_{X/Y} \circ \pi_{1*} \circ \psi_{\Gamma_f}$ gives

$$\text{Res}_{X/Y}(f^{\bullet, k}(u_i)) \leq C_3 \frac{\|u_i\|_{N^k(X)_\mathbb{R}}}{(H_X^n)} \times \text{Res}_{X/Y}(f^{\bullet, k}(H_X^k)),$$

where C_3 is a positive constant which depends only on the choice of big nef divisors. This implies by intersecting with H_X^{e-k} the inequality:

$$\|((f, g)^{\bullet, k}(\alpha_i))\|_{N_{e-k}(X/Y)_\mathbb{R}} \leq C_3 \frac{\|u_i\|_{N^k(X)_\mathbb{R}}}{(H_X^n)} \|(f, g)^{\bullet, k}(\varphi(H_X^k))\|_{N_{e-k}(X/Y)_\mathbb{R}}.$$

In particular we have shown that:

$$1 \leq \frac{\|(f, g)^{\bullet, k} \alpha\|_{N_{e-k}(X/Y)_\mathbb{R}}}{\|(f, g)^{\bullet, k} \varphi(H_X^k)\|_{N_{e-k}(X/Y)_\mathbb{R}}} \leq \frac{2C_1 C_2 C_3}{(H_X^n)},$$

which concludes the proof. □

6. SEMI-CONJUGATION BY DOMINANT RATIONAL MAPS

In this section, we consider a more general situation than in the previous section. We still suppose that the varieties X_i and Y_i are of dimension n and l respectively such that the relative dimension is $e = n - l$, but we suppose the maps $q_i : X_i \dashrightarrow Y_i$ merely *rational* and dominant: they may exhibit indeterminacy points. Recall also that H_{X_i} and H_{Y_i} are again big and nef Cartier divisors on X_i and Y_i respectively.

Definition 6.0.4. *Let $f : X_1 \dashrightarrow X_2$, $g : Y_1 \dashrightarrow Y_2$, $q_1 : X_1 \dashrightarrow Y_1$ and $q_2 : X_2 \dashrightarrow Y_2$ be four dominant rational maps such that $q_2 \circ f = g \circ q_1$. We define the k -th relative dynamical degree of f (still denoted $\text{reldeg}_k(f)$) as the relative degree $\text{reldeg}_k(\tilde{f})$ with respect to the rational map $\Gamma_{q_1}/Y_1 \xrightarrow[\tilde{g}]{\tilde{f}} \Gamma_{q_2}/Y_2$ where Γ_{q_i} are the normalization of the graphs of q_i in $X_i \times Y_i$ for each integer $i \in \{1, 2\}$ respectively and $\tilde{f} : \Gamma_{q_1} \dashrightarrow \Gamma_{q_2}$ is the rational map induced by f .*

Theorem 6.0.5. (1) *Consider now the following commutative diagram:*

$$\begin{array}{ccccc} X_1 & \xrightarrow{f_1} & X_2 & \xrightarrow{f_2} & X_3 \\ \downarrow q_1 & & \downarrow q_2 & & \downarrow q_3 \\ Y_1 & \xrightarrow{g_1} & Y_2 & \xrightarrow{g_2} & Y_3 \end{array}$$

where $f_i : X_i \dashrightarrow X_{i+1}$, $g_i : Y_i \dashrightarrow Y_{i+1}$, $q_1 : X_1 \dashrightarrow Y_1$, $q_2 : X_2 \dashrightarrow Y_2$, $q_3 : X_3 \dashrightarrow Y_3$ are dominant rational maps for any integer $i \in \{1, 2, 3\}$ such that $q_{i+1} \circ f_i = g_i \circ q_i$ for any integer $i \in \{1, 2\}$. Then there exists a constant $C > 0$ which depends only e, k and the choice of big nef Cartier divisors such that:

$$\text{reldeg}_k(f_2 \circ f_1) \leq C \text{reldeg}_k(f_2) \text{reldeg}_k(f_1).$$

(2) Consider now the following commutative diagram:

$$\begin{array}{ccccc}
 & & X'_1 & \overset{\tilde{f}}{\dashrightarrow} & X'_2, \\
 & & \swarrow \varphi_1 & & \searrow \varphi_2 \\
 X_1 & \overset{f}{\dashrightarrow} & X_2 & & \\
 \downarrow q_1 & & \downarrow q_2 & & \downarrow \\
 Y_1 & \overset{g}{\dashrightarrow} & Y_2 & & \\
 & & \swarrow \phi_1 & & \searrow \phi_2 \\
 & & Y'_1 & \overset{\tilde{g}}{\dashrightarrow} & Y'_2 \\
 & & \downarrow & & \downarrow
 \end{array}$$

where $f : X_1 \dashrightarrow X_2$, $g : Y_1 \dashrightarrow Y_2$, $q_1 : X_1 \dashrightarrow Y_1$, $q_2 : X_2 \dashrightarrow Y_2$ are four dominant rational maps such that $q_2 \circ f = g \circ q_1$. We consider some birational maps $\varphi_i : X'_i \dashrightarrow X_i$ and $\phi_i : Y'_i \dashrightarrow Y_i$ for $i = 1, 2$ such that $\tilde{f} = \varphi_2^{-1} \circ f \circ \varphi_1$ and $\tilde{g} = \phi_2^{-1} \circ g \circ \phi_1$. Then for any integer $0 \leq k \leq e$, there exists a constant $C > 0$ which depends on e, k and on the choice of big nef Cartier divisors such that:

$$\frac{1}{C} \operatorname{reldeg}_k(f) \leq \operatorname{reldeg}_k(\tilde{f}) \leq C \operatorname{reldeg}_k(f). \quad (25)$$

Proof. (1) Let $\tilde{f}_i : \Gamma_{q_i} \dashrightarrow \Gamma_{q_{i+1}}$ be the rational maps induced by f_i on the graph Γ_{q_i} of q_i for $i \in \{1, 2\}$ respectively. Then (1) results directly from Theorem 5.2.1 applied to the composition $\Gamma_{q_1}/Y_1 \xrightarrow[\tilde{g}_1]{\tilde{f}_1} \Gamma_{q_2}/Y_2 \xrightarrow[\tilde{g}_2]{\tilde{f}_2} \Gamma_{q_3}/Y_3$.

(2) Let us suppose first that the maps $q_i : X_i \rightarrow Y_i$ and $q'_i : X'_i \rightarrow Y'_i$ are all regular for $i = 1, 2$.

Let us apply successively Theorem 5.2.1 to the composition $X'_1/q'_1 Y'_1 \xrightarrow[\phi_1]{\varphi_1} X_1/q_1 Y_1 \xrightarrow[g]{f} X_2/q_2 Y_2 \xrightarrow[\phi_2^{-1}]{\varphi_2^{-1}} X'_2/q'_2 Y'_2$. We obtain :

$$\operatorname{reldeg}_k(\varphi_2^{-1} \circ f \circ \varphi_1) \leq C_2 \operatorname{reldeg}_k(f \circ \varphi_1) \operatorname{reldeg}_k(\varphi_2^{-1}) \leq C_1 C_2 \operatorname{reldeg}_k(f) \operatorname{reldeg}_k(\varphi_1) \operatorname{reldeg}_k(\varphi_2^{-1}), \quad (26)$$

where $C_1 = (e - k + 1)^k / (H_{X_1}^e \cdot q_1^* H_{Y_1}^l)$ and $C_2 = (e - k + 1)^k / (H_{X_2}^e \cdot q_2^* H_{Y_2}^l)$. This proves that:

$$\operatorname{reldeg}_k(\varphi_2^{-1} \circ f \circ \varphi_1) \leq C \operatorname{reldeg}_k(f),$$

where

$$C = \frac{(e - k + 1)^{2k} \operatorname{reldeg}_k(\varphi_1) \operatorname{reldeg}_k(\varphi_2^{-1})}{(H_{X_1}^e \cdot q_1^* H_{Y_1}^l)(H_{X_2}^e \cdot q_2^* H_{Y_2}^l)}.$$

The proof follows easily from the regular case since the maps $\Gamma_{q'_1} \dashrightarrow \Gamma_{q_1}$ and $\Gamma_{q'_2} \dashrightarrow \Gamma_{q_2}$ are birational where $\Gamma_{q'_i}$ are the graphs of q'_i in $X'_i \times Y'_i$ for $i = 1, 2$. \square

Proof of Theorem 1: (1) We apply Theorem 5.2.1 to $Y_1 = Y_2 = Y_3 = \operatorname{Spec}(\kappa)$, $X_1 = X_2 = X_3 = X$ and $H_{X_1} = H_{X_2} = H_{X_3} = H_X$, we get thus the desired conclusion:

$$\operatorname{deg}_k(g \circ f) \leq \frac{(n - k + 1)^k}{(H_X^n)} \operatorname{deg}_k(f) \operatorname{deg}_k(g).$$

(2) Applying Theorem 6.0.5.(2) to the varieties $X'_1 = X'_2 = X_1 = X_2 = X$, $Y'_1 = Y'_2 = Y_1 = Y_2 = \operatorname{Spec}(\kappa)$, to the choice of big nef divisors $H_{X'_1} = H_{X'_2} = H'_X$, $H_{Y'_1} = H_{Y'_2} = H'_Y$, $H_{X_1} = H_{X_2} = H_X$ and to the rational maps $\varphi_1 = \varphi_2 = \operatorname{Id}_X$, $\phi_1 = \phi_2 = g = \operatorname{Id}_{\operatorname{Spec}(\kappa)}$, $f : X \dashrightarrow X$ yields the desired result.

7. MIXED DEGREE FORMULA

Let us consider three dominant rational maps $f : X \dashrightarrow X$, $q : X \dashrightarrow Y$, $g : Y \dashrightarrow Y$ such that $q \circ f = g \circ q$. Theorem 6.0.5.(1) implies that for any integer $k \leq e$ the sequence $\text{reldeg}_k(f^n)$ is submultiplicative. Define k -th relative dynamical degree as follows.

$$\lambda_k(f, X/Y) := \lim_{p \rightarrow +\infty} (\text{reldeg}_k(f^p))^{1/p}.$$

When Y is reduced to a point, then we simply write $\lambda_k(f) := \lambda_k(f, X/\{pt\})$.

Remark 7.0.6. Since $\text{reldeg}_k(f^p) \in \mathbb{N}$ is an integer, one has that $\lambda_k(f, X/Y) \geq 1$.

Remark 7.0.7. Theorem 6.0.5.(2) implies that $\lambda_k(f, X/Y)$ does not depend on the choice of big nef Cartier divisors and on any choice of varieties X' and Y' which are birational to X and Y respectively.

Our aim in this section is to prove Theorem 4. To that end, we follow the approach from [DNT12]. The main ingredient (Corollary 7.1.5) is an inequality relating basepoint free classes which generalizes to arbitrary fields (see [DN11, Proposition 2.3] and [DNT12, Proposition 2.5]). This inequality is a direct consequence of Theorem 7.1.1 which estimates the positivity of the diagonal in a quite general setting. After this, we prove in Theorem 7.2.3 the submultiplicativity formula for the mixed degrees. Once the submultiplicativity of these mixed degrees holds, the proof follows from a linear algebra argument.

7.1. Positivity estimate of the diagonal. In this section, we prove the following theorem.

Theorem 7.1.1. *Let $q : X \rightarrow Y$ be a surjective morphism such that $\dim Y = l$ and such that q is of relative dimension e . There exists a constant $C > 0$ such that for any surjective generically finite morphism $\pi : X' \rightarrow X$ and any class $\gamma \in \text{BPF}^{l+e}(X' \times X')$:*

$$(\gamma \llcorner [\Delta_{X'}]) \leq C \times (\gamma \cdot (\pi \times \pi)^*(H_X^e \cdot H_Y^l)), \quad (27)$$

where p_1 and p_2 are the projections from $X' \times X'$ to the first and the second factor respectively, $H_X = p_1^*H_X + p_2^*H_X$ and $H_Y = p_1^*q^*H_Y + p_2^*q^*H_Y$, and where $\Delta_{X'}$ (resp. Δ_X) is the diagonal of X' (resp. of X) in $X' \times X'$ (resp. in $X \times X$).

Remark 7.1.2. The fact that the constant $C > 0$ does not depend on π but only on H_X , H_Y is crucial in the applications. Moreover, we believe that the following conjecture, which is a stronger version of (27), should be true:

$$[\Delta_{X'}] \leq C \psi_{X' \times X'}((\pi \times \pi)^*(H_X^e \cdot H_Y^l)) \in N_{l+e}(X' \times X')_{\mathbb{R}}. \quad (28)$$

We shall use several times the following lemma.

Lemma 7.1.3. *Let $X_1/q_1 Y_1 \xrightarrow[f]{g} X_2/q_2 Y_2$ be two dominant rational maps where $\dim Y_1 = \dim Y_2 = l$ and $\dim X_1 = \dim X_2 = e + l$. We denote by Γ_f and Γ_g the normalizations of the graph of f and g in $X_1 \times X_2$ and $Y_1 \times Y_2$ respectively, $\pi_1, \pi_2, \pi'_1, \pi'_2$ are the projections from Γ_f and Γ_g on the first and the second factor respectively. Then there exists a constant $C > 0$ such that for any surjective generically finite morphism $\pi : X' \rightarrow \Gamma_f$, any integer $0 \leq j \leq l$ and any class $\beta \in \text{BPF}^{e+l-j}(X')$, one has:*

$$(\beta \cdot \pi^* \pi_2^* q_2^* H_{Y_2}^j) \leq C \frac{\text{deg}_j(g)}{(H_{Y_1}^l)} \times (\beta \cdot \pi^* \pi_1^* q_1^* H_{Y_1}^j),$$

where $\text{deg}_j(g)$ is the j -th degree of the rational map g with respect to the divisors H_{Y_1} and H_{Y_2} .

Proof of Theorem 7.1.1. By Siu's inequality, we can suppose that both the classes H_X and H_Y are ample in X and Y respectively. We proceed in three steps. Fix $\pi : X' \rightarrow X$.

Step 1: We suppose first that $X = \mathbb{P}^l \times \mathbb{P}^e$, $Y = \mathbb{P}^l$ and q is the projection onto the first factor. Since $X \times X$ is smooth, the pullback $(\pi \times \pi)^*$ is well-defined in $N_{l+e}(X \times X)_{\mathbb{R}}$ because the morphism $\psi_{X \times X} : N^{l+e}(X \times X)_{\mathbb{R}} \rightarrow N_{l+e}(X \times X)_{\mathbb{R}}$ is an isomorphism. Our objective is to prove that there exists a constant $C_1 > 0$ such that

$$[\Delta_{X'}] \leq C_1 \times \psi_{X' \times X'}((\pi \times \pi)^*(H_X^e \cdot H_Y^l)) \in N_{l+e}(X' \times X')_{\mathbb{R}}.$$

As $X \times X$ is homogeneous, we apply the following lemma analogous to [Tru16, Lemma 4.4] which we prove at the end of the section.

Lemma 7.1.4. *Let X be a homogeneous projective variety of dimension n and let $\pi : X' \rightarrow X$ be a surjective generically finite morphism. Then one has that :*

$$[\Delta_{X'}] \leq (\pi \times \pi)^*[\Delta_X] \in N_n(X' \times X')_{\mathbb{R}}.$$

We denote by p'_1, p'_2 (resp. p''_1, p''_2) the projections from $Y \times Y$ (resp. from $X \times X$) onto the first and the second factor respectively. Since the basepoint free cone has a non-empty interior by Theorem 3.3.3.(1) and since the class $p_1^* H_Y + p_2^* H_Y$ is ample on $Y \times Y$, there exists a constant $C_2 > 0$ such that the class $-[\Delta_Y] + C_2(p_1^* H_Y + p_2^* H_Y)^l \in N^l(Y \times Y)_{\mathbb{R}}$ is basepoint free. Since $\Delta_X = \Delta_Y \times \Delta_{\mathbb{P}^e}$ and by intersection and by pullback, we have that the class:

$$-[\Delta_X] + C_2 \times H_Y^l \cdot p^*[\Delta_{\mathbb{P}^e}] \in N^{e+l}(X \times X)$$

is basepoint free where p denotes the projection from $X \times X$ to $\mathbb{P}^e \times \mathbb{P}^e$. By the same argument, there exists a constant $C_3 > 0$ such that the class $-p^*[\Delta_{\mathbb{P}^e}] + C_3 H_X^e \in N^e(X \times X)_{\mathbb{R}}$ is basepoint free. We have proved that the class:

$$-[\Delta_X] + C_2 C_3 H_Y^l \cdot H_X^e \in N^{e+l}(X \times X)_{\mathbb{R}}$$

is basepoint free. Since the basepoint free cone is stable by pullback, we have thus:

$$[\Delta_{X'}] \leq (\pi \times \pi)^*[\Delta_X] \leq C_1 \times \psi_{X' \times X'}((\pi \times \pi)^*(H_Y^l \cdot H_X^e)) \in N_{l+e}(X' \times X')_{\mathbb{R}},$$

where $C_1 = C_2 \times C_3$ as required.

Step 2: We now suppose that $X = Y \times \mathbb{P}^e$. Since Y is projective, there exists a dominant rational map $\phi : Y \dashrightarrow \mathbb{P}^l$. Let Y' be the normalization of the graph of ϕ in $Y \times \mathbb{P}^e \times \mathbb{P}^l$ and we denote by ϕ_1 and φ_1 the projections from Y' onto the first and the second factor respectively. Let $\varphi_2 : Y' \times \mathbb{P}^e \rightarrow \mathbb{P}^l \times \mathbb{P}^e$ (resp. $\phi_2 : Y' \times \mathbb{P}^e \rightarrow X$) the map induced by φ_1 (resp. ϕ_1). Let X'' be the fibred product of X' with $Y' \times \mathbb{P}^e$ so that ϕ_3, π' are the projections from X'' onto X' and $Y' \times \mathbb{P}^e$ respectively. We obtain the following commutative diagram:

$$\begin{array}{ccccc}
 X' & \xleftarrow{\phi_3} & X'' & & \\
 \downarrow \pi & & \downarrow \pi' & & \\
 Y \times \mathbb{P}^e & \xleftarrow{\phi_2} & Y' \times \mathbb{P}^e & \xrightarrow{\varphi_2} & \mathbb{P}^l \times \mathbb{P}^e \\
 \downarrow q & & \downarrow p_{Y'} & & \downarrow p_{\mathbb{P}^l} \\
 Y & \xleftarrow{\phi_1} & Y' & \xrightarrow{\varphi_1} & \mathbb{P}^l
 \end{array}$$

where $p_{Y'}$ and $p_{\mathbb{P}^l}$ are the projections from $Y' \times \mathbb{P}^e$ and $\mathbb{P}^l \times \mathbb{P}^e$ onto Y' and \mathbb{P}^l respectively and where the horizontal arrows are birational maps. Let us prove that there exists a constant $C_4 > 0$ which does not depend on the morphism $\pi : X' \rightarrow X$ such that for any basepoint free class $\gamma' \in \text{BPF}^{e+l}(X'' \times X'')$, one has:

$$(\gamma' \lrcorner [\Delta_{X''}]) \leq C_4 (\gamma' \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^*(H_X^e \cdot H_Y^l)).$$

Fix a class $\gamma' \in \text{BPF}^{e+l}(X'' \times X'')$. We apply the conclusion of the first step to the surjective generically finite morphism $\pi'' := \varphi_2 \circ \pi' : X'' \rightarrow \mathbb{P}^l \times \mathbb{P}^e$. There exists a constant $C_1 > 0$ such that

$$[\Delta_{X''}] \leq C_1 \psi_{X'' \times X''}((\pi'' \times \pi'')^*(H_{\mathbb{P}^l}^l \cdot H_{\mathbb{P}^l \times \mathbb{P}^e}^e)) \in N_{l+e}(X'' \times X'')_{\mathbb{R}}, \quad (29)$$

where $H_{\mathbb{P}^l \times \mathbb{P}^e}$ is an ample Cartier divisor in $(\mathbb{P}^l \times \mathbb{P}^e)^2$ and $H_{\mathbb{P}^l}$ is the pullback by $p_{\mathbb{P}^l} \times p_{\mathbb{P}^l}$ of an ample Cartier divisor in $\mathbb{P}^l \times \mathbb{P}^l$. Let us apply Theorem 3.4.4 to the class $(\pi'' \times \pi'')^* H_{\mathbb{P}^l \times \mathbb{P}^e}^e$ and to the class $(\pi' \times \pi')^*(\phi_2 \times \phi_2)^* H_X$, there exists a constant $C_5 > 0$ such that:

$$\begin{aligned} (\pi'' \times \pi'')^* H_{\mathbb{P}^l \times \mathbb{P}^e}^e &\leq C_5 \frac{((\pi' \times \pi')^*((\phi_2 \times \phi_2)^* H_X^{2l+e} \cdot (\varphi_2 \times \varphi_2)^* H_{\mathbb{P}^l \times \mathbb{P}^e}^e))}{((\pi' \times \pi')^*(\phi_2 \times \phi_2)^* H_X^{2(l+e)})} \\ &\quad \times (\pi' \times \pi')^*(\phi_2 \times \phi_2)^* H_X^e \in N^e(X'' \times X'')_{\mathbb{R}}. \end{aligned}$$

Since $((\pi' \times \pi')^* \alpha) = \deg(\pi')(\alpha)$ for any class $\alpha \in N^{2l+2e}((Y' \times \mathbb{P}^e)^2)_{\mathbb{R}}$, we have thus:

$$(\pi'' \times \pi'')^* H_{\mathbb{P}^l \times \mathbb{P}^e}^e \leq C_6 (\pi' \times \pi')^*(\phi_2 \times \phi_2)^* H_X^e \in N^e(X'' \times X'')_{\mathbb{R}}, \quad (30)$$

where $C_6 = C_5((\phi_2 \times \phi_2)^* H_X^{2l+e} \cdot (\varphi_2 \times \varphi_2)^* H_{\mathbb{P}^l \times \mathbb{P}^e}^e) / ((\phi_2 \times \phi_2)^* H_X^{2l+2e}) > 0$ does not depend on $\pi : X' \rightarrow X$. Using (30) and (29), we obtain:

$$[\Delta_{X''}] \leq C_7 \times \psi_{X'' \times X''}((\pi'' \times \pi'')^* H_{\mathbb{P}^l}^l \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^* H_X^e) \in N_{l+e}(X'' \times X'')_{\mathbb{R}}, \quad (31)$$

where $C_7 = C_6 \times C_1$. Since the basepoint free cone is contained in the nef cone by Theorem 3.3.3.(5), we have thus:

$$(\gamma' \lrcorner [\Delta_{X''}]) \leq C_7 (\gamma' \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^* H_X^e \cdot (\pi'' \times \pi'')^* H_{\mathbb{P}^l}^l). \quad (32)$$

Let us denote by $X_1 = (Y \times \mathbb{P}^e)^2$, $X_2 = (\mathbb{P}^l \times \mathbb{P}^e)^2$, $Y_1 = Y \times Y$, $Y_2 = \mathbb{P}^l \times \mathbb{P}^l$ and let $f := (\varphi_2 \circ \phi_2^{-1} \times \varphi_2 \circ \phi_2^{-1}) : X_1 \dashrightarrow X_2$ and $g := (\varphi_1 \circ \phi_1^{-1} \times \varphi_1 \circ \phi_1^{-1}) : Y_1 \dashrightarrow Y_2$ be the corresponding dominant rational maps. Let us apply Lemma 7.1.3 to the class $(\pi' \times \pi')^*(\varphi_2 \times \varphi_2)^* H_{\mathbb{P}^l}^l$ and to the class $(\pi' \times \pi')^*(\phi_2 \times \phi_2)^* H_Y^l$, there exists a constant $C_8 > 0$ which is independent of the morphism $\pi' \times \pi' : X'' \times X'' \rightarrow (Y' \times \mathbb{P}^e)^2$ such that for any class $\beta \in \text{BPF}^{2e+l}(X'' \times X'')$:

$$(\beta \cdot (\pi' \times \pi')^*(\varphi_2 \times \varphi_2)^* H_{\mathbb{P}^l}^l) \leq C_8 \frac{\deg_l(g)}{(H_Y^{2l})} (\beta \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^* H_Y^l). \quad (33)$$

Using (32) and (33) to the class $\beta = \gamma' \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^* H_X^e \in \text{BPF}^{l+2e}(X'' \times X'')$, we obtain:

$$(\gamma' \lrcorner [\Delta_{X''}]) \leq C_4 (\gamma' \cdot (\phi_3 \times \phi_3)^*(\pi \times \pi)^*(H_X^e \cdot H_Y^l)),$$

where $C_4 = C_4 \times C_8(\deg_l(g))/(H_Y^{2l}) > 0$ does not depend on π . The conclusion of the theorem follows from the projection formula and from the fact that $\phi_3 \times \phi_3$ is a birational map. Indeed, we apply the previous inequality to $\gamma' = (\phi_3 \times \phi_3)^* \gamma$ where $\gamma \in \text{BPF}^{l+e}(X' \times X')$, we obtain

$$(\gamma \lrcorner [\Delta_{X'}]) = ((\phi_3 \times \phi_3)^* \gamma \lrcorner [\Delta_{X''}]) \leq C_4 (\gamma \cdot (\pi \times \pi)^*(H_X^e \cdot H_Y^l))$$

as required.

Step 3: We prove the theorem in the general case. Suppose $q : X \rightarrow Y$ is a surjective morphism of relative dimension e and fix a class $\beta \in \text{BPF}^{l+e}(X' \times X')$. Since X is projective over Y , there exists a closed immersion $i : X \rightarrow Y \times \mathbb{P}^N$ such that $q = p'_Y \circ i$ where p'_Y is the projection of $Y \times \mathbb{P}^N$

onto Y . Let us choose a projection $Y \times \mathbb{P}^N \dashrightarrow Y \times \mathbb{P}^e$ so that the composition with i gives a dominant rational map $f : X \dashrightarrow Y \times \mathbb{P}^e$. Let us denote by Γ_f the normalization of the graph of f in $X \times Y \times \mathbb{P}^e$ and π_1, π_2 the projections of Γ_f onto the first and the second factor respectively. We set X'' the fibred product of X' with Γ_f and we denote by π' and ϕ the projection of X'' to Γ_f and X' respectively. We get the following commutative diagram:

$$\begin{array}{ccc} X' & \longleftarrow & X'' \\ \downarrow \pi & & \downarrow \pi' \\ X & \longleftarrow \pi_1 & \Gamma_f \\ \downarrow q & \searrow f & \downarrow \pi_2 \\ Y & \longleftarrow p_Y & Y \times \mathbb{P}^e \end{array}$$

where p_Y is the projection of $Y \times \mathbb{P}^e$ onto Y . We apply the result of Step 2 to the class $(\phi \times \phi)^* \beta \in \text{BPF}^{l+e}(X'' \times X'')$ and to the diagonal of X'' . There exists a constant $C_4 > 0$ which does not depend on π such that:

$$((\phi \times \phi)^* \beta \llcorner [\Delta_{X''}]) \leq C_4 ((\phi \times \phi)^* (\beta \cdot (\pi \times \pi)^* H_Y^l) \cdot ((\pi_2 \circ \pi') \times (\pi_2 \circ \pi'))^* H_{Y \times \mathbb{P}^e}^e). \quad (34)$$

Let us apply Theorem 3.4.4 to the class $((\pi_2 \circ \pi') \times (\pi_2 \circ \pi'))^* H_{Y \times \mathbb{P}^e}^e$ and to the class $(\phi \times \phi)^* (\pi \times \pi) H_X$. There exists a constant $C_9 > 0$ such that:

$$((\pi_2 \circ \pi') \times (\pi_2 \circ \pi'))^* H_{Y \times \mathbb{P}^e}^e \leq C_9 \frac{((\pi' \times \pi')^* ((\pi_2 \times \pi_2)^* H_{Y \times \mathbb{P}^e}^e \cdot (\pi_1 \times \pi_1)^* H_X^{2l+2e}))}{((\pi' \times \pi')^* (\pi_1 \times \pi_1)^* H_X^{2l+2e})} (\phi \times \phi)^* (\pi \times \pi) H_X^e \in \mathbb{N}^e(X'' \times X'')_{\mathbb{R}}.$$

Since $((\pi' \times \pi')^* ((\pi_2 \times \pi_2)^* H_{Y \times \mathbb{P}^e}^e \cdot (\pi_1 \times \pi_1)^* H_X^{2l+2e})) / ((\pi' \times \pi')^* (\pi_1 \times \pi_1)^* H_X^{2l+2e}) = \deg_e(f \times f) / (H_X^{2l+2e})$ and using (34), we obtain:

$$((\phi \times \phi)^* \beta \llcorner [\Delta_{X''}]) \leq C ((\phi \times \phi)^* (\beta \cdot (\pi \times \pi)^* (H_X^e \cdot H_Y^l))),$$

where $C = C_4 C_9 \deg_e(f \times f) / (H_X^{2l+2e})$. Since the morphism $\pi_1 : \Gamma_f \rightarrow X$ is birational, the map $\phi : X'' \rightarrow X'$ is also birational and we conclude using the projection formula and since $(\phi \times \phi)_* [\Delta_{X''}] = [\Delta_{X'}]$:

$$(\beta \llcorner [\Delta_{X'}]) \leq C (\beta \cdot (\pi \times \pi)^* (H_X^e \cdot H_Y^l)).$$

□

Corollary 7.1.5. *Let $q : X \rightarrow Y$ be a surjective morphism of relative dimension e where $\dim Y = l$. Then there exists a constant $C > 0$ such that for any surjective generically finite morphism $\pi : X' \rightarrow X$ such that for any class $\alpha \in \text{BPF}^k(X')$ and any class $\beta \in \text{BPF}^{l+e-k}(X')$, one has:*

$$(\beta \cdot \alpha) \leq C \sum_{\max(0, k-l) \leq j \leq \min(k, e)} U_j(\pi_* \psi_{X'}(\alpha)) \times (\beta \cdot \pi^*(q^* H_Y^{k-j} \cdot H_X^j)), \quad (35)$$

where $U_j(\pi_* \psi_{X'}(\alpha)) = (H_X^{e-j} \cdot q^* H_Y^{l-k+j} \llcorner \pi_* \psi_{X'}(\alpha))$.

Remark 7.1.6. Note that when $k \leq e$, then the inequality is already a consequence of Siu's inequality (Theorem 3.4.4). Indeed, the term on the right hand side of (35) with $j = k$ corresponds exactly to the term $C(\pi^* H_X^{n-k} \cdot \alpha) \times \pi^* H_X^k$.

Remark 7.1.7. Equation (35) proves that the class

$$-\psi_{X'}(\alpha) + C \sum_{\max(0, k-l) \leq j \leq \min(k, e)} U_j(\pi_* \psi_{X'}(\alpha)) \times \psi_{X'}(\pi^*(q^* H_Y^{k-j} \cdot H_X^j)) \in \mathbb{N}_{n-k}(X')_{\mathbb{R}}$$

is in the dual of the basepoint free cone $\text{BPF}^{n-k}(X')$. Moreover, if (28) is satisfied, then this class is pseudo-effective.

Proof. We apply Theorem 7.1.1 to the class $\gamma = p_1^*\beta \cdot p_2^*\alpha \in \text{BPF}^n(X' \times X')$. There exists a constant $C_1 > 0$ such that for any surjective generically finite morphism $\pi : X' \rightarrow X$ and any class $\gamma \in \text{BPF}^n(X' \times X')$, one has:

$$(\gamma \lrcorner [\Delta_{X'}]) \leq C_1(\gamma \cdot (\pi \times \pi)^*(H_X^e \cdot H_Y^l)).$$

We denote by p_1 and p_2 the projections of $X' \times X'$ onto the first and the second factors respectively. Fix $\alpha \in \text{BPF}^k(X')$ and $\beta \in \text{BPF}^{n-k}(X')$. Let us apply the previous inequality to $\gamma = p_1^*\beta \cdot p_2^*\alpha \in \text{BPF}^n(X' \times X')$. We obtain:

$$(\beta \cdot \alpha) = (p_1^*\beta \cdot p_2^*\alpha \lrcorner [\Delta_{X'}]) \leq C_1(p_1^*\beta \cdot p_2^*\alpha \cdot (\pi \times \pi)^*(H_X^e \cdot H_Y^l)).$$

Since $(p_1^*\pi^*(H_X^i \cdot q^*H_Y^j) \cdot p_2^*(\pi^*(q^*H_Y^{l-i} \cdot H_X^{e-j}) \cdot \gamma)) = 0$ when $i + j \neq k$, we obtain :

$$(\beta \cdot \alpha) \leq C \sum_{\max(0, k-l) \leq j \leq \min(e, k)} (\pi^*(q^*H_Y^{l-k+j} \cdot H_X^{e-j}) \cdot \alpha)(\pi^*(q^*H_Y^{k-j} \cdot H_X^j) \cdot \beta).$$

where $C = C_1 \left(1 + \max \left(\binom{e}{j} \binom{l}{k-j} \right) \right)$. Hence by the projection formula, we have proved the required inequality:

$$(\beta \cdot \alpha) \leq C \sum_{\max(0, k-l) \leq j \leq \min(e, k)} U_j(\pi_*\psi_{X'}(\alpha)) \times (\beta \cdot \pi^*(q^*H_Y^{k-j} \cdot H_X^j)).$$

□

Proof of Lemma 7.1.4: (see [Tru16, Lemma 4.4]) Since X is homogeneous, it is smooth. Let G be the automorphism group of $X \times X$, we denote by \cdot the (transitive) action of G on $X \times X$. By generic flatness (see [FGI⁺05, Theorem 5.12]), there exists a non empty open subset $V \subset X \times X$ such that the restriction of $\pi \times \pi$ to $U := (\pi \times \pi)^{-1}(V)$ is flat over V . Recall that two subvarieties $V \subset X \times X$ and $W \subset X \times X$ intersect properly in $X \times X$ if $\dim(V \cap W) = \dim V + \dim W - 2n$. Since G acts transitively on $X \times X$, there exists by [Ful98, Lemma B.9.2] a Zariski dense open subset $O \subset G$ such that for any point $g \in O$, the cycle $g \cdot [\Delta_X]$ intersects properly every component of $X \times X \setminus V$. In particular, there exists a one parameter subgroup $\tau : \mathbb{G}_m \rightarrow G$ such that $\tau(1) = \text{Id} \in G$ and such that τ maps the generic point of \mathbb{G}_m to a point in O . Let S be the closure in $X' \times X' \times \mathbb{P}^1$ of the set $\{(x', t) \in U \times \mathbb{G}_m \mid (\pi \times \pi)(x') \in \tau(t) \cdot \Delta_X\}$. Let $p : X' \times X' \times \mathbb{P}^1 \rightarrow X' \times X'$ be the projection onto $X' \times X'$ and let $f : S \rightarrow \mathbb{P}^1$ be the morphism induced by the projection of $X' \times X' \times \mathbb{P}^1$ onto \mathbb{P}^1 . As in [Ful98, Section 1.6], we denote by $S_t := p_*[f^{-1}(t)] \in Z_n(X' \times X')$ for any $t \in \mathbb{G}_m$. By construction the cycle $S_1 \in Z_n(X' \times X')$ is effective and its support contains the diagonal $\Delta_{X'}$ in $X' \times X'$, hence:

$$[\Delta_{X'}] \leq S_1 \in N_n(X' \times X')_{\mathbb{R}}.$$

Let $t \in \mathbb{G}_m$ such that $\tau(t) \in O$. Since $S_1 = S_t \in A_n(X' \times X')$ for any $t \in \mathbb{P}^1$, we have thus:

$$[\Delta_{X'}] \leq S_t \in N_n(X' \times X')_{\mathbb{R}}.$$

Since the cycle $\tau(t) \cdot [\Delta_X]$ intersects properly every component of $X \times X \setminus V$, we apply the result in [Ful98, Example 11.4.8.(b)] to the morphism $\pi \times \pi : X' \times X' \rightarrow X \times X$ so that

$$S_t = \overline{[(\pi \times \pi)^{-1}(\tau(t) \cdot \Delta_X)]} = (\pi \times \pi)^*[\Delta_X] \in A_n(X' \times X').$$

We have thus proved:

$$[\Delta_{X'}] \leq (\pi \times \pi)^*[\Delta_X] \in N_n(X' \times X')_{\mathbb{R}}.$$

□

Proof of Lemma 7.1.3. Observe that one has the following commutative diagram:

$$\begin{array}{ccccc}
 & & X' & & \\
 & & \downarrow \pi & & \\
 & & \Gamma_f & & \\
 & \swarrow \pi_1 & & \searrow \pi_2 & \\
 X_1 & \overset{f}{\dashrightarrow} & & X_2 & \\
 \downarrow q_1 & & & & \downarrow q_2 \\
 Y_1 & \overset{g}{\dashrightarrow} & & Y_2 & \\
 \swarrow \pi'_1 & & \Gamma_g & & \searrow \pi'_2
 \end{array}$$

Fix a class $\beta \in \text{BPF}^{e+l-j}(X')$. By linearity and by Proposition 2.1.8, we can suppose that the class β is induced by a product of nef divisors $D_1 \cdots D_{e_1+e+l-j}$ where D_i are nef divisors on X'_1 where $p : X'_1 \rightarrow X'$ is a flat morphism of relative dimension e_1 . The intersection $(\beta \cdot \pi^* \pi_2^* q_2^* H_{Y_2}^j)$ is thus given by the formula:

$$(\beta \cdot \pi^* \pi_2^* q_2^* H_{Y_2}^j) = (D_1 \cdots D_{e_1+e+l-j} \cdot p^* \pi^* \pi_2^* q_2^* H_{Y_2}^j).$$

Take A an ample Cartier divisor on X'_1 and set $\alpha_\epsilon = (D_1 + \epsilon A) \cdots (D_{e_1+e} + \epsilon A) \in \mathbb{N}^{e_1+e}(X'_1)_{\mathbb{R}}$ for any $\epsilon > 0$. Since the class α_ϵ is a complete intersection, there exists a cycle $V_\epsilon \in Z_l(X'_1)_{\mathbb{R}}$ such that $\psi_{X'_1}(\alpha_\epsilon) = \{V_\epsilon\} \in N_l(X'_1)_{\mathbb{R}}$ and such that the restrictions of the morphisms $\pi_1 \circ \pi \circ p$ and $\pi_2 \circ \pi \circ p$ to the support of V_ϵ are surjective and generically finite onto Y_1 and Y_2 respectively. We apply Theorem 3.4.4 to the class $(p^* \pi^* \pi_2^* q_2^* H_{Y_2}^j)|_{V_\epsilon}$ and to $(p^* \pi^* \pi_1^* q_1^* H_{Y_1}^j)|_{V_\epsilon}$, we get:

$$p^* \pi^* \pi_2^* q_2^* H_{Y_2}^j \cdot \alpha_\epsilon \leq C \frac{(p^* \pi^* (\pi_2^* q_2^* H_{Y_2}^j \cdot \pi_1^* H_{Y_1}^{l-j}) \llcorner \{V_\epsilon\})}{(p^* \pi^* \pi_1^* q_1^* H_{Y_1}^l \llcorner \{V_\epsilon\})} \times p^* \pi^* \pi_1^* q_1^* H_{Y_1}^j \cdot \alpha_\epsilon \in \mathbb{N}^{j+e_1+e}(X'_1)_{\mathbb{R}}.$$

By the projection formula applied to the morphism $\pi \circ p$, we have that $(p^* \pi^* (\pi_2^* q_2^* H_{Y_2}^j \cdot \pi_1^* H_{Y_1}^{l-j}) \llcorner \{V_\epsilon\}) / (p^* \pi^* \pi_1^* q_1^* H_{Y_1}^l \llcorner \{V_\epsilon\}) = \text{deg}_j(g) / (H_{Y_1}^l)$, hence:

$$p^* \pi^* \pi_2^* q_2^* H_{Y_2}^j \cdot \alpha_\epsilon \leq C \frac{\text{deg}_j(g)}{(H_{Y_1}^l)} p^* \pi^* \pi_1^* q_1^* H_{Y_1}^j \cdot \alpha_\epsilon \in \mathbb{N}^{j+e_1+e}(X'_1)_{\mathbb{R}}.$$

We intersect with the class $(D_{e_1+e+1} \cdots D_{e_1+e+l-j}) \in \mathbb{N}^{l-j}(X'_1)_{\mathbb{R}}$ and take the limit as ϵ tends to zero. We obtain:

$$(\beta \cdot \pi^* \pi_2^* q_2^* H_{Y_2}^j) = (D_1 \cdots D_{e_1+e+l-j} \cdot p^* \pi^* \pi_2^* q_2^* H_{Y_2}^j) \leq C \frac{\text{deg}_j(g)}{(H_{Y_1}^l)} (\beta \cdot \pi^* \pi_1^* q_1^* H_{Y_1}^j),$$

as required. \square

7.2. Submultiplicativity of mixed degrees.

Definition 7.2.1. Let $X_1/q_1 Y_1 \xrightarrow{f} X_2/q_2 Y_2$ be rational maps where $e = \dim X_i - \dim Y_i$ and $l = \dim Y_i$ for $i = 1, 2$. We fix some ample divisors H_{X_i} and H_{Y_i} on each variety respectively. We define for any integer $0 \leq k \leq n$:

$$a_{k,j}(f) := \begin{cases} ((H_{Y_1}^{l-j} \cdot H_{X_1}^{e+j-k}) \llcorner f^{\bullet k}(H_{X_2}^k)) & \text{if } \max(0, k-e) \leq j \leq l, \\ 0 & \text{otherwise.} \end{cases}$$

Remark 7.2.2. For $j = 0$, it is the k -th relative degree $a_{k,0}(f) = \text{reldeg}_k(f)$ and when $j = l$, it corresponds to the k -th degree of f , $a_{k,l}(f) = \text{deg}_k(f)$.

Theorem 7.2.3. *Let $q_1 : X_1 \rightarrow Y_1$, $q_2 : X_2 \rightarrow Y_2$, $q_3 : X_3 \rightarrow Y_3$ be three surjective morphisms such that $\dim X_i = e + l$ and $\dim Y_i = l$ for all $i \in \{1, 2, 3\}$. Then there exists a constant $C > 0$ such that for any rational maps $X_1/q_1 Y_1 \xrightarrow{f_1/g_1} X_2/q_2 Y_2$, $X_2/q_2 Y_2 \xrightarrow{f_2/g_2} X_3/q_3 Y_3$ and for all integers $0 \leq j_0 \leq l$:*

$$a_{k,j_0}(f_2 \circ f_1) \leq C \sum_{\max(0,k-l) \leq j \leq \min(e,k)} \text{deg}_{k-j}(g_1) a_{k,k-j}(f_2) a_{j,j_0-k}(f_1).$$

Proof. Since we are in the same situation as Theorem 5.2.1, we can consider the diagram (19) and we keep the same notations. We denote by $n = e + l$ the dimension of X_i .

Let us denote by d the topological degree of the map f_2 . We apply Corollary 7.1.5 to the pliant class $\alpha := (1/d)v^*\pi_4^*H_{X_3}^k \in \text{BPF}^k(\Gamma)$, to the class $\beta := u^*\pi_1^*(H_{X_1}^{e-k+j_0} \cdot q_1^*H_{Y_1}^{l-j_0}) \in \text{BPF}^{n-k}(\Gamma)$ and to the morphism $\pi = \varphi \circ \pi_3 \circ v$. There exists a constant $C_1 > 0$ which depends only on the choice of divisors $H_{Y_2 \times \mathbb{P}^e}$ and H_{Y_2} such that:

$$a_{k,j_0}(f_2 \circ f_1) \leq C_1 \sum_{\max(0,k-l) \leq j \leq \min(e,k)} U_j(\pi_*\psi_\Gamma(\alpha))(\beta \cdot \pi^*(H_{X_2}^j \cdot q_2^*H_{Y_2}^{k-j})),$$

where $U_j(\gamma) = (H_{X_2}^{e-j} \cdot q_2^*H_{Y_2}^{l-k+j} \lrcorner \gamma)$ for any class $\gamma \in \mathbb{N}_{n-k}(X_2)_{\mathbb{R}}$. We observe that $U_j(\pi_*\psi_\Gamma(\alpha)) = a_{k,k-j}(f_2)$. We have thus:

$$a_{k,j_0}(f_2 \circ f_1) \leq C_1 \sum_{\max(0,k-l) \leq j \leq \min(e,k)} a_{k,k-j}(f_2) (u^*(\pi_1^*(H_{X_1}^{e-k+j_0} \cdot q_1^*H_{Y_1}^{l-j_0}) \cdot \pi_2^*(H_{X_2}^j \cdot q_2^*H_{Y_2}^{k-j}))). \quad (36)$$

Applying Lemma 7.1.3 to the class $u^*\pi_2^*q_2^*H_{Y_2}^{k-j} \in \text{BPF}^{k-j}(\Gamma)$ and to $\beta' = \beta \cdot u^*\pi_2^*H_{X_2}^j \in \text{BPF}^{n-k+j}(\Gamma)$, there exists a constant $C_2 > 0$ such that :

$$(\beta' \cdot u^*\pi_2^*q_2^*H_{Y_2}^{k-j}) \leq C_2 \text{deg}_{k-j}(g_1) (u^*(\pi_1^*(H_{X_1}^{e-k+j_0} \cdot q_1^*H_{Y_1}^{l-j_0+k-j}) \cdot \pi_2^*H_{X_2}^j)).$$

Since the map $u : \Gamma \rightarrow \Gamma_{f_1}$ is birational, we have that:

$$(u^*(\pi_1^*(H_{X_1}^{e-k+j_0} \cdot q_1^*H_{Y_1}^{l-j_0}) \cdot \pi_2^*(H_{X_2}^j \cdot q_2^*H_{Y_2}^{k-j}))) \leq C_2 \text{deg}_{k-j}(g_1) a_{j,j_0+j-k}(f_1). \quad (37)$$

Finally, (36) and (37) imply:

$$a_{k,j_0}(f_2 \circ f_1) \leq C \sum_{\max(0,k-l) \leq j \leq \min(e,k)} a_{k,k-j}(f_2) a_{j,j_0+j-k}(f_1) \text{deg}_{k-j}(g_1),$$

where $C = C_2 C_1 > 0$ is a constant which is independent of f_1 and f_2 as required. \square

7.3. Proof of Theorem 4. By definition of the relative degrees, we are reduced to prove the theorem when $q : X \rightarrow Y$ is a proper surjective morphism. Recall that $\dim X = n$ and $\dim Y = l$ such that $q : X \rightarrow Y$ has relative dimension $e = n - l$. Let us consider the following commutative

diagram:

$$\begin{array}{ccc}
 & \Gamma_f & \\
 \pi_1 \swarrow & & \searrow \pi_2 \\
 X & \overset{f}{\dashrightarrow} & X \\
 q \downarrow & & \downarrow q \\
 Y & \overset{g}{\dashrightarrow} & Y \\
 \pi'_1 \swarrow & & \searrow \pi'_2 \\
 & \Gamma_g &
 \end{array} \tag{38}$$

where $f : X \dashrightarrow X$, $g : Y \dashrightarrow Y$ are dominant rational maps, Γ_f, Γ_g are the normalization of the graph of f and g respectively, $\pi_1, \pi_2, \pi'_1, \pi'_2$ are the projections from Γ_f and Γ_g onto the first and second factor respectively and $\varpi : \Gamma_f \rightarrow \Gamma_g$ is the restriction of $q \times q$ to Γ_f . The following lemma proves that $\max_{j \leq k} (\lambda_j(f, X/Y) \lambda_{k-j}(g)) \leq \lambda_k(f)$.

Lemma 7.3.1. *For any integer $\max(0, k-l) \leq j \leq \min(k, e)$, we have that $\lambda_j(f, X/Y) \lambda_{k-j}(g) \leq \lambda_k(f)$.*

Proof. We are reduced to show that for any $\max(0, k-l) \leq j \leq \min(k, e)$, there exists a constant $C > 0$ such that:

$$\deg_{k-j}(g) \operatorname{reldeg}_j(f) \leq C \deg_k(f).$$

For this, it suffices to consider the product $(\pi_1^*(H_X^{e-j} \cdot q^* H_Y^{l-k+j}) \cdot \pi_2^*(H_X^j \cdot q^* H_Y^{k-j}))$. Since $\pi_i \circ q = \varpi \circ \pi'_i$ for $i \in \{1, 2\}$, we obtain:

$$(\pi_1^*(H_X^{e-j} \cdot q^* H_Y^{l-k+j}) \cdot \pi_2^*(H_X^j \cdot q^* H_Y^{k-j})) = (\varpi^*(\pi_1'^* H_Y^{l-k+j} \cdot \pi_2'^* H_Y^{k-j}) \cdot \pi_1^* H_X^{e-j} \cdot \pi_2^* H_X^j).$$

Moreover, one has that $\pi_1'^* H_Y^{l-k+j} \cdot \pi_2'^* H_Y^{k-j} = (\pi_1'^* H_Y^{l-k+j} \cdot \pi_2'^* H_Y^{k-j}) [p_0] = \deg_{k-j}(g) [p_0]$ where p_0 is a general point in Γ_g . We can hence apply Proposition 4.3.1 to the morphism $\varpi : \Gamma_f \rightarrow \Gamma_g$ and obtain:

$$(\pi_1^*(H_X^{e-j} \cdot q^* H_Y^{l-k+j}) \cdot \pi_2^*(H_X^j \cdot q^* H_Y^{k-j})) = \deg_{k-j}(g) (\pi_1^* H_X^{e-j} \cdot \pi_2^* H_X^j \lrcorner [\Gamma_{f p_0}]).$$

Since π_1' is a birational morphism, a general fiber of ϖ is equal to a general fiber of $\pi_1' \circ \varpi$. In other words, we have that $\operatorname{Res}_{\Gamma_f/\Gamma_g} = \operatorname{Res}_{\Gamma_f/Y}$ and since $\pi_1^* H_X^{e-j} \cdot \pi_2^* H_X^j \lrcorner [\Gamma_{f p_0}] = \operatorname{Res}_{\Gamma_f/\Gamma_g} (\pi_1^* H_X^{e-j} \cdot \pi_2^* H_X^j)$, we obtain:

$$(\pi_1^*(H_X^{e-j} \cdot q^* H_Y^{l-k+j}) \cdot \pi_2^*(H_X^j \cdot q^* H_Y^{k-j})) = \deg_{k-j}(g) \times \operatorname{reldeg}_j(f).$$

As H_X is ample, we apply Theorem 3.3.3 to the classes $\pi_2^* q^* H_Y$ and $\pi_2^* H_X$:

$$\pi_2^* q^* H_Y^{k-j} \leq (n-k+j+1)^{k-j} \frac{(\pi_2^* q^* H_Y^{k-j} \cdot \pi_2^* H_X^{n-k+j})}{(\pi_2^* H_X^n)} \pi_2^* H_X^{k-j} = C_1 \pi_2^* H_X^{k-j} \in \mathbb{N}^{k-j}(X)_{\mathbb{R}},$$

where $C_1 = (n-k+j+1)^{k-j} (q^* H_Y^{k-j} \cdot H_X^{n-k+j}) / (H_X^n)$ depends only on n, k and the choice of big nef Cartier divisors. Intersecting with $\pi_1^* H_X^{n-k} \cdot \pi_2^* H_X^j$, one obtains:

$$\deg_{k-j}(g) \cdot \operatorname{reldeg}_j(f) \leq C_1 (\pi_2^* H_X^k \cdot \pi_1^*(H_X^{e-j} \cdot q^* H_Y^{l-k+j})).$$

By the same argument, there exists a constant $C_2 > 0$ which depends only on H_Y, H_X and k such that:

$$\pi_1^* q^* H_Y^{l-k+j} \leq C_2 \pi_1^* H_X^{l-k+j}.$$

Hence, we obtain:

$$\deg_{k-j}(g) \operatorname{reldeg}_j(f) \leq C \deg_k(f),$$

where $C = C_1 C_2$. □

Let us prove the converse inequality. We fix an integer $0 \leq k \leq n$. Let us apply Theorem 7.2.3 to $f_1 = f$, $f_2 = f^p$, $g_1 = g$ and $g_2 = g^p$, we can rewrite the inequality as:

$$a_{k,j_0}(f^{p+1}) \leq C \sum_{\max(0,k-e) \leq j \leq \min(k,l)} \deg_j(g) a_{k-j,j_0-j}(f) a_{k,j}(f^p). \quad (39)$$

Let us denote by $U_k(f)$ the column vector given by:

$$U_k(f) = (a_{k,i}(f))_{0 \leq i \leq l} = \begin{pmatrix} a_{k,0}(f) \\ \dots \\ a_{k,l}(f) \end{pmatrix}.$$

Let us also denote by $M_k(f)$ the $(l+1) \times (l+1)$ lower-triangular matrix given by:

$$M_k(f) := (\deg_j(g) a_{k-j,i-j}(f) \times \mathbb{1}_{[k-e, \min(k,l)]}(j))_{0 \leq i \leq l, 0 \leq j \leq l},$$

where $\mathbb{1}_A$ denotes the characteristic function of the set A . Therefore, (39) can be rewritten as:

$$U_k(f^{p+1}) \leq C M_k(f) \cdot U_k(f^p),$$

where \cdot denotes the linear action on \mathbb{Z}^{l+1} . A simple induction proves:

$$U_k(f^p) \leq C^p (M_k(f))^{p-1} \cdot U_k(f)$$

Since the $(l+1)$ -th entry of the vector $U_k(f^p)$ corresponds to $\deg_k(f^p)$, we deduce that:

$$\deg_k(f^p)^{1/p} \leq C \langle e_l, (M_k(f))^p \cdot U_k(f) \rangle^{1/p}, \quad (40)$$

where (e_0, \dots, e_l) denotes the canonical basis of \mathbb{Z}^{l+1} . In particular, $\deg_k(f^p)^{1/p}$ is controlled up to a constant by the eigenvalues of the matrix $M_k(f)$ which are $\deg_j(g) \operatorname{reldeg}_{k-j}(f)$ for $\max(0, k-e) \leq j \leq \min(k, l)$ since $M_k(f)$ is lower-triangular. Applying (40) to f^r , we get:

$$\deg_k(f^{pr})^{1/(pr)} \leq C^{1/r} \|U_k(f^r)\|^{1/pr} \max_{\max(0,k-e) \leq j \leq \min(k,l)} (\deg_j(g^r) \operatorname{reldeg}_{k-j}(f^r))^{1/r}.$$

We conclude by taking the limsup as $r \rightarrow +\infty, p \rightarrow +\infty$:

$$\lambda_k(f) \leq \max_{\max(0,k-l) \leq j \leq \min(k,e)} \lambda_{k-j}(g) \lambda_j(f, X/Y).$$

Remark 7.3.2. Note that the previous theorem gives information only on the dynamical degrees of f . Lemma 7.3.1 provides a lower bound on the degree of f^p . However, one cannot find an upper bound for $\deg_k(f^p)$ which would only depend on the relative degrees and the degree on the base. If $X = E \times E$ is a product of two elliptic curves and if $f : (z, w) \in E \times E \rightarrow (z, z + w)$ is an automorphism of X , then the degree growth of f^p is equivalent to p^2 whereas the degree on the base and on any fiber are trivial.

8. KÄHLER CASE

We prove the submultiplicativity of the k -th degrees in the case where (X, ω) is a complex compact Kähler manifold. For any closed smooth (p, q) -form α on X , we denote by $\{\alpha\}$ its class in the Dolbeault cohomology $H^{p,q}(X)_{\mathbb{R}}$.

Definition 8.0.3. *Let (X, ω) be a compact Kähler manifold. A class $\alpha \in H^{1,1}(X)_{\mathbb{R}}$ is nef if for any $\epsilon > 0$, the class $\alpha + \epsilon\{\omega\}$ is represented by a Kähler metric.*

Definition 8.0.4. *Let (X, ω) be a compact Kähler manifold of dimension n . A class α of degree (k, k) is pseudo-effective if it can be represented by a closed positive current T . Moreover, one says that α is big if there exists a constant $\delta > 0$ such that $T - \delta\omega$ is a closed positive current and we write $T \geq \delta\omega^k$.*

Theorem 8.0.5. (cf [Xia15, Remark 3.1]) *Let (X, ω) be a compact Kähler manifold of dimension n . Let k be an integer and α, β be two nef classes in $H^{1,1}(X)$ such that $\alpha^k \in H^{k,k}(X)$ is big and such that $\int_X \alpha^n - \binom{n}{k} \int_X \alpha^{n-k} \wedge \beta^k > 0$. Then the class $\alpha^k - \beta^k$ is big.*

Recall that the degree of a meromorphic selfmap $f : X \dashrightarrow X$ when (X, ω) is given by:

$$\deg_k(f) := \int_{\Gamma_f} \pi_1^* \omega^{n-k} \wedge \pi_2^* \omega^k,$$

where Γ_f is the desingularization of the graph of f and π_i are the projections from Γ_f onto the first and the second factor respectively.

Corollary 8.0.6. *Let (X_1, ω_{X_1}) , (X_2, ω_{X_2}) and (X_3, ω_{X_3}) be some compact Kähler manifolds of dimension n . Then there exists a constant $C > 0$ which depends only on the choice of the Kähler classes ω_{X_i} such that for any dominant meromorphic maps $f_1 : X_1 \dashrightarrow X_2$ and $f_2 : X_2 \dashrightarrow X_3$, one has:*

$$\deg_k(f_2 \circ f_1) \leq C \deg_k(f_1) \deg_k(f_2).$$

Moreover, the constant C may be chosen to be equal to $\binom{n}{k} / (\int_{X_2} \omega_{X_2}^n)$.

Proof. The previous theorem gives that for any big nef class $\beta^k \in H^{k,k}(X)$, for any nef class $\alpha \in H^{1,1}(X)$, one has:

$$\alpha^k \leq \binom{n}{k} \frac{\int_X \alpha^k \wedge \beta^{n-k}}{\int_X \beta^n} \times \beta^k. \quad (41)$$

Then, the proof is formally the same as Theorem 5.2.1. Indeed, one only needs to consider the diagram (19) where $Y_1 = Y_2 = Y_3$ are reduced to a point and where $\Gamma_{f_1}, \Gamma_{f_2}, \Gamma$ are the desingularizations of the graph of f_1, f_2 and $\pi_3^{-1} \circ f_1 \circ \pi_1$ respectively. We apply (41) to $\alpha = v^* \pi_4^* \omega_{X_3}$ and $\beta = v^* \pi_3^* \omega_{X_2}$ to obtain:

$$v^* \pi_4^* \omega_{X_3}^k \leq \binom{n}{k} \frac{\deg_k(f_2)}{\int_{X_2} \omega_{X_2}^n} \times v^* \pi_3^* \omega_{X_2}^k.$$

By intersecting the previous inequality with the class $u^* \pi_1^* \omega_{X_1}^{n-k}$, we finally get:

$$\deg_k(f_2 \circ f_1) \leq \binom{n}{k} \frac{\deg_k(f_2) \deg_k(f_1)}{\int_{X_2} \omega_{X_2}^n}.$$

□

APPENDIX A. COMPARISON WITH FULTON'S APPROACH

In [Ful98, Chapter 19], a cycle $z \in Z_k(X)$ on a variety X is defined to be numerically trivial if $(c_\perp z)$ for any product $c = c_{i_1}(E_1) \cdot \dots \cdot c_{i_p}(E_p) \in A^k(X)$ of Chern classes $c_{i_j}(E_j)$ where E_j is a vector bundle on X and $i_1 + \dots + i_p = k$. This appendix is devoted to the proof of the following result:

Theorem A.1. *Let X be a normal projective variety of dimension n . For any $z \in Z_k(X)$, the following conditions are equivalent:*

- (i) *For any product of Chern classes $c = c_{i_1}(E_1) \cdot \dots \cdot c_{i_p}(E_p) \in A^k(X)_\mathbb{R}$ where E_j are vector bundles on X and $i_1 + \dots + i_p = k$, we have $(c_\perp z) = 0$.*
- (ii) *For any integer e , any flat morphism $p_1 : X_1 \rightarrow X$ of relative dimension e where X_1 is a projective scheme and any Cartier divisors D_1, \dots, D_{e+k} on X_1 , we have $(D_1 \cdot \dots \cdot D_{e+k} \lrcorner p_1^* z) = 0$.*

- (iii) For any integer e , any flat morphism $p_1 : X_1 \rightarrow X$ of relative dimension e between normal projective varieties and any Cartier divisors D_1, \dots, D_{e+k} on X_1 , we have $(D_1 \cdots \cdots D_{e+k} \lrcorner p_1^* z) = 0$.

The implication (ii) \Rightarrow (i) follows immediately from the definition of Chern classes. The implication (ii) \Rightarrow (iii) is also straightforward. For the converse implications (i) \Rightarrow (ii) and (i) \Rightarrow (iii), we rely on the following proposition.

Proposition A.2. *Let $q : X \rightarrow Y$ be a flat morphism of relative dimension e where X is a projective scheme and Y is a normal projective variety. For any Cartier divisors D_1, \dots, D_{e+k} be some ample Cartier divisors on X , there exist vector bundles E_j , and a homogeneous polynomial $c = P(c_{i_1}(E_1), \dots, c_{i_p}(E_p))$ of degree k with respect to the weight (i_1, \dots, i_p) , with rational coefficients such that for any cycle $z \in Z_k(X)$, $(c \cdot z) = (D_1 \cdots \cdots D_{e+k} \cdot q^* z)$.*

Proof. We take some ample Cartier divisors D_1, \dots, D_{e+k} on X . We denote by \mathcal{L}_i the line bundle $\mathcal{O}_X(D_i)$. By Grauert's Theorem (cf [Har77, Corollary 12.9]), the sheaves $R^i q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})$ are locally free. By [Har77, Theorem 8.8], we have that $R^i q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}) = 0$ for $i > 0$ and m_i large enough since the line bundle \mathcal{L}_i are ample. So the sheaf $q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})$ is locally free and we have in $K_0(Y)$:

$$q_*[\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}] = \sum (-1)^i [R^i q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})] = [q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})]. \quad (42)$$

Lemma A.3. *For any $j \leq k$:*

- (1) *The function $(m_1, \dots, m_{e+k}) \rightarrow \text{ch}_j(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})) \in N^j(Y)_{\mathbb{R}}$ is a polynomial of degree $e + j$ with coefficients in $N^j(Y)$.*
- (2) *For any cycle $z \in Z_j(Y)$, the coefficient in $m_1 \cdots \cdots m_{e+k}$ in $(\text{ch}_j(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})) \lrcorner z)$ is $((D_1 \cdots \cdots D_{e+k}) \lrcorner q^* z)$.*

Proof. Let us set $\mathcal{F} = \mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}$. We prove the result by induction on $0 \leq j \leq k$.

For $j = 0$, choosing a point $y \in Y(\kappa)$, the number $\text{ch}_0(q_*(\mathcal{F}))$ is equal to $h^0(X_y, \mathcal{F}|_{X_y})$. By asymptotic Riemann-Roch, for m_1, \dots, m_{e+k} large enough, it is a polynomial of degree $\dim X_y = e$. Moreover, Snapper's theorem (see [Deb01, Definition 1.7]) states that the coefficient in $m_1 \cdots \cdots m_{e+k}$ is the number $(D_1 \cdots \cdots D_{e+k} \lrcorner [X_y])$.

We suppose by induction that $\text{ch}_i(q_*(\mathcal{F}))$ is a polynomial of degree $e + i$ for any $i \leq j$ where $j \leq k - 1$. For any subvariety V of dimension $j + 1$ in Y , we denote by W its scheme-theoretic preimage by q .

For any scheme V , let us denote by τ_V the morphisms:

$$\tau_V : K_0(V) \otimes \mathbb{Q} \rightarrow A_{\bullet}(V) \otimes \mathbb{Q}.$$

We refer to [Ful98, Theorem 18.3] for the construction of this morphism and its properties. We apply Grothendieck-Riemann-Roch's theorem for singular varieties (see [Ful98, Theorem 18.3.(1)]) and using (42), we get in $A_{\bullet}(Y)_{\mathbb{Q}}$:

$$\text{ch}(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})) \lrcorner \tau_V(\mathcal{O}_V) = q_*(\text{ch}(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}) \lrcorner \tau_W(\mathcal{O}_W)). \quad (43)$$

The term in $A_0(Y)_{\mathbb{Q}}$ in the left handside of the previous equation is equal to:

$$\text{ch}_{j+1}(q_*(\mathcal{F})) \lrcorner [V] + \sum_{i \leq j} \text{ch}_i(q_*(\mathcal{F})) \lrcorner \tau_{V,i}(\mathcal{O}_V),$$

where $\tau_{V,i}(\mathcal{O}_V)$ is the term in $A_i(Y)$ of $\tau_V(\mathcal{O}_V)$. By the induction hypothesis, every $\text{ch}_i(q_*(\mathcal{F}))$ is a polynomial of degree $e + i$, and the right hand side of equation (43) is a polynomial of degree $e + j + 1$, so $\text{ch}_{j+1}(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}))$ is also a polynomial of degree $e + j + 1$. Now we

identify the coefficients in $m_1 \cdot \dots \cdot m_{e+k}$ of the term in $N_0(Y)$ in equation (43). It follows from [Ful98, example 18.3.11] that $\tau_W(\mathcal{O}_W) = [W] + R_W$ where R_W is a linear combination of cycles of dimension $< e + k$. Therefore, the coefficient in $m_1 \cdot \dots \cdot m_{e+k}$ of the right hand side of equation (43) in $N_0(Y)$ is $((D_1 \cdot \dots \cdot D_{e+k}) \lrcorner [W])$ if $j + 1 = k$ or 0 otherwise.

We have proved that the coefficient of $\text{ch}_{j+1}(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}})) \lrcorner [V]$ is $((D_1 \cdot \dots \cdot D_{e+k}) \lrcorner [W])$ if $\dim V = k$ or 0 otherwise. Extending it by linearity, one gets the desired result. \square

We have that $\text{ch}_k(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}))$ is by definition a polynomial in Chern classes of vector bundles on Y . Using the previous lemma, the coefficient $U(D_1, \dots, D_{e+k})$ in $m_1 \cdot \dots \cdot m_{e+k}$ of $\text{ch}_k(q_*(\mathcal{L}_1^{m_1} \otimes \dots \otimes \mathcal{L}_{e+k}^{m_{e+k}}))$ is equal to $P(c_{i_1}(E_1), \dots, c_{i_p}(E_p))$ where P is a homogeneous polynomial with rational coefficients of degree k with respect to the weight (i_1, \dots, i_p) and E_i are vector bundles on Y . We have proven that for any cycle $z \in Z_k(Y)$:

$$(P(c_{i_1}(E_1), \dots, c_{i_p}(E_p)) \lrcorner z) = ((D_1 \cdot \dots \cdot D_{e+k}) \lrcorner q^* z).$$

As any Cartier divisor can be written as a difference of ample Cartier divisors. The proposition provides a proof for the implication (i) \Rightarrow (ii) of Theorem A.1. \square

Remark A.4. In codimension 1, the intersection product $(D_1 \cdot \dots \cdot D_{e+1} \lrcorner q^* z)$ is represented by Deligne's product $I_X(\mathcal{O}_X(D_1), \dots, \mathcal{O}_X(D_{e+1})) \in N^1(X)_{\mathbb{R}}$ (see [Gar00] for a reference). Indeed, one has by [Gar00, Section 6] that for any cycle $z \in N_1(X)$:

$$c_1(I_X(\mathcal{O}_X(D_1), \dots, \mathcal{O}_X(D_{e+1}))) \lrcorner z = D_1 \cdot \dots \cdot D_{e+1} \lrcorner q^* z.$$

This gives an answer to the question of numerical pullback formulated in [FL14b, section 1.2].

Corollary A.5. *Let $q : X \rightarrow Y$ be a flat morphism of relative dimension e between normal projective varieties. Then the morphism $q^* : A_{\bullet}(Y)_{\mathbb{Q}} \rightarrow A_{e+\bullet}(X)_{\mathbb{Q}}$ induces a morphism of abelian groups $q^* : N_{\bullet}(Y)_{\mathbb{Q}} \rightarrow N_{e+\bullet}(X)_{\mathbb{Q}}$. By duality, the morphism $q_* : A^{\bullet}(X)_{\mathbb{Q}} \rightarrow A^{\bullet-e}(Y)_{\mathbb{Q}}$ induces a morphism of abelian groups $q_* : N^{\bullet}(X)_{\mathbb{Q}} \rightarrow N^{\bullet-e}(Y)_{\mathbb{Q}}$.*

REFERENCES

- [BC16] Jérémy Blanc and Serge Cantat, *Dynamical degrees of birational transformations of projective surfaces*, J. Amer. Math. Soc. **29** (2016), no. 2, 415–471. MR 3454379
- [BD15] Jérémy Blanc and Julie Déserti, *Degree growth of birational maps of the plane*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **14** (2015), no. 2, 507–533. MR 3410471
- [BdFFU15] Sébastien Boucksom, Tomasso de Fernex, Charles Favre, and Stefano Urbinati, *Valuation spaces and multiplier ideals on singular varieties*, Recent advances in algebraic geometry, London Math. Soc. Lecture Note Ser., vol. 417, Cambridge Univ. Press, Cambridge, 2015, pp. 29–51. MR 3380442
- [BFJ08] Sébastien Boucksom, Charles Favre, and Mattias Jonsson, *Degree growth of meromorphic surface maps*, Duke Math. J. **141** (2008), no. 3, 519–538. MR 2387430
- [BS92] Eric Bedford and John Smillie, *Polynomial diffeomorphisms of \mathbb{C}^2 . III. Ergodicity, exponents and entropy of the equilibrium measure*, Math. Ann. **294** (1992), no. 3, 395–420. MR 1188127
- [Can11] Serge Cantat, *Sur les groupes de transformations birationnelles des surfaces*, Ann. of Math. (2) **174** (2011), no. 1, 299–340. MR 2811600
- [CC15] Dawei Chen and Izzet Coskun, *Extremal higher codimension cycles on moduli spaces of curves*, Proc. Lond. Math. Soc. (3) **111** (2015), no. 1, 181–204. MR 3404780
- [CLO16] Izzet Coskun, John Lesieutre, and John Christian Ottem, *Effective cones of cycles on blow-ups of projective space*, arXiv preprint 1603.04808 (2016).
- [Cut15] Steven Dale Cutkosky, *Teissier's problem on inequalities of nef divisors*, J. Algebra Appl. **14** (2015), no. 9, 1540002, 37. MR 3368254
- [CZ12] Serge Cantat and Abdelghani Zeghib, *Holomorphic actions, Kummer examples, and Zimmer program*, Ann. Sci. Éc. Norm. Supér. (4) **45** (2012), no. 3, 447–489. MR 3014483

- [Dan94] V. I. Danilov, *Algebraic varieties and schemes*, Algebraic geometry, I, Encyclopaedia Math. Sci., vol. 23, Springer, Berlin, 1994, pp. 167–297. MR 1287420
- [DDG11] Jeffrey Diller, Romain Dujardin, and Vincent Guedj, *Dynamics of meromorphic mappings with small topological degree II: Energy and invariant measure*, Comment. Math. Helv. **86** (2011), no. 2, 277–316. MR 2775130
- [Deb01] Olivier Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, New York, 2001. MR 1841091
- [DELV11] Olivier Debarre, Lawrence Ein, Robert Lazarsfeld, and Claire Voisin, *Pseudoeffective and nef classes on abelian varieties*, Compos. Math. **147** (2011), no. 6, 1793–1818. MR 2862063
- [DF01] Jeffrey Diller and Charles Favre, *Dynamics of bimeromorphic maps of surfaces*, Amer. J. Math. **123** (2001), no. 6, 1135–1169. MR 1867314
- [DN11] Tien-Cuong Dinh and Viêt-Anh Nguyễn, *Comparison of dynamical degrees for semi-conjugate meromorphic maps*, Comment. Math. Helv. **86** (2011), no. 4, 817–840. MR 2851870
- [DNT12] Tien-Cuong Dinh, Viêt-Anh Nguyễn, and Tuyen Trung Truong, *On the dynamical degrees of meromorphic maps preserving a fibration*, Communications in Contemporary Mathematics **14** (2012), no. 06, 1250042.
- [DNT16] Tien-Cuong Dinh, Viet-Anh Nguyen, and Tuyen Trung Truong, *Growth of the number of periodic points for meromorphic maps*, arXiv preprint 1601.03910 (2016).
- [DS04a] Tien-Cuong Dinh and Nessim Sibony, *Groupes commutatifs d'automorphismes d'une variété kählérienne compacte*, Duke Math. J. **123** (2004), no. 2, 311–328. MR 2066940
- [DS04b] ———, *Regularization of currents and entropy*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 6, 959–971. MR 2119243
- [DS05a] ———, *Dynamics of regular birational maps in \mathbb{P}^k* , J. Funct. Anal. **222** (2005), no. 1, 202–216. MR 2129771
- [DS05b] ———, *Une borne supérieure pour l'entropie topologique d'une application rationnelle*, Ann. of Math. (2) **161** (2005), no. 3, 1637–1644. MR 2180409
- [DS10] ———, *Dynamics in several complex variables: endomorphisms of projective spaces and polynomial-like mappings*, Holomorphic dynamical systems, Lecture Notes in Math., vol. 1998, Springer, Berlin, 2010, pp. 165–294. MR 2648690
- [DTV10] Henry De Thélin and Gabriel Vigny, *Entropy of meromorphic maps and dynamics of birational maps*, Mém. Soc. Math. Fr. (N.S.) (2010), no. 122, vi+98. MR 2752759
- [Eis95] David Eisenbud, *Commutative algebra*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry. MR 1322960
- [FGI⁺05] B. Fantechi, L. Göttsche, L. Illusie, S. Kleiman, N. Nitsure, and A. Vistoli, *Fundamental algebraic geometry: Grothendieck's fga explained*, Mathematical Surveys and Monographs, 2005.
- [FL14a] Mihai Fulger and Brian Lehmann, *Kernels of numerical pushforwards*, arXiv preprint 1407.6455 (2014).
- [FL14b] ———, *Positive cones of dual cycle classes*, arXiv preprint 1408.5154 (2014).
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry. MR 1234037
- [Ful98] ———, *Intersection theory*, second ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 2, Springer-Verlag, Berlin, 1998. MR 1644323
- [Gar00] Elvira Muñoz García, *Fibrés d'intersection*, Compositio Math. **124** (2000), no. 3, 219–252. MR 1809336
- [GGJ⁺16] José Ignacio Burgos Gil, Walter Gubler, Philipp Jell, Klaus Kuennemann, and Florent Martin, *Differentiability of non-archimedean volumes and non-archimedean monge-amp\ere equations (with an appendix by robert lazarsfeld)*, arXiv preprint 1608.01919 (2016).
- [Giz80] Marat Kharisovitch Gizatullin, *Rational G-surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), no. 1, 110–144, 239. MR 563788
- [Gro87] Misha Gromov, *Entropy, homology and semialgebraic geometry*, Astérisque (1987), no. 145-146, 5, 225–240, Séminaire Bourbaki, Vol. 1985/86. MR 880035
- [Gue05] Vincent Guedj, *Ergodic properties of rational mappings with large topological degree*, Ann. of Math. (2) **161** (2005), no. 3, 1589–1607. MR 2179389
- [Har77] Robin Hartshorne, *Algebraic geometry*, vol. 52, Springer Science & Business Media, 1977.
- [Jon96] Aise Johan De Jong, *Smoothness, semi-stability and alterations*, Publications Mathématiques de l'IHÉS (1996).

- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. MR 1658959
- [Laz04] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series. MR 2095471
- [LX15] Brian Lehmann and Jian Xiao, *Convexity and zariski decomposition structure*.
- [Mil13] James S. Milne, *Lectures on étale cohomology (v2.21)*, 2013, Available at www.jmilne.org/math/.
- [Pop16] Dan Popovici, *Sufficient bigness criterion for differences of two nef classes*, Math. Ann. **364** (2016), no. 1-2, 649–655. MR 3451400
- [RS97] Alexander Russakovskii and Bernard Shiffman, *Value distribution for sequences of rational mappings and complex dynamics*, Indiana Univ. Math. J. **46** (1997), no. 3, 897–932. MR 1488341
- [Sib99] Nessim Sibony, *Dynamique des applications rationnelles de \mathbf{P}^k* , Dynamique et géométrie complexes (Lyon, 1997), Panor. Synthèses, vol. 8, Soc. Math. France, Paris, 1999, pp. ix–x, xi–xii, 97–185. MR 1760844
- [Tru15] Tuyen Trung Truong, *(relative) dynamical degrees of rational maps over an algebraic closed field*, arXiv preprint 1501.01523 (2015).
- [Tru16] ———, *Relative dynamical degrees of correspondance over a field of arbitrary characteristic*, arXiv (2016).
- [Xia15] Jian Xiao, *Weak transcendental holomorphic Morse inequalities on compact Kähler manifolds*, Ann. Inst. Fourier (Grenoble) **65** (2015), no. 3, 1367–1379. MR 3449182
- [Yom87] Yosef Yomdin, *Volume growth and entropy*, Israel Journal of Mathematics **57** (1987), no. 3, 285–300.
- [Zha14] De-Qi Zhang, *Compact Kähler manifolds with automorphism groups of maximal rank*, Trans. Amer. Math. Soc. **366** (2014), no. 7, 3675–3692. MR 3192612

(Nguyen-Bac Dang) CMLS, ÉCOLE POLYTECHNIQUE, CNRS, UNIVERSITÉ PARIS-SACLAY, 91128 PALAISEAU CEDEX, FRANCE

E-mail address: nguyen-bac.dang@polytechnique.edu