

Addressing high variability in LCA with Global Sensitivity Assessment: from a single parameterized model to multi-parameterized clustered models

Paula Perez-Lopez, Alessio Mastrucci, Enrico Benetto, Isabelle Blanc

▶ To cite this version:

Paula Perez-Lopez, Alessio Mastrucci, Enrico Benetto, Isabelle Blanc. Addressing high variability in LCA with Global Sensitivity Assessment: from a single parameterized model to multi-parameterized clustered models. 22nd SETAC Europe LCA Case Study Symposium, SETAC, Sep 2016, Montpellier, France. hal-01447625

HAL Id: hal-01447625 https://hal.science/hal-01447625

Submitted on 27 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Addressing high variabilityin LCA with Global Sensitivity Assessment: from a single parameterized modelto multi parameterized clustered models.

Pérez-López P, Mastrucci A, Benetto E, Blanc I.

The application of Life Cycle Assessment (LCA) has been traditionally based on deterministic models that provide estimates of the environmental impacts of complex processes through sets of single-value parameters. Too often, this approach neglects to provide the potential distribution of the response of specific systems linked either orboth to uncertainty (due to lack of knowledge in parameters, models and practitioner's decisions) and to variability (caused by inherent differences between locations, technologies and temporal frames for example).

Global sensitivity analysis (GSA)is an efficient method to quantify the impact of both causes of LCA results variance from each input's variationandto support LCA model simplification. However, uncertainty and variability have different implicationswhen refining LCA parameterized models to enhance their quality, so the identification of the main cause of variationis an essential step.

If uncertainty is the major source of variation, the improvement of the LCA model may be obtained by collecting additional information of the system that allows:i) a deeper knowledge of the model parameters, ii) the development of a more detailed and relevant model or iii) a change in arbitrary decisions affecting LCA results.On the contrary, the predominance of variability reveals the existence of potential significantly different alternatives of the modeled process. Consequently better qualified estimates of the environmental impacts cannot be obtained by acquiring more precise input data.

In this work, we propose a method to improve the quality of LCA results linked to variabilitybased variation in complex models by

- 1) Identifying whether variability is actually the main cause of variation or not,
- 2) Grouping the different alternatives into a relevant set of system typologies.Such clustering step does reduce the range of LCA results for each typology,

3) For each cluster, setting a simplified parametrized model based on key parameters responsible for most of the cluster'svariability.

We apply this approach to the quantification of the greenhouse gas (GHG) emissions from the building sector by taking as an example the residential building stock of a representativetown in Luxembourg. The methodreduces the GHG estimates variabilityinduced by agenericsingle parameterized LCA model for all existing types of buildings by establishing multi parameterizedclustered modelsbased onbuilding type-specific key parameters.