



# Effects of pyrolytic and petrogenic polycyclic aromatic hydrocarbons on 1 swimming and metabolic performance of zebrafish contaminated by 2 ingestion 3 4

Julie Lucas, Isabelle Percelay, Thibault Larcher, Christel Lefrancois

## ► To cite this version:

Julie Lucas, Isabelle Percelay, Thibault Larcher, Christel Lefrancois. Effects of pyrolytic and petrogenic polycyclic aromatic hydrocarbons on 1 swimming and metabolic performance of zebrafish contaminated by 2 ingestion 3 4. *Ecotoxicology and Environmental Safety*, 2016, 132, pp.145-152. <10.1016/j.ecoenv.2016.05.035>. <hal-01451518>

**HAL Id: hal-01451518**

**<https://hal.science/hal-01451518v1>**

Submitted on 1 Feb 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

**Effects of pyrolytic and petrogenic polycyclic aromatic hydrocarbons on swimming and metabolic performance of zebrafish contaminated by ingestion**

Lucas J. <sup>a,b,\*</sup>, Percelay I. <sup>a</sup>, Larcher T. <sup>c</sup>, Lefrançois C. <sup>a</sup>

<sup>a</sup> UMR 7266 Littoral Environnement Sociétés (LIENSs), Institut du Littoral et de l'Environnement, 2 rue Olympe de Gouges, 17000 La Rochelle, France

<sup>b</sup> Ifremer, Place Gaby Coll, BP7, 17137 L'Houmeau, France

<sup>c</sup> INRA UMR 703, APEX, Oniris, La Chantrerie, 44300 Nantes, France

\*Author to whom correspondence should be addressed. Tel.: +33 5 46 45 72 17; e-mail:

[julie.lucas@laposte.net](mailto:julie.lucas@laposte.net)

## Abstract

Depending on their origins, polycyclic aromatic hydrocarbons (PAH) are characterized by different chemical properties. Petrogenic PAH (e.g. from fossil fuels) and pyrolytic PAH (e.g. those produced by incineration processes) are therefore expected to affect organisms differently. The impact of trophic exposure to these PAH was investigated on swimming and metabolic performance of zebrafish *Danio rerio*. Two-month-old juveniles and six-month-old adults were individually challenged following a swimming step protocol. While pyrolytic exposure did not affect fish whatever the duration of exposure, it appeared that petrogenic PAH impaired adults' performance. Indeed, the active metabolic rate in petrogenic PAH-contaminated adults was significantly reduced by 35%, and critical swimming speed by 26.5%. This was associated with cardiac abnormalities, which are expected to contribute to the reduction of oxygen transport, particularly during intensive effort. These results may be due to the different composition and toxicity of PAH mixtures.

**Keywords:** trophic exposition, petroleum hydrocarbons, critical swimming speed, aerobic metabolic rate, sub-lethal concentration

## 1. Introduction

Locomotion performance in fish is considered to be a relevant biological function contributing to survival (e.g. Graham et al., 1990; Jones et al., 1974; Plaut, 2001; Stobutzki and Bellwood, 1994; Swanson et al., 1998), since it is related to predator–prey interactions, food research and migration (Drucker and Jensen, 1996; Hammer, 1995; Plaut, 2001; Reidy et al., 1995; Videler, 1993; Walker et al., 2005; Watkins, 1996;). Critical swimming speed (i.e. the maximal velocity a fish can reach during a swimming step protocol,  $U_{crit}$ ; Hammer, 1995) is frequently employed as an indicator of swimming capacities in a varying environment (e.g. Beamish, 1978). Since  $U_{crit}$  notably depends on the maximal ability of fish to provide energy during sustained swimming activity, these studies are often associated with the assessment of metabolic performance, which contribute to a better view of the environmental impact. Aerobic metabolic scope (AMS; Brett, 1964; Fry, 1947, 1971) is defined as the difference between (i) the active metabolic rate (AMR), which is the highest metabolic rate the organism can sustain under maximal activity and (ii) the standard metabolic rate (SMR), the metabolic rate necessary to maintain vital functions and measured under resting conditions. Metabolic and swimming performance are known to be modulated by a set of environmental parameters, such as temperature (Blier et al., 1997; Farrell, 2007; Rome et al., 2007), dissolved oxygen (Diaz, 2001; Diaz et al., 2004) or pollutants (Johansen and Jones, 2011; Lefrançois and Claireaux, 2003; Marit and Weber, 2012; Shingles et al., 2001; Schurmann and Steffensen, 1997; Thomas et al., 2013).

This study aimed at investigating responses of fish exposed to environmentally relevant concentrations of PAH through the assessment of AMS and  $U_{crit}$  as indicators of the physiological state of the organism (Fry, 1947). The increase of anthropogenic activities in coastal areas induces discharges of PAH in aquatic ecosystems. Due to their high

liposolubility, PAH are typically adsorbed by organic matter or marine sediments, bioaccumulated by organisms at the lowest trophic levels (e.g. invertebrates; Bustamante et al., 2012; O'Connor and Lauenstein, 2006) and transferred through trophic chains (Hylland, 2006; Vignet et al., 2014). Past studies on fish have demonstrated that hydrocarbon exposure causes a decrease of  $U_{crit}$ . For instance, *Danio rerio* showed a 15% and 22% reduction of  $U_{crit}$  while exposed to crude oil (Hicken et al., 2011) and TCDD (2,3,7,8-tetrachlorodibenzo-p-dioxin; Marit and Weber, 2012), respectively. Moreover, PAH could induce damage of tissues and/or organs involved in oxygen transport (e.g. fish gills, Fasulo et al., 2012; mussel gills, Cappello et al., 2013). This could induce a decrease of AMR as observed by Davoodi and Claireaux (2007) after acute PAH exposure of *Solea solea*. In addition, the implementation of detoxification processes which are energy costly have already been demonstrated. These processes are likely to increase costs of maintenance and, consequently, SMR. The consequent potential reduction of AMS would reflect a decrease in the capacity of fish to sustain energy-demanding activities.

The effects of PAH mixtures were tested on zebrafish *Danio rerio* (F. Hamilton, 1822) contaminated by ingestion. Both pyrolytic PAH (PY-PAH, resulting from the combustion of organic matter) and petrogenic PAH (HO-PAH, originated from fossil fuels; Hylland, 2006) were considered in the present study. Present as complex mixtures in the environment, PY-PAH and HO-PAH show different chemical properties and may induce various effects on aquatic life. The main hypothesis was that chronic exposure to PY-PAH and HO-PAH would impair AMS by increasing SMR and/or reducing AMR. The consequent potential reduction of AMS would indicate a decrease in the capacity of the fish to support oxygen-demanding activities beyond SMR and consequently a reduction of  $U_{crit}$ . In addition, PAH may induce anatomic impairments due to teratogenic effects (Hawkins et al., 1990; Myers et al., 1991). Swimming and metabolic functions were explored, as well as the cardiac and skeletal muscle

structures, as two key tissues involved the tested functions.

## 2. Materials and methods

### 2.1. General production of zebrafish

Pairs of zebrafish (wild-type Tuebingen strain, TU) were reared together in 10 L tanks in the laboratory. Aquaria were filled with a mix of osmosis and filtered tap water which was characterized by the following parameters: temperature  $28 \pm 0.5$  °C, conductivity  $300 \pm 50$   $\mu\text{S}\cdot\text{cm}^{-1}$ , oxygen saturation  $\geq 80\%$ , pH  $7.5 \pm 0.5$ . Fish were reared with a photoperiod of 14 h light/10 h dark. Ammonia, nitrite and nitrate remained within recommended ranges (Lawrence, 2007). The fish were fed twice daily with uncontaminated commercial dry food (INICIO Plus, BioMar, France) and once by *Artemia* sp. nauplii (Ocean Nutrition, Belgium), occasionally supplemented with red sludge worms (Boschetto, Poland).

Over a period of one month, spawn was obtained from pairs weekly following the protocol described in Lucas et al. (2014a) and Vignet et al. (2014). Briefly, the day before spawn was collected, these fish couples were isolated in 1 L breeding tanks (Aqua Schwarz, Germany). After collection, fertilized and normally developed eggs were selected using a binocular microscope. Spawn was then mixed to avoid any parental influence. From 2 weeks onwards, fish were kept in groups of 30 individuals in 10 L aquaria in the same specific rearing system described above.

### 2.2. Food contamination protocol

Artificial dry food (INICIO Plus, BioMar France) was spiked with mixtures of PAH as

described in detail by Vignet et al. (2014). Briefly, the pyrolytic fraction (*PY*) was extracted from natural sediment sampled at the site of Oissel (Seine Bay, France), while the petrogenic fraction (*HO*) was extracted from heavy Erika fuel oil. For both types of *PY* and *HO* mixtures, quantification of the 16 individual PAH of the US EPA list was made, according to a protocol described in Budzinski et al. (1997) and Cachot et al. (2007). The food was then spiked with the extracted *PY* or *HO* fractions according to a protocol adapted from Vicquelin et al. (2011). The target concentration was 15000 ng.g<sup>-1</sup> of food and based on the sum of the 16 PAH from the US EPA list in dry weight. This corresponds to three times the PAH concentration organisms may be exposed to in their natural environment (Cachot et al., 2006; Cailleaud et al., 2007; Payne et al., 2008). This high concentration was used to highlight the toxicological effects on the physiological performances investigated in this study. It is called *X* in the rest of the manuscript. Food size was adapted to the developmental stage of *D. rerio*. A *control* treatment was also considered, and consisted of feeding groups of fish with uncontaminated food, treated only by the solvent dichloromethane. This solvent was used to improve PAH incorporation into the food, and then evaporated. PAH compositions and concentrations in the diet are described in detail in Table 1.

**Table 1.** Concentration of 21 US-EPA PAH in contaminated dry food. Values are expressed as mean±standard deviation. n.d.=not detected

PAHs	Molecular weight (g.mol <sup>-1</sup> )	Concentration in dry food (ng.g <sup>-1</sup> )			
		pyrolytic PAH		petrogenic PAH	
		Solvent	3X	Solvent	3X
Naphtalene	128.2	4.01±2.21	199.13±32.06	3.79±0.97	405.02±73.41
Acenaphthylene	152.2	0.87±0.26	124.82±12.38	1.31±0.51	13.10±12.77
Acenaphthene	154.2	1.91±0.64	99.49±10.80	27.29±26.07	190.11±17.26
Fluorene	166.2	2.62±0.48	148.22±18.02	2.34±1.05	312.01±25.07
Dibenzothiophenes	184.26	0.68±0.46	117.95±10.61	0.34±0.16	545.94±53.56
Phenanthrene	178.2	9.36±3.73	1029.78±88.76	6.11±2.73	1278.88±51.23
Antracene	178.2	0.78±0.31	598.14±53.81	0.37±0.22	219.76±9.75
Fluoranthenes	202.3	1.37±0.39	1968±179.78	1.92±0.71	145.08±17.70
Pyrenes	202.3	1.62±0.21	1648.54±179.45	1.45±0.27	709.13±64.10
Benzo[a]anthracene	228.3	0.76±0.26	2286.28±200.96	n.d.	542.78±29.08
Chrysene+Triphenylene	228.3	0.77±0.26	2984.46±240.64	n.d.	1073.29±78.69
Benzonaphthothiophene	252.3	6.53±0.73	660.11±56.98	n.d.	572.96±35.67
Benzo[b,k,l]fluoranthenes	252.3	1.54±0.56	3164.65±269.51	n.d.	362.75±17.62
Benzo[e]pyrene	252.3	0.89±0.37	1267.09±108.51	n.d.	535.54±25.67
Benzo[a]pyrene	252.3	0.72±0.55	1395.49±146.63	n.d.	342.12±10.42
Perylene	252.3	0.96±0.48	437.37±39.40	n.d.	172.01±13.89
Indeno[1,2,3-cd]pyrene	276.3	n.d.	1218.26±123.95	n.d.	68.42±10.88
Dibenzo[a]anthracene + Dibenzo[a]chrysene	276.3	2.22±1.74	369.38±53.74	n.d.	112.55±6.26
Benzo[g,h,i]perylene	276.3	n.d.	913.77±88.29	n.d.	480.53±12.21
Sum PAHs		34 ± 6	17305 ± 4798	39 ± 30	8082.5 ± 305

### 2.3. Zebrafish PAH exposure protocol

Contamination through food was conducted by feeding zebrafish larvae twice a day with one of the treatments (i.e. *control* and *X* for both types of contamination *PY* and *HO*). The diet started from the first meal (5 days post fertilization, dpf) until the age at the test: 2 months and 6 months for juveniles and adults, respectively. In accordance with the protocols in Vignet et al. (2014), larvae were fed ad libitum and then, starting from 2 months old, the ration of food was decreased by 5% to 2% of the biomass in each tank in order to maintain constant growth. For all fish, brine shrimp were distributed as complementary food once a day. *PY* exposure took place during the second semester of 2011 and *HO* exposure was carried out in summer 2013. Quantification of hydroxylated metabolites in larvae at 15 dpf indicated a dose-dependent increase of metabolites, confirming successful contamination (total

concentrations of hydroxylated metabolites for *PY*-PAH: *control* = 9.1 ng.g<sup>-1</sup> of tissue, *X* = 275 ng.g<sup>-1</sup>; Vignet et al., 2014).

For the *X* treatment tested, the average concentrations found were 136 ng.g<sup>-1</sup> for *PY* larvae and 42 ng.g<sup>-1</sup> for *HO* larvae, while it was 2 or 4 ng.g<sup>-1</sup> in the *PY* and *HO control* fish, respectively (Vignet et al., 2014). It appears that PAH concentration is higher for *PY* than for *HO*.

## 2.4. Aerobic metabolic scope and critical swimming speed

### 2.4.1. Experimental set-up

For both types of contamination (i.e. *PY*- and *HO*-PAH), two identical 170 mL swimming respirometers (Loligo Systems, Denmark) were used to assess the swimming and metabolic performance of fish. Each swimming respirometer was composed of (a) a swimming chamber, where the fish was placed to be tested, (b) a motor fitted with a three-bladed propeller to control water flow and (c) honeycomb placed at each side of the swimming chamber to laminarize the water flow. Each swimming respirometer was submerged in a 20 L buffer tank, and filled with temperature-controlled (i.e. 28 °C) and oxygenated mixed water as in the rearing system. The oxygen consumption (MO<sub>2</sub> in mg O<sub>2</sub>.g<sup>-1</sup>.h<sup>-1</sup>) associated with the activity of fish was measured by intermittent-flow respirometry (Steffensen, 1989). Water supply in each swimming respirometer was provided by flush pumps, which controlled water flow from the buffer tank to the swimming respirometer. This allowed alternation between phases of oxygen renewal and phases of MO<sub>2</sub> measurement with a cycle of 5:20 min. In addition, a peristaltic pump was used to create a continuous water flow from the respirometer to the oxygen probe (flow through an oxygen minisensor, PreSens, Germany). The probe was connected to an oxymeter (Microx, PreSens, Germany) to record the level of dissolved

oxygen in the water every 5 s. The probe was calibrated once at the beginning of the swim test using 0% and 100% air saturation for a controlled temperature of 28 °C.

#### 2.4.2. *Experimental protocol*

The fish were starved 24 h before being transferred individually to a swimming respirometer. They were then individually submitted to a swimming challenge. Each experimental trial consisted of challenging two fish in parallel exposed to one of three treatments: (i) fish fed by the *control* food and (ii) fish exposed to *PY*- or *HO*-PAH. Each individual was submitted to a short acclimation swimming period (~5 min) during which the velocity was increased to 2 body lengths per second ( $\text{BL.s}^{-1}$ ). Then, fish were left undisturbed with a water flow of 0.5  $\text{BL.s}^{-1}$  for 2 h to permit recovery from handling stress. The swimming challenge started by increasing water flow by steps of 1.5  $\text{BL.s}^{-1}$  from 0.5 to 6.5  $\text{BL.s}^{-1}$  and steps of 0.75  $\text{BL.s}^{-1}$  to the end of the test. Each step lasted 20 min, during which fish  $\text{MO}_2$  was measured. The experiment stopped when the fish fatigued, i.e. when it did not manage to swim against the current and stay on the honeycomb. Speed was then decreased to 0.5  $\text{BL.s}^{-1}$  for a recovery period of 1.5 h. Fish were then removed from the swimming respirometer and anaesthetized using benzocaine (50  $\text{mg.L}^{-1}$ ). Standard and total length, mass and sex of each individual were determined; characteristics of the fish tested in the swim tunnel are described in Table 2. Before and after each trial, a blank measurement was performed to quantify microbial oxygen consumption in the swimming respirometer. The average of these two values was subtracted from the measured oxygen consumption. Each fish was tested once.

**Table 2.** Biometry of fish for each PAHs fractions tested (Mean±SE). SL represents standard length and TL total length

	Treatment	Lifestage	Fish	Weight (g)	SL (cm)	TL (cm)
Pyrolytic PAH	Control	juveniles	n=21	0.19±0.06	2.20±0.19	2.68±0.35
		adults	n=16	0.68±0.21	3.13±0.25	3.88±0.35
	3X	juveniles	n=22	0.17±0.05	2.10±0.26	2.59±0.26
		adults	n=15	0.48±0.13	2.81±0.24	3.50±0.22
Petrogenic PAH	Control	juveniles	n=15	0.09±0.03	1.82±0.28	2.26±0.34
		adults	n=19	0.30±0.06	2.62±0.19	3.29±0.23
	3x	juveniles	n=12	0.07±0.03	1.55±0.26	1.97±0.29
		adults	n=11	0.12±0.04	1.79±0.33	2.27±0.32

196

### 197 2.4.3. Calculations

#### 198 2.4.3.1. Critical swimming speed

199 The critical swimming speed  $U_{crit}$  was calculated according to the formula of Brett (1964):

200

$$201 \quad U_{crit} = U_t + t_1 \cdot t^{-1} \cdot U_1 \quad \text{Equation (1)}$$

202

203 Where  $U_t$  (in  $BL \cdot s^{-1}$ ) is the highest velocity maintained for an entire step,  $t_1$  (in min) is the  
 204 time spent until the exhaustion of fish at the last step,  $t$  (in min) is the swimming period for  
 205 each step (i.e. 20 min in the present study) and  $U_1$  is the increment velocity (0.75 or 1.5  
 206  $BL \cdot s^{-1}$ ).

207

#### 208 2.4.3.2. Oxygen consumption ( $MO_2$ )

209 Oxygen consumption  $MO_2$  is expressed in  $mg \ O_2 \cdot g^{-1} \cdot h^{-1}$  and calculated using the following

formula:

$$MO_{2\text{meas}} = \Delta[O_2] \cdot V \cdot \Delta t^{-1} \cdot M_{\text{meas}}^{-1} \quad \text{Equation (2)}$$

where  $\Delta[O_2]$  (in mg  $O_2 \cdot L^{-1}$ ) is the variation in oxygen concentration during the measurement period  $\Delta t$  (in h),  $V$  (in L) is the volume of the respirometer minus the volume of the fish and  $M_{\text{meas}}$  (in g) is the fish mass measured.

An allometric relationship exists between oxygen consumption and body mass, which permits correction of  $MO_{2\text{meas}}$  using the following formula:

$$MO_{2\text{cor}} = MO_{2\text{meas}} \cdot (M_{\text{meas}} \cdot M_{\text{cor}}^{-1})^{1-b} \quad \text{Equation (3)}$$

where  $MO_{2\text{cor}}$  (in mg  $O_2 \cdot g^{-1} \cdot h^{-1}$ ) is the oxygen consumption related to a standard fish of 1 g ( $M_{\text{cor}}$ ),  $MO_{2\text{meas}}$  (in mg  $O_2 \cdot g^{-1} \cdot h^{-1}$ ) is the oxygen consumption estimated for experimental fish whose mass was  $M_{\text{meas}}$  (in g) and  $b$  is the allometric scaling exponent describing the relationship between oxygen consumption and body mass. A previous study of Lucas et al. (2014a) estimated  $b$  in zebrafish *D. rerio* from the relationship between AMR and body mass, i.e.  $b_{\text{AMR}}$ , and from the relationship between SMR and body mass, i.e.  $b_{\text{SMR}}$ . We used  $b_{\text{AMR}} = 0.926$  for the correction of AMR and  $b_{\text{SMR}} = 0.965$  for the correction of SMR.

#### 2.4.3.3. Standard metabolic rate, active metabolic rate and aerobic metabolic scope

Oxygen consumption is known to increase exponentially with swimming speed (Brett, 1964). The following equation can be therefore fitted on data and allows the assessment of SMR:

$$MO_2 = \text{SMR} \cdot \exp^{bU} \quad \text{Equation (4)}$$

where  $MO_2$  is the oxygen consumption (in  $mg\ O_2.g^{-1}.h^{-1}$ ), SMR is the intercept (i.e. the  $MO_2$  when  $U = 0\ BL.s^{-1}$ ),  $b$  is a constant and  $U$  is the swimming speed (in  $BL.s^{-1}$ ).

AMR is evaluated as the maximum oxygen consumption measured during the swimming challenge. AMS is the difference between AMR and SMR.  $U_{crit}$ , SMR, AMR and AMS were assessed for each fish except for HO-PAH-exposed juveniles. No SMR data were indeed recorded for these individuals, because of the small size of the fish (relative to the swim tunnel volume) and the low quality of the oxygen signal. The HO exposure impaired the growth of fish exposed to (Vignet et al. 2014) and consequently HO-PAH-exposed juveniles were smaller than PY-PAH-exposed juveniles (Table 2).

## 2.5. Histological analysis

Analysis occurred after the swimming challenge, in older fish: 3-month-old juveniles for both types of contamination, 9-month-old adults for PY-PAH and 7-month-old adults for HO-PAH. A total of 20 juveniles per treatment (i.e. *control* and *X*) were analysed for PY- and HO-PAH exposure. In adults, 30 fish per treatment were observed for PY-PAH, and 25 fish were observed for HO-PAH. Fish were euthanized with a lethal dose of benzocaine ( $100\ mg.l^{-1}$ ; Sigma-Aldrich), fixed in 4 % formalin for 2 weeks, dehydrated and embedded in paraffin wax (Daouk et al., 2011). For each fish, 4 serial  $4\ \mu m$ -thick sections spaced from from about  $250\ \mu m$ , were routinely stained with haematoxylin-eosin-saffron (HES). The histological appearance of cardiac and skeletal muscles of fish was then observed using a bright-field microscope (Nikon Eclipse 90i, Champigny, France) equipped with 2X, 4X, 10X, 20X and 40X objectives and a SPOT digital camera (AZ microscope, London, UK). Lesions were systematically recorded by a skilled veterinary pathologist (member of the the European College of Veterinary Pathologists).


## 2.6. Statistical analysis

Statistical analysis was carried out using GraphPad Prism software. For AMR, SMR, AMS and  $U_{crit}$ , normality and homoscedasticity were checked using Kolmogorov–Smirnov and Bartlett tests, respectively, for each duration of exposure and for each type of contamination (i.e. *PY* and *HO*). Student's t-tests were used to compare individuals exposed to the *control* treatment with individuals exposed to PAH. Chi-square tests were used to compare histological anomalies (i.e. cardiac and skeletal muscle structures) between *control* fish and those exposed to *PY*-PAH or *HO*-PAH. For all tests, the results were considered to be significantly different when  $p < 0.05$ .

## 3. Results

### 3.1. $U_{crit}$


Whatever the duration of exposure, no significant difference was found between fish exposed to *control* or *PY*-PAH (Student's t-test, degrees of freedom (dof) = 41 and  $p = 0.34$  for juveniles; dof = 26 and  $p = 0.40$  for adults; Fig. 1A). On the contrary,  $U_{crit}$  decreased significantly in both *HO*-PAH-contaminated juveniles and adults compared to *control*-exposed individuals (Student's t-test: dof = 24 and  $p = 0.011$ , dof = 28 and  $p = 0.01$ , respectively; Fig. 1B).



**Fig. 1** Critical swimming speed ( $U_{crit}$ ) of zebrafish exposed to control (white bars) or to PAHs from pyrolytic (A) or petrogenic (B) origins (in grey). PAHs concentrations in food was 3X; the environmental reference concentrations  $X$  corresponding  $X_{py}=17305\pm4798\mu\text{g.g}^{-1}$  and  $X_{pe}=8082.5\pm305\mu\text{g.g}^{-1}$  of dry food (table 1).  $U_{crit}$  was evaluated in juvenile and adults. Results are expressed as mean values  $\pm$  standard deviation. For a given type of PAHs (PY or HO) and a given duration, \* indicates a significant difference in  $U_{crit}$  between control and 3X treatment.

### 3.2. Aerobic metabolic scope, standard and active metabolic rate

For PY-PAH contamination, no significant difference was found between contaminated fish and the control whatever the metabolic rate observed: SMR (Fig. 2A; Student's t-test: dof = 32 and  $p=0.40$ , dof = 26 and  $p=0.46$ , for juveniles and adults, respectively), AMR (Fig. 2A; Student's t-test: dof = 34 and  $p=0.29$ , dof = 26 and  $p=0.44$  for juveniles and adults, respectively) and AMS (Student's t-test: dof = 34 and  $p=0.29$ , dof = 26 and  $p=0.44$  for juveniles and adults, respectively).



300

301

302

303

304

305

306

307

308

**Fig. 2.** Standard metabolic rate (SMR), active metabolic rate (AMR) and aerobic metabolic scope (AMS) of zebrafish *Danio rerio* exposed to control food (white bars) containing, pyrolytic (A) or petrogenic (B) PAHs (grey bars). PAH concentrations in food was X; the environmental reference concentrations corresponding to  $X_{py}=17305\pm4798\mu\text{g.g}^{-1}$  and  $X_{pe}=8082.5\pm305\mu\text{g.g}^{-1}$  of dry food (table 1). Aerobic metabolic rate was estimated at two life stages: juvenile and adults. Results are expressed as mean values  $\pm$  standard deviation. For a given type of PAHs (PY or HO) and a given duration, \* indicates a significant difference in AMR between control and X treatment.

309

310

311

312

313


314

315

In *D. rerio* which ingested HO-PAH-contaminated food, the lack of SMR data for juveniles did not allow estimation of SMR and AMS for this stage. In adults, SMR did not differ significantly between control and contaminated fish (Fig. 2B; Student's t-test: dof = 27 and p = 0.35). On the contrary, AMR was significantly impaired, but only in adults (Student's t-test: dof = 20 and p = 0.43, dof = 27 and p = 0.006, for juveniles and adults, respectively; Fig. 2B), as was AMS (Student's t-test: dof = 27 and p = 0.003).

### 3.3. Histological analysis

During histopathological examination, no anomalies were observed in skeletal muscle for either *PY*- or *HO*-PAH exposure. While no heart damage was observed for *PY*-PAH exposure, 4 out of 25 fish exposed to *HO*-PAH for 7 months displayed generalized oedema that severely thickened the peritoneum all around the abdominal cavity. Significant heart impairments were associated with this oedema: the pericardium appeared dilated with fluid and the heart was very small and rounded with a thin ventricular wall (Fig. 3; chi-square test, dof = 1 and  $p = 0.002$ ). These lesions are characteristic of cardiac insufficiency with secondary collection of fluid in connective tissue and body cavities.



**Fig. 3** Heart of zebrafish exposed to control condition of food (A) or to PAHs from petrogenic origin (B). Compared to a control fish, some fish exposed to PAHs displayed a large pericardial cavity with a small rounded heart. at: atrium; ba: bulbus arteriosus; p: pericardium; sv: sinus venosus; vt: ventricle. Hemalun-Eosin-Saffron. Bars = 200  $\mu$ m.

#### 4. Discussion

Under these experimental conditions and at the two life stages tested, trophic exposure to PAH differently impaired the fish depending on the type of PAH, i.e. PY or HO, and the considered traits. AMR and  $U_{crit}$  were indeed influenced by HO-PAH, but SMR was not.

First, for a general comparison, swimming performance data were in the range of what other authors have observed in *D. rerio*, since average  $U_{crit}$  ranged between 10 and 16.5 BL.s<sup>-1</sup> in *control* individuals (e.g.  $U_{crit} = 13$  BL.s<sup>-1</sup> in Plaut and Gordon, 1994;  $U_{crit} = 15.5$  BL.s<sup>-1</sup> in Plaut, 2001);  $U_{crit} = 18$  BL.s<sup>-1</sup> in Palstra et al., 2010).

AMR values in *control* fish ranged between 0.9 and 1.6 mg O<sub>2</sub>.h<sup>-1</sup>.kg<sup>-1</sup>, i.e. 30 to 60 lower compared with other studies in the same species (Thomas and Janz, 2011, Lucas et al., 2014a, 2016). Such differences observed in AMR estimation could be explained by differences in the general physiological status of the tested zebrafish (e.g. related to age, thermal acclimatization, rearing conditions), as well as in the protocols used. It is also worth noting that Peake and Farrell (2006) underlined the importance of behavioural regulations in assessing fish performance through swimming challenges. They suggested that behaviour, rather than physiological exhaustion, participates in the fatigue observed in fish tested in a swim flume, thereby increasing the variability of the measured parameters (i.e. AMR and  $U_{crit}$ ).

Focussing on contamination, HO-PAH exposure induced a significant decrease of 15 and 26.5% of  $U_{crit}$  in zebrafish exposed for 2 and 6 months, respectively. A similar pattern was observed in the literature regarding pollutant effects (Cheng and Farrell, 2007; Hicken et al., 2011; Howard, 1975; Marit and Weber, 2011, 2012; Thomas and Janz, 2011). In HO-PAH adults, this was associated with a significant reduction of AMR, cardiac lesions (i.e. a smaller

and more rounded heart with a thin ventricular wall), generalized oedema and collection of fluid in body cavities indicating a functional cardiac insufficiency. These lesions may induce a reduction of cardiac performance and capacity to transport oxygen. This may limit aerobic metabolism, especially during high levels of activity, such as occurs during swimming (Farrell 1997, 2007; Fry, 1947, 1971; Claireaux et al., 2005; Priede, 1985).

Even if  $U_{crit}$  was reduced in HO-PAH juveniles, no impairment of AMR or cardiac function was observed. It is worth noting that  $U_{crit}$  depends on both aerobic and anaerobic metabolism (Webb, 1998), which could explain the results observed. However, the anaerobic part of the metabolism required to attain  $U_{crit}$  was not assessed in the present study, nor was oxygen debt after the swimming test (e.g. Vagner et al., 2008). Indeed, it would have been interesting to assess whether reduction of  $U_{crit}$  was associated to a different solicitation of anaerobic metabolism in juveniles and adults'

Contrary to the HO-PAH mixture, exposure to PY-PAH did not impair fish metabolic and swimming performance or cardiac anatomy. The lack of effect could be explained by the fact that organisms that suffer long-term chronic environmental stress can present physiological adaptations to maintain their homeostasis. Chronic exposure to PY-PAH may have induced such adaptations in *D. rerio*.

Whatever the type of exposure, no effect was observed on SMR. The range of observed values are in agreement with previous studies carried out for the same species and with a similar experimental approach ( $SMR = 0.19 \text{ mg O}_2 \cdot \text{g}^{-1} \cdot \text{h}^{-1}$  in Barrionuevo and Burggren, 1999; mean  $\pm$  s.d.,  $SMR = 0.31 \pm 0.11$  and  $0.35 \pm 0.16 \text{ mg O}_2 \cdot \text{g}^{-1} \cdot \text{h}^{-1}$  in juveniles and adults, respectively, in Lucas et al., 2014a, 2016).

Even though the concentration of PAH and their metabolite compounds was not assessed in juveniles and adults in this study, they were estimated in 15 dpf larvae exposed to HO- or PY-

PAH, and ranged between 42 ng.g<sup>-1</sup> and 136 ng.g<sup>-1</sup>, respectively (Vignet et al., 2014).

It is likely that PAH metabolite concentration increased proportionally with the duration of exposure. However, the present results suggest that the level of contamination tested was not sufficiently extreme to induce significant variation in SMR in our experimental conditions., PAH exposure did not induce supplementary costs of maintenance due to detoxification processes in zebrafish (Lannig et al., 2006; Lucas et al., 2016; Sokolova and Lannig, 2008; Wilson et al., 1994), which is contrary to one of the initial hypotheses.

The different impairment of the types of PAH mixture the fish were exposed to could be explained by their different chemical properties depending on the composition of individual PAH (Vignet et al., 2014). It is worth noting that toxicity depends on individual intrinsic properties of each component and their interactions with enzymatic systems in organisms. Mainly characterized by four- to six-ring PAH and 5% of methylated compounds, PY-PAH could induce less toxic effects than HO-PAH despite its higher concentration (Table 1). Indeed, HO-PAH mixtures were characterized by three- and four-ring PAH, and 47% of methylated compounds, which are considered to be more toxic for organisms (Di Giulio and Hinton, 2008; Hall and Oris, 1991; Hylland, 2006; Tuvikene, 1995). In addition, adults that were exposed to PAH for 6 months were therefore expected to contain more PAH metabolites than juveniles, which coped with PAH exposure for only 2 months. This may contribute to the fact that exposure to HO-PAH induced a significant reduction of AMR in adults and not in juveniles. However, this pattern was not observed in juveniles and adults exposed to PY-PAH (Fig. 1B) as well as in the similar study of Lucas et al. (2016).

## 5. Conclusion

Our results suggest that, contrary to HO-PAH, PY-PAH have no direct effect on aerobic metabolism and swimming performance. Lucas et al. (2014b) have observed impairment of metabolic performance and cardiac activity on the progeny of PY-PAH-contaminated *D. rerio* parents. The parental effect of PAH exposure appears to be an important subject to study. This study on swimming performance and associated energetic costs will be completed by investigations regarding escape performance. This type of swimming depends on anaerobic metabolism, as well as on sensorial and locomotion performance. Moreover, interfering with some physiological mechanisms, PAH could be narcotic, inducing lethargic behaviour and impairing the swimming performance of fish (Wassenberg and Di Giulio, 2004). All these studies will permit a global view of the impact of this type of trophic contamination across zebrafish generations.

## **Acknowledgements**

The authors are grateful to Didier Leguay and Michel Prineau for their help during the experiment. All experiments were carried out at Ifremer (Plateforme d'Ecophysiologie des Poissons), La Rochelle station, France. This study was financially supported by the ANR project ConPhyPoP (CES 09\_002) and JL received a doctoral grant from the Regional Council of Poitou-Charentes. This study was conducted under the approval of the Animal Care Committee of France under the official licence to Marie-Laure Bégout (17-010).

## References

- Barrionuevo, W.R., Burggren, W.W., 1999. O<sub>2</sub> consumption and heart rate in developing zebrafish (*Danio rerio*): influence of temperature and ambient O<sub>2</sub>. *Am. J. Physiol.* 276, 505–513.
- Beamish, F.W.H., 1978. Swimming capacity, in: Hoar, W.S., Randall, D.J. (Eds.), *Fish Physiology*, Vol. 7. Academic Press, New York, London, pp. 101–187.
- Blier, P.U., Pelletier, D., Dutil, J.D., 1997. Does aerobic capacity set a limit on fish growth rate? *Rev. Fish. Sci.* 5(4), 323–340.
- Brett, J.R., 1964. The respiratory metabolism and swimming performance of young sockeye salmon. *J. Fish. Res. Board Can.* 21, 1183–1226.
- Budzinski, H., Jones, I., Bellocq, J., Pierard, C., Garrigues, P., 1997. Evaluation of sediment contamination by polycyclic aromatic hydrocarbons in the Gironde estuary. *Mar. Chem.* 58, 85–97.
- Bustamante, P., Luna-Acosta, A., Clemens, S., Cassi, R., Thomas-Guyon, H., Warnau, M. (2012). Bioaccumulation and metabolism of 14C-pyrene by the Pacific oyster *Crassostrea gigas* exposed via seawater. *Chem.* 87, 938–944.
- Cachot, J., Geffard, O., Augagneur, S., Lacroix, S., Le Menach, K., Peluhet, L., Budzinski, H., 2006. Evidence of genotoxicity related to high PAH content of sediments in the upper part of the Seine estuary (Normandy, France). *Aquat. Toxicol.* 79, 257–267.
- Cachot, J., Law, M., Pottier, D., Peluhet, L., Norris, M., Budzinski, H., Winn, R., 2007. Characterization of toxic effects of sediment-associated organic pollutants using the lambda transgenic medaka. *Environ. Sci. Technol.* 41, 7830–7836.
- Cailleaud, K., Forget-Leray, J., Souissi, S., Hilde, D., LeMenach, K., Budzinski, H., 2007. Seasonal variations of hydrophobic organic contaminant concentrations in the water-

column of the Seine Estuary and their transfer to a planktonic species *Eurytemora*  
*affinis* (Calanoida, copepoda). Part 1: PCBs and PAHs. Chemosphere 70(2), 270–280.

Cannas, M., Atzori, F., Rupsard, F., Bustamante, P., Loizeau, V., Lefrançois, C., 2013. PCB  
contamination does not alter aerobic metabolism and tolerance to hypoxia of juvenile  
sole (*Solea solea*). Aquat. Toxicol. 127, 54–60.

Cappello, T., Mauceri, A., Corsaro, C., Maisano, M., Parrino, V., Lo Paro, G., Messina, G.,  
Fasulo, S., 2013. Impact of environmental pollution on caged mussels *Mytilus*  
*galloprovincialis* using NMR-based metabolomics. Mar. Pollut. Bull. 77, 132–139.

Cheng, W.W., Farrell, A.P., 2007. Acute and sublethal toxicities of rotenone in juvenile  
rainbow trout (*Oncorhynchus mykiss*): swimming performance and oxygen consumption.  
Arch. Environ. Contam. Toxicol. 52, 388–396.

Claireaux, G., McKenzie, D.J., Genge, A.G., Chatelier, A., Aubin, J., Farrell, A.P., 2005.  
Linking swimming performance, cardiac pumping ability and cardiac anatomy in  
rainbow trout. J. Exp. Biol. 208, 1775–1784.

Clark, T.D., Donaldson, M.R., Pieperhoff, S., Drenner, S.M., Lotto, A., Cooke, S.J., Hinch,  
S.G., Patterson, D.A., Farrell, A.P., 2012. Physiological benefits of being small in a  
changing world: responses of coho salmon (*Oncorhynchus kisutch*) to an acute thermal  
challenge and a simulated capture event. PLoS One 7, e39079.

Daouk, T., Larcher T., Rupsard, F., Lyphout, L., Rigaud, C., Ledevin, M., Loizeau, V.,  
Cousin, X., 2011. Long-term food-exposure of zebrafish to PCB mixtures mimicking  
some environmental situations induces ovary pathology and impairs reproduction ability.  
Aquat. Toxicol. 105(3–4), 270–278.

Davoodi, F., Claireaux, G., 2007. Effects of exposure to petroleum hydrocarbons upon the  
metabolism of the common sole *Solea solea*. Mar. Pollut. Bull. 54, 928–934.

Diaz, R.J., 2001. Overview of hypoxia around the world. J. Environ. Qual. 30, 275–281.

479 Diaz, R.J., Nestlerode, J., Diaz, M.L., 2004. A global perspective on the effects of  
 480 eutrophication and hypoxia on aquatic biota, in: Rupp, G.L., White, M.D. (Eds.),  
 481 Proceedings of the 7th International Symposium on Fish Physiology, Toxicology and  
 482 Water Quality, Tallinn, Estonia, May 12–15, 2003. EPA 600/R-04/049, U.S.  
 483 Environmental Protection Agency, Ecosystems Research Division, Athens, Georgia, pp.  
 484 1–33.

485 Drucker, E.G., Jensen, J.S., 1996. Pectoral fin locomotion in the striped surfperch. I.  
 486 Kinematics effects of swimming speed and body size. *J. Exp. Biol.* 199, 2235–2242.

487 Farrell, A.P., 1997. Effects of temperature on cardiovascular performance, in: Wood, C.M.,  
 488 McDonald, D.G. (Eds.), *Global Warming Implications for Freshwater and Marine Fish*.  
 489 S.E.B. Seminar Series 61. Cambridge University Press, Cambridge, pp. 135–153.

490 Farrell, A.P., 2007. Cardiorespiratory performance during prolonged swimming tests with  
 491 salmonids: a perspective on temperature effects and potential analytical pitfalls. *Philos.*  
 492 *Trans. R. Soc., B* 362, 2017–2030.

493 Fasulo, S., Maisano, M., Sperone, E., Mauceri, A., Bernabo, I., Cappello, T., D’Agata, A.,  
 494 Tripepi, S., Bruneli, E., 2012. Toxicity of Foroozan crude oil to ornate wrasse  
 495 (*Thalassoma pavo*, Osteichthyes, Labridae): ultrastructure and cellular biomarkers. *Ital.*  
 496 *J. Zool.* 79(2), 182–199.

497 Fry, F.E.J., 1947. The effects of the environment on animal activity. University of Toronto  
 498 Studies in Biology Series 55, 1–62.

499 Fry, F.E.J., 1971. The effect of environmental factors on the physiology of fish, in: Hoar, W.S.,  
 500 Randall, D.J. (Eds.), *Fish Physiology Vol. VI*. Academic Press, New York, pp. 1–98.

501 Graham, J.B., Dewar, H., Lai, N.C., Lowell, W.R., Arce, S.M., 1990. Aspects of shark  
 502 swimming performance determined using a large water tunnel. *J. Exp. Biol.* 151, 175–  
 503 192.

504 Hall, T.A., Oris, J.T., 1991. Anthracene reduces reproductive potential and is maternally  
 505 transferred during long-term exposure in fathead minnows. *Aquat. Toxicol.* 19, 249–  
 506 264.

507 Hammer, C., 1995. Fatigue and exercise tests with fish. *Comp. Biochem. Physiol.* 112, 1–20.

508 Hawkins, W.E., Walker, W.W., Overstreet, R.M., Lytle, J.S., Lytle, T.F., 1990. Carcinogenic  
 509 effects of some polycyclic aromatic hydrocarbons on the Japanese medaka and guppy in  
 510 waterborne exposures. *Sci. Total Environ.* 94(1–2), 155–167.

511 Hicken, C.E., Linbo, T.L., Baldwin, D.H., Willis, M.L., Myers, M.S., Holland, L., Incardona,  
 512 J.P., 2011. Sub-lethal exposure to crude oil during embryonic development alters cardiac  
 513 morphology and reduces aerobic capacity in adult fish. *Proc. Natl. Acad. Sci. U. S. A.*  
 514 108, 7086–7090.

515 Howard, T.E., 1975. Swimming performance of juvenile coho salmon (*Oncorhynchus kisutch*)  
 516 exposed to bleached kraft pulpmill effluent. *J. Fish. Res. Board Can.* 32, 789–793.

517 Hylland, K., 2006. Polycyclic aromatic hydrocarbon (PAH) ecotoxicology in marine  
 518 ecosystems. *J. Toxicol. Environ. Health, Part A* 69(1–2), 109–23.

519 Johansen, J.L., Jones, G.P., 2011. Increasing ocean temperature reduces the metabolic  
 520 performance and swimming ability of coral reef damselfishes. *Global Change Biol.* 17,  
 521 2971–2979.

522 Jones, D.R., Kiceniuk, J.W., Bamford, O.S., 1974. Evaluation of the swimming performance  
 523 of several fish species from the Mackenzie River. *J. Fish. Res. Board Can.* 31, 1641–  
 524 1647.

525 Jourdan-Pineau, H., Dupont-Prinet, A., Claireaux, G., McKenzie, D.J., 2010. An investigation  
 526 of metabolic prioritization in the European sea bass, *Dicentrarchus labrax*. *Physiol.*  
 527 *Biochem. Zool.* 83, 68–77.

528 Lannig, G., Flores, J.F., Sokolova, I.M., 2006. Temperature-dependent stress response in

529 oysters, *Crassostrea virginica*: pollution reduces temperature tolerance in oysters. *Aquat.*  
530 *Toxicol.* 79, 278–287.

531 Lawrence, C., 2007. The husbandry of zebrafish (*Danio rerio*): a review. *Aquaculture* 269, 1–  
532 20.

533 Lefrançois, C., Claireaux, G., 2003. Influence of ambient oxygenation and temperature on  
534 metabolic scope and scope for heart rate in the common sole *Solea solea*. *Mar. Ecol.: Prog. Ser.* 259, 273–284.

536 Lucas, J., Bonnieux, A., Lyphout, L., Cousin, X., Miramand, P., Lefrancois, C., 2016. Trophic  
537 contamination by pyrolytic polycyclic aromatic hydrocarbons does not affect aerobic  
538 metabolic scope in zebrafish *Danio rerio*. *Ecotoxicol. Environ. Saf.* 88, 433–442

539 Lucas, J., Schouman, A., Lyphout, L., Cousin, X., Lefrancois, C., 2014a. Allometric  
540 relationship between body mass and aerobic metabolism in zebrafish *Danio rerio*. *J.*  
541 *Fish Biol.* 84(4), 1171–1178.

542 Lucas J., Perrichon P., Nouhaud M., Audras A., Le Menach, K., Budzinski H. , Lefrancois C.  
543 2014b Aerobic metabolism and cardiac activity in the descendants of zebrafish exposed  
544 to pyrolytic polycyclic aromatic hydrocarbons. *Environ. Sci. Pollut. Res.* 21:13888–  
545 13897

546 Marit, J.S., Weber, L.P., 2011. Acute exposure to 2,4-dinitrophenol alters zebrafish swimming  
547 performance and whole body triglyceride levels. *Comp. Biochem. Physiol., Part C:*  
548 *Toxicol. Pharmacol.* 154, 14–18.

549 Marit, J.S., Weber, L.P., 2012. Persistent effects on adult swim performance and energetics in  
550 zebrafish developmentally exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Aquat.*  
551 *Toxicol.* 106–107, 131–139.

552 Myers, M.S., Landah, J.T., Krahn, M.M., McCain, B.B., 1991. Related lesions and exposure  
553 to relationships between hepatic neoplasms toxic chemicals in marine fish from the U.S.

554 West Coast. Environ. Health Perspect. 90, 7–15.

555 O'Connor, T.P. & Lauenstein, G.G. (2006). Trends in chemical concentrations in mussels and  
556 oysters collected along the US coast: update to 2003. Mar. Environ. Res. 62, 261–285.

557 Palstra, A.P., Tudorache, C., Rovira, M., Brittiijn, S.A., Burgerhout, E., van den Thillart, G. E.,  
558 Spaink, H.P., Planas, J.V., 2010. Establishing zebrafish as a novel exercise model:  
559 swimming economy, swimming-enhanced growth and muscle growth marker gene  
560 expression. PLoS One 5(12), e14483.

561 Payne, J.R., Driskell, W.B, Short, J.W., Larsen, M.L., 2008. Long term monitoring for oil in  
562 the Exxon Valdez spill region. Mar. Pollut. Bull. 56(12), 2067–2081.

563 Peake, S.J., Farrell, A.P., 2006. Fatigue is a behavioural response in respirometer confined  
564 smallmouth bass. J. Fish Biol. 68, 1742–1755.

565 Plaut, I., 2001. Critical swimming speed: its ecological relevance. Comp. Biochem. Physiol.  
566 131, 41–50.

567 Plaut, I., Gordon, M.S., 1994. Swimming metabolism of wild-type and cloned zebrafish  
568 *Brachydanio rerio*. J. Exp. Biol. 194, 209–223.

569 Priede, I.G., 1985. Metabolic scope in fishes, in: Tytler, P., Calow, P. (Eds.), Fish Energetics:  
570 New Perspectives. Croom Helm, London and Sydney, pp. 33–63.

571 Reidy, S.P., Nelson, J.A., Tang, Y., Kerr, S.R., 1995. Postexercise metabolic rate in Atlantic  
572 cod (*Gadus morhua*) and its dependence upon the method of exhaustion. J. Fish Biol. 47,  
573 377–386.

574 Roche, D.G., Binning, S.A., Bosiger, Y., Johansen, J.L., Rummer, J.L., 2013. Finding the best  
575 estimates of metabolic rates in a coral reef fish. J. Exp. Biol. 216(11), 2103–2110.  
576 doi:10.1242/jeb.082925

577 Rome, L.C., 2007. The effect of temperature and thermal acclimation on the sustainable  
578 performance of swimming scup. Philos. Trans. R. Soc., B 362, 1995–2016.

579 Schurmann, H., Steffensen, J.F., 1997. Effects of temperature, hypoxia and activity on the  
580 metabolism of juvenile Atlantic cod. *J. Fish Biol.* 50, 1166–1180.

581 Shingles, A., McKenzie, D.J., Taylor, E.W., Moretti, A., Butler, P.J., Ceradini, S., 2001.  
582 Effects of sublethal ammonia exposure on swimming performance in rainbow trout  
583 (*Oncorhynchus mykiss*). *J. Exp. Biol.* 204, 2691–2698.

584 Sokolova, I.M, Lannig, G., 2008. Interactive effects of metal pollution and temperature on  
585 metabolism in aquatic ectotherms: implications of global climate change. *Clim. Res.* 37,  
586 181–201.

587 Steffensen, J.F. (1989). Some errors in respirometry of aquatic breathers: how to avoid and  
588 correct for them. *Fish Physiol. Biochem.* 6, 49–59.

589 Stobutzki, I.C., Bellwood, D.R., 1994. An analysis of the sustained swimming abilities of pre-  
590 and post-settlement coral reef fishes. *J. Exp. Biol. Ecol.* 175, 275–286.

591 Swanson, C., Young, P.S., Cech, J.J., Jr., 1998. Swimming performance of Delta smelt:  
592 maximum performance, and behavioral and kinematic limitations on swimming at  
593 submaximal velocities. *J. Exp. Biol.* 201, 333–345.

594 Thomas, J.K., Janz, D.M., 2011. Dietary selenomethionine exposure in adult zebrafish alters  
595 swimming performance, energetics and the physiological stress response. *Aquat.*  
596 *Toxicol.* 102, 79–86.

597 Thomas, J.K., Wiseman, S., Giesy, J.P., Janz, D.M., 2013. Effects of chronic dietary  
598 selenomethionine exposure on repeat swimming performance, aerobic metabolism and  
599 methionine catabolism in adult zebrafish (*Danio rerio*). *Aquat. Toxicol.* 130–131, 112–  
600 122.

601 Tuvikene, A., 1995. Responses of fish to polycyclic aromatic hydrocarbons. *Ann. Zool. Fenn.* 32,  
602 295–309.

603 Vagner, M., Lefrançois, C., Ferrari, R.S., Satta, A., Domenici, P., 2008. The effect of acute

hypoxia on swimming stamina at optimal swimming speed in flathead grey mullet  
*Mugil cephalus*. Mar. Biol. 155, 183–190.

Vicquelin, L., Leray-Forget, J., Peluhet, L., LeMenach, K., Deflandre, B., Anschutz, P.,  
 Etcheber, H., Morin, B., Budzinski, H., Cachot, J., 2011. A new spiked sediment assay  
 using embryos of the Japanese medaka specifically designed for a reliable toxicity  
 assessment of hydrophobic chemicals. Aquat. Toxicol. 105, 235–245.

Videler, J.J., 1993. Fish Swimming. Chapman and Hall, London.

Vignet, C., LeMenach, K., Mazurais, D., Lucas, J., Perrichon, P., Le Bihanic, F., Lyphout, L.,  
 Frère, L., Bégout, M.-L., Zambonio-Infante, J.-L., Budzinski, H., Cousin, X., 2014.  
 Chronic exposures to pyrolytic and petrogenic mixtures of PAHs through diet produce  
 physiological disruptions in zebrafish - Part I: Survival and growth. Environ. Sci. Pollut.  
 Res.

Waiwood, K.G., Beamish, F.W.H., 1978. Effects of copper, pH and hardness on the critical  
 swimming performance of rainbow trout (*Salmo gairdneri* Richardson). Water Res. 12,  
 611–619.

Walker, J.A., Ghalambor, C.K., Griset, O.L., McKenny, D., Reznick, D.N., 2005. Do faster  
 starts increase the probability of evading predators? Funct. Ecol. 19(5), 808–815.

Wassenberg, D.M., Di Giulio, R.T., 2004. Synergistic embryotoxicity of polycyclic aromatic  
 hydrocarbon aryl hydrocarbon receptor agonists with cytochrome P4501A inhibitors in  
*Fundulus heteroclitus*. Environ. Health Perspect. 112(17), 1658–1664.  
 doi:10.1289/ehp.7168

Watkins, T.B., 1996. Predator-mediated selection on burst swimming performance in tadpoles  
 of the Pacific tree frog, *Pseudacris regilla*. Physiol. Zool. 69, 154–167.

Webb, P.W., 1998. Swimming, in: Evans, D.H. (Ed.), The Physiology of Fishes. CRC Marine  
 Science Series, New York, pp. 3–24.

629 Wilson, R.W., Bergman, H.L., Wood, C.M., 1994. Metabolic costs and physiological  
630 consequences of acclimation to aluminum in juvenile rainbow trout (*Oncorhynchus*  
631 *mykiss*). 1: Acclimation specificity, resting physiology, feeding, and growth. Can. J. Fish.  
632 Aquat. Sci. 51, 527–535.  
633  
634