



HAL
open science

Invasive water primroses in France become a weed problem in meadows

Jacques J. Haury, Julie Coudreuse, Michel M. Bozec

► **To cite this version:**

Jacques J. Haury, Julie Coudreuse, Michel M. Bozec. Invasive water primroses in France become a weed problem in meadows. International Conference on Ecological Sciences, Oct 2016, Marseille, France. , 2016, Sféologie 2016. hal-01453899

HAL Id: hal-01453899

<https://hal.science/hal-01453899>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Invasive Water Primroses in France become a weed problem in meadows

Jacques Haury*¹, Julie Coudreuse , and Michel Bozec

¹Agrocampus Ouest, UMR Ecologie et Santé des Ecosystèmes (AO UMR ESE) – Ministère de l'Agriculture, de l'Agroalimentaire et de la Forêt – 65 rue de Saint Briec, CS 84215, F35042 RENNES Cedex, France

Abstract

Invasive Water Primroses (*Ludwigia grandiflora* and *L. peploides*) are amphibious Onagraceae native in tropical and subtropical American areas. They become a weed problem in wet meadows along rivers and marshes. They affect pasture productivity, biodiversity and land use for cattle breeders.

Considering the distribution in France and results of enquiries, we present the problem and focus on local risk analysis of invasion by these plants.

We have performed detailed studies in Western France since 2001 at the site level. In many areas, terrestrial forms appear on winter-submersed meadows. They can be as productive in terrestrial habitats than in aquatic ones, around 2 kg dry weight per m². Both species can grow altogether but *L. grandiflora* is more competitive. Colonization begins when flooding with cuttings that settle when native plants do not form dense canopy. Aquatic plants are dense, longer with poorly branched stems, while terrestrial ones in wet meadows are shorter, with a bushy form. They are highly competitive over native plants. At the beginning or colonization, populations do not form fruits, at least for *L. grandiflora*, but after some years, fruits are observed. *Ex situ* germination studies point out seed fertility. Seedlings begin to appear in some sites.

Effects on agricultural exploitations are very bad: loose of EU grants, reduction of grazing area and impossibility to use the colonized pastures, leading to land abandonment and marsh agriculture decrease. Some recommendations to reduce colonization and restore pastures conclude the presentation.

Keywords: Jussie, Water Primrose, Wetlands, Agriculture, Adaptation, Biological Invasion

*Speaker