



**HAL**  
open science

# Total Domination, Connected Vertex Cover and Steiner Tree with Conflicts

Alexis Cornet, Christian Laforest

► **To cite this version:**

Alexis Cornet, Christian Laforest. Total Domination, Connected Vertex Cover and Steiner Tree with Conflicts. [Research Report] LIMOS (UMR CNRS 6158), universit  Clermont Auvergne, France. 2017. hal-01455072v1

**HAL Id: hal-01455072**

**<https://hal.science/hal-01455072v1>**

Submitted on 3 Feb 2017 (v1), last revised 19 Dec 2017 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin e au d p t et   la diffusion de documents scientifiques de niveau recherche, publi s ou non,  manant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv s.

# Total Domination, Connected Vertex Cover and Steiner Tree with Conflicts

Alexis Cornet

Christian Laforest

February 2, 2017

## Abstract

Total dominating set, connected vertex cover and Steiner tree are well-known graph problems. Despite the fact that they are NP-complete to optimize, it is easy (even trivial) to find solutions, regardless of their size. In this paper, we study a variant of these problems by adding conflicts, that are pairs of vertices that cannot be both in a solution. This new constraint leads to situations where it is NP-complete to decide if there exists a solution avoiding conflicts. This paper proposes NP-completeness proofs of the existence of a solution for different restricted classes of graphs and conflicts, improving recent results. We also propose polynomial time constructions in several restricted cases and we introduce a new parameter, the *stretch*, to capture the locality of the conflicts.

## 1 Introduction

In the field of discrete optimization, graphs problems have been extensively studied. An instance of a problem consists of a graph  $G$  (together with other parameters), and the goal is to compute a structure  $S$  which satisfies some constraints and whose value is optimized for a criterion. It is the case for domination problems, connected vertex cover, and Steiner tree for which approximation algorithm is a field of active research [2]. However, deciding the existence of solutions (and finding one), regardless of its size, can easily be done in polynomial time.

In the real world, there can be incompatibilities between some vertices of  $G$  because of various reasons, for example structural incompatibilities between components of a system, security reasons, mutually exclusives funding, interface incompatibilities and so on. To model these situations, we say that two elements  $u$  and  $v$  are in *conflict*, if  $u$  and  $v$  cannot be both in a solution.  $G = (V, E)$  will be called the *support graph*, and the set of conflicts will be interpreted as a graph on the same vertices, called the *conflict graph*  $C$ . No solution can contain both ends of an edge of  $C$ , i.e. any solution must be an independent set of  $C$ . Given this additional constraint, deciding the existence of a solution becomes harder. In fact, we show that deciding the existence of a solution for total dominating set with no conflicts, connected vertex cover with no conflict and Steiner tree with no conflicts is NP-complete even for restricted graph classes.

Problems with conflicts, also known as forbidden pairs, have been studied in numerous problems. In a recent serie of papers [6–8, 11, 12], authors deal with conflicts between pairs of edges, in problems such as path, spanning tree, Hamiltonian path and Hamiltonian cycle. Most of the results are NP-completeness theorems on the existence of such objects. Conflicts between vertices have also been studied until recently by several authors. Most of the works in [5, 9, 10, 14] study the complexity of finding paths without conflicts. In [4], authors prove the NP-completeness of deciding the existence of the solution in some problems with conflicts, including domination problems, connected vertex cover and Steiner tree. However, their reductions use complex graphs, and a lot of conflicts. In an ongoing work [3], we prove the NP-completeness of deciding the existence of a dominating set with no conflicts and independent dominating set with no conflicts in very restricted classes of graphs.

In this paper, in sections 2, 3 and 4, we investigate the complexity of total dominating set with no conflicts, connected vertex cover with no conflicts and Steiner tree with no conflicts for some graph classes.

Support graph $G$		Conflict graph $C$		Complexity
Class	max degree	Class	Stretch	
TOTAL DOMINATING SET				
Bipartite	4	Maximum degree 1	= 2	NPC
-	2	-	-	P
Caterpillar	3	Maximum degree 1	Finite	NPC
CONNECTED VERTEX COVER				
Bipartite	4	Maximum degree 1	= 2	NPC
Minimal degree $(1/2 - \epsilon) \cdot n$	-	-	-	NPC
Tree	-	-	Finite	P
Split	-	-	-	P
STEINER TREE				
Bipartite	4	Maximum degree 1	= 2	NPC
Tree	-	-	Finite	P
Planar bipartite	3	Maximum degree 1	-	NPC
Planar chordal	4	$\cup$ complete bipartite $\leq 4$	Finite	NPC
Split	-	Maximum degree 1	= 1	NPC
Dirac	-	$P_1, P_2$ and $P_3$	$\leq 2$	NPC
Dirac	-	Dirac	$\leq 2$	NPC

Figure 1: Our results

We aim to prove NP-completeness for the smallest possible classes, for both the support graph and the conflict graph. Therefore, we refine results of [4] by drawing a more accurate picture of the NP-completeness of the problems restricted to specific graph classes, both sparse or dense. Since we obtain NP-completeness results for restricted classes of graphs, we introduce a new parameter to capture the locality of conflicts: the *stretch*. For any conflict  $ab$ , the stretch of  $ab$  is the distance between  $a$  and  $b$  in the support graph. We consider the maximum stretch among all conflicts.

Most of the following NP-completeness proofs use reductions from a version of 3-SAT. Let us define the problem formally.

- Instance:  $(X, Cl)$  where  $X$  is a set of boolean variables and  $Cl$  a set of disjunctive 3-clauses over  $X$ .
- Question: Is there an assignment on  $X$  satisfying  $Cl$ ?

The 3 SAT problem is NP-complete, even if each variable appears in at most 4 clauses [13] (in a positive or a negative form). This result will be useful to reduce the maximum degree of support graph or conflict graph in several reductions. Moreover, one can suppose without loss of generality that each literal appears in at most 3 clauses: otherwise its negation cannot appear and the literal can be set to *true* and the variable removed from the instance together with clauses in which it appears.

In this paper we need several additional notations and graph classes. If  $G = (V, E)$  is a graph and  $X \subseteq V$  then  $G[X]$  is the graph induced by  $X$  in  $G$ . Two vertices linked by an edge are neighbors. The neighborhood of  $u$  in  $G$  is noted  $N_G(u)$  or  $N(u)$  when there is no ambiguities.  $P_n$  is a path on  $n$  vertices.

A graph of  $n$  vertices is a *Dirac graph* if each vertex has degree at least  $n/2$ . A graph is a *split graph* if it can be partitioned in an independent set and a clique. A *caterpillar* is a tree that has a dominating path. Figure 1 summarizes the results of this paper.

## 2 Total Dominating Set

Given  $(G, C)$  where  $G = (V, E)$  is the support graph and  $C$  the conflict graph, a *total dominating set with no conflicts* (TDSwnc) is a subset of vertices  $S \subseteq V$  such that:






Figure 3: Graph equivalent to the 3 SAT formula  $(\bar{a} \vee b \vee c) \wedge (a \vee \bar{b} \vee \bar{c}) \wedge (a \vee b \vee c)$




Figure 4: Gadget used to split  $K_{1,2}$ (left),  $K_{2,2}$  (middle) and  $K_{1,3}$  (right) of conflict into graphs of maximum degree 1

star, leaves are dominated by the center. Moreover, each clause is satisfied thus for each star, a leaf belongs to  $S$  and the center is dominated. Thus  $S$  is a TDSwnc of  $G$ .

Let  $S$  be a TDSwnc of  $(G, C)$ . Let  $A$  be the following assignment :  $l_i = 1$  if  $l_i \in S$ ,  $l_i = 0$  otherwise. A conflict exists for each pair  $a, \bar{a}$ , thus a literal and its negation cannot be set simultaneously to 1, hence the assignment is consistent. Moreover, for each clause  $c_i$ , there is a star which center can only be dominated by a vertex representing one of its literals, thus each clause is satisfied.

□

In the above reduction, the conflict graph is a disjoint union of small complete bipartite graphs. Next lemma presents gadgets to split these bipartites into graphs of maximum degree 1.

**Lemma 2.** *Given  $(G, C)$  a graph with conflicts where  $G$  is a disjoint union of caterpillars of maximum degree  $\delta > 1$  and  $C$  an union of complete bipartite graph of at most 4 vertices, it is possible to construct  $(G', C')$  where  $G$  is a disjoint union of caterpillars of maximum degree  $\delta$  and  $C$  a graph of maximum degree 1 such that  $(G, C)$  has a TDSwnc if and only if  $(G', C')$  has a TDSwnc.*

*Proof.* Let  $(G, C)$  a graph with conflicts where  $G$  is a disjoint union of caterpillars of maximum degree  $\delta > 1$  and  $C$  a disjoint union of complete bipartite graphs of at most 4 vertices. Split  $K_{1,2}$ ,  $K_{1,3}$  and  $K_{2,2}$  of conflicts into graphs of maximum degree 1 using gadgets from Figure 4. One can see by exhaustive search that the choice of a vertex  $a$  forbids all the vertices  $\bar{a}$  (and conversely), and that the new paths can be dominated regardless of the choice of  $a$  or  $\bar{a}$ . Only paths were created in this transformation, hence the maximum degree of the graph did not changed.

□

The conflict graph is now of maximum degree 1 but the support graph is not connected. The next result shows how to connect the disjoint caterpillars using gadgets.




Figure 5: Gadget used to connect two connected components without changing the existence of a solution.

**Lemma 3.** *Given  $(G, C)$  a graph with conflicts where  $G$  is a disjoint union of caterpillars of maximum degree  $\delta > 2$  and  $C$  a graph of maximum degree 1, it is possible to construct  $(G', C')$  where  $G'$  is a connected caterpillar of maximum degree  $\delta$  and  $C'$  a graph of maximum degree 1 such that  $(G, C)$  has a TDSwnC if and only if  $(G', C')$  has a TDSwnC.*

*Proof.* Let  $(G, C)$  be a graph with conflicts where  $G$  is a disjoint union of caterpillars of maximum degree  $\delta > 2$  and  $C$  is a graph of maximum degree 1. If  $G$  is a single caterpillar, the lemma is true. Otherwise, let  $p_1$  and  $p_2$  be two caterpillars of  $G$  and let  $(G_1, C_1)$  be the graph with conflict where  $p_1$  and  $p_2$  are connected by  $p$ , where  $p$  is the caterpillar shown in Figure 5. More specifically, connect vertices 1 and 4 of  $P$  to the extremities of the longest path of the caterpillars  $p_1$  and  $p_2$ . Thus, the graph is still an union of caterpillars. The extremities of paths are leaves, their degrees change from 1 to 2, hence the maximum degree of the graph does not change. Moreover,  $p$  can be dominated only if 1 and 4 do not belong to the TDSwnC. This can be proved by exhaustive search of the TDSwnC of the gadget. Thus it can be used to connect two caterpillars without changing the existence of a solution. Moreover,  $G_1$  has one less caterpillar than  $G$ . Repeat this transformation until there is only one caterpillar. □

**Theorem 2.** *Given  $(G, C)$  a graph with conflicts, deciding whether there exists a TDSwnC is NP-complete, even if  $G$  is a caterpillar of maximum degree 3 and  $C$  a graph of maximum degree 1.*

*Proof.* Theorem 2 follows from lemmas 1, 2 and 3. □

Previous results proved that TDSwnC is NP-complete in caterpillar of maximum degree 3. We know prove that this result is in some way the strongest possible, since the problem becomes polynomial when the maximum degree is 2.

**Theorem 3.** *If  $G$  is a graph of maximum degree 2 then deciding whether  $(G, C)$  has a TDSwnC can be done in polynomial time.*

*Proof.* We reduce the problem to 2 SAT for which polynomial algorithms are known (see [1]). Let  $(G = (V, E), C)$  be an instance of TDSwnC where  $G$  has maximum degree 2. Suppose without loss of generality that there is no isolated vertex (otherwise the graph has no TDSwnC). Construct  $(X, Cl)$  the following SAT instance:  $X = V$  and  $Cl = \bigwedge_{x \in V} (\bigvee_{y \in N(x)} y) \bigwedge_{ab \in C} (\bar{a} \vee \bar{b})$ . Each vertex has at most 2 neighbors hence we obtain a 2 SAT instance.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Then  $S = \{x | x = 1\}$  is without conflict in  $C$  since for each conflict  $ab$  the clause  $\bar{a} \vee \bar{b}$  exists. Moreover, for each  $x \in V$ , the clause  $\bigvee_{y \in N(x)} y$  exists, hence  $x$  is dominated by one of its neighbors. Thus  $S$  is a TDSwnC.

Let  $S$  be a TDSwnC of  $(G, C)$ . Construct  $A$  an assignment on  $X$  as follows:  $x$  is set to 1 if  $x \in S$ ,  $x$  is set to 0 otherwise. For each clause  $c$ , there exists a vertex of  $V$  which can only be dominated by vertices representing literals of the clause. One of these vertices belongs to  $S$ , thus the clause is satisfied. For each clause  $\bar{a} \vee \bar{b}$ , there exists a conflict  $ab$ , thus at most one vertex belongs to  $S$ , hence the clause is satisfied. Thus  $A$  satisfies  $Cl$ . □




Figure 6: Graph equivalent to the 3 SAT formula  $(a \vee \bar{b} \vee c) \wedge (a \vee b \vee \bar{d})$

### 3 Connected Vertex Cover

Given  $(G, C)$  where  $G = (V, E)$ , a *connected vertex cover with no conflicts* (CVCwnC) is a subset of vertices  $S \subseteq V$  such that:

- for each  $xy \in E$ ,  $x \in S$  or  $y \in S$
- for each  $xy \in C$ ,  $x \notin S$  or  $y \notin S$
- $G[S]$  is connected.

**Theorem 4.** *Given  $(G, C)$  a graph with conflicts, deciding whether there exists a CVCwnC is NP-complete even if  $G$  is a bipartite graph of maximum degree 4 and  $C$  is a graph of maximum degree 1 of stretch 2.*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance where each variable is in at most 4 clauses. Construct  $(G, C)$  an instance of CVCwnC as follows: for each variable  $\alpha$ , vertices  $\alpha, \bar{\alpha}$  and  $r_\alpha$  are created, along with the edges  $\alpha r_\alpha$  and  $\bar{\alpha} r_\alpha$  and the conflict  $\alpha \bar{\alpha}$ . Vertices  $r_\alpha$  are connected by intermediate vertices  $r_i$ . For each clause  $c_i = (a \vee b \vee c)$ , vertices  $c_i$  and  $c'_i$  are created, along with the edges  $c_i c'_i, c_i a, c_i b, c_i c$ . Thus  $G$  is bipartite of maximum degree 4 and conflicts are a graph of maximum degree 1 of stretch 2. An example is shown at Figure 6.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Construct  $S$  as follows:  $S$  contains all vertices corresponding to positive literals of  $A$ . Vertices  $r_\alpha$  and  $r_i$  and  $c_i$  are also added to  $S$ . Thus  $S$  is a Vertex Cover. Moreover, it is without conflict since a literal and its negation can not be set to 1 simultaneously. Vertices  $r_\alpha, r_i$  and vertices corresponding to literals are connected, and for each clause  $c_i = (a \vee b \vee c)$ ,  $a, b$  or  $c$  belong to  $S$ , thus  $c_i$  is connected.

Let  $S$  be a CVCwnC of  $(G, C)$ . Literals corresponding to vertices of  $S$  are set to 1, other variables are set to 0. Since conflicts exist between each literal and its negation, we obtain an assignment. Moreover, vertices  $c_i$  necessarily belong to  $S$ , and each needs to be connected by a vertex corresponding to a literal of the associated clause. Thus, each clause is satisfied. □

Let us point out two graph classes in which deciding the existence of a CVCwnC is polynomial.

**Remark 1.** *Deciding the existence of CVCwnC is polynomial in trees. There exists a unique Connected Vertex Cover minimal for inclusion, the set of internal vertices of the tree: it is sufficient to test if it is without conflict.*

**Theorem 5.** *Given  $(G, C)$  a graph with conflicts, deciding whether there exists a CVCwnC is polynomial if  $G$  is a split graph.*

*Proof.* Let  $G = (V, E)$  be a split graph where  $V = K \cup I$  where  $K$  is a clique and  $I$  is an independent set and  $C$  the conflict graph.

- If  $K$  is without conflicts, then  $K$  is a CVCwnC of  $(G, C)$ .




Figure 7: Construction of the dense graph

- If  $C[K]$  is a star of center  $a$ , then if  $S = (K - a) \cup N_G(a)$  is without conflicts,  $S$  is a CVCwnC. (Two possibilities for the center  $a$  if the star is an edge). If  $(K - a) \cup N_G(a)$  has a conflict, then  $(G, C)$  does not have a CVCwnC.

- If  $C[K]$  is not a star, then at least 2 vertices  $a$  and  $b$  of  $K$  cannot be in the solution: the edge  $ab$  will not be covered and thus there is no CVCwnC.  $\square$

In Theorem 4 we proved NP-completeness for bipartite graphs. We are now interested in dense graphs, and prove NP-completeness for dense graphs.

**Theorem 6.** *For all  $\epsilon > 0$ , given  $(G, C)$  a graph with conflicts, deciding whether there exist a CVCwnC is NP-complete even if  $G$  is of minimum degree  $(1/2 - \epsilon) \cdot n$ .*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance of  $m$  clauses over  $n$  variables. Construct  $(G, C)$  an instance of CVCwnC as follows. Let  $d$  be the least integer greater than  $(m + 2n)/2\epsilon$ . For each variable  $x \in X$  create the vertices  $x, \bar{x}$ . For each clause  $c_i$  create vertex  $c_i$ . Let  $G_X$  be the set of vertices representing literals.  $G_X$  has  $2n$  vertices. Let  $G_{Cl}$  be the set of  $m$  vertices representing clauses. Construct  $G_K$  a clique of size  $d$  and  $G_I$  an independent set of size  $d$ . Complete bipartite are created between  $G_K$  and  $G_X$ ,  $G_{Cl}$  and  $G_I$ ,  $G_K$  and  $G_I$ . For each clause  $c_i = (a \vee b \vee d)$  edges  $c_i a, c_i b, c_i d$  are created. Construct  $C$  as follow. Each pair  $\{x, \bar{x}\}$  is in conflict. Each vertex of  $G_I$  is in conflict with all the other vertices of  $G$ . We present a scheme of the support graph used in the reduction at Figure 7. To ensure readability, conflicts are not represented.

$G$  has  $m + 2n + 2d$  vertices and  $d \geq (m + 2n)/2\epsilon$ . Because of the complete bipartite graphs, vertices of  $G_X, G_{Cl}, G_I$  and  $G_K$  has degree at least  $d$ . It can be derived by basic arithmetic on inequalities that  $d > (1/2 - \epsilon)(m + 2n + 2d)$ , hence the graph satisfies the assumption of the theorem.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ , and let  $A_1$  be the set of positive literals of  $A$ . Set  $VC = A_1 \cup G_K \cup G_{Cl}$ .  $VC$  does not contain simultaneously vertices representing a literal and its negation, neither vertices of  $G_i$ , thus it is a set without conflicts in  $C$ . Edges between  $G_K$  and  $G_X$ ,  $G_{Cl}$  and  $G_I$ ,  $G_K$  and  $G_I$ ,  $G_X$  and  $G_{Cl}$  are covered. Thus  $VC$  is a vertex cover of  $G$ . Vertices of  $G_K$  and  $G_X$  are connected. Moreover, for each vertex  $c_i \in Cl$  there exists a vertex  $x$  of  $A_1$  corresponding to a positive literal of  $c_i$ . Thus  $VC$  is connected.

Let  $VC$  be a CVCwnC of  $(G, C)$ . Let  $A$  be the following assignment: for each literal  $x$ , if  $x$  is represented by a vertex of  $VC$ , then  $x = 1$ , otherwise,  $x = 0$ .  $VC$  is not reduced to a single vertex, hence  $VC$  cannot contain vertices of  $G_I$  because of conflicts between  $G_I$  and all the vertices.  $VC$  must contain the




Figure 8: Graph equivalent to the 3 SAT formula  $(a \vee \bar{b} \vee c) \wedge (a \vee b \vee \bar{d})$ . Vertices of  $M$  are squares.

neighborhood of  $G_I$ :  $G_{Cl} \cup G_K$ . Moreover, each vertex of  $G_{Cl}$  must be connected to  $G_K$ . It can only be done via vertices of  $G_X$ . For each vertex representing a clause  $(a \vee b \vee c)$  there exist a vertex representing one of its literals in  $VC$ . Hence,  $A$  satisfies  $Cl$ . Moreover, a vertex representing a variable and a vertex representing its negation can not be set to 1 simultaneously, thus the assignment is consistent.  $\square$

## 4 Steiner tree

A Steiner tree of  $(G = (V, E), M)$  where  $M \subseteq V$  is a subtree of  $G$  that includes all the vertices of  $M$ .

Given  $(G, M, C)$  where  $G = (V, E)$  and  $M \subseteq V$ , a *Steiner Tree without conflict* (STwnc) is a subset of vertices  $S \subseteq V$  such that :

- for each  $xy \in C$ ,  $x \notin S$  or  $y \notin S$
- $M \subseteq S$
- $G[S]$  is connected.

If  $G[S]$  is connected and without conflict, it is easy to extract a covering tree of  $S$ , hence we are interested only in the set of vertices of the tree.

First, we prove NP-completeness when the conflicts are local.

**Theorem 7.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exists a STwnc is NP-complete even if  $G$  is a bipartite graph of maximum degree 4 and  $C$  a graph of maximum degree 1 and of stretch 2.*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance where each variable is in at most 4 clauses. Suppose without loss of generality that each literal is in at most 3 clauses. Construct  $(G, M, C)$  an instance of STwnc as follows: for each variable  $\alpha$  vertices  $\alpha, \bar{\alpha}$  and  $r_\alpha$  are created, along with the edges  $\alpha r_\alpha$  and  $r_\alpha \bar{\alpha}$  and the conflict  $\alpha \bar{\alpha}$ . Vertices  $r_\alpha$  are connected by intermediate vertices  $m_i$ . For each clause  $c_i = (a \vee b \vee c)$ , a vertex  $c_i$  is created along with the edges  $c_i a, c_i b, c_i c$ . The graph is of maximum degree 5, and the conflict graph is a graph of maximum degree 1 and of stretch 2. The set  $M$  is composed of all the vertices  $m_i$  and  $c_i$ . An example is shown at Figure 8.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Set  $S = M \cup_\alpha \{r_\alpha\} \cup P$  where  $P$  is the set of vertices representing positive literals of  $A$ . Since a variable cannot be set to one simultaneously with its negation,  $S$  is without conflicts. Vertices  $m_i$  and  $r_\alpha$  are connected. Moreover, for each vertex  $c_i$ , there exists a clause  $c_i$  in which one of the literals is set to 1 in  $A$ , thus  $c_i$  is connected to the other vertices of  $S$ . Thus,  $S$  is connected in  $G$ .

Let  $S$  be a STwnc of  $(G, M, C)$ . The value of literals of  $X$  which vertices are in  $S$  is set to 1. Since conflicts exist between variables and their negation, we obtain an assignment. Moreover, for each vertex  $c_i$  representing a clause  $a \vee b \vee c$ , one of the vertices  $a, b, c$  must belong to  $S$ , hence the clause is satisfied.  $\square$




Figure 9: Graph equivalent to the 3 SAT formula  $(a \vee b \vee c) \wedge (\bar{a} \vee b \vee \bar{c}) \wedge (\bar{a} \vee b \vee \bar{c}) \wedge (\bar{a} \vee b \vee c)$ . Vertices of  $M$  are squares. In the black rectangle, gadget corresponding to the clause  $(\bar{a} \vee b \vee \bar{c})$ .

The next theorems are NP-completeness results for more restricted class of support graph, but the conflicts are no longer local (i.e. stretch is not bounded).

**Theorem 8.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exist a  $STwnc$  is NP-complete even if  $G$  is a planar bipartite graph of maximum degree 3 and  $C$  a graph of maximum degree 1.*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance where each variable is in at most 4 clauses. Construct  $(G, M, C)$  an instance of  $STwnc$  where  $G$  is a planar bipartite graph of maximum degree 3 and  $C$  a graph of maximum degree 1 such that there exists an assignment on  $X$  satisfying  $Cl$  if and only if there exist a  $STwnc$  of  $(G, M, C)$ .

For each clause, create a 15 vertices gadget composed of 3 non-disjoint paths having the same extremities, each corresponding to a literal. One can see this gadget in Figure 9. These gadgets are connected linearly, and the set  $M$ , composed of only two vertices, is the first vertex of the first gadget and the last vertex of the last gadget. For each pair of literals  $(x, \bar{x})$ , add a conflict between a vertex with no conflict of the path representing  $x$  and a vertex with no conflict of the path representing  $\bar{x}$ . The graph is planar bipartite of maximum degree 3 and the conflict graph is of maximum degree 1.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Construct  $S$  a subset of  $G$  as follows. For each gadget  $g$  representing a clause  $(a \vee b \vee c)$ , choose in  $S$  a path representing a positive literal. Each gadget is passed through, hence vertices of  $M$  are connected. The set  $S$  is the path of vertices connecting the two vertices of  $M$ . Moreover, this path does not pass through path representing variable and their negation, hence it is without conflicts.

Let  $S$  be a  $STwnc$  of  $(G, M, C)$ . Suppose without loss of generality that  $S$  is minimal for inclusion (otherwise, it can be minimalized). Construct  $A$  an assignment on  $X$  satisfying  $Cl$  as follows: for each gadget representing a clause  $(a \vee b \vee c)$ ,  $S$  contains a path representing one of its literals. This one is set to 1 in  $A$ . Since there exist conflicts between paths representing variable and their negation, the assignment  $A$  is consistent. Moreover, for each clause, a literal is set to 1 hence  $A$  satisfies  $Cl$ . □

**Theorem 9.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exists a  $STwnc$  is NP-complete even if  $G$  is planar chordal graph of maximum degree 4 and  $C$  a disjoint union of complete bipartite graphs of at most 4 vertices.*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance where each variable is in at most 4 clauses. Construct  $(G, M, C)$  the following instance of  $STwnc$ . For each clause  $c_i = (a \vee b \vee c)$ , vertices  $c_i, c'_i, a, b, c, m_i$  and edges  $c'_i c_i, c_i a, c_i b, c_i c, ab, ac, am_i, bm_i, bc, cm_i$  are created. The set  $M$  is the set of all vertices  $m_i$  and  $c'_i$ . Vertices  $c'_i$  are connected to form a path. For each pair of vertices  $\{\alpha, \bar{\alpha}\}$  representing a literal and its negation, the conflict  $\alpha \bar{\alpha}$  is created. The graph is chordal planar of maximum degree 4, and the conflict graph is an union of complete bipartite graphs of at most 4 vertices. An example is shown at Figure 10.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Construct  $S = M \cup_i \{c_i\} \cup P$  where  $P$  is the set of vertices representing positive literals of  $A$ . Then  $\cup_i \{c_i\} \cup_i \{c'_i\}$  is connected. Moreover, for each clause  $c_i$ , a vertex  $x$  corresponding to a positive literal belongs to  $S$ . Hence, the vertex  $m_i$  is connected and  $S$  is connected in




Figure 10: Graph equivalent to the 3 SAT formula  $(a \vee b \vee c) \wedge (a \vee \bar{b} \vee \bar{c}) \wedge (\bar{a} \vee b \vee c)$ . Vertices of  $M$  are squares.

$G$ . Moreover, conflicts are only between variables and their negation, which can not be simultaneously set to 1 in  $A$ . Thus,  $S$  is without conflicts.

Let  $S$  be a STwnc of  $(G, M, C)$ . Literals corresponding to vertices of  $S$  are set to 1, the other variables are set to 0. Since there exist conflicts between vertices representing variables and their negation, we obtain an assignment. Moreover, each vertex  $m_i$  must be connected to the other vertices of  $S$  by a vertex representing a literal of the associated clause, which is satisfied.  $\square$

**Remark 2.** *Deciding the existence of a STwnc is polynomial in trees. There exists an unique Steiner tree minimal for inclusion, hence it is sufficient to test if it is without conflicts.*

**Theorem 10.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exists a STwnc is NP-complete even if  $G$  is a split graph and  $C$  a graph of maximum degree 1 and of stretch 1.*

*Proof.* Let  $(X, Cl)$  be a 3 SAT instance. Suppose without loss of generality that  $Cl$  contain several clauses. Create  $(G = (K, I, E), M, C)$  an instance of STwnc as follows. Each literal becomes a vertex of  $K$  and each clause a vertex of  $I$ . Conflicts are added between variable and their negations, and edges between clauses and their literals. The graph is a split graph, and the conflict graph is a graph of maximum degree 1 and of stretch 1. Set  $M = I$ . An example is shown at Figure 11.

Let  $A$  be an assignment on  $X$  satisfying  $Cl$ . Construct  $S = I \cup P$  where  $P$  is the set of vertices corresponding to positive literals of  $A$ . Since a literal and its negation cannot both be set to 1,  $S$  is without conflicts. Moreover, for each  $c_i \in I$ , the associated clause is satisfied, thus there exists a neighbor in  $K \cap S$ . Hence, each vertex of  $I$  is connected to  $K \cap S$  and since  $K$  is a clique,  $S$  is connected.

Let  $S$  be a STwnc of  $(G = (K, I, E), M, C)$ . The value of literals corresponding to vertices of  $S$  is set to 1, and the value 0 is set to other variables. Since there exist conflict between vertices representing literals and vertices representing their negation, we have an assignment. Moreover, each vertex  $c_i$  must be connected to  $K$  by a vertex representing a literal of the associated clause, thus the clause is satisfied.  $\square$

Previous theorems aimed to prove NP-completeness for sparse support and conflict graphs. We now prove that the problem remains NP-complete in a class of dense graphs: Dirac graphs (which are graphs of minimum degree  $n/2$ ).

**Theorem 11.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exists a STwnc is NP-complete even if  $G$  is a Dirac graph and  $C$  a disjoint union of  $P_1, P_2$  and  $P_3$ .*




Figure 11: Graph equivalent to the 3 SAT formula  $(a \vee \bar{c} \vee b) \wedge (\bar{a} \vee b \vee \bar{c})$ . Vertices of  $M$  are squares.

*Proof.* Let  $(G_1 = (V_1, E_1), M_1, C_1)$  be an instance of STwnc where  $C_1$  is a graph of maximum degree 1. This problem is NP-complete from Theorem 7. Construct  $(G_2 = (V_2, E_2), M_2, C_2)$  a graph with conflicts such that there exists a STwnc of  $(G_2, M_2, C_2)$  if and only if there exists a STwnc of  $(G_1, M_1, C_1)$ . Let  $n = |V_1|$ . Set  $V_2 = V_1 \cup A \cup B$  where  $A$  is a path of length  $n$  and  $B$  an independent set of size  $2n$ .  $M_2 = M_1 \cup A$ .  $A$  is connected to an arbitrary vertex of  $M_1$ . A complete bipartite graph is added between  $B$  and  $V_2 - B$ . Conflicts of  $C_1$  are added to  $C_2$ . Moreover, each vertex of  $A$  is in conflict with 2 distinct vertices of  $B$ . By construction,  $G_2$  is a Dirac graph and  $C_2$  a disjoint union of  $P_1, P_2$  and  $P_3$ .

Let  $T_1$  be a STwnc of  $G_1$ . Then  $T_2 = T_1 \cup A$  is a STwnc of  $G_2$ .

Suppose there exists  $T_2$  a STwnc of  $(G_2, M_2, C_2)$ . Then  $T_2 \cap B = \emptyset$ . Moreover, vertices of  $A$  do not connect vertices of  $V_1$ . Hence,  $T_1 = T_2 - A$  is a STwnc of  $(G_1, M_1, C_1)$ .  $\square$

**Theorem 12.** *Given  $(G, M, C)$  a graph with conflicts and a subset of vertices, deciding whether there exist a STwnc is NP-complete even if  $G$  is a Dirac and  $C$  a Dirac graph.*

*Proof.* Let  $(G_1 = (V_1, E_1), M_1, C_1)$  be an instance of STwnc. Construct  $(G_2 = (V_2, E_2), M_2, C_2)$  a Dirac graph with conflicts such that there exists a STwnc of  $(G_2, M_2, C_2)$  if and only if there exists a STwnc of  $(G_1, M_1, C_1)$ . Let  $n = |V_1|$ . Set  $V_2 = V_1 \cup A$  where  $A$  is an independent set of size  $n$ .  $M_2 = M_1$ . A complete bipartite graph is created between  $V_1$  and  $A$ . Conflicts of  $C_1$  exist in  $C_2$ . Moreover, each vertex of  $A$  is in conflict with all vertices of  $V_1$ . By construction,  $G_2$  and  $C_2$  are Dirac graphs.

Let  $T$  be a STwnc of  $(G_1, M_1, C_1)$ . Thus it is a STwnc of  $G_2$ .

Suppose that there exist  $T$  a STwnc of  $(G_2, M_2, C_2)$ . Then  $T \cap A = \emptyset$ . Hence,  $T$  is a STwnc of  $(G_1, M_1, C_1)$ .  $\square$

## 5 Conclusion

Our paper strengthens the results of [4] for CVCwnc and STwnc, by proving NP-completeness in smaller graph classes. More exactly, we proved that these problems remain NP-complete for some classes of sparse graphs. For STwnc, we proved that the problem is also NP-complete in dense graphs. We also extended the notion of conflicts to a new problem: TDSwnc and proved its NP-completeness in a very sparse graph class: caterpillars of maximum degree 3. Moreover, we proved that this result is in some way the strongest possible since the problem becomes polynomial in graph of maximum degree 2. Furthermore, we proved that all these problems are NP-complete even when the stretch of the conflicts is two at most, i.e. when the conflicts remain local.

A natural extension of this work would be to work on the complexity of CVCwnc and TDSwnc in dense graphs, and the complexity of CVCwnc in other classes of sparse graphs, for example planar graphs. It would also be interesting to study other graphs problems with conflicts.

## References

- [1] Bengt Aspvall, Michael F Plass, and Robert Endre Tarjan. A linear-time algorithm for testing the truth of certain quantified boolean formulas. *Information Processing Letters*, 8(3):121–123, 1979.
- [2] Giorgio Ausiello, Pierluigi Crescenzi, Giorgio Gambosi, Viggo Kann, Alberto Marchetti-Spaccamela, and Marco Protasi. *Complexity and approximation: Combinatorial optimization problems and their approximability properties*. Springer Science & Business Media, 2012.
- [3] Alexis Cornet and Christian Laforest. Domination problems with conflicts. *writing in progress*.
- [4] Francois Delbot, Christian Laforest, and Raksmeay Phan. Hardness Results and Approximation Algorithms for Discrete Optimization Problems with Conditional and Unconditional Forbidden Vertices. Research report, HAL, <https://hal.archives-ouvertes.fr/hal-01257820>, January 2016.
- [5] Harold N Gabow, Shachindra N Maheshwari, and Leon J Osterweil. On two problems in the generation of program test paths. *Software Engineering, IEEE Transactions on*, (3):227–231, 1976.
- [6] Mamadou Moustapha Kanté, Christian Laforest, and Benjamin Momège. An exact algorithm to check the existence of (elementary) paths and a generalisation of the cut problem in graphs with forbidden transitions. In *SOFSEM 2013*, pages 257–267, 2013.
- [7] Mamadou Moustapha Kanté, Christian Laforest, and Benjamin Momège. Trees in graphs with conflict edges or forbidden transitions. In *TAMC 2013, Hong Kong*, pages 343–354, 2013.
- [8] Mamadou Moustapha Kanté, Fatima Zahra Moataz, Benjamin Momège, and Nicolas Nisse. Finding paths in grids with forbidden transitions. In *WG 2015*, pages 154–168, 2015.
- [9] Petr Kolman and Ondřej Pangrác. On the complexity of paths avoiding forbidden pairs. *Discrete Applied Mathematics*, 157(13):2871–2876, 2009.
- [10] Jakub Kováč. Complexity of the path avoiding forbidden pairs problem revisited. *Discrete Applied Mathematics*, 161(10):1506–1512, 2013.
- [11] Christian Laforest and Benjamin Momège. Some hamiltonian properties of one-conflict graphs. In *IWOCA 2014*, pages 262–273, 2014.
- [12] Christian Laforest and Benjamin Momège. Nash-williams-type and chvátal-type conditions in one-conflict graphs. In *SOFSEM 2015*, pages 327–338, 2015.
- [13] Craig A. Tovey. A simplified np-complete satisfiability problem. *Discrete Applied Mathematics*, 8(1):85–89, 1984.
- [14] Hananya Yinnone. On paths avoiding forbidden pairs of vertices in a graph. *Discrete applied mathematics*, 74(1):85–92, 1997.