

Editorial: Perceptual Grouping-The State of The Art

Birgitta Dresp, Stephen Grossberg, Adam Reeves

▶ To cite this version:

Birgitta Dresp, Stephen Grossberg, Adam Reeves. Editorial: Perceptual Grouping-The State of The Art. Frontiers in Psychology, 2017, 10.3389/fpsyg.2017.00067. hal-01455558

HAL Id: hal-0145558

https://hal.science/hal-01455558

Submitted on 10 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

6

9

14

24

26

Editorial: Perceptual Grouping - The State Of The Art

2 Birgitta Dresp-Langley, Stephen Grossberg, Adam Reeves

3 ICube UMR 7357 CNRS, Strasbourg University, Strasbourg, FRANCE

Perceptual neuroscience has identified mechanisms of perceptual grouping which account for the 4

ways in which visual sensitivity to ordered structure and regularities expresses itself, in behavior and 5

in the brain. The need to actively construct order, notably representations of objects in depth, is

mandated as soon as visual signals reach the retina, given the occlusion of retinal signals by retinal 7

veins and other retinal elements or blur. Multiple stages of neural processing transform fragmented 8

signals into visual key representations of 3D scenes that can be used to control effective behaviors.

Since our survival depends on our abilities to pick up order in the physical world, and since we 10

conceive the physical world as an ordered one, our perception of it must somehow be sensitive to 11

order. The articles from this collection illustrate how texture dissimilarity, boundary completion, 12

surface filling-in, and figure-ground segregation generate perceptual order. Structural regularities are 13

required for a visual texture to be discriminated from random noise by human observers, as shown in

https://doi.org/10.3389/fpsyg.2015.00734, where Katkov, Harris and Sagi compare the performance

15 of human observers to those of an Ideal Observer and an Order Observer model. The Order Observer 16

model, based on the capture of statistical regularities in textures, is shown to closely reflect the 17

sensitivity of the human visual system. Spatiotemporal boundary formation is critical to the 18

perception of shape boundaries and global motion https://doi.org/10.3389/fpsyg.2016.00910, as 19

shown here by Erlikhman and Kellman, who demonstrate boundary integration from spatially and 20

temporally sparse transformations of texture. The visual system uses positions and times of element 21

22 transformations to extract local structural regularities such as edge fragments, which then connect by

interpolation processes to produce larger shapes with a spatiotemporal integration constraint of ~80 23

ms between elements. Structural ambiguity and uncertainty pose a specific problem here. Perceptual

25 grouping not only accounts for the ways in which physically defined order is processed by the visual

system, but also how such order may be inferred in cases of uncertainty, and how this may give rise

to the perception of "illusory" or "phantom" shapes. Roncato, Guidi, Parlangeli and Battaglini 27

discuss a new phenomenon https://doi.org/10.3389/fpsyg.2016.00959 in which illusory streaks 28

irradiate from the vertexes of angular contours, connecting pairs of figures and forming phantom net 29

shapes that are only clearly visible in a specific figural context, as discussed here in the light of 30

several theories of object contour detection. The real world conditions or configurative order to 31

which *perceptual grouping* is sensitive also determine our aesthetic perception, as shown already 32

elsewhere in a large body of previous work on aesthetic preferences for structural regularities. Here, 33

Pinna, Porcheddu and Deiana https://doi.org/10.3389/fpsyg.2016.01051 discuss a novel variant of 34 perceptual grouping called accentuation for perceptual coupling. New conditions for grouping that 35

cannot be explained in terms of the classical Gestalt principles (similarity etc) are shown. This new 36

work reveals the supremacy of dissimilarity as an important figural complement to similarity in the 37

genesis of aesthetic structure, involving novel dynamics of perceptual grouping they call *coupling*. 38

The visual system organizes incoming information early on and interprets a 2D image in terms of the 39

real world, producing configurative order that provides perceptual continuity and enables object-

40

based attention. In https://doi.org/10.3389/fpsyg.2015.02054 Grossberg reviews the Form-And-41

Color-And-DEpth (FACADE) model and its laminar cortical embodiment and extension by the 3D 42

LAMINART models. These have explained, simulated, and predicted many perceptual and 43

44 neurobiological data about how the visual cortex carries out 3D vision and figure-ground perception.

and how these cortical mechanisms generate 3D percepts from 2D pictures of occluding and 45

occluded objects. The models provide a unified explanation of neurophysiological data about cortical 46

area V2 cells reported in a remarkable series of articles from the von der Heydt laboratory. Properties 47

48 of border ownership, contrast preference, binocular stereoscopic information, selectivity to side-offigure, Gestalt rules, strength of their modulation by attention and temporal properties are explained. 49 The segregation of image parts into foreground and background requires border ownership 50 assignment by the visual system and recordings from monkey visual cortex show that many neurons, 51 especially in area V2, are selective to border ownership. These neurons are edge selective and have 52 classical receptive fields (CRF). Their responses are modulated (enhanced or suppressed) depending 53 54 on the location of a 'figure' relative to the edge in their receptive field. Each neuron has a fixed preference for location on one side or the other. This selectivity is derived from image context far 55 beyond the CRF. In https://doi.org/10.3389/fpsyg.2015.01695 von der Heydt reviews evidence 56 57 indicating that border ownership selectivity reflects the formation of early object representations, which he terms 'proto-objects'. The evidence includes experiments showing (1) reversal of border 58 ownership signals with change of perceived object structure, (2) border ownership specific 59 enhancement of responses in object-based selective attention, (3) persistence of border ownership 60 signals in accordance with continuity of object perception, and (4) remapping of border ownership 61 signals across saccades and object movements. Grouping, persistence, and remapping are shown to 62 be manifest in low-level visual areas, which challenges some of the traditional views and top-down-63 only models of *perceptual grouping*. To generate perceptual judgments relative to figure and ground, 64 polarity-invariant and polarity-specific object properties are used by the visual brain. Dresp-Langley 65 and Grossberg https://doi.org/10.3389/fpsyg.2016.01102 demonstrate conditions under which figure-66 ground is determined by the orientation of contrasts, not by their relative sign (Dresp, 1998; Dresp & 67 Fischer, 2001; Dresp & Langley, 2005). Psychophysical data from human observers responding to 68 69 structurally ambiguous, fragmented 2D shapes with varying local contrast signs reveal polarityinvariant perceptual grouping predicted by the earliest version of the FACADE theory and, more 70 recently, by the 3D LAMINART models. Research into processes of perceptual grouping has 71 important societal and technological aspects, for visual interface design and, in particular, image-72 guided surgery platforms and simulators. Such implications, brought forward here by Dresp-Langley 73 in https://doi.org/10.3389/fpsyg.2015.01565 illustrate why boundary completion and figure-ground 74 segregation are not only essential for realistic scene perception, but also for visually guided action 75 76 planning (Bonnet & Dresp, 2001), visual consciousness (Dresp-Langley & Durup, 2009; 2012) and 77 behavior.

References

78

82

83

86 87

88

89

- 79 Dresp, B. (1998). The effect of practice on the visual detection of near-threshold lines. *Spatial Vision*, **11**, 1-13.
- 80 Bonnet, C., & Dresp, B. (2001). Investigations of sensory magnitude and perceptual processing with reaction times. 81 *Psychologica*, **25**, 63-86.
 - Dresp, B., & Fischer, S. (2001). Asymmetrical contrast effects induced by luminance and colour configurations. *Perception & Psychophysics*, **63**, 1262-1270.
- Dresp, B., & Langley, O. K. (2005). Long-range spatial integration across contrast signs: a probabilistic mechanism? *Vision Research*, **45**, 275-284.
 - Dresp-Langley, B, & Durup, J (2009) A plastic temporal code for conscious state generation in the brain. *Neural Plasticity*, 1-15. http://dx.doi.org/10.1155/2009/482696
 - Dresp-Langley, B, & Durup, J (2012) Does consciousness exist independently of present time and present time independently of consciousness? *Open Journal of Philosophy*, 2, 45-49.
- 90 <u>http://dx.doi.org/10.4236/ojpp.2012.21007</u>