



HAL
open science

Selecting on fat and protein content instead of milk yield would not improve dairy cows' fertility

Nicolas Bedere, Catherine Disenhaus, Vincent Ducrocq, Ségolène Colette, Luc Delaby

► To cite this version:

Nicolas Bedere, Catherine Disenhaus, Vincent Ducrocq, Ségolène Colette, Luc Delaby. Selecting on fat and protein content instead of milk yield would not improve dairy cows' fertility. 67. Annual Meeting of the European Federation of Animal Science (EAAP), Aug 2016, Belfast, United Kingdom. Annual Meeting of the European Association for Animal Production, 22, 2016, Annual Meeting of the European Association for Animal Production. hal-01455921

HAL Id: hal-01455921

<https://hal.science/hal-01455921>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Selecting on fat and protein content instead of milk yield would not improve dairy cows' fertility

N. Bedere¹, C. Disenhaus¹, V. Ducrocq², S. Leurent-Colette³ and L. Delaby¹

¹INRA, AGROCAMPUS OUEST, PEGASE, Saint-Gilles, 35590, France, ²INRA, AgroParisTech, Université Paris-Saclay, GABI, Jouy-en-Josas, 78352, France, ³INRA, Domaine Experimental du Pin-au-Haras, Exmes, 61310, France; nicolas.bedere@rennes.inra.fr

To compare reproductive performances of cows with high genetic merit for fat and protein content vs high genetic merit for milk yield, an experiment was conducted on 500 cows at the INRA farm of Le Pin-au-Haras. The herd was composed of Holstein and Normande cows. In each breed, cows with high genetic merit for milk yield were classified in a 'Milk Group' (MG) and those with high genetic merit for fat and protein content were classified in a 'Content Group' (CG) with equivalent milk solids production over the lactation. Two grazing-based feeding systems were used, a 'High' (maize silage / grazing plus concentrate) and a 'Low' (grass silage / grazing with no concentrate) system. To study the different steps of the reproductive process, milk progesterone information (3/week), oestrous behaviour recording and pregnancy diagnosis (ultrasonography) were combined. Within breeds, cows in the CG had an earlier resumption of luteal activity than cows in the MG (-6 d, $P=0.01$). Occurrence of prolonged luteal phases was not related to genetic groups. Both genetic groups had similar ovulation detection rate and proportion of standing behaviour. No difference in fertility was found between genetic groups in Normande cows. Holstein cows in the CG showed more non-fertilization/early embryo mortality in the Low system (+31 points, $P=0.002$) and more late embryo mortality/foetal death in the High system (+18 points, $P=0.06$). The MG was exporting more energy in early lactation because of a higher milk yield peak and the CG was exporting more energy in mid-lactation because of a better persistency. Trade-off between milk production and reproduction was not found: cows in the MG had a degraded cyclicity and those in the CG had a degraded fertility. Selecting dairy cows on fat and protein content may not be a promising way to maintain good productive and reproductive performances.