



HAL
open science

A correction of a characterization of planar partial cubes

Rémi Desgranges, Kolja Knauer

► **To cite this version:**

Rémi Desgranges, Kolja Knauer. A correction of a characterization of planar partial cubes. *Discrete Mathematics*, 2017. hal-01457179

HAL Id: hal-01457179

<https://hal.science/hal-01457179>

Submitted on 6 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A correction of a characterization of planar partial cubes

Rémi Desgranges* Kolja Knauer†

January 7, 2017

Abstract

In this note we determine the set of expansions such that a partial cube is planar if and only if it arises by a sequence of such expansions from a single vertex. This corrects a result of Peterin.

1 Introduction

A graph is a *partial cube* if it is isomorphic to an isometric subgraph G of a hypercube graph Q_d , i.e., $\text{dist}_G(v, w) = \text{dist}_{Q_n}(v, w)$ for all $v, w \in G$. Any isometric embedding of a partial cube into a hypercube leads to the same partition of edges into so-called Θ -classes, where two edges are equivalent, if they correspond to a change in the same coordinate of the hypercube. This can be shown using the Djoković-Winkler-relation Θ which is defined in the graph without reference to an embedding, see [5, 6].

Let G^1 and G^2 be two isometric subgraphs of a graph G that (edge-)cover G and such that their intersection $G' := G^1 \cap G^2$ is non-empty. The *expansion* H of G with respect to G^1 and G^2 is obtained by considering G^1 and G^2 as two disjoint graphs and connecting them by a matching between corresponding vertices in the two resulting copies of G' . A result of Chepoi [3] says that a graph is a partial cube if and only if it can be obtained from a single vertex by a sequence of expansions. An equivalence class of edges with respect to Θ in a partial cube is an inclusion minimal edge cut. The inverse operation of an expansion in partial cubes is called *contraction* and consists in taking a Θ -class of edges E_f and contracting it. The two disjoint copies of the corresponding G^1 and G^2 are just the two components of the graph where E_f is deleted.

2 The flaw and the result

Let H be an expansion of a planar graph G with respect to G^1 and G^2 . Then H is a *2-face expansion* of G if G^1 and G^2 have plane embeddings such that $G' := G^1 \cap G^2$ lies on a face in both the respective embeddings. Peterin [4] proposes a theorem stating that a graph is a planar partial cube if and only

*ENS Cachan, Université Paris-Saclay

†Laboratoire d'Informatique Fondamentale, Aix-Marseille Université and CNRS, Faculté des Sciences de Luminy, F-13288 Marseille Cedex 9, France

if it can be obtained from a single vertex by a sequence of 2-face expansions. However, his argument has a flaw, since G' lying on a face of G^1 and G^2 does not guarantee that the expansion H be planar. Indeed, Figure 1 shows an example of such a 2-face expansion H of a planar graph G that is non-planar.




Figure 1: Left: A 2-face expansion H of a planar partial cube G , where G^1 and G^2 are drawn as crosses and circles, respectively. Right: A subdivision of $K_{3,3}$ (bold) in H , certifying that H is not planar.

The correct concept are non-crossing 2-face expansions: We call an expansion H of a planar graph G with respect to subgraphs G^1 and G^2 a *non-crossing 2-face expansion* if G^1 and G^2 have plane embeddings such that $G' := G^1 \cap G^2$ lies on the outer face of both the respective embeddings, such that the orderings on G' obtained from traversing the outer faces of G^1 and G^2 in the clockwise order, respectively, are opposite.

Lemma 1. *For a partial cube $H \not\cong K_1$ the following are equivalent:*

- (i) H is planar,
- (ii) H is a non-crossing 2-face expansion of a planar partial cube G ,
- (iii) if H is an expansion of G , then G is planar and H is a non-crossing 2-face expansion of G .

Proof.

(ii) \implies (i): Let G be a planar partial cube and G^1 and G^2 two subgraphs satisfying the preconditions for doing a non-crossing 2-face expansion. We can thus embed G^1 and G^2 disjointly into the plane such that the two copies of $G' := G^1 \cap G^2$ appear in opposite order around their outer face, respectively. Connecting corresponding vertices of the two copies of G' by a matching E_f does not create crossings, because the 2-face expansion is non-crossing, see Figure 2. Thus, if H is a non-crossing 2-face expansion of G , then H is planar.

(i) \implies (iii): Let H be a planar partial cube, that is an expansion of G . Thus, there is a Θ -class E_f of H such that $G = H/E_f$. In particular, since contraction preserves planarity, G is planar.

Consider now H with some planar embedding. Since H is a partial cube, E_f is an inclusion-minimal edge cut of H . Thus, $H \setminus E_f$ has precisely two components corresponding to G^1 and G^2 , respectively. Since E_f is a minimal cut its planar dual is a simple cycle C_f . It is well-known, that any face of a planar embedded graph can be chosen to be the outer face without changing the combinatorics of the embedding. We change the embedding of H , such




Figure 2: Two disjoint copies of subgraphs G^1 and G^2 in a planar partial cube H .

that some vertex v of C_f corresponds to the outer face of the embedding, see Figure 2.

Now, without loss of generality C_f has G^1 and G^2 in its interior and exterior, respectively. Since C_f is connected and disjoint from G^1 and G^2 it lies in a face of both. By the choice of the embedding of H it is their outer face. Moreover, since every vertex from a copy of G' in G^1 can be connected by an edge of E_f to its partner in G^2 crossing an edge of C_f but without introducing a crossing in H , the copies of G' in G^1 and G^2 lie on this face, respectively.

Furthermore, following E_f in the sense of clockwise traversal of C_f gives the same order on the two copies of G' , corresponding to a clockwise traversal on the outer face of G^1 and a counter-clockwise traversal on the outer face of G^2 . Thus, traversing both outer faces in clockwise order the obtained orders on the copies of G' are opposite. Hence H is a non-crossing 2-face expansion of G .

(iii) \implies (ii): Since $H \not\cong K_1$, it is an expansion of some partial cube G . The rest is trivial. \square

Lemma 1 yields our characterization.

Theorem 2. *A graph H is a planar partial cube if and only if H arises from a sequence of non-crossing 2-face expansions from K_1 .*

Proof.

\implies : Since H is a partial cube by the result of Chepoi [3] it arises from a sequence of expansions from K_1 . Moreover, all these sequences have the same length corresponding to the number of Θ -classes of H . We proceed by induction on the length ℓ of such a sequence. If $\ell = 0$ the sequence is empty and there is nothing to show. Otherwise, since $H \not\cong K_1$ is planar we can apply Lemma 1 to get that H arises by a non-crossing 2-face expansions from a planar partial cube G . The latter has a sequence of expansions from K_1 of length $\ell - 1$ which by induction can be chosen to consist of non-crossing 2-face expansions. Together with the expansion from G to H this gives the claimed sequence from H .

\impliedby : Again we induct on the length ℓ of the sequence. If $\ell = 0$ we are fine since K_1 is planar. Otherwise, consider the graph G in the sequence such that H is its non-crossing 2-face expansion. Then G is planar by induction and H is planar by Lemma 1, since it is a non-crossing 2-face expansion of G . \square

3 Remarks

We have characterized planar partial cubes graphs by expansions. Planar partial cubes have also been characterized in a topological way as dual graphs of non-separating pseudodisc arrangements [1]. There is a third interesting way of characterizing them. The class of planar partial cubes is closed under *partial cube minors*, see [2], i.e., contraction of G to G/E_f where E_f is a Θ -class and restriction to a component of $G \setminus E_f$. What is the family of minimal obstructions for a partial cube to being planar, with respect to this notion of minor? The answer will be an infinite list, since a subfamily is given by the set $\{G_n \square K_2 \mid n \geq 3\}$, where G_n denotes the *gear graph* (also known as *cogwheel*) on $2n + 1$ vertices and \square is the Cartesian product of graphs. See Figure 3 for the first three members of the family.




Figure 3: The first three members of an infinite family of minimal obstructions for planar partial cubes.

Acknowledgements:

We wish to thank Iztok Peterin for discussing this result with us and two referees for helpful comments.

References

- [1] M. ALBENQUE AND K. KNAUER, *Convexity in partial cubes: the hull number.*, Discrete Math., 339 (2016), pp. 866–876.
- [2] V. CHEPOI, K. KNAUER, AND T. MARC, *Partial cubes without Q_3^- minors*, (2016). arXiv:1606.02154.
- [3] V. CHEPOJ, *Isometric subgraphs of Hamming graphs and d -convexity.*, Cybernetics, 24 (1988), pp. 6–11.
- [4] I. PETERIN, *A characterization of planar partial cubes.*, Discrete Math., 308 (2008), pp. 6596–6600.
- [5] D. Ž. DJOKOVIĆ, *Distance-preserving subgraphs of hypercubes.*, J. Comb. Theory, Ser. B, 14 (1973), pp. 263–267.
- [6] P. M. WINKLER, *Isometric embedding in products of complete graphs.*, Discrete Appl. Math., 7 (1984), pp. 221–225.