



HAL
open science

CAESAR: a new tool to assess relatives' experience of dying and death in the ICU

Nancy Kentish-Barnes, Valerie Seegers, Stephane Legriél, Alain Cariou, Samir Jaber, Jean Yves Lefrant, Bernard Floccard, Anne Renault, Isabelle Vinatier, Armelle Mathonnet, et al.

► To cite this version:

Nancy Kentish-Barnes, Valerie Seegers, Stephane Legriél, Alain Cariou, Samir Jaber, et al.. CAESAR: a new tool to assess relatives' experience of dying and death in the ICU. *Intensive Care Medicine*, 2016, 42 (6), pp.995-1002. 10.1007/s00134-016-4260-4 . hal-01458388

HAL Id: hal-01458388

<https://hal.science/hal-01458388>

Submitted on 5 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CAESAR: a new tool to assess relatives' experience of dying and death in the ICU

Nancy Kentish-Barnes¹, Valérie Seegers^{2,3}, Stéphane Legriel⁴, Alain Cariou^{5,6}, Samir Jaber^{7,8}, Jean-Yves Lefrant^{9,10}, Bernard Floccard¹¹, Anne Renault¹², Isabelle Vinatier¹³, Armelle Mathonnet¹⁴, Danielle Reuter¹, Olivier Guisset¹⁵, Christophe Cracco¹⁶, Amélie Seguin¹⁷, Jacques Durand-Gasselín¹⁸, Béatrice Éon¹⁹, Marina Thirion²⁰, Jean-Philippe Rigaud²¹, Bénédicte Philippon-Jouve²², Laurent Argaud^{11,23}, Renaud Chouquer²⁴, Mélanie Adda²⁵, Laurent Papazian^{25,33}, Céline Dedrie²⁶, Hugues Georges²⁷, Eddy Lebas²⁸, Nathalie Rolin²⁹, Pierre-Edouard Bollaert^{30,31}, Lucien Lecuyer³², Gérald Viquesnel¹⁷, Marc Léone^{25,33}, Ludivine Chalumeau-Lemoine³⁴, Zoé Cohen-Solal¹, Maité Garrouste-Orgeas³⁵, Fabienne Tamion³⁶, Bruno Falissard², Sylvie Chevret^{37,38} and Elie Azoulay^{1,37*}

Abstract

Purpose: To develop an instrument designed specifically to assess the experience of relatives of patients who die in the intensive care unit (ICU).

Methods: The instrument was developed using a mixed methodology and validated in a prospective multicentre study. Relatives of patients who died in 41 ICUs completed the questionnaire by telephone 21 days after the death, then completed the Hospital Anxiety and Depression Scale, Impact of Event Scale-Revised and Inventory of Complicated Grief after 3, 6, and 12 months.

Results: A total of 600 relatives were included, 475 in the main cohort and 125 in the reliability cohort. The 15-item questionnaire, named CAESAR, covered the patient's preferences and values, interactions with/around the patient and family satisfaction. We defined three groups based on CAESAR score tertiles: lowest (≤ 59 , $n = 107$, 25.9%), middle ($n = 185$, 44.8%) and highest (≥ 69 , $n = 121$, 29.3%). Factorial analysis showed a single dimension. Cronbach's alpha in the main and reliability cohorts was 0.88 (0.85–0.90) and 0.85 (0.79–0.89), respectively. Compared to a high CAESAR score, a low CAESAR score was associated with greater risks of anxiety and depression at 3 months [1.29 (1.13–1.46), $p = 0.001$], post-traumatic stress-related symptoms at 3 [1.34 (1.17–1.53), $p < 0.001$], 6 [OR = 1.24 (1.06–1.44), $p = 0.008$] and 12 [OR = 1.26 (1.06–1.50), $p = 0.01$] months and complicated grief at 6 [OR = 1.40 (1.20–1.63), $p < 0.001$] and 12 months [OR = 1.27 (1.06–1.52), $p = 0.01$].

*Correspondence: elie.azoulay@sls.aphp.fr

¹ Assistance Publique-Hôpitaux de Paris, Famiréa Research Group, Saint-Louis University Hospital, Paris, France

Full author information is available at the end of the article

Take-home message: A 15-item questionnaire (CAESAR) proved reliable for assessing the experience of relatives with the dying and death of a loved one in the intensive care unit. This instrument should prove useful as the primary endpoint in trials of interventions designed to improve family care.

Conclusions: The CAESAR score 21 days after death in the ICU is strongly associated with post-ICU burden in the bereaved relatives. The CAESAR score should prove a useful primary endpoint in trials of interventions to improve relatives' well-being.

Keywords: Relatives, Intensive care unit, Death, Bereavement, Complicated grief

Introduction

An increasing number of deaths occur in hospitals, particularly in intensive care units (ICUs) [1], where the burden on relatives is heavy [2–5]. This fact, together with the high mortality rates in ICUs, has prompted tremendous efforts to understand the experience of dying and death in the ICU [6]. Studies have identified quality markers [7], targets for improving end-of-life care, and metrics for assessing the effectiveness of ICU interventions. For instance, a communication strategy based on a proactive family conference was shown to diminish the post-ICU burden on relatives, who were less affected by post-traumatic stress disorder (PTSD), anxiety and depression [4]. Identifying meaningful outcomes for patients who die in the ICU has been the focus of specific studies. The components of the post-ICU burden are not necessarily specific to the ICU. Characterising the specific aspects of dying and death in the ICU that are associated with the subsequent burden on relatives is a prerequisite to developing preventive strategies.

A tool designed to measure the quality of dying and death (QODD) in any setting (e.g. hospital, nursing home, home or hospice) from the relatives' perspective has been developed by a group that defined QODD as 'the degree to which a person's preferences for dying and the moment of death agree with observations of how the person actually died as reported by others' [8]. This group used a literature review, patient interviews, focus groups, prior research, existing instruments, and a list of desirable measurement properties to identify six conceptual domains and 31 specific issues [8]. Of these 31 issues, 23 were selected as relevant to a QODD measurement tool for ICU patients [9]. The ICU-QODD has shown valuable psychometric properties [10]. After the death, the ICU-QODD is completed by the physicians and nurses in charge of the patient within a few days or weeks and by the relatives after several months.

The QODD tool centres on the patient. The relatives, however, also require specific attention to ensure that their suffering during and after the death is alleviated to the extent possible. ICU healthcare workers must be able to identify the personal characteristics that affect relatives' coping style and ability, such as the nature of their relationship with the patient, ability to understand medical information, communication style with the ICU staff, beliefs, support system and usual degree of psychological

well-being. The first step towards achieving this objective is to understand the experience of relatives of patients dying in the ICU.

Here, our objective was to develop and validate a tool specifically designed to assess the overall experience of relatives of patients who die in the ICU. We used a mixed methodology to create the questionnaire then validated it in a multicentre prospective study in 41 French ICUs.

Patients and methods

The study was approved by the institutional review board of the Paris-North Hospitals (IRB00006477, approval #11-019), Paris 7 University. Informed consent was obtained from each relative before study inclusion.

Development of the CAESAR instrument

We used three sources of data to develop the initial questionnaire: a review of the medical, nursing and social-science literature; the experience accumulated by our research group; and in-depth qualitative interviews by a sociologist (NKB) with relatives of patients expected to die in the ICU, bereaved relatives, physicians and nurses.

A panel of ten investigators (five physicians, two nurses and three sociologists/psychologists) held several meetings to discuss the data collected from these three sources. They identified eight domains and 50 items about the experience of relatives (Table S1). The items were then tested twice with relatives of patients, as well as with physicians and nurses, in one ICU (Saint Louis Hospital). The distribution of each item was examined, and a floor or ceiling effect was defined as having more than 85 % of respondents with the lowest or highest item score [11], a characteristic potentially associated with poor discrimination. Items with a large number of missing answers were then identified as potentially unreliable. Finally, a factorial analysis was performed to identify items with a weak representation. On the basis of the results of these analyses, the panel eliminated 17 items. The remaining 33 items (Table S2) fell into three domains: the patient (preparation for death, whole person concerns, symptoms, personal care, and treatment preferences); interactions with and around the patient (quality of communication between the ICU team and the patient and between the ICU team and the relatives, particularly whether conflicts arose), and family needs and satisfaction. The items are not arranged by domain on the questionnaire form but

Table 1 Final 15-item CAESAR questionnaire

Items	Mean score (standard deviation)
1. Was your loved one's pain well controlled throughout the ICU stay? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.2 (1.2)
2. Do you feel that your loved one's dignity was maintained? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.6 (0.9)
3. Do you feel the ICU team was attentive to your loved one? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.4 (0.9)
4. Are you satisfied with the quality of medical care received by your loved one? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.4 (1.1)
5. During the days before the death, were you clearly informed that your loved one was dying? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	3.2 (1.3)
6. Are you satisfied with the quality of the communication between you and the physicians? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.4 (1.1)
7. Are you satisfied with the quality of the communication between you and the nurses? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.7 (0.8)
8. Were you in conflict with the ICU team? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.6 (0.8)
9. Were you given the opportunity to discuss your loved one's wishes, as well as your own preferences, with the ICU team? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.1 (1.1)
10. Did your loved one refuse any of the suggested treatments? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.1 (0.8)
11. Do you believe the ICU team went too far in the treatment given to your loved one? Do you believe the ICU team used unnecessary treatments? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	3.9 (1.3)
12. Were you able to say goodbye and express important feelings to your loved one? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	3.9 (1.3)
13. Were you present when your loved one died? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.5 (0.9)
14. Are you satisfied with the support you received while your loved one was dying? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	3.9 (0.6)
15. During your loved one's stay in the ICU, did you receive counselling, for instance from a psychologist? Please rate this experience: 1 Traumatic, 2 Painful, 3 Difficult, 4 Acceptable, 5 Comforting	4.2 (0.8)

are instead asked in the order believed to be easiest and least traumatic for the relatives. Each item is answered by both a written description and a score on a 5-point scale (1, traumatic; 2, painful; 3, difficult; 4, acceptable; and 5, comforting). A questionnaire was also developed for the physicians and nurses and used for a separate analysis. Finally, 18 additional items were eliminated because they were redundant, as determined on the basis of inter-item Spearman's correlation coefficients [12]. This produced the 15-item questionnaire shown in Table 1 and designated CAESAR. Relatives need approximately 20 min to complete the instrument.

Validation of the CAESAR questionnaire

The 33-item questionnaire was validated in a multicentre prospective study conducted from July 2011 to July 2013 in 41 French ICUs belonging to the FAMIREA network, as described elsewhere [13]. In each ICU, one of the intensivists included consecutive adults who died after at least 48 h in the ICU. For each patient, we included the

relative who served as the surrogate at the time of death (designated proxy > spouse > adult child > sibling > other relative, in that order). Eligibility criteria for relatives were having visited the patient at least once in the ICU and having sufficient knowledge of French to complete the assessment tools.

The data shown in the tables and figures were collected prospectively. The relatives completed the 33-item questionnaire during a telephone interview 21 days after the patient's death. To validate the questionnaire, a reliability cohort of relatives of patients who died in the same 41 ICUs was recruited. During a telephone interview 3 months after the death, each relative in the cohort completed the Hospital Anxiety and Depression Scale (HADS) and Impact of Event Scale-Revised (IES-R) for PTSD symptoms. All telephone interviews were conducted by the same sociologist with extensive interviewing experience (ZCS). Finally, 6 and 12 months after the death, a questionnaire including the Inventory of Complicated Grief (ICG) and IES-R was mailed to each relative.




Fig. 1 Recruitment flow chart for the development and reliability cohorts

Factorial validity was assessed in the main sample by determining the dimensional structure of the 33-item questionnaire and final 15-item CAESAR questionnaire. To this end, we used maximum likelihood factor analysis with varimax rotation. The number of factors was determined from the observation of the screen plots and value of simulations [14–16]. Internal consistency was deemed acceptable when Cronbach’s alpha [17] was in the 0.70–0.95 range [18]. Confidence intervals (CIs) were computed on the basis of 1000 bootstrap replicates. Factorial validity was assessed in the reliability sample using the same methodology.

For the final 15-item tool, the item scores (range 1–5) were summed to obtain a global score (15–75).

Statistical analysis

Statistical analyses were performed using the R 3.1 package with the “psy” package [19]. Quantitative data were described as median (interquartile range) and binary and categorical data as number (%). All statistical tests were two-sided and p values of 0.05 or less were considered significant.

Results

Study population

Figure 1 is the study flow chart. Among the 4607 patients admitted to the 41 participating ICUs during the study period, 875 (19 %) died, including 228 who met exclusion criteria, 104 for whom the opportunity for inclusion was missed and 68 whose relatives refused participation. For

each of the remaining 475 (54 %) patients, one relative was included. On day 21, 430 (90.5 %) relatives completed the 33-item instrument. A different group of 232 relatives was included in the reliability cohort; 116 (93 %) of these relatives completed the 33-item instrument on day 21.

Table 2 reports the main characteristics of the relatives and end-of-life circumstances according to CAESAR scores.

CAESAR scores

Of the 430 33-item questionnaires completed on day 21, 413 (96 %) had no missing data for the 15 CAESAR items and were therefore used to determine the global score. The median score was 66 (range 21–75, interquartile range 59–69). Of the score values, 107 (25.9 %) were in the lowest tertile (≤ 59), 185 (44.8 %) in the middle tertile and 121 (29.3 %) in the highest tertile (≥ 69). Table 3 reports the prevalence of symptoms of post-ICU burden according to CAESAR score tertiles.

Figure S1A depicts the distribution of individual item scores and Fig. S1B the distribution of the global CAESAR score. The factorial analysis of the main sample, assessed by screen plots, was consistent with a single dimension, as 42.1 % of the variance was explained by the first principal component (Fig. S2A). Cronbach’s alpha was 0.88 (95 % CI 0.85–0.90). Similarly, in the reliability sample, the results suggested a one-dimensional structure, with 35.9 % of the variance explained by the first principal component

Table 2 Characteristics of the relatives, end-of-life, and ICU by CAESAR score tertile

	Overall <i>n</i> = 413 (%)	Lowest tertile <i>n</i> = 107	Middle tertile <i>n</i> = 185	Highest tertile <i>n</i> = 121	<i>p</i> value
Relatives, <i>n</i> (%)					
Gender, <i>n</i> (%)					
Male	128 (31.0 %)	38 (35.5 %)	51 (27.6 %)	39 (32.2 %)	0.346
Female	285 (69.0 %)	69 (64.5 %)	134 (72.4 %)	82 (67.8 %)	
Relationship to patient					
Spouse	140 (33.9 %)	38 (35.5 %)	69 (37.3 %)	33 (27.3 %)	0.199
Adult child	188 (45.5 %)	50 (46.7 %)	82 (44.3 %)	56 (46.3 %)	
Parent	13 (3.1 %)	5 (4.7 %)	3 (1.6 %)	5 (4.1 %)	
Brother/sister	37 (9.0 %)	10 (9.3 %)	16 (8.6 %)	11 (9.1 %)	
Other	35 (8.5 %)	4 (3.7 %)	15 (8.1 %)	16 (13.2 %)	
Religious/spiritual beliefs					
Practice of a religion	280 (67.8 %)	71 (66.4 %)	122 (65.9 %)	87 (71.9 %)	0.496
	107 (25.9 %)	22 (20.6 %)	48 (25.9 %)	37 (30.6 %)	0.243
End-of-life					
Length of ICU stay (days), median (IQR)	7 (4–17)	11 (6–24)	6 (3–16)	6.5 (4–12)	0.008
Use of vasopressors, <i>n</i> (%)	314 (76. %)	91 (85 %)	140 (75.7 %)	83 (68.6 %)	0.016
Decision to withhold or withdraw life-sustaining therapy, <i>n</i> (%)					
None	80 (19.4 %)	26 (24.3 %)	36 (19.5 %)	18 (14.9 %)	0.494
Withholding decision	134 (32.4 %)	31 (29 %)	61 (33 %)	42 (34.7 %)	
Withdrawing decision	199 (48.2 %)	50 (46.7 %)	88 (47.6 %)	61 (50.4 %)	
Family participation in the decision-making process, <i>n</i> (%)					
Actively involved	198 (47.9 %)	40 (37.4 %)	92 (49.7 %)	66 (54.5 %)	0.103
Informed of decisions	135 (32. %7)	41 (38.3 %)	57 (30.8 %)	37 (30.6 %)	
Neither involved nor informed	80 (19.4 %)	26 (24.3 %)	36 (19.5 %)	18 (14.9 %)	
How the death was disclosed to the family, <i>n</i> (%)					
At the bedside	228 (55.2 %)	65 (60.7 %)	93 (50.3 %)	70 (57.9 %)	0.066
Over the phone	136 (32.9 %)	27 (25.2 %)	72 (38.9 %)	37 (30.6 %)	
Upon arrival at the ICU	49 (11.9 %)	15 (14.9 %)	20 (10.8 %)	14 (11.5 %)	
Mechanical ventilation at the time of death, <i>n</i> (%)					
Intubated	273 (66.1 %)	75 (70.1 %)	125 (67.6 %)	73 (60.3 %)	0.572
Recently extubated	58 (14.0 %)	13 (12.1 %)	24 (13 %)	21 (17.4 %)	
Never intubated	82 (19.9 %)	19 (17.8 %)	36 (19.5 %)	27 (22.3 %)	
Involvement of external consultants, <i>n</i> (%)					
Palliative-care team	12 (2.9 %)	4 (3.7 %)	4 (2.2 %)	4 (3.3 %)	0.702
Clergyman/chaplain	51 (12.3 %)	17 (15.9 %)	21 (11.4 %)	13 (10.7 %)	0.372
ICU characteristics, <i>n</i> (%)					
24-hour visiting	151 (36.6 %)	32 (29.9 %)	72 (38.9 %)	47 (38.8 %)	0.252
Psychologist available	150 (36.3 %)	38 (35.5 %)	69 (37.3 %)	43 (35.5 %)	0.933

ICU intensive care unit, IQR interquartile range

(Fig. S2). Cronbach's alpha was 0.85 (95 % CI 0.79–0.89). Spearman's correlation coefficients between each item and the global score ranged from 0.39 to 0.79 (Table S3).

Factors associated with the CAESAR score

A longer ICU stay length was significantly associated with a lower CAESAR score indicating a more difficult experience with dying and death [11 (6–24), 6 (3–16) and 6.5 (4–12) days in the low, middle and high tertiles, respectively, $p = 0.008$]. Vasopressor therapy was also

associated with a lower CAESAR score (85, 75.7 and 68.6 % of patients received vasopressors in the low, middle and high tertiles, respectively, $p = 0.016$).

At 3 months, 370 (86 %) relatives completed the HADS and IES-R. Among them, 190 (51.4 %) had symptoms of depression and 129 (34.9 %) symptoms of anxiety. At 6 and 12 months, 268 (64.9 %) and 209 (50.6 %) relatives completed the ICG and IES-R. Among them, 139 (51.9 %) met criteria for complicated grief at 6 months and 113 (54.1 %) at 12 months. Complicated grief at 6 months was

Table 3 Post-ICU burden according to CAESAR score tertile

Median (IQR) or numbers (%)	Overall	Lowest tertile <i>n</i> = 107	Middle tertile <i>n</i> = 185	Highest tertile <i>n</i> = 121	<i>p</i> value
At 3 months (<i>n</i> = 385)					
HADS score	13 (7–22)	18 (9.5–24)	13 (7–20)	10 (5–17.75)	<0.001
Relatives with symptoms of anxiety	198 (51.4 %)	58 (61.1 %)	83 (50.3 %)	49 (44.5 %)	0.058
Relatives with symptoms of depression	137 (35.6 %)	49 (51.6 %)	57 (34.5 %)	23 (20.9 %)	<0.001
IES-R score	30 (13–45)	38 (25–54)	28 (14–45)	17 (5.25–17.75)	<0.001
Relatives with PTSD symptoms	173 (44.9 %)	57 (60.0 %)	74 (44.8 %)	34 (30.9 %)	<0.001
At 6 months (<i>n</i> = 267)					
ICG score	27 (16–40)	37 (24.5–49)	25 (17–39.75)	22 (10.5–35.5)	<0.001
Relatives with complicated grief	140 (52.4 %)	49 (73.1 %)	54 (49.1 %)	36 (39.6 %)	<0.001
IES-R score	29 (16–43)	35 (21–49.5)	28 (17–45.75)	25 (12–36)	0.004
Relatives with PTSD symptoms	116 (43.4 %)	37 (55.2 %)	47 (42.7 %)	31 (34.1 %)	0.029
At 12 months (<i>n</i> = 207)					
ICG score	27 (14–41)	31 (21–45)	29 (19–42)	22 (9.75–34.25)	0.003
Relatives with complicated grief	112 (54.1 %)	29 (64.4 %)	53 (60.2 %)	31 (40.8 %)	0.013
IES-R score	26 (13–43)	31 (21–46)	30 (13.75–45)	20.5 (10–30.25)	0.002
Relatives with PTSD symptoms	76 (36.7 %)	21 (46.7 %)	38 (43.2 %)	18 (23.7 %)	0.011

Lowest, middle and highest CAESAR tertiles were scores ≤ 59 , between 60 and 68, and ≥ 69 , respectively

ICU intensive care unit, IQR interquartile range, HADS Hospital Anxiety and Depression Scale, IES-R Impact of Event Scale-Revised, PTSD post-traumatic stress disorder, ICG Inventory of Complicated Grief

more common in the lowest CAESAR score tertile than in the middle tertile [odds ratio (OR) 1.27, 95 % CI 1.10–1.47, $p = 0.002$] or highest tertile (OR 1.40, 95 % CI 1.20–1.63, $p < 0.001$) (Fig. 2). Corresponding figures at 12 months were as follows: OR 1.04, 95 % CI 0.87–1.24, $p = 0.64$; and OR 1.27, 95 % CI 1.06–1.52, $p = 0.011$, respectively. Table S4 reports associations between CAESAR score tertiles and clinical endpoints at 3, 6 and 12 months.

The prevalence of PTSD-related symptoms as assessed by the IES-R decreased over time, from 44.6 % ($n = 165/370$) at 3 months to 42.9 % ($n = 115/268$) at 6 months and 36.8 % ($n = 77/209$) at 12 months. At 6 months, relatives in the lowest CAESAR score tertile were at higher risk for PTSD-related symptoms compared to those in the middle tertile (OR 1.13, 95 % CI 0.98–1.32, $p = 0.102$) or highest tertile (OR 1.24, 95 % CI 1.06–1.44, $p = 0.008$) (Fig. 2). Corresponding figures at 12 months were as follows: OR 1.04, 95 % CI 0.87–1.23, $p = 0.69$; and OR 1.26, 95 % CI 1.06–1.50, $p = 0.011$, respectively.

Discussion

This study performed in 41 ICUs belonging to the FAMIREA network allowed us to develop and validate a new tool, the 15-item CAESAR questionnaire, designed to measure the self-reported experience of relatives with the dying and death of a family member in the ICU. The CAESAR score 21 days after the patient's death correlated strongly with the presence in the following months of symptoms of anxiety, depression, PTSD and complicated grief.

This tool should prove useful in assessing the efficacy of interventions designed to improve the quality of relatives' experience with dying and death of ICU patients. Furthermore, it can serve to identify areas where improvements are most needed. These expected benefits are of considerable importance, as high levels of post-ICU burden continue to be documented despite recent advances in communication strategies, training, incorporation of palliative care in the ICU and recognition of relatives' vulnerability [20–22].

To determine how to improve end-of-life care in the ICU, we need objective and subjective information on the process of dying, the management of this process in the ICU, and the effects of specific features of the process on relatives' well-being. The items of the CAESAR questionnaire were derived from studies in the field of social science and from qualitative interviews of relatives of ICU patients and of ICU workers. CAESAR provides information on the relatives' experience with ICU practices rather than on the practices themselves. This approach sheds light on the aspects of end-of-life care that are important to families, as opposed to physicians and nurses. Moreover, as shown in a previous publication [13], our study identified features of end-of-life care that correlated with major components of the post-ICU burden such as PTSD and complicated grief. Thus, an individualized determination of the part each specific patient wants to play and the opportunity for the relatives to say goodbye and to be present at the time of death are particularly important to the relatives [13].




Fig. 2 Association between the CAESAR score and clinical endpoints 3, 6 and 12 months after the patient's death, assessed by logistic regression

The CAESAR scale adds to the ICU-QODD, which relies primarily on data from family members and not from healthcare workers [23]. Furthermore, we focused on relatives and not patients. When developing the CAESAR tool, we eliminated items that often went unanswered in the test administrations. ICU-QODD and CAESAR include shared variables that influence their scores, such as pain control, patient's dignity, family presence at the time of death and discussion of the patient's wish to withdraw life support [24]. Other variables specific to CAESAR, such as quality of communication, are considered by relatives to be among the most highly valued aspects of care [25]. Studies comparing the information supplied by the two tools are warranted. However, discrepancies can be expected, as family members give more favourable ratings of dying and death compared to nurses [26].

The global CAESAR score provides a picture of relatives' experience with end-of-life care in the ICU. By determining this score at regular intervals, ICUs can monitor their performance over time. However, further work is needed to assess the sensitivity of the CAESAR score to interventions designed to improve dying and death in the ICU. The median CAESAR score was 66/75 (21–75), and only 25.9 % of relatives were in the lowest tertile. Whether CAESAR can serve as a benchmarking tool needs further exploration.

This study has several limitations. First, all participating ICUs were in France and the findings may not be applicable to other countries. However, the large number of ICUs and relatives and the validation of the results in a reliability cohort support the robustness of the data. Second, instead of asking the relatives to complete the questionnaire immediately after the death, we waited 3 weeks. We believe that completing the questionnaire immediately may be difficult or even distressful. Compared to earlier studies, the evaluation of the relatives' experience was performed far earlier and using a more uniform approach. Moreover, a single highly experienced sociologist administered the questionnaire during a telephone interview, offering support whenever the items activated distressing emotions. Such support is known to be much appreciated by families [27]. Last, not all relatives could be contacted at 6 and 12 months. However, the response rates at these time points are at the high end of the range reported in earlier studies [4, 9].

In summary, we describe and validate a new instrument for assessing the experience of relatives of patients who die in the ICU. Our study suggests new targets for interventions designed to improve family care in the ICU. Only a quarter of the relatives had low CAESAR scores indicating a distressing experience. However, these low scores were associated with a greater post-ICU burden. The CAESAR score could serve as a primary endpoint in studies of

interventions aimed at improving the experience of relatives with the dying and death of a loved one in the ICU.

Author details

¹ Assistance Publique-Hôpitaux de Paris, Famiréa Research Group, Saint-Louis University Hospital, Paris, France. ² Paris-Sud Innovation Group in Mental Health, UMR 669, INSERM, Paris-Sud University, Paris, France. ³ Data Management Research Department DRCl, Angers Hospital and SFR ICAT, University of Angers, Angers, France. ⁴ Versailles Hospital, Versailles, France. ⁵ Assistance Publique-Hôpitaux de Paris, Cochin University Hospital, Paris, France. ⁶ Paris Descartes University, Paris, France. ⁷ Saint Eloi University Hospital, Montpellier, France. ⁸ Montpellier 1 University, Montpellier, France. ⁹ Carêmeau University Hospital, Nîmes, France. ¹⁰ Nîmes University, Nîmes, France. ¹¹ Hospices Civils de Lyon, Edouard Herriot University Hospital, Lyon, France. ¹² Cavale Blanche University Hospital, Brest, France. ¹³ Les Oudairies Hospital, La Roche Sur Yon, France. ¹⁴ Hospital de la Source, Orléans, France. ¹⁵ Saint André University Hospital, Bordeaux, France. ¹⁶ Angoulême Hospital, Angoulême, France. ¹⁷ Caen University Hospital, Caen, France. ¹⁸ Sainte Musse Hospital, Toulon, France. ¹⁹ La Timone University Hospital, Marseille, France. ²⁰ Victor Dupouy Hospital, Argenteuil, France. ²¹ Dieppe Hospital, Dieppe, France. ²² Roanne Hospital, Roanne, France. ²³ Lyon Est University, Lyon, France. ²⁴ Annecy Hospital, Annecy, France. ²⁵ Hôpital Nord University Hospital, Marseille, France. ²⁶ Roubaix Hospital, Roubaix, France. ²⁷ Chatilliez Hospital, Tourcoing, France. ²⁸ Bretagne Atlantique Hospital, Vannes, France. ²⁹ Marc Jacquet Hospital, Melun, France. ³⁰ Nancy University Hospital, Nancy, France. ³¹ Lorraine University, Nancy, France. ³² Sud Francilien Hospital, Evry, France. ³³ Aix-Marseille University, Marseille, France. ³⁴ Gustave Roussy Institut, Villejuif, France. ³⁵ Saint Joseph Hospital, Paris, France. ³⁶ Rouen University Hospital, Rouen, France. ³⁷ Biostatistics and Clinical Epidemiology research team, U1153, INSERM, Paris Diderot Sorbonne University, Paris, France. ³⁸ Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Service de Biostatistique et Information Médicale, Paris, France.

Acknowledgments

These contributors participated to the study: Michel Badet, Chambéry Hospital, Chambéry, France; Julie Carr, Saint Eloi University Hospital, Montpellier, France; Sophie Cayot-Constantin, Estaing University Hospital, Clermont Ferrand, France; Vincent Das, André Grégoire Hospital, Montreuil, France; Fabienne Fieux, Saint Louis University Hospital, Paris, France; Emmanuelle Hammad, Hôpital Nord University Hospital, Marseille, France; Mercé Jourdain, Roger Salengro University Hospital, Lille, France; Véronique Leray, La Croix Rousse Hospital, Lyon, France; Djamel Mokart, Institut Paoli Calmettes, Marseille, France; Sami Hreich Hôpital Nord University Hospital, Marseille, France; Michel Ramakers, Saint Lô Hospital, Saint Lô, France; Jean-Michel Robert, Edouard Herriot University Hospital, Lyon, France; Antoine Roquilly, Hôtel Dieu University Hospital, Nantes, France.

Supported by a Grant from the French Ministry of Health (PHRC 10 104). Medical Intensive Care Unit, Hôpital Saint-Louis, ECSTRA team, Biostatistics and clinical epidemiology, UMR 1153 (Center of Epidemiology and Biostatistics Sorbonne Paris Cité, CRESS), INSERM, Paris Diderot Sorbonne University, Paris, France.

Compliance with ethical standards

Conflicts of interest

None of the authors declare any conflict of interest in relation to this manuscript.

References

1. Angus DC, Barnato AE, Linde-Zwirble WT et al (2004) Use of intensive care at the end of life in the United States: an epidemiologic study. *Crit Care Med* 32:638–643

- Azoulay E, Pochard F, Kentish-Barnes N et al (2005) Risk of post-traumatic stress symptoms in family members of intensive care unit patients. *Am J Respir Crit Care Med* 171:987–994. doi:10.1164/rccm.200409-1295OC
- Cuthbertson BH, Hull A, Strachan M, Scott J (2003) Post-traumatic stress disorder after critical illness requiring general intensive care. *Intensive Care Med* 30:450–455. doi:10.1007/s00134-003-2004-8
- Lautrette A, Darmon M, Megarbane B et al (2007) A communication strategy and brochure for relatives of patients dying in the ICU. *N Engl J Med* 356:469–478. doi:10.1056/NEJMoa063446
- Siegel MD, Hayes E, Vanderwerker LC et al (2008) Psychiatric illness in the next of kin of patients who die in the intensive care unit. *Crit Care Med* 36:1722–1728. doi:10.1097/CCM.0b013e318174da72
- Nelson JE, Azoulay E, Curtis JR et al (2012) Palliative care in the ICU. *J Palliat Med* 15:168–174. doi:10.1089/jpm.2011.9599
- Clarke EB, Luce JM, Curtis JR et al (2004) A content analysis of forms, guidelines, and other materials documenting end-of-life care in intensive care units. *J Crit Care* 19:108–117
- Patrick DL, Engelberg RA, Curtis JR (2001) Evaluating the quality of dying and death. *J Pain Symptom Manage* 22(3):717–726
- Mularski RA, Heine CE, Osborne ML et al (2005) Quality of dying in the ICU: ratings by family members. *Chest* 128:280–287. doi:10.1378/chest.128.1.280
- Curtis JR, Patrick DL, Engelberg RA et al (2002) A measure of the quality of dying and death. Initial validation using after-death interviews with family members. *J Pain Symptom Manage* 24:17–31
- Terwee CB, Bot SDM, de Boer MR et al (2006) Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 60:34–42. doi:10.1016/j.jclinepi.2006.03.012
- Falissard B (2008) Mesurer la subjectivité en santé. Elsevier Masson, Issy-les-Moulineaux
- Kentish-Barnes N, Chaize M, Seegers V et al (2015) Complicated grief after death of a relative in the intensive care unit. *Eur Respir J* 45:1341–1352. doi:10.1183/09031936.00160014
- Lance CE (2006) The sources of four commonly reported cut-off criteria: what did they really say? *Organ Res Methods*. doi:10.1177/1094428105284919
- Cattell RB (1966) The scree test for the number of factors. *Multivar Behav Res* 1:245–276
- Guttman L (1954) Some necessary conditions for common-factor analysis. *Psychometrika* 19:149–161
- Cronbach LJ, Warrington WG (1951) Time-limit tests: estimating their reliability and degree of speeding. *Psychometrika* 16:167–188
- Nunnally JC, Bernstein IH, Berge J (1967) *Psychometric theory*. McGraw-Hill, New York
- Falissard B (2012) *Psy: various procedures used in psychometry*. R package version 1.1. <http://CRAN.R-project.org/package=psy>. Accessed 1 Jan 2015
- Curtis JR (2015) Palliative care in critical illness: challenges for research and practice. *Palliat Med* 29:291–292. doi:10.1177/0269216315573901
- Long AC, Curtis JR (2014) Quality of dying in the ICU: understanding ways to make it better. *Intensive Care Med* 40:1793. doi:10.1007/s00134-014-3512-4
- Aslakson RA, Curtis JR, Nelson JE (2014) The changing role of palliative care in the ICU. *Crit Care Med* 42:2418–2428. doi:10.1097/CCM.0000000000000573
- Curtis JR, Downey L, Engelberg RA (2013) The quality of dying and death: is it ready for use as an outcome measure? *Chest* 143:289–291. doi:10.1378/chest.12-1941
- Glavan BJ, Engelberg RA, Downey L, Curtis JR (2008) Using the medical record to evaluate the quality of end-of-life care in the intensive care unit. *Crit Care Med* 36:1138–1146. doi:10.1097/CCM.0b013e318168f301
- Nelson JE, Mulkerin CM, Adams LL, Pronovost PJ (2006) Improving comfort and communication in the ICU: a practical new tool for palliative care performance measurement and feedback. *Qual Saf Health Care* 15:264–271. doi:10.1136/qshc.2005.017707
- Levy CR, Ely EW, Payne K et al (2005) Quality of dying and death in two medical ICUs: perceptions of family and clinicians. *Chest* 127:1775–1783. doi:10.1378/chest.127.5.1775
- Kentish-Barnes N, McAdam JL, Kouki S et al (2015) Research participation for bereaved family members: experience and insights from a qualitative study. *Crit Care Med*. doi:10.1097/CCM.0000000000001092