

#### Preliminary results on plant Farmland papers Audrey Alignier

#### ▶ To cite this version:

Audrey Alignier. Preliminary results on plant Farmland papers. Farmland Autumn Writing Workshop, Nov 2015, Cassagnabère, France. 20 p. hal-01458631

#### HAL Id: hal-01458631 https://hal.science/hal-01458631

Submitted on 5 Jun2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# Preliminary results on plant Farmland paper

Audrey Alignier, INRA SAD-Paysage

Farmland Autumn Writing workshop – Cassagnabère – 03/11/2015



## A brief summary of previsional papers using plant data

**Paper 1**: How landscape composition and configuration influence plant diversity <u>at the field scale</u> ?

- Same general hypotheses on landscape metrics as the joint paper

- Focusing on landscape composition :

Which crops are delivering most of the plant diversity (or species richness)? Can we increase plant diversity by increasing the number of crops?

- Focusing on landscape configuration :

Are responses between field centre and field margin different ?

**Paper 2**: How landscape composition and configuration influence plant functional groups <u>at the crop scale</u>?

Focus on species composition and functional groups

Can be enhanced certain functional group by sowing a certain crop? What crop can we plant to enhance rare/damaging weeds ?

## A brief summary of previsional papers using plant data

**Paper 1**: How landscape composition and configuration influence plant diversity at the field scale ?


response on pollinators and insect-pollinated plants response on pollinators and insect-pollinated plants response on pollinators and fine the paper on grass multifunction and fine the plants of the paper of the plants the plants the plants

Focus on species composition and functional groups

*Can be enhanced certain functional group by sowing a certain crop?* What crop can we plant to enhance rare/damaging weeds?


## Paper 1 – Species accumulation curves per region

Using « 2013 » data:




## Paper 1 – Species richness per field

ANOVA test: p < 0.001



Note that <u>all</u> plant species from all crop types were included in analyses.

## Paper 1 – Sampling design and sampling unit



### Partitioning diversity:

	at the FIELD scale	at the TRANSECT scale
Gamma	Total species richness	Total species richness
Alpha	Mean species richness (=averaging species richness over the two transects per field)	-
Beta	Gamma - Alpha	-

### Partitioning diversity:

	at the FIELD scale	at the TRANSECT scale
Gamma	Total species richness	Total species richness
Alpha	Mean species richness (=averaging species richness over the two transects per field)	-
Beta	Gamma - Alpha	-
Beta_JAC	Jaccard index (=differences in species composition between the two transects per field)	

• Use of simple linear models as:

Response variable ~ Crop\_SHDI + Crop\_TBL + Crop type + Region + Crop\_SHDI:Region + Crop\_TBL:Region + Crop\_SDHI:Crop + Crop\_TBL:Crop

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	-	-

• Use of simple linear models as:

```
Response variable ~ Crop_SHDI + Crop_TBL + Crop type + Region +
Crop_SHDI:Region + Crop_TBL:Region + Crop_SDHI:Crop + Crop_TBL:Crop
```

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	-	-

• Testing the robustness of models by adding 'non crop cover' and 'woody boundary' :



• Use of simple linear models as:

```
Response variable ~ Crop_SHDI + Crop_TBL + Crop type + Region +
Crop_SHDI:Region + Crop_TBL:Region + Crop_SDHI:Crop + Crop_TBL:Crop
```

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	-	-

• Testing the robustness of models by adding 'non crop cover ' and 'woody boundary' :



• Adding random effects in more complex models (LMMs) as:

Response variable ~ Crop\_SHDI + Crop\_TBL + Crop type + Region + non crop cover + woody boundary + Crop\_SHDI:Region + Crop\_TBL:Region + (1| Landscape) + (1| Year)

• Use of simple linear models as:

```
Response variable ~ Crop_SHDI + Crop_TBL + Crop type + Region +
Crop_SHDI:Region + Crop_TBL:Region + Crop_SDHI:Crop + Crop_TBL:Crop
```

		$\frown$		
	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	ns	-
			$\mathbf{\nabla}$	

• Testing the robustness of models by adding 'non crop cover ' and 'woody boundary' :



• Adding random effects in more complex models (LMMs) as:

Response variable ~ Crop\_SHDI + Crop\_TBL + Crop type + Region + non crop cover + woody boundary + Crop\_SHDI:Region + Crop\_TBL:Region + (1| Landscape) + (1| Year)

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	- / ns	-

More diverse landscapes have less diversity:

...more resistant to weed invasion ?

... less habitats for specialist plants ?

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	- / ns	-

More diverse landscapes have less diversity:

...more resistant to weed invasion ?

... less habitats for specialist plants ?

• Adding « region » as random effect in mixed models (LMMs) :

	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	+	+	+	ns
Crop_TBL	ns	ns	ns	-


	Gamma	Alpha	Beta	Beta_JAC
Crop_SDHI	-	-	ns	ns
Crop_TBL	ns	ns	- / ns	-

More diverse landscapes have less diversity:

...more resistant to weed invasion?

... less habitats for specialist plants ?


• Adding « region » as random effect in mixed models (LMMs) :



		Gamma	Alpha	Beta	Beta_JAC
	Crop_SDHI	-	-	ns	ns
	Crop_TBL	ns	ns	-1 ns ti	ie · resul
More	e diverse land	scapes have le	ess diversity:		
				nt to weed inv	
		• • •	less habitats	for specialist	plants?
• Adding	« region » as	random offor	t in mixed m	odole (LNANAE)	•
Auuing	« region » as	random enec			•
		Gamma	Alpha	Beta	Beta_JAC
	Crop_SDHI	+	+	+	ns

## Paper 1 – Methods and results


ii) To investigate the contribution of crop diversity to species diversity



sites

## Paper 1 – Methods and results

ii) To investigate the contribution of crop diversity to species diversity



Number of crops

iii) To test differences in species diversity response between field border and field centre

Use of mixed effect models (LMMs) as:

Response variable ~ Crop\_SHDI + Crop\_TBL + Crop type + Region + non crop cover + woody boundary + Crop\_SHDI:Region + Crop\_TBL:Region + (1| Landscape) + (1| Year)

	Gamma field border	Gamma field centre
	Total species richness per transect « field border »	Total species richness per transect « field centre »
Crop_SDHI	-	ns
Crop_TBL	ns	ns

Include region as random effect in mixed effect models : TO DO

#### • Grouping crop types and work at the 'crop type' scale

Some crops appear only in few regions *e.g.* almonds

Avoid singleton in models; may help to elucidate the strong regional effect

Proposals:

- 1) arable crops, 2) perennial non woody crops, 3) perennial woody crops ... etc
- 1) winter cereals, 2) summer cereals, 3) grasslands, 4) orchards, ... etc

#### • Differentiating species diversity between field border and field centre

Separate data from field border and field centre as they experienced different disturbance level

#### • Going further in landscape structure characterization

Deepen the effect of landscape composition/configuration by including % grasslands, % hedges, etc.

#### • Separating flora into functional groups

Separate annual from perennial plant species Categorize species as rare, harmful, ... etc