



# Incidence de l'âge et de l'épaisseur de lard dorsal a 100 kg sur la carrière reproductive des truies Large White

R. Gueblez, J.M. Gestin, Genevieve Le Henaff

## ► To cite this version:

R. Gueblez, J.M. Gestin, Genevieve Le Henaff. Incidence de l'âge et de l'épaisseur de lard dorsal a 100 kg sur la carrière reproductive des truies Large White. Annales de zootechnie, 1985, 34, pp.365-365. hal-01460791

**HAL Id: hal-01460791**

**<https://hal.science/hal-01460791>**

Submitted on 2 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

herd having farrowed twice (272 litters) of 969 crossbred *Pietrain* × *Large White* piglets born in the 2nd litters of the 1st generation (group 1) and 579 *Large White* piglets born in the 1st litters of the 2nd generation (group 2). We did not observe any significant effect of sow haplotypes on the number of live born, still born or weaned piglets. We found a significant reduction in the size of the litters in which 25 p. 100 of the piglets should be homozygous for SLA, because their parents had the same haplotype (— 2 piglets born per litter,  $P < 0.01$ ). In group 1, as far as haplotype 2 was concerned, this reduction was due to a significant deficit in homozygous piglets (8.9 instead of 25 p. 100,  $P < 0.01$ ). Thus, there may be a lethal gene associated to this haplotype. In the two groups, the presence of haplotype 1 was associated with a lower mortality rate before weaning. In group 1, it seems that the presence of haplotype 1 was associated with a higher mortality after weaning due to diarrhoea, but this was not the case in group 2. It may be concluded that the main effect of SLA is the unfavourable influence of homozygosity on prolificacy.

### Effect of age and backfat thickness on reproductive performance of 100 kg *Large White* sows

R. GUEBLEZ \*, J.M. GESTIN \*, Geneviève LE HENAFF \*\*

\* *Institut Technique du Porc, B.P. 3, 35650 Le Rheu*

\*\* *I.N.R.A., Station de Génétique quantitative et appliquée  
78350 Jouy-en-Josas*

The age and backfat thickness of 5000 *Large White* selection gilts of 100 kg live weight tested in 1981 and 1982 were related to their reproductive performance as multipliers until the 4th litter. Performances at 100 kg were found to have a marked effect on longevity. Cullings before the 3rd litter were all the more frequent as performances at 100 kg were high. Thus, 45 p. 100 of the animals exhibiting the lowest performances (age at 100 kg  $\geq 170$  days or backfat thickness at 100 kg  $\geq 20$  mm or standard index  $< 80$  points) reached the fourth litter versus 28 p. 100 of the animals with the highest performances (age at 100 kg  $< 140$  days, backfat at 100 kg  $< 14$  mm or standard index  $\geq 120$  points). Prolificacy was not affected. However, the 1st and 2nd litter of the leanest sows at 100 kg tended to be 0.3 to 0.5 piglet larger than those of the fattest sows of the same weight.

This study showed that longevity has to be taken into account in any experiment on sow herd management. Moreover, it is necessary to study the possibilities of improving longevity of purebred herds using other techniques than genetic ones before reconsidering selection goals in female lines.