



HAL
open science

Multiplying a Gaussian Matrix by a Gaussian Vector (and the Gauss-Laplace Transmutation)

Pierre-Alexandre Mattei

► **To cite this version:**

Pierre-Alexandre Mattei. Multiplying a Gaussian Matrix by a Gaussian Vector (and the Gauss-Laplace Transmutation). 2017. hal-01462941v1

HAL Id: hal-01462941

<https://hal.science/hal-01462941v1>

Preprint submitted on 9 Feb 2017 (v1), last revised 5 Apr 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTIPLYING A GAUSSIAN MATRIX BY A GAUSSIAN VECTOR (AND THE GAUSS-LAPLACE TRANSMUTATION)

Pierre-Alexandre Mattei

pierre-alexandre.mattei@parisdescartes.fr

Laboratoire MAP5, UMR CNRS 8145

Université Paris Descartes

45 rue des Saints-Pères, 75006 Paris, France

Abstract

We provide a new simple characterization of the multivariate generalized Laplace distribution. In particular, our characterization implies that the product of a Gaussian matrix with independent and identically distributed columns and an independent isotropic Gaussian vector follows a *symmetric* multivariate generalized Laplace distribution.

1. Introduction

Wishart and Bartlett (1932) proved that the inner product of two independent bidimensional standard gaussian vectors follows a Laplace distribution. This result is deeply linked to the fact that the Laplace distribution can be represented as an infinite scale mixture of Gaussians with gamma mixing distribution. Specifically, if σ^2 follows a Gamma(1, 1/2) distribution and $x|\sigma \sim \mathcal{N}(0, \sigma^2)$, then x follows a standard Laplace distribution¹. This representation – which was recently named the *Gauss-Laplace transmutation* by Ding and Blitzstein (2015) following a blog post by Christian P. Robert² – is particularly useful if one wants to simulate a Laplace random variable: its use constitutes for example the cornerstone of the Gibbs sampling scheme for the Bayesian lasso of Park and Casella (2008).

In this short paper, we are interested in studying links between multivariate counterparts of these two characterizations. More specifically, we give a new simple characterization of the *multivariate generalized Laplace distribution* of Kotz, Kozubowski, and Podgórski (2001). As a corollary, we show that the product of a zero-mean Gaussian matrix with i.i.d. columns and a zero-mean isotropic Gaussian vector follows a symmetric multivariate generalized Laplace distribution, a result that has useful applications for Bayesian principal component analysis (Bouveyron, Latouche, and Mattei, 2016).

In the remainder of this paper, p and d are two positive integers.

2. The multivariate generalized Laplace distribution

While the definition of the univariate Laplace distribution is widely undisputed, there exist several different generalizations of this distribution to higher dimensions – a comprehensive review of such generalizations can be found in the monograph of Kotz, Kozubowski, and Podgórski (2001). In particular, Anderson (1992) introduced a zero-mean elliptically contoured p -dimensional Laplace distribution with univariate Laplace marginals by considering

1. The shape-rate parametrization of the gamma distribution is used through this paper. Note also that a standard Laplace distribution is centered with variance two.
2. <https://xianblog.wordpress.com/2015/10/14/gauss-to-laplace-transmutation/>

characteristic functions of the form

$$\forall \mathbf{u} \in \mathbb{R}^p, \phi(\mathbf{u}) = \frac{1}{1 + \frac{1}{2}\mathbf{u}^T \boldsymbol{\Sigma} \mathbf{u}},$$

where $\boldsymbol{\Sigma} \in \mathcal{S}_p^+$. This distribution was notably promoted by Eltoft, Kim, and Lee (2006) and is arguably the most popular multivariate generalization of the Laplace distribution (Kotz, Kozubowski, and Podgórski, 2001, p. 229). Among its advantages, this distribution can be slightly generalized to model skewness, by building on characteristic functions of the form

$$\forall \mathbf{u} \in \mathbb{R}^p, \phi(\mathbf{u}) = \frac{1}{1 + \frac{1}{2}\mathbf{u}^T \boldsymbol{\Sigma} \mathbf{u} - i\boldsymbol{\mu}^T \mathbf{u}},$$

where $\boldsymbol{\mu} \in \mathbb{R}^p$ accounts for asymmetry. Similarly to the univariate Laplace distribution, this asymmetric multivariate generalization is infinitely divisible (Kotz, Kozubowski, and Podgórski, 2001, p. 256). Therefore, it can be associated with a specific Lévy process (Kyprianou, 2014, p. 5), whose increments will follow yet another generalization of the Laplace distribution, the *multivariate generalized asymmetric Laplace distribution*. This distribution, introduced by Kotz, Kozubowski, and Podgórski (2001, p. 257) and further studied by Kozubowski, Podgórski, and Rychlik (2013), will be the cornerstone of our analysis of multivariate characterizations of Laplace and Gaussian distributions.

Definition 1 *A random variable $\mathbf{z} \in \mathbb{R}^p$ is said to have a **multivariate generalized asymmetric Laplace distribution** with parameters $s > 0$, $\boldsymbol{\mu} \in \mathbb{R}^p$ and $\boldsymbol{\Sigma} \in \mathcal{S}_p^+$ if its characteristic function is*

$$\forall \mathbf{u} \in \mathbb{R}^p, \phi_{\text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s)}(\mathbf{u}) = \left(\frac{1}{1 + \frac{1}{2}\mathbf{u}^T \boldsymbol{\Sigma} \mathbf{u} - i\boldsymbol{\mu}^T \mathbf{u}} \right)^s.$$

It is denoted by $\mathbf{z} \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s)$.

General properties of the generalized asymmetric Laplace distribution are discussed by Kozubowski, Podgórski, and Rychlik (2013). We list here a few useful ones.

Proposition 2 *Let $s > 0$, $\boldsymbol{\mu} \in \mathbb{R}^p$ and $\boldsymbol{\Sigma} \in \mathcal{S}_p^+$. If $\mathbf{z} \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s)$, we have $\mathbb{E}(\mathbf{z}) = s\boldsymbol{\mu}$ and $\text{Cov}(\mathbf{z}) = s(\boldsymbol{\Sigma} + \boldsymbol{\mu}\boldsymbol{\mu}^T)$. Moreover, if $\boldsymbol{\Sigma}$ is positive definite, the density of \mathbf{z} is given by*

$$\forall \mathbf{x} \in \mathbb{R}^p, f_{\mathbf{z}}(\mathbf{x}) = \frac{2e^{\boldsymbol{\mu}^T \boldsymbol{\Sigma}^{-1} \mathbf{x}}}{(2\pi)^{p/2} \Gamma(s) \sqrt{\det \boldsymbol{\Sigma}}} \left(\frac{Q(\mathbf{x})}{C(\boldsymbol{\Sigma}, \boldsymbol{\mu})} \right)^{s-p/2} K_{s-p/2}(Q(\mathbf{x})C(\boldsymbol{\Sigma}, \boldsymbol{\mu})),$$

where $Q_{\boldsymbol{\Sigma}}(\mathbf{x}) = \sqrt{\mathbf{x}^T \boldsymbol{\Sigma}^{-1} \mathbf{x}}$, $C(\boldsymbol{\Sigma}, \boldsymbol{\mu}) = \sqrt{2 + \boldsymbol{\mu}^T \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu}}$ and $K_{s-p/2}$ is the modified Bessel function of the second kind of order $s - p/2$.

Note that the $\text{GAL}_1(2b^2, 0, 1)$ case corresponds to a centered univariate Laplace distribution with scale parameter $b > 0$. In the symmetric case ($\boldsymbol{\mu} = 0$) and when $s = 1$, we recover the multivariate generalization of the Laplace distribution of Anderson (1992).

An appealing property of the multivariate generalized Laplace distribution is that it is also endowed with a multivariate counterpart of the Gauss-Laplace transmutation.

Theorem 3 (Generalized Gauss-Laplace transmutation) *Let $s > 0$, $\boldsymbol{\mu} \in \mathbb{R}^p$ and $\boldsymbol{\Sigma} \in \mathcal{S}_p^+$. If $u \sim \text{Gamma}(s, 1)$ and $\mathbf{x} \sim \mathcal{N}(0, \boldsymbol{\Sigma})$ is independent of u , we have*

$$u\boldsymbol{\mu} + \sqrt{u}\mathbf{x} \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s). \quad (1)$$

A proof of this result can be found in Kotz, Kozubowski, and Podgórski (2001, chap. 6). This generalized transmutation explains why the multivariate generalized Laplace distribution can also be seen as a multivariate generalization of the *variance-gamma distribution* which is widely used in the field of quantitative finance (Madan, Carr, and Chang, 1998). Infinite mixtures similar to (1) are called *variance-mean mixtures* (Barndorff-Nielsen, Kent, and Sørensen, 1982) and are discussed for example by Yu (2017).

Another useful property of the multivariate generalized Laplace distribution is that, under some conditions, it is closed under convolution.

Proposition 4 *Let $s_1, s_2 > 0$, $\boldsymbol{\mu} \in \mathbb{R}^p$ and $\boldsymbol{\Sigma} \in \mathcal{S}_p^+$. If $\mathbf{z}_1 \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_1)$ and $\mathbf{z}_2 \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_2)$ are independent random variables, then*

$$\mathbf{z}_1 + \mathbf{z}_2 \sim \text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_1 + s_2). \quad (2)$$

Proof Since \mathbf{z}_1 and \mathbf{z}_2 are independent, we have for all $\mathbf{u} \in \mathbb{R}^p$,

$$\phi_{\mathbf{z}_1 + \mathbf{z}_2}(\mathbf{u}) = \phi_{\text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_1)}(\mathbf{u})\phi_{\text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_2)}(\mathbf{u}) = \left(\frac{1}{1 + \frac{1}{2}\mathbf{u}^T \boldsymbol{\Sigma} \mathbf{u} - i\boldsymbol{\mu}^T \mathbf{u}} \right)^{s_1 + s_2}$$

which is the characteristic function of the $\text{GAL}_p(\boldsymbol{\Sigma}, \boldsymbol{\mu}, s_1 + s_2)$ distribution. ■

3. A new characterization involving a product between a Gaussian matrix and a Gaussian vector

We now state our main theorem, which gives a new characterization of multivariate generalized Laplace distributions with half-integer shape parameters.

Theorem 5 *Let \mathbf{W} be a $p \times d$ random matrix with i.i.d. columns following a $\mathcal{N}(0, \boldsymbol{\Sigma})$ distribution, $\mathbf{y} \sim \mathcal{N}(0, \mathbf{I}_d)$ be a Gaussian vector independent from \mathbf{W} and let $\boldsymbol{\mu} \in \mathbb{R}^p$. We have*

$$\mathbf{W}\mathbf{y} + \|\mathbf{y}\|_2^2 \boldsymbol{\mu} \sim \text{GAL}_p(2\boldsymbol{\Sigma}, 2\boldsymbol{\mu}, d/2). \quad (3)$$

Proof For each $k \in \{1, \dots, d\}$ let \mathbf{w}_k be the k -th column of \mathbf{W} , $u_k = y_k^2$ and $\boldsymbol{\xi}_k = y_k \mathbf{w}_k + y_k^2 \boldsymbol{\mu}$. To prove the theorem, we will prove that $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_d$ follow a GAL distribution and use the decomposition

$$\mathbf{W}\mathbf{y} + \|\mathbf{y}\|_2^2 \boldsymbol{\mu} = \sum_{k=1}^d \boldsymbol{\xi}_k.$$

Let $k \in \{1, \dots, d\}$. Since \mathbf{y} is standard Gaussian, $u_k = y_k^2$ follows a $\chi^2(1)$ distribution, or equivalently a $\text{Gamma}(1/2, 1/2)$ distribution. Therefore, $u_k/2 \sim \text{Gamma}(1/2, 1)$.

Moreover, note that $\sqrt{u_k}\mathbf{w}_k = |y_k|\mathbf{w}_k = y_k\text{sign}(y_k)\mathbf{w}_k \stackrel{d}{=} y_k\mathbf{w}_k$ since $|y_k|$ and $\text{sign}(y_k)$ are independent and $\text{sign}(y_k)\mathbf{w}_k \stackrel{d}{=} \mathbf{w}_k$. Therefore, according to the generalized Gauss-Laplace transmutation, we have

$$\boldsymbol{\xi}_k = \sqrt{\frac{u_k}{2}}\sqrt{2}\mathbf{w}_k + \frac{u_k}{2}2\boldsymbol{\mu} \sim \text{GAL}_p(2\boldsymbol{\Sigma}, 2\boldsymbol{\mu}, 1/2).$$

Since $\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_d$ are i.i.d. and following a $\text{GAL}_p(2\boldsymbol{\Sigma}, 2\boldsymbol{\mu}, 1/2)$ distribution, we can use Proposition 4 to conclude that

$$\mathbf{W}\mathbf{y} + \|\mathbf{y}\|_2^2\boldsymbol{\mu} = \sum_{k=1}^d \boldsymbol{\xi}_k \sim \text{GAL}_p(2\boldsymbol{\Sigma}, 2\boldsymbol{\mu}, d/2).$$

■

In the symmetric case ($\boldsymbol{\mu} = 0$), this result gives the distribution of the product between a Gaussian matrix with i.i.d. columns and a isotropic Gaussian vector.

Corollary 6 *Let \mathbf{W} be a $p \times d$ random matrix with i.i.d. columns following a $\mathcal{N}(0, \boldsymbol{\Sigma})$ distribution and let $\mathbf{y} \sim \mathcal{N}(0, \alpha\mathbf{I}_d)$ be a Gaussian vector independent from \mathbf{W} . Then*

$$\mathbf{W}\mathbf{y} \sim \text{GAL}_p(2\alpha\boldsymbol{\Sigma}, 0, d/2). \quad (4)$$

Moreover, if u is a standard Gamma variable with shape parameter $d/2$ and if $\mathbf{x} \sim \mathcal{N}(0, 2\alpha\boldsymbol{\Sigma})$ is a Gaussian vector independent of u , then

$$\mathbf{W}\mathbf{y} \stackrel{d}{=} \sqrt{u}\mathbf{x}. \quad (5)$$

Less general versions of Theorem 5 have been proved in the past, dating back to the derivation of the inner product of two i.i.d. standard Gaussian vectors by Wishart and Bartlett (1932). In particular, the unidimensional case ($p = 1$) was recently proven by Gaunt (2014) in order to obtain bounds for the convergence rate of random sums involving Gaussian products. The multivariate symmetric isotropic case ($\boldsymbol{\mu} = 0$ and $\boldsymbol{\Sigma}$ proportional to \mathbf{I}_p) was proven by Bouveyron, Latouche, and Mattei (2016) in order to derive the marginal likelihood of the noiseless probabilistic PCA model of Roweis (1998). While the proof of Bouveyron, Latouche, and Mattei (2016) relied on characteristic functions and the properties of Bessel functions, the proof that we presented here is closer in spirit to the one of Gaunt (2014), based on representations of distributions.

Acknowledgements

I thank Charles Bouveyron, Pierre Latouche and Christian P. Robert for fruitful advices and discussions. Part of this work was made during a visit to University College Dublin, funded by the Fondation Sciences Mathématiques de Paris (FSMP).

References

- D. N. Anderson. A multivariate Linnik distribution. *Statistics & Probability Letters*, 14(4): 333–336, 1992.
- O. Barndorff-Nielsen, J. Kent, and M. Sørensen. Normal variance-mean mixtures and z distributions. *International Statistical Review/Revue Internationale de Statistique*, pages 145–159, 1982.
- C. Bouveyron, P. Latouche, and P.-A. Mattei. Bayesian variable selection for globally sparse probabilistic PCA. Technical report, HAL-01310409, 2016.
- P. Ding and J. K. Blitzstein. Representation for the Gauss-Laplace transmutation. *arXiv preprint arXiv:1510.08765*, 2015.
- T. Eltoft, T. Kim, and T.-W. Lee. On the multivariate Laplace distribution. *IEEE Signal Processing Letters*, 13(5):300–303, 2006.
- R. E. Gaunt. Variance-gamma approximation via Stein’s method. *Electronic Journal of Probability*, 19(38):1–33, 2014.
- S. Kotz, T. Kozubowski, and K. Podgórski. *The Laplace distribution and generalizations: a revisit with applications to communications, economics, engineering, and finance*. Number 183. Springer Science & Business Media, 2001.
- T. Kozubowski, K. Podgórski, and I. Rychlik. Multivariate generalized Laplace distribution and related random fields. *Journal of Multivariate Analysis*, 113:59–72, 2013.
- A. Kyprianou. *Fluctuations of Lévy processes with applications: Introductory Lectures*. Springer Science & Business Media, 2014.
- D. B. Madan, P. P. Carr, and E. C. Chang. The variance gamma process and option pricing. *European finance review*, 2(1):79–105, 1998.
- T. Park and G. Casella. The Bayesian lasso. *Journal of the American Statistical Association*, 103(482):681–686, 2008.
- S. Roweis. EM algorithms for PCA and SPCA. *Advances in neural information processing systems*, pages 626–632, 1998.
- J. Wishart and M. S. Bartlett. The distribution of second order moment statistics in a normal system. *Mathematical Proceedings of the Cambridge Philosophical Society*, 28, 10 1932.
- Y. Yu. On normal variance–mean mixtures. *Statistics & Probability Letters*, 121:45–50, 2017.