



**HAL**  
open science

## Geometric side of a local relative trace formula

Patrick Delorme, Pascale Harinck, Sofiane Souaifi

► **To cite this version:**

Patrick Delorme, Pascale Harinck, Sofiane Souaifi. Geometric side of a local relative trace formula. Transactions of the American Mathematical Society, 2019, 371 (3), pp.1815-1857. 10.1090/tran/7360 . hal-01467988

**HAL Id: hal-01467988**

**<https://hal.science/hal-01467988>**

Submitted on 15 Feb 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Geometric side of a local relative trace formula

P. Delorme\*, P. Harinck, S. Souaifi

## Abstract

Following a scheme suggested by B. Feigon, we investigate a local relative trace formula in the situation of a reductive  $p$ -adic group  $G$  relative to a symmetric subgroup  $H = \underline{H}(F)$  where  $\underline{H}$  is split over the local field  $F$  of characteristic zero and  $G = \underline{G}(F)$  is the restriction of scalars of  $\underline{H}_{/E}$  relative to a quadratic unramified extension  $E$  of  $F$ . We adapt techniques of the proof of the local trace formula by J. Arthur in order to get a geometric expansion of the integral over  $H \times H$  of a truncated kernel associated to the regular representation of  $G$ .

*Mathematics Subject Classification 2000:* 11F72, 22E50.

*Keywords and phrases:*  $p$ -adic reductive groups, symmetric spaces, local relative trace formula, truncated kernel, orbital integrals.

## Contents

<b>Introduction</b>	<b>2</b>
<b>1 Preliminaries</b>	<b>8</b>
1.1 Reductive $p$ -adic groups	8
1.2 The symmetric space $H \backslash G$	12
1.3 Weyl integration formula and orbital integrals	14
<b>2 Geometric side of the local relative trace formula</b>	<b>21</b>
2.1 Truncation	21
2.2 The truncated kernel	24
2.3 Preliminaries to estimates	27
2.4 Proof of Theorem 2.3	33
2.5 The function $J^T(f)$	41

---

\*The first author was supported by a grant of Agence Nationale de la Recherche with reference ANR-13-BS01-0012 FERPLAY.

## Introduction

In this article, we investigate a local relative trace formula in the situation of  $p$ -adic groups relative to a symmetric subgroup. This work is inspired by the recent results of B. Feigon ([F]), where she investigated what she called a local relative trace formula on  $PGL(2)$  and a local Kuznetsov trace formula for  $U(2)$ .

Before we describe our setting and results, we would to explain on the toy model of finite groups the framework of the formulas of B. Feigon. We even start with the more general framework of the relative trace formula initiated by H. Jacquet ([J], see also [O] for an account of some applications of this relative trace formula).

Let  $G$  be a finite group and let  $H, H', \Gamma$  be subgroups of  $G$ . We endow any finite set with the counting measure. We denote by  $r$  the right regular representation of  $G$  on  $L^2(\Gamma \backslash G)$  and we consider the  $H$ -fixed linear form  $\xi$  on  $L^2(\Gamma \backslash G)$  defined by

$$\xi = \sum_{h \in H \cap \Gamma \backslash H} \delta_{\Gamma h} \tag{0.1}$$

where  $\delta_{\Gamma h}$  is the Dirac measure of the coset  $\Gamma h$ , or in other words

$$\xi(\psi) = \int_{H \cap \Gamma \backslash H} \psi(\Gamma h) dh, \quad \psi \in L^2(\Gamma \backslash G).$$

We define similarly  $\xi'$  relative to  $H'$ .

We view  $\xi, \xi'$  as elements of  $L^2(\Gamma \backslash G)$  and we form the coefficient  $c_{\xi, \xi'}(g) = (r(g)\xi, \xi')$ . Integrating over functions on  $G$ , it defines a "distribution"  $\Theta$  on  $G$  which is right invariant by  $H$  and left invariant by  $H'$ . The relative trace formula in this context gives two expressions of  $\Theta(f)$  for  $f$  a function on  $G$ , the first one, called the geometric side, in terms of orbital integrals, and the second one, called the spectral side, in terms of irreducible representations of  $G$ .

First we deal with the geometric side. For this purpose we introduce suitable orbital integrals. For  $\gamma \in \Gamma$ , we set  $[\gamma] := (H' \cap \Gamma)\gamma(H \cap \Gamma)$  and one introduces two subgroups of  $H' \times H$

$$(H' \times H)_\gamma = \{(h', h) | h'\gamma h^{-1} = \gamma\}, (H' \cap \Gamma \times H \cap \Gamma)_\gamma = (H' \times H)_\gamma \cap (\Gamma \times \Gamma).$$

Then, we define the orbital integral of a function  $f$  on  $G$  by

$$I([\gamma], f) = \int_{(H' \times H)_\gamma \backslash (H' \times H)} f(h'\gamma h^{-1}) dh' dh.$$

Let  $f$  be a function on  $G$ . Since  $r(g)\delta_{\Gamma h} = \delta_{\Gamma hg^{-1}}$ , the definition of  $\xi$  and  $\xi'$  gives

$$\Theta(f) = \sum_{g \in G} f(g) \Theta(g) = \sum_{g \in G} f(g) \frac{1}{\text{vol}(\Gamma \cap H)} \frac{1}{\text{vol}(\Gamma \cap H')} \sum_{h \in H} \sum_{h' \in H'} (\delta_{\Gamma h g^{-1}}, \delta_{\Gamma h'}).$$

Changing  $g$  in  $g^{-1}h$  and using the fact that  $(\delta_{\Gamma g}, \delta_{\Gamma h'})$  is equal to 1 for  $g \in \Gamma h'$  and to zero otherwise, one gets

$$\Theta(f) = \frac{1}{\text{vol}(\Gamma \cap H)} \frac{1}{\text{vol}(\Gamma \cap H')} \sum_{h \in H} \sum_{h' \in H'} \sum_{\gamma \in \Gamma} f(h' \gamma h). \quad (0.2)$$

A simple computation of volumes leads to the geometric expression of  $\Theta$  in terms of orbital integrals

$$\Theta(f) = \sum_{[\gamma] \in H' \cap \Gamma \backslash \Gamma / \Gamma \cap H} \text{vol}((H' \cap \Gamma \times H \cap \Gamma)_{\gamma} \backslash (H' \times H)_{\gamma}) I([\gamma], f). \quad (0.3)$$

Let us turn to the spectral side. We decompose  $L^2(\Gamma \backslash G)$  into isotypic components  $\oplus_{\pi \in \hat{G}} \mathcal{H}_{\pi}$ . The restriction of  $\xi$  and  $\xi'$  to  $\mathcal{H}_{\pi}$  will be denoted  $\xi_{\pi}$  and  $\xi'_{\pi}$  respectively. The spectral formula for  $\Theta$  is the simple equality

$$\Theta = \sum_{\pi \in \hat{G}} c_{\xi_{\pi}, \xi'_{\pi}}. \quad (0.4)$$

Notice that it might be also interesting to decompose further the representation into irreducible representations and the restriction of  $\xi$  to each of them will be called a period.

There is a third interpretation of the distribution  $\Theta$ . If  $f$  is a function on  $G$ , then the operator  $r(f)$  on  $L^2(\Gamma \backslash G)$  is an integral operator whose kernel  $K_f$  is the function on  $\Gamma \backslash G \times \Gamma \backslash G$  given by

$$K_f(x, y) = \sum_{\gamma \in \Gamma} f(x^{-1} \gamma y).$$

By (0.2), one gets easily the following expression of  $\Theta(f)$

$$\Theta(f) = \int_{(H' \cap \Gamma \backslash H') \times (H \cap \Gamma \backslash H)} K_f(h', h) dh' dh. \quad (0.5)$$

This point of view is probably the best one. But it is important to have the representation theoretic meaning of  $\Theta$ .

The toy model for the local relative trace formula of B. Feigon appears as a particular case of the above relative trace formula. In that case, the groups  $G$ ,  $H$  and  $H'$  are products  $G_1 \times G_1$ ,  $H_1 \times H_1$  and  $H'_1 \times H'_1$  respectively and  $\Gamma$  is the diagonal of  $G_1 \times G_1$ . Then  $\Gamma \backslash G$  identifies with  $G_1$  and the right representation corresponds

to the representation  $R$  of  $G_1 \times G_1$  on  $L^2(G_1)$  given by  $[R(x, y)\phi](g) = \phi(x^{-1}gy)$ . Then, we have

$$\xi(\psi) = \int_{H_1} \psi(h)dh, \quad \psi \in L^2(G_1).$$

The spectral side is more concrete. If  $(\pi_1, \mathcal{H}_{\pi_1})$  is an irreducible unitary representation of  $G_1$  then  $G_1 \times G_1$  acts on  $\text{End}(\mathcal{H}_{\pi_1})$  by an irreducible representation denoted by  $\pi$ . It is unitary if we use the scalar product associated to the Hilbert-Schmidt norm. Moreover  $L^2(G_1)$  is canonically isomorphic to the direct sum  $\bigoplus_{\pi_1 \in \hat{G}_1} \text{End}(\mathcal{H}_{\pi_1})$ . Let  $P_\pi$  be the orthogonal projector onto the space of invariant vectors under  $H_1$ . Then, the period map  $\xi_\pi$ , which is a linear form on  $\text{End}(\mathcal{H}_{\pi_1})$ , is given by

$$\xi_\pi(T) = \int_{H_1} \text{Tr}(\pi_1(h)T)dh = (T, P_\pi), \quad T \in \text{End}(\mathcal{H}_{\pi_1}).$$

One further decomposes  $\xi_\pi$  by using an orthonormal basis  $(\eta_{\pi_1, i})$  of the space of  $H_1$ -invariant vectors. We will use the identification of  $\text{End}(\mathcal{H}_{\pi_1})$  with the tensor product of  $\mathcal{H}_{\pi_1}$  with its conjugate complex vector space. In this identification, one has

$$P_\pi = \sum_i \eta_{\pi_1, i} \otimes \eta_{\pi_1, i}.$$

We define similar notations for  $\xi'$  relative to  $H'$ . Then, for two functions  $f_1, f_2$  on  $G_1$ , the spectral side (0.4) can be written

$$\Theta(f_1 \otimes f_2) = \sum_{\pi_1 \in \hat{G}_1} \sum_{i, i'} c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_1) c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_2).$$

For the geometric side, we define the integral orbital of a function  $f$  on  $G_1$  by

$$I(g, f) = \int_{(H'_1 \times H_1)_g \backslash H'_1 \times H_1} f(h'gh^{-1})dhdh'$$

which depends only on the double coset  $H'_1 g H_1$ . Then one gets by (0.3) the equality

$$\Theta(f_1 \otimes f_2) = \sum_{g \in H'_1 \backslash G_1 / H_1} v(g) I(g, f_1) I(g, f_2)$$

where the  $v(g)$ 's are positive constants depending on volumes. Hence the final form of the local relative trace formula is:

$$\sum_{g \in H'_1 \backslash G_1 / H_1} v(g) I(g, f_1) I(g, f_2) = \sum_{\pi_1 \in \hat{G}_1} \sum_{i, i'} c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_1) c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_2).$$

This formula allows to invert the orbital integrals  $I(g, f_1)$ . For this purpose, one chooses  $g_1 \in G_1$  and takes for  $f_2$  the Dirac measure at  $g_1$ . Then  $I(g_1, f_2) = 1$  and the other orbital integrals of  $f_2$  are zero. Hence

$$v(g_1) I(g_1, f_1) = \sum_{\pi_1 \in \hat{G}_1} \sum_{i, i'} c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_1) c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_2).$$

In order to make the formula more precise, one needs to compute the constants  $c_{\eta_{\pi_1, i}, \eta'_{\pi_1, i'}}(f_2)$ .

The inversion of orbital integrals is one of our motivations to investigate a local relative trace formula in the situation of  $p$ -adic groups relative to a symmetric subgroup  $H$  and we will take  $H = H'$ .

In this article, we consider a reductive algebraic group  $\underline{H}$  defined over a non archimedean local field  $F$  of characteristic 0. We fix a quadratic unramified extension  $E$  of  $F$  and we consider the group  $\underline{G} := \text{Res}_{E/F}\underline{H}$  obtained by restriction of scalars of  $\underline{H}$ , where here  $\underline{H}$  is considered as a group defined over  $E$ . We denote by  $H$  and  $G$  the group of  $F$ -points of  $\underline{H}$  and  $\underline{G}$  respectively. Then  $G$  is isomorphic to  $\underline{H}(E)$  and  $H$  appears as the fixed points of  $G$  under the involution of  $G$  induced by the nontrivial element of the Galois group of  $E/F$ . We assume that  $\underline{H}$  is split over  $F$  and we fix a maximal split torus  $A_0$  of  $H$ . The groups  $G$  and  $H$  correspond to  $G_1$  and  $H_1 = H'_1$  respectively in our example of a local relative trace formula for finite groups.

The starting point of our study is the analogue to the expression (0.5). We consider the regular representation  $R$  of  $G \times G$  on  $L^2(G)$  given by  $(R(g_1, g_2)\psi)(x) = \psi(g_1^{-1}xg_2)$ . Then for  $f = f_1 \otimes f_2$  where  $f_1$  and  $f_2$  are two smooth compactly supported functions on  $G$ , the corresponding operator  $R(f)$  is an integral operator on  $L^2(G)$  with smooth kernel

$$K_f(x, y) = \int_G f_1(xg)f_2(gy)dg = \int_G f_1(g)f_2(x^{-1}gy)dg.$$

As  $H$  may be not compact, even modulo the split component  $A_H$  of the center of  $H$ , we have to truncate this kernel to integrate it. We multiply this kernel by a product of functions  $u(x, T)u(y, T)$  where  $u(\cdot, T)$  is the characteristic function of a large compact subset in  $A_H \backslash H$  depending on a parameter  $T \in a_0 = \text{Rat}(A_0) \otimes_{\mathbb{Z}} \mathbb{R}$  ( $\text{Rat}(A_0)$  is the group of  $F$ -rational characters of  $A_0$ ) as in [Ar3] (cf. (2.7)). As  $H$  is split, we have  $A_H = A_G$ . Hence the kernel  $K_f$  is left invariant by the diagonal  $\text{diag}(A_H)$  of  $A_H$  and we can integrate the truncated kernel over  $\text{diag}(A_H) \backslash H \times H$ . We set

$$K^T(f) := \int_{\text{diag}(A_H) \backslash (H \times H)} K_f(x_1, x_2)u(x_1, T)u(x_2, T)d\overline{(x_1, x_2)}.$$

In [Ar3], J. Arthur studies the integral of  $K_f(x, x)u(x, T)$  over  $A_G \backslash G$  to obtain its local trace formula on reductive groups.

We study the geometric expression of the distribution  $K^T(f)$  and its dependence on the parameter  $T$ . Our main results (Theorem 2.3 and Corollary 2.11) assert that  $K^T(f)$  is asymptotic as  $T$  approaches infinity to another distribution  $J^T(f)$  of the form

$$J^T(f) = \sum_{k=0}^N p_{\xi_k}(T, f)e^{\xi_k(T)} \tag{0.6}$$

where  $\xi_0 = 0, \dots, \xi_N$  are distinct points of the dual space  $ia_0^*$  and each  $p_{\xi_k}(T, f)$  is a polynomial function in  $T$ . Moreover, the constant term  $\tilde{J}(f) := p_0(0, f)$  of  $J^T(f)$  is well-defined and uniquely determined by  $K^T(f)$ . We give an explicit expression of this constant term in terms of weighted orbital integrals.

These results are analogous to those of [Ar3] for the group case. Our proof follows closely the study by J. Arthur of the geometric side of his local trace formula which we were able to adapt under our assumptions to the case of double truncations.

In the first section, we introduce notation on groups and on symmetric spaces according to [RR]. The starting point of our study is the Weyl integration formula established in [RR], which takes into account the  $(H, H)$ -double classes of  $\sigma$ -regular elements of  $G$  (cf. (1.30) and (1.32)). These double classes are express in terms of  $\sigma$ -torus, that is torus whose elements are anti-invariant by  $\sigma$ . Under our assumptions, there is a bijective correspondence  $S \rightarrow S_\sigma$  between maximal tori of  $H$  and maximal  $\sigma$ -tori of  $G$  which preserves  $H$ -conjugacy classes.

Then the Weyl integration formula can be written in terms of Levi subgroups  $M \in \mathcal{L}(A_0)$  of  $H$  containing  $A_0$  and  $M$ -conjugacy classes of maximal anisotropic tori of  $M$  (cf. (1.33)):

$$\int_G f(g)dg = \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{diag(A_M) \backslash H \times H} f(h^{-1} x_m \gamma l) d(\overline{h, l}) d\gamma$$

where  $\kappa_S$  is a finite subset of  $G$ ,  $c_M$  and  $c_{S, x_m}$  are positive constants,  $\mathcal{T}_M$  is a suitable set of anisotropic tori of  $M$  and  $\Delta_\sigma$  is a jacobian.

A fundamental result for our proofs concerns the orbital integral  $\mathcal{M}(f)$  of a compactly smooth function  $f$  on  $G$ . It is defined on  $\sigma$ -regular points by

$$\mathcal{M}(f)(x_m \gamma) = |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/4} \int_{diag(A_S) \backslash H \times H} f(h^{-1} x_m \gamma l) d(\overline{h, l}),$$

where  $S$  is a maximal torus of  $H$ ,  $x_m \in \kappa_S$  and  $\gamma \in S_\sigma$  such that  $x_m \gamma$  is  $\sigma$ -regular. As in the group case using the exponential map and the property that each root of  $S_\sigma$  has multiplicity 2 in the Lie algebra of  $G$ , we prove that the orbital integral is bounded on the subset of  $\sigma$ -regular points of  $G$  (cf. Theorem 1.2).

In the second section, we explain the truncation process based on the notion of  $(H, M)$ -orthogonal sets and prove our main results. Using the Weyl integration formula, we can write

$$K^T(f) = \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} K^T(x_m, \gamma, f) d\gamma$$

where

$$K^T(x_m, \gamma, f) = |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{diag(A_M) \backslash H \times H} \int_{diag(A_M) \backslash H \times H} f_1(y_1^{-1} x_m \gamma y_2)$$

$$\times f_2(x_1^{-1}x_m\gamma x_2)u_M(x_1, y_1, x_2, y_2, T)d(x_1, x_2)d(y_1, y_2)$$

and

$$u_M(x_1, y_1, x_2, y_2, T) = \int_{A_H \backslash A_M} u(y_1^{-1}ax_1, T)u(y_2^{-1}ax_2, T)da.$$

The function  $J^T(f)$  is obtained in a similar way to  $K^T(f)$  where we replace the weight function  $u_M(x_1, y_1, x_2, y_2, T)$  by another weight function  $v_M(x_1, y_1, x_2, y_2, T)$ .

The weight function  $v_M$  is given by

$$v_M(x_1, y_1, x_2, y_2, T) := \int_{A_H \backslash A_M} \sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T))da$$

where  $\sigma_M(\cdot, \mathcal{Y})$  is the function of [Ar3] depending on a  $(H, M)$ -orthogonal set  $\mathcal{Y}$  and  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  is a  $(H, M)$ -orthogonal set obtained as the "minimum" of two  $(H, M)$ -orthogonal sets  $\mathcal{Y}_M(x_1, y_1, T)$  and  $\mathcal{Y}_M(x_2, y_2, T)$  (cf. (2.4), Lemma 2.2 and (2.11)). If  $\mathcal{Y}_1$  and  $\mathcal{Y}_2$  are two  $(H, M)$ -orthogonal positive sets then the "minimum"  $\mathcal{Z}$  of  $\mathcal{Y}_1$  and  $\mathcal{Y}_2$  satisfies the property that the convex hull  $\mathcal{S}_M(\mathcal{Z})$  in  $a_H \backslash a_M$  of the points of  $\mathcal{Z}$  is the intersection of the convex hulls  $\mathcal{S}_M(\mathcal{Y}_1)$  and  $\mathcal{S}_M(\mathcal{Y}_2)$  in  $a_H \backslash a_M$  of the points of  $\mathcal{Y}_1$  and  $\mathcal{Y}_2$  respectively.

If  $\|T\|$  is large relative to  $\|x_i\|, \|y_i\|, i = 1, 2$  then  $\sigma_M(\cdot, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T))$  is just the characteristic function of  $\mathcal{S}_M(\mathcal{Y}_M(x_1, y_1, x_2, y_2, T))$ . In that case, this function is equal to the product of  $\sigma_M(\cdot, \mathcal{Y}_M(x_1, y_1, T))$  and  $\sigma_M(\cdot, \mathcal{Y}_M(x_2, y_2, T))$ .

Our proofs consist to establish good estimates of  $|u_M((x_1, y_1, x_2, y_2, T) - v_M(x_1, y_1, x_2, y_2, T)|$  when  $x_i, y_i, i = 1, 2$  satisfy  $f_1(y_1^{-1}x_m\gamma y_2)f_1(x_1^{-1}x_m\gamma x_2) \neq 0$  for some  $\gamma \in S_\sigma$  and  $x_m \in \kappa_S$ . Then, using that orbital integrals are bounded, we deduce our result on  $|K^T(f) - J^T(f)|$ .

This work is a first step towards a local relative trace formula. For the spectral side, we have to prove that  $K^T(f)$  is asymptotic to a distribution  $k^T(f)$  which is of general form (0.6) and constructed from spectral data. We hope that we can express the constant term of  $k^T(f)$  in terms of regularized local period integrals introduced by B. Feigon in [F] in the same way than Jacquet-Lapid-Rogawski regularized period integrals for automorphic forms in [JLR]. We plan to explicit such a local relative trace formula for  $PGL(2)$ .

**Acknowledgments.** We thank warmly Bertrand Lemaire for his answers to our many questions on algebraic groups. We thank Bertrand Rémy and David Renard for our helpful discussions. We thank also Guy Henniart for providing us a proof of (1.5).



# 1 Preliminaries

## 1.1 Reductive $p$ -adic groups

Let  $F$  be a non archimedean local field of characteristic 0 and odd residual characteristic  $q$ . Let  $|\cdot|_F$  denote the normalized valuation on  $F$ .

For an algebraic variety  $\underline{M}$  defined over  $F$ , we identify  $\underline{M}$  with  $\underline{M}(\overline{F})$  where  $\overline{F}$  is an algebraic closure of  $F$  and we set  $M := \underline{M}(F)$ .

We will use conventions like in [W2]. One considers various algebraic groups  $\underline{J}$  defined over  $F$ , and sentences like

” let  $M$  be an algebraic group” will mean ” let  $M$  be the  $F$ -points of an algebraic group  $\underline{M}$  defined over  $F$ ” and ” let  $A$  be a split torus ” will mean ” let  $A$  be the group of  $F$ -points of a torus,  $\underline{A}$ , defined and split over  $F$  .” (1.1)

If  $J$  is an algebraic group, one denotes by  $\text{Rat}(J)$  the group of its rational characters defined over  $F$ . If  $V$  is a vector space,  $V^*$  will denote its dual. If  $V$  is real,  $V_{\mathbb{C}}$  will denote its complexification.

Let  $\underline{G}$  be an algebraic reductive group defined over  $F$ . We fix a maximal split torus  $A_0$  of  $G$  and we denote by  $M_0$  its centralizer in  $G$ .

We denote by  $A_G$  the maximal split torus of the center of  $G$  and we define

$$a_G := \text{Hom}_{\mathbb{Z}}(\text{Rat}(G), \mathbb{R}).$$

One has the canonical map  $h_G : G \rightarrow a_G$  which is defined by

$$e^{\langle h_G(x), \chi \rangle} = |\chi(x)|_F, \quad x \in G, \chi \in \text{Rat}(G). \quad (1.2)$$

The restriction of rational characters from  $G$  to  $A_G$  induces an isomorphism

$$\text{Rat}(G) \otimes_{\mathbb{Z}} \mathbb{R} \simeq \text{Rat}(A_G) \otimes_{\mathbb{Z}} \mathbb{R}. \quad (1.3)$$

Notice that  $\text{Rat}(A_G)$  appears as a generating lattice in the dual space  $a_G^*$  of  $a_G$  and

$$a_G^* \simeq \text{Rat}(G) \otimes_{\mathbb{Z}} \mathbb{R}. \quad (1.4)$$

The kernel of  $h_G$ , which is denoted by  $G^1$ , is the intersection of the kernels of  $|\chi|_F$  for all character  $\chi \in \text{Rat}(G)$  of  $G$ . The groupe  $G^1$  is distinguished in  $G$  and contains the derived group  $G_{der}$  of  $G$ . Moreover, it is well-known that

the group  $G^1$  is generated by the compact subgroups of  $G$ . (1.5)

G. Henniart has communicated to us an unpublished proof of this result by N. Abe, F. Herzig, G. Henniart and M.F. Vigneras.

One denotes by  $a_{G,F}$  (resp.,  $\tilde{a}_{G,F}$ ) the image of  $G$  (resp.,  $A_G$ ) by  $h_G$ .  
Then  $G/G^1$  is isomorphic to the lattice  $a_{G,F}$ . (1.6)

If  $P$  is a parabolic subgroup of  $G$  with Levi subgroup  $M$ , we keep the same notation with  $M$  instead of  $G$ .

The inclusions  $A_G \subset A_M \subset M \subset G$  determine a surjective morphism  $a_{M,F} \rightarrow a_{G,F}$  (resp., an injective morphism,  $\tilde{a}_{G,F} \rightarrow \tilde{a}_{M,F}$ ) which extends uniquely to a surjective linear map  $h_{MG}$  from  $a_M$  to  $a_G$  (resp., injective map between  $a_G$  and  $a_M$ ). The second map allows to identify  $a_G$  with a subspace of  $a_M$  and the kernel of the first one,  $a_M^G$ , satisfies

$$a_M = a_M^G \oplus a_G. \tag{1.7}$$

For  $M = M_0$ , we set  $a_0 := a_{M_0}$  and  $a_0^G := a_{M_0}^G$ . We fix a scalar product  $(\cdot, \cdot)$  on  $a_0$  which is invariant under the Weyl group  $W(G, A_0)$  of  $(G, A_0)$ . Then  $a_G$  identifies with the fixed point set of  $a_0$  by  $W(G, A_0)$  and  $a_0^G$  is an invariant subspace of  $a_0$  under  $W(G, A_0)$ . Hence, it is the orthogonal subspace to  $a_G$  in  $a_0$ . The space  $a_G^*$  might be viewed as a subspace of  $a_0^*$  by (1.7). Moreover, by definition of the surjective map  $a_0 \rightarrow a_G$ , one deduces that

$$\text{if } m_0 \in M_0 \text{ then } h_G(m_0) \text{ is the orthogonal projection of } h_{M_0}(m_0) \text{ onto } a_G. \tag{1.8}$$

From (1.7) applied to  $(M, M_0)$  instead of  $(G, M)$ , one obtains a decomposition  $a_0 = a_0^M \oplus a_M$ . From the  $W(G, A_0)$  invariance of the scalar product, one gets:

$$\begin{aligned} &\text{The decomposition } a_0 = a_0^M \oplus a_M \text{ is an orthogonal decomposition.} \\ &\text{The space } a_M^* \text{ appears as a subspace of } a_0^* \text{ and, in the identification of } a_0 \text{ with } a_0^* \text{ given by the scalar product, } a_M^* \text{ identifies with } a_M. \end{aligned} \tag{1.9}$$

The decomposition  $a_M = a_M^G \oplus a_G$  is orthogonal relative to the restriction to  $a_M$  of the  $W(G, A_0)$ -invariant inner product on  $a_0$  and the natural map  $h_{MG}$  is identified with the orthogonal projection of  $a_M$  onto  $a_G$ .

$$\text{In particular, } a_{G,F} \text{ is the orthogonal projection of } a_{M,F} \text{ onto } a_G. \text{ Moreover, we have } \tilde{a}_{G,F} = a_G \cap \tilde{a}_{M,F} \text{ (cf. [Ar3] (1.4)).} \tag{1.10}$$

By a Levi subgroup of  $G$ , we mean a group  $M$  containing  $M_0$  which is the Levi component of a parabolic subgroup of  $G$ . If  $P$  is a parabolic subgroup containing  $M_0$  then it has a unique Levi subgroup denoted by  $M_P$  which contains  $M_0$ . We will denote by  $N_P$  the unipotent radical of  $P$ .

For a Levi subgroup  $M$ , we write  $\mathcal{L}(M)$  for the finite set of Levi subgroups of  $G$  which contain  $M$  and we also let  $\mathcal{P}(M)$  denote the finite set of parabolic subgroups  $P$  with  $M_P = M$ .

Let  $K$  be the fixator of a special point in the apartment of  $A_0$  in the Bruhat-Tits building. We have the Cartan decomposition

$$G = KM_0K. \quad (1.11)$$

If  $P = M_P N_P$  is a parabolic subgroup of  $G$  containing  $M_0$ , then

$$G = PK = M_P N_P K. \quad (1.12)$$

If  $x \in G$ , we can write

$$x = m_P(x)n_P(x)k_P(x), \quad m_P(x) \in M_P, n_P(x) \in N_P, k_P(x) \in K. \quad (1.13)$$

We set

$$h_P(x) := h_{M_P}(m_P(x)). \quad (1.14)$$

The point  $m_P(x)$  is defined up an element of  $K$  but  $h_P(x)$  does not depend of this choice.

We introduce a norm  $\|\cdot\|$  on  $G$  as in ([W2] §I.1.) (called height function in ([Ar3])). Let  $\Lambda_0 : G \rightarrow GL_n(\mathbb{F})$  be an algebraic embedding. For  $g \in G$ , we write

$$\Lambda_0(g) = (a_{i,j})_{i,j=1\dots n}, \quad \Lambda_0(g^{-1}) = (b_{i,j})_{i,j=1\dots n}.$$

We set

$$\|g\| := \sup_{i,j} \sup(|a_{i,j}|_{\mathbb{F}}, |b_{i,j}|_{\mathbb{F}}). \quad (1.15)$$

If  $\Lambda : G \rightarrow GL_d(\mathbb{F})$  is another algebraic embedding then the norm  $\|\cdot\|_{\Lambda}$  attached to  $\Lambda$  as above is equivalent to  $\|\cdot\|$  in the following sense: there are a positive constant  $C_{\Lambda}$  and a positive integer  $d_{\Lambda}$  such that

$$\|g\|_{\Lambda} \leq C_{\Lambda} \|g\|^{d_{\Lambda}}.$$

This allows us to use results of [W2] for estimates on norms.

The following properties of  $\|\cdot\|$  are immediate consequences of definition:

$$1 \leq \|x\| = \|x^{-1}\|, \quad x \in G, \quad (1.16)$$

$$\|xy\| \leq \|x\| \|y\|, \quad x, y \in G. \quad (1.17)$$

In order to have estimates, we introduce the following notation. Let  $r$  be a positive integer. Let  $f$  and  $g$  be two positive functions defined over a subset  $W$  of  $G^r$ .

$$\text{We write } f(x) \leq g(x), x \in W \text{ if and only if there are a positive constant } c \text{ and a positive integer } d \text{ such that } f(x) \leq cg(x)^d \text{ for all } x \in W. \quad (1.18)$$

We write  $f(x) \approx g(x), x \in W$  if  $f(x) \leq g(x), x \in W$  and  $g(x) \leq f(x), x \in W$ . (1.19)

If  $f_1, f_2$  and  $f_3$  are positive functions on  $G^r$ , we clearly have

if  $f_1(x) \leq f_2(x), x \in W$  and  $f_2(x) \leq f_3(x), x \in W$  then  $f_1(x) \leq f_3(x), x \in W$ ,  
if  $f_1(x) \approx f_2(x), x \in W$  and  $f_2(x) \approx f_3(x), x \in W$  then  $f_1(x) \approx f_3(x), x \in W$ .

Moreover, if  $f_1, f_2, g_1$  and  $g_2$  are positive functions on  $G^r$  which take values greater or equal to 1, we obtain easily the following properties:

1. for all positive integer  $d$ , we have  $f_1(x) \approx f_1(x)^d, x \in W$ ,
2. if  $f_1(x) \leq g_1(x), x \in W$  and  $f_2(x) \leq g_2(x), x \in W$  then  
 $(f_1 f_2)(x) \leq (g_1 g_2)(x), x \in W$ , (1.20)
3. if  $f_1(x) \approx g_1(x), x \in W$  and  $f_2(x) \approx g_2(x), x \in W$  then  
 $(f_1 f_2)(x) \approx (g_1 g_2)(x), x \in W$ .

Since  $\|x\| = \|xyy^{-1}\| \leq \|xy\|\|y\|$  and  $\|xy\| \leq \|x\|\|y\|$ , we obtain

If  $\Omega$  is a compact subset of  $G$ , then  $\|x\| \approx \|x\omega\|, x \in G, \omega \in \Omega$ . (1.21)

Let  $P = M_P N_P$  be a parabolic subgroup of  $G$  containing  $M_0$ . Then, each  $x \in G$  can be written  $x = m_P(x)n_P(x)k$  where  $m_P(x) \in M_P, n_P(x) \in N_P$  and  $k \in K$ . By ([W2] Lemma II.3.1), we have

$$\|m_P(x)\| + \|n_P(x)\| \leq \|x\|, x \in G. (1.22)$$

Recall that  $G^1$  is the kernel of  $h_G : G \rightarrow a_G$ . Let us prove that

$$\|xa\| \approx \|x\|\|a\|, x \in G^1, a \in A_G. (1.23)$$

According to the Cartan decomposition (1.11), if  $g \in G$ , we denote by  $m_0(g)$  an element of  $M_0$  such that there exist  $k, k' \in K$  with  $g = km_0(g)k'$ . Notice that  $\|h_{M_0}(m_0(g))\|$  does not depend on our choice of  $m_0(g)$ . By (1.21), one has

$$\|g\| \approx \|m_0(g)\|, g \in G, (1.24)$$

and by ([W2] 1.1.(6)) we have

$$\|m_0\| \approx e^{\|h_{M_0}(m_0)\|}, m_0 \in M_0. (1.25)$$

Let  $x \in G^1$  and  $a \in A_G$ . Then  $m_0(x) \in G^1 \cap M_0$  and  $m_0(xa) = m_0(x)a$ . Thus, one has  $h_G(m_0(x)) = 0$ . We deduce from (1.8) that  $h_{M_0}(m_0(x))$  belongs to  $a_0^G$ . Since

$h_{M_0}(m_0(x)a) = h_{M_0}(m_0(x)) + h_{M_0}(a)$  and  $h_{M_0}(a) \in a_G$ , we obtain by orthogonality that

$$\frac{1}{2}(\|h_{M_0}(m_0(x))\| + \|h_{M_0}(a)\|) \leq \|h_{M_0}(m_0(x)a)\| \leq \|h_{M_0}(m_0(x))\| + \|h_{M_0}(a)\|.$$

Hence (1.23) follows from (1.24) and (1.25).

We denote by  $C_c^\infty(G)$  the space of smooth functions on  $G$  with compact support. We normalize Haar measures according to [Ar3] §1. Unless otherwise stated, the Haar measure on a compact group will be normalized to have total volume 1.

Let  $M$  be a Levi subgroup of  $G$ . We fix a Haar measure on  $a_M$  so that the volume of the quotient  $a_M/\tilde{a}_{M,F}$  equals 1.

Let  $P = MN_P \in \mathcal{P}(M)$ . We denote by  $\delta_P$  the modular function of  $P$  given by

$$\delta_P(mn) = e^{2\rho_P(h_M(m))}, m \in M, n \in N_P,$$

where  $2\rho_P$  is the sum of roots, with multiplicity, of  $(P, A_M)$ . Let  $\bar{P} = MN_{\bar{P}}$  be the parabolic subgroup which is opposite to  $P$ . If  $dn$  is a Haar measure on  $N_P$  then the number

$$\gamma(P) = \int_{N_P} e^{2\rho_{\bar{P}}(h_{\bar{P}}(n))} dn$$

is finite. Moreover, the measure  $\gamma(P)^{-1}dn$  is independent of the choice of  $dn$  and thus defines a canonical Haar measure on  $N_P$ .

If  $dm$  is a Haar measure on  $M$  then there exists a unique Haar measure  $dg$  on  $G$ , independent of the choice of the parabolic subgroup  $P$ , such that

$$\int_G f(g)dg = \frac{1}{\gamma(P)\gamma(\bar{P})} \int_{N_P} \int_M \int_{N_{\bar{P}}} f(nm\bar{n})\delta_P(m)^{-1}d\bar{n} dm dn,$$

for  $f \in C_c^\infty(G)$ . We say that  $dm$  and  $dg$  are compatible. Compatibility has the obvious transitivity property relative to Levi subgroups of  $M$ . Using the Iwasawa decomposition (1.12), these measures satisfy

$$\int_G f(g)dg = \frac{1}{\gamma(P)} \int_K \int_M \int_{N_P} f(mnk)dn dm dk.$$

## 1.2 The symmetric space $H \backslash G$

Let  $E$  be an unramified quadratic extension of  $F$ . Thus  $E = F[\tau]$  where  $\tau^2$  is not a square in  $F$ . We denote by  $\sigma$  the nontrivial element of the Galois group  $\mathcal{G}al(E/F)$  of  $E/F$ . The normalized valuation  $|\cdot|_E$  on  $E$  satisfies  $|x|_E = |x|_F^2$  for  $x \in F$ .

If  $\underline{J}$  is an algebraic group defined over  $F$ , as usual we denote by  $J$  its group of points over  $F$ . Let  $\underline{J} \times_F E$  be the group, defined over  $E$ , obtained from  $\underline{J}$  by extension of scalars. We consider the group

$$\tilde{\underline{J}} := \text{Res}_{E/F}(\underline{J} \times_F E)$$

defined over  $F$ , obtained by restriction of scalars.

With our convention, one has  $\tilde{J} = \tilde{J}(F)$  and  $\tilde{J}$  is isomorphic to  $\underline{J}(E)$ .

Let  $\underline{H}$  be a reductive group defined over  $F$ . In all this article, we assume that  $\underline{H}$  is split over  $F$  and we set  $\underline{G} := \tilde{H}$  and  $G := \tilde{H}$ . We fix a maximal split torus  $A_0$  of  $H$ . Then  $A_0$  is also a maximal split torus of  $G$  and we have  $A_H = A_G$ .

The nontrivial element  $\sigma$  of  $\mathcal{G}al(E/F)$  induces an involution of  $\underline{G}$  defined over  $F$ , which we denote by the same letter. This automorphism  $\sigma$  extends to an  $E$ -automorphism  $\sigma_E$  on  $\underline{G} \times_F E$ .

We consider the canonical map  $\varphi$  defined over  $F$  from  $\underline{G}$  to  $(\underline{H} \times_F E) \times (\underline{H} \times_F E)$  by  $\varphi(g) = (g, \sigma(g))$ .

Then,  $\varphi$  extends uniquely to an isomorphism  $\Psi$  defined over  $E$  from  $\underline{G} \times_F E$  to  $(\underline{H} \times_F E) \times (\underline{H} \times_F E)$  such that  $\Psi(g) = (g, \sigma(g))$  for all  $g \in \underline{G}$  (1.26) and if  $\Psi(g) = (g_1, g_2)$  then  $\Psi(\sigma_E(g)) = (g_2, g_1)$ .

Now we turn to the description of the geometric structure of the symmetric space  $\mathcal{S} = H \backslash G$  according to [RR] sections 2 and 3.

Let  $\underline{\mathfrak{g}}$  be the Lie algebra of  $\underline{G}$  and  $\mathfrak{g}$  be the Lie algebra of its  $F$ -points. We will say that  $\mathfrak{g}$  is the Lie algebra of  $G$  and the Lie algebra  $\mathfrak{h}$  of  $H$  consists of the elements of  $\mathfrak{g}$  invariant by  $\sigma$ . We denote by  $\mathfrak{q}$  the space of antiinvariant elements of  $\mathfrak{g}$  by  $\sigma$ . Thus, one has  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q}$  and  $\mathfrak{g}$  may be identified with  $\mathfrak{h} \otimes_F E$ .

As in ([RR] §2.), we say that a subspace  $\mathfrak{c}$  of  $\mathfrak{q}$  is a Cartan subspace of  $\mathfrak{q}$  if  $\mathfrak{c}$  is a maximal abelian subspace of  $\mathfrak{q}$  made of semisimple elements. As  $E = F[\tau]$ , the multiplication by  $\tau$  induces an isomorphism between the set of Cartan subspaces of  $\mathfrak{q}$  and the set of Cartan subalgebras of  $\mathfrak{h}$  which preserves  $H$ -conjugacy classes.

We denote by  $\underline{\mathcal{P}}$  the connected component of 1 in the set of  $x$  in  $\underline{G}$  such that  $\sigma(x) = x^{-1}$ . Then the map  $\underline{p}$  from  $\underline{G}$  to  $\underline{\mathcal{P}}$  defined by  $\underline{p}(x) = x^{-1}\sigma(x)$  induces an isomorphism of affine varieties  $\underline{p} : \underline{H} \backslash \underline{G} \rightarrow \underline{\mathcal{P}}$ .

A torus  $\underline{A}$  of  $\underline{G}$  is called a  $\sigma$ -torus if  $\underline{A}$  is a torus defined over  $F$  contained in  $\underline{\mathcal{P}}$ . Notice that such torus are called  $\sigma$ -split torus in [RR]. We prefer change the terminology as  $\sigma$ -tori are not necessarily split over  $F$ . Each  $\sigma$ -torus is the centralizer in  $\underline{\mathcal{P}}$  of a Cartan subspace of  $\mathfrak{q}$ , or equivalently of a Cartan subalgebra of  $\mathfrak{h}$ .

Let  $S$  be a maximal torus of  $H$ . We denote by  $\underline{S}_\sigma$  the connected component of  $\tilde{S} \cap \underline{\mathcal{P}}$ . Then  $\underline{S}_\sigma$  is a  $\sigma$ -torus defined over  $F$  which identifies with the antidiagonal  $\{(s, s^{-1}); s \in \underline{S}\}$  of  $\underline{S} \times \underline{S}$  by the isomorphism (1.26). Thus,  $\underline{S}_\sigma$  is a maximal  $\sigma$ -torus and each maximal  $\sigma$ -torus arises in this way. The  $H$ -conjugacy classes of maximal tori of  $H$  are in bijective correspondence with the  $H$ -conjugacy classes of maximal  $\sigma$ -tori of  $G$  by the map  $S \mapsto S_\sigma$ . The roots of  $\underline{S}$  (resp.;  $\underline{S}_\sigma$ ) in  $\underline{\mathfrak{h}} = Lie(\underline{H})$  (resp.;

$\underline{\mathfrak{q}} \otimes_{\mathbb{F}} \bar{\mathbb{F}})$  are the restrictions of the roots of  $\tilde{\underline{S}}$  in  $\underline{\mathfrak{g}} = \text{Lie}(\underline{G})$ .

Therefore, each root of  $\underline{S}$  (resp.;  $\underline{S}_\sigma$ ) in  $\underline{\mathfrak{g}}$  has multiplicity two. If  $\tilde{\underline{S}}$  splits over a finite extension  $\mathbb{F}'$  of  $\mathbb{F}$ , we denote by  $\Phi(S'_\sigma, \mathfrak{g}')$  (resp.;  $\Phi(S', \mathfrak{h}')$ ) the set of roots of  $\underline{S}_\sigma(\mathbb{F}')$  in  $\underline{\mathfrak{g}} \otimes_{\mathbb{F}} \mathbb{F}'$  (resp.;  $\underline{S}(\mathbb{F}')$  in  $\underline{\mathfrak{h}} \otimes_{\mathbb{F}} \mathbb{F}'$ ).  
Let  $\tilde{\underline{\mathfrak{g}}}$  be the Lie algebra of  $\tilde{\underline{S}}$ . Then, the differential of each root  $\alpha$  of  $\Phi(\tilde{S}', \mathfrak{g}')$  defines a linear form on  $\tilde{\underline{\mathfrak{g}}} \otimes_{\mathbb{F}} \mathbb{F}'$  which we denote by the same letter. (1.27)

Let  $\mathcal{G}al(\bar{\mathbb{F}}/\mathbb{F})$  be the Galois group of  $\bar{\mathbb{F}}/\mathbb{F}$ . By ([RR] §3), the set of  $(H, S_\sigma)$ -double cosets in  $\underline{HS}_\sigma \cap G$  are parametrized by the finite set  $I$  of cohomology classes in  $H^1(\mathcal{G}al(\bar{\mathbb{F}}/\mathbb{F}), \underline{H} \cap \underline{S}_\sigma)$  which split in both  $\underline{H}$  and  $\underline{S}_\sigma$ . To each such classe  $m$ , we attach an element  $x_m \in G$  of the form  $x_m = h_m a_m^{-1}$  with  $h_m \in \underline{H}$  and  $a_m \in \underline{S}_\sigma$  such that  $m_\gamma = h_m^{-1} \gamma(h_m) = a_m^{-1} \gamma(a_m)$  for all  $\gamma \in \mathcal{G}al(\bar{\mathbb{F}}/\mathbb{F})$ .

**1.1 Lemma.** *Let  $x \in G$  such that  $x = hs$  with  $h \in \underline{H}$  and  $s \in \tilde{\underline{S}}$ . Then,  $xSx^{-1}$  is a maximal torus of  $H$  and there exists  $h' \in H$  such that  $x' = h'x$  centralizes the split connected component  $A_S$  of  $S$ .*

*Proof :*

Replace  $S$  by a  $H$ -conjugate if necessary, we may assume that  $A := A_S$  is contained in the fixed maximal split torus  $A_0$  of  $H$ . Since  $H$  is split,  $A_0$  is also a maximal split torus of  $G$ .

Since  $x = hs \in G$ , the torus  $\underline{S}' := x\underline{S}x^{-1}$  is equal to  $h\underline{S}h^{-1} \subset \underline{H}$ . Thus  $\underline{S}'$  is defined over  $\mathbb{F}$  and contained in  $\underline{H}$  and we obtain the first assertion.

Let  $S' := \underline{S}'(\mathbb{F})$  and let  $A'$  be the split connected component of  $S'$ . There exists  $h_1 \in H$  such that  $h_1 A' h_1^{-1} \subset A_0$ . We set  $x_1 = h_1 x$ , thus we have  $A_1 := x_1 A x_1^{-1} \subset A_0$ .

Let  $M = Z_G(A)$  and  $M_1 = Z_G(A_1) = x_1 M x_1^{-1}$ . Then  $A_0$  and  $x_1 A_0 x_1^{-1}$  are maximal split tori of  $M_1$ . Therefore, there exists  $y_1 \in M_1$  such that  $y_1 x_1 A_0 x_1^{-1} y_1^{-1} = A_0$ . As  $H$  is split, the Weyl group of  $A_0$  in  $G$  coincides with the Weyl group of  $A_0$  in  $H$ . Thus, there exist  $h_2 \in N_H(A_0)$  and  $v \in Z_G(A_0)$  such that  $z := y_1 x_1 = h_2 v$ .

For  $a \in A \subset A_0$ , one has  $z a z^{-1} = h_2 a h_2^{-1} = y_1 x_1 a x_1^{-1} y_1^{-1} = x_1 a x_1^{-1}$  since  $x_1 a x_1^{-1} \in A_1$  and  $y_1 \in M_1$ . One deduces that  $x' := h_2^{-1} h_1 x$  centralizes  $A$ .  $\square$

Thus, for each maximal torus  $S$  of  $H$ , we can fix a finite set of representatives  $\kappa_S = \{x_m\}_{m \in I}$  of the  $(H, S_\sigma)$ -double cosets in  $\underline{HS}_\sigma \cap G$  such that each element  $x_m$  may be written  $x_m = h_m a_m^{-1}$  where  $h_m \in \underline{H}$  centralizes  $A_S$  and  $a_m \in \underline{S}_\sigma$ . Hence  $x_m$  centralizes  $A_S$ . (1.28)

### 1.3 Weyl integration formula and orbital integrals

We first recall basic notions on the symmetric space according to ([RR], §3). An element  $x$  in  $\underline{G}$  is called  $\sigma$ -semisimple if the double coset  $\underline{H}x\underline{H}$  is Zariski closed. This is equivalent to say that  $\underline{p}(x)$  is a semisimple point of  $\underline{G}$ . We say that a semisimple

element  $x$  is  $\sigma$ -regular if this closed double coset  $\underline{H}x\underline{H}$  is of maximal dimension. This is equivalent to say that the centralizer of  $\underline{p}(x)$  in  $\mathfrak{q}$  (resp.;  $\underline{\mathcal{P}}$ ) is a Cartan subspace of  $\mathfrak{q}$  (resp.; a maximal  $\sigma$ -torus of  $\underline{G}$ ).

We denote by  $G^{\sigma\text{-reg}}$  the set of  $\sigma$ -regular elements of  $G$ .

For  $g \in G$ , we denote by  $D_G(g)$  the coefficient of the least power of  $t$  appearing nontrivially in  $\det(t + 1 - \text{Ad}(g))$ . We define the  $H$ -biinvariant function  $\Delta_\sigma$  on  $G$  by  $\Delta_\sigma(x) = D_G(\underline{p}(x))$ . Then by ([RR], Lemma 3.2. and Lemma 3.3), the set of  $g \in G$  such that  $\Delta_\sigma(g) \neq 0$  coincides with  $G^{\sigma\text{-reg}}$ .

Let  $S$  be a maximal torus of  $H$  with Lie algebra  $\mathfrak{s}$ . Then  $\tilde{\mathfrak{s}} := \mathfrak{s} \otimes_{\mathbb{F}} \mathbb{E}$  identifies with the Lie algebra of  $\tilde{S}$ . For  $g \in x_m S_\sigma$  with  $x_m \in \kappa_S$ , one has

$$\Delta_\sigma(g) = D_G(\underline{p}(g)) = \det(1 - \text{Ad}(\underline{p}(g)))_{\mathfrak{g}/\tilde{\mathfrak{s}}}. \quad (1.29)$$

By ([RR] Theorem 3.4 (1)), the set  $G^{\sigma\text{-reg}}$  is a disjoint union

$$G^{\sigma\text{-reg}} = \bigcup_{\{S\}_H} \bigcup_{x_m \in \kappa_S} H((x_m S_\sigma) \cap G^{\sigma\text{-reg}})H, \quad (1.30)$$

where  $\{S\}_H$  runs the  $H$ -conjugacy classes of maximal tori of  $H$ .

If  $x_m \in \kappa_S$  then  $x_m = h_m a_m$  for some  $h_m \in \underline{H}$  and  $a_m \in \underline{S}_\sigma$ , hence  $\underline{p}(x_m) = a_m^{-2}$  commutes with  $S$  and  $S_\sigma$ . Therefore for  $\gamma \in S_\sigma$ , we have

$$\underline{p}(x_m \gamma) = \underline{p}(x_m) \gamma^{-2} \quad \text{and} \quad H x_m \gamma S = H x_m \gamma.$$

We have the following Weyl integration formula (cf. [RR] Theorem 3.4 (2)):

Let  $f$  be a compactly supported smooth function on  $G$ , then we have

$$\int_G f(y) dy = \sum_{\{S\}_H} \sum_{x_m \in \kappa_S} c_{S, x_m}^0 \int_{S_\sigma} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{S \setminus H} \int_H f(h x_m \gamma l) dh d\bar{l} d\gamma, \quad (1.31)$$

where the constants  $c_{S, x_m}^0$  are explicitly given in ([RR] Theorem 3.4 (1)).

For our purposes, we need another version of this Weyl integration formula. Let  $S$  be a maximal torus of  $H$ . We denote by  $A_S$  its split connected component. Since the quotient  $A_S \setminus S$  is compact, by our choice of measure, the integration over  $S \setminus H$  in the Weyl formula above can be replaced by an integration over  $A_S \setminus H$ . Moreover, it is convenient to change  $h$  into  $h^{-1}$ . As every  $x_m \in \kappa_S$  commutes with  $A_S$  (cf. (1.28)), one can replace the integration over  $(A_S \setminus H) \times H$ , by an integration over  $\text{diag}(A_S) \setminus (H \times H)$  where  $\text{diag}(A_S)$  is the diagonal of  $A_S$ . This gives the following Weyl integration formula equivalent to (1.31):

$$\int_G f(y) dy = \sum_{\{S\}_H} \sum_{x_m \in \kappa_S} c_{S, x_m}^0 \int_{S_\sigma} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_S) \setminus (H \times H)} f(h^{-1} x_m \gamma l) d(\overline{h, l}) d\gamma. \quad (1.32)$$



We will now describe the  $H$ -conjugacy classes of maximal tori of  $H$  in terms of Levi subgroups  $M$  in  $\mathcal{L}(A_0)$  and  $M$ -conjugacy classes of some tori of  $M$ .

Let  $M \in \mathcal{L}(A_0)$ . We denote by  $N_H(M)$  its normalizer in  $H$ . If  $S$  is a maximal torus of  $M$ , we denote by  $W(M, S)$  (resp.  $W(H, S)$ ) its Weyl group in  $M$  (resp.  $H$ ). We choose a set  $\mathcal{T}_M$  of representatives for the  $M$ -conjugacy classes of maximal tori  $S$  in  $M$  such that  $A_M \backslash S$  is compact. For  $M, M' \in \mathcal{L}(A_0)$ , we write  $M \sim M'$  if  $M$  and  $M'$  are conjugate under  $H$ .

Let  $S$  be a maximal torus of  $H$  whose split connected component  $A_S$  is contained in  $A_0$ . Then, the centralizer  $M$  of  $A_S$  belongs to  $\mathcal{L}(A_0)$  and  $S$  is a maximal torus of  $M$  such that  $A_M \backslash S$  is compact. If  $S'$  is a maximal torus conjugated to  $S$  by  $H$  such that  $A_{S'}$  is contained in  $A_0$ , then the centralizer  $M'$  of  $A_{S'}$  in  $H$  belongs to  $\mathcal{L}(A_0)$  and  $M' \sim M$ .

Since each maximal torus of  $H$  is  $H$ -conjugate to a maximal torus  $S$  such that  $A_S \subset A_0$ , we obtain a surjective map  $S \mapsto \{S\}_H$  from the set of  $S$  in  $\mathcal{T}_M$  where  $M$  runs a system of representatives of  $\mathcal{L}(A_0)_{/\sim}$  to the set of  $H$ -conjugacy classes of maximal tori of  $H$ .

Let  $M \in \mathcal{L}(A_0)$ . By ([Ko ] (7.12.3)), the cardinal of the class of  $M$  in  $\mathcal{L}(A_0)_{/\sim}$  is equal to

$$\frac{|W(H, A_0)|}{|W(M, A_0)||N_H(M)/M|}$$

where  $N_H(M)$  is the normalizer of  $M$  in  $H$ .

By ([Ko ] Lemma 7.1), if  $S$  is a maximal torus of  $M$ , then the number of  $M$ -conjugacy classes of maximal torus  $S'$  in  $M$  such that  $S'$  is  $H$ -conjugate to  $S$  is equal to

$$\frac{|N_H(M)/M||W(M, S)|}{|W(H, S)|}.$$

Therefore, we can rewrite (1.32) as follows:

$$\int_G f(g)dg = \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_M) \backslash H \times H} f(h^{-1} x_m \gamma l) d\overline{(h, l)} d\gamma \quad (1.33)$$

where

$$c_M = \frac{|W(M, A_0)|}{|W(H, A_0)|} \quad \text{and} \quad c_{S, x_m} = \frac{|W(H, S)|}{|W(M, S)|} c_{S, x_m}^0.$$

Let  $f \in C_c^\infty(G)$ . We define the orbital integral  $\mathcal{M}(f)$  of  $f$  on  $G^{\sigma\text{-reg}}$  as follows. Let  $S$  a maximal torus of  $H$ . For  $x_m \in \kappa_S$  and  $\gamma \in S_\sigma$  with  $x_m \gamma \in G^{\sigma\text{-reg}}$ , we set

$$\begin{aligned} \mathcal{M}(f)(x_m \gamma) &:= |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/4} \int_{\text{diag}(A_S) \backslash (H \times H)} f(h^{-1} x_m \gamma l) d\overline{(h, l)} \\ &= |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/4} \int_{S \backslash H} \int_H f(h x_m \gamma l) dh d\bar{l}. \end{aligned} \quad (1.34)$$

Our definition corresponds, up to a positive constant, to Definition 3.8 of [RR]. Indeed, by definition of  $\Delta_\sigma$ , we have  $\Delta_\sigma(x_m\gamma) = D_G(\underline{p}(x_m\gamma))$ . Since we can write  $x_m = h_m a_m$  with  $h_m \in \underline{H}$  and  $a_m \in \underline{S}_\sigma$ , we have  $\underline{p}(x_m\gamma) = \underline{p}(x_m)\gamma^{-2} = a_m^{-2}\gamma^{-2}$  for  $\gamma \in S_\sigma$ . Let  $F'$  be an extension of  $E$  such that  $\tilde{S}$  splits over  $F'$  and  $a_m \in \underline{S}_\sigma(F')$ . Since each root  $\alpha$  of  $\underline{S}_\sigma(F')$  in  $\mathfrak{g} \otimes F'$  have multiplicity  $m(\alpha) = 2$ , using notation of (1.27), we obtain:

$$\Delta_\sigma(x_m\gamma) = \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}')} (1 - \underline{p}(x_m)^\alpha \gamma^{-2\alpha})^2 = \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}')} (\gamma^\alpha - \underline{p}(x_m)^\alpha \gamma^{-\alpha})^2,$$

hence

$$\begin{aligned} |\Delta_\sigma(x_m\gamma)|_{F'}^{1/4} &= \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}')} |(\gamma^\alpha - \underline{p}(x_m)^\alpha \gamma^{-\alpha})^{m(\alpha)-1}|_{F'}^{1/2}, \\ &= \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}')} |(\gamma^\alpha - \underline{p}(x_m)^\alpha \gamma^{-\alpha})|_{F'}^{1/2}. \end{aligned}$$

Then, the Weyl integration formula (1.31) in terms of orbital integrals is given as in ([RR] page 126) by

$$\int_G f(y) dy = \sum_{\{S\}_H} \sum_{x_m \in \kappa_S} c_{S, x_m}^0 \int_{S_\sigma} |\Delta_\sigma(x_m\gamma)|_{F'}^{1/4} \mathcal{M}(f)(x_m\gamma) d\gamma.$$

**1.2 Theorem.** *Let  $f \in C_c^\infty(G)$  and  $S$  be a maximal torus of  $H$ . Let  $x_m \in \kappa_S$ .*

1. *There exists a compact set  $\Omega$  in  $S_\sigma$  such that, for any  $\gamma$  in the complementary of  $\Omega$  with  $x_m\gamma \in G^{\sigma\text{-reg}}$ , one has  $\mathcal{M}(f)(x_m\gamma) = 0$ .*

2.

$$\sup_{\gamma \in S_\sigma; x_m\gamma \in G^{\sigma\text{-reg}}} |\mathcal{M}(f)(x_m\gamma)| < +\infty.$$

*Proof :*

The proof follows that of the group case (cf. [HC3] proof of Theorem 14). We write it for convenience of the reader.

1. Let  $\omega$  be the support of  $f$ . We consider the set  $\omega_S$  of elements  $\gamma$  in  $S_\sigma$  such that  $x_m\gamma$  is in the closure of  $H\omega H$ . For  $g \in G$ , we consider the polynomial function

$$\det(1 - t - \text{Ad } \underline{p}(g)) = t^n + q_{n-1}(g)t^{n-1} + \dots + q_l(g)t^l \quad (1.35)$$

where  $l$  is the rank of  $G$  and  $n$  its dimension. Each  $q_j$  is a  $H \times H$  biinvariant regular function on  $G$ , thus it is bounded on  $x_m\omega_S$ . Therefore, the roots of  $\det(1 - t - \text{Ad } \underline{p}(g))$  are bounded on  $x_m\omega_S$ .

For  $\gamma \in S_\sigma$ , we have  $\underline{p}(x_m\gamma) = \underline{p}(x_m)\gamma^{-2}$ . We choose a finite extension  $F'$  of  $F$  such that  $\tilde{S}$  splits over  $F'$  and  $\underline{p}(x_m) \in \underline{S}_\sigma(F')$ . Using notation of (1.27), the roots of  $\det(1 - t - \text{Ad } \underline{p}(x_m\gamma))$  are the numbers  $(1 - \underline{p}(x_m)^\alpha \gamma^{-2\alpha})$  for  $\alpha \in \Phi(S'_\sigma, \mathfrak{g}')$ . Since

these roots are bounded on  $x_m\omega_S$ , we deduce that the maps  $\gamma \rightarrow \gamma^\alpha$ ,  $\alpha \in \Phi(S'_\sigma, \mathfrak{g}')$ , are bounded on  $\omega_S$ . This implies that  $\omega_S$  is bounded. Then, the closure  $\Omega$  of  $\omega_S$  satisfies the first assertion.

2. By 1., if  $\gamma \notin \Omega$  then  $\mathcal{M}(f)(x_m\gamma) = 0$ . Thus, it is enough to prove that for each  $\gamma_0 \in S_\sigma$ , there exists a neighborhood  $V_{\gamma_0}$  of  $\gamma_0$  in  $S_\sigma$  such that

$$\sup_{\gamma \in V_{\gamma_0}, x_m\gamma \in G^{\sigma-reg}} |\mathcal{M}(f)(x_m\gamma)| < +\infty. \quad (1.36)$$

Let  $y_0 := \underline{p}(x_m\gamma_0)$ . We first assume that  $y_0$  is central in  $G$ . Then, we have  $\Delta_\sigma(x_m\gamma_0\gamma) = D_G(y_0\gamma^{-2}) = D_G(\gamma^{-2})$  for  $\gamma \in S_\sigma$  and  $x_m\gamma_0h(x_m\gamma_0)^{-1} \in H$  for  $h \in H$ . We define the function  $f_0$  on  $G$  by  $f_0(g) := f(x_m\gamma_0g)$ . Then, we have  $\mathcal{M}(f_0)(\gamma) = \mathcal{M}(f)(x_m\gamma_0\gamma)$  for  $\gamma \in S_\sigma \cap G^{\sigma-reg}$ . Thus, we are reduced to the case  $y_0 = 1$ . As in the group case, we use the exponential map "exp" which is well defined in a neighborhood of 0 in  $\mathfrak{g}$  since the characteristic of  $\mathbb{F}$  is equal to zero (cf. [HC4] §10). As in ([HC1] proof of Lemma 15), we can choose a  $H$ -invariant open neighborhood  $V_0$  of 0 in  $\mathfrak{h}$  such that the map  $X \in V_0 \mapsto \exp(\tau X)$  is an isomorphism and an homeomorphism onto its image and there is a  $H$ -invariant function  $\varphi \in C_c^\infty(\mathfrak{h})$  such that  $\varphi(X) = 1$  for  $X \in V_0$ . We define  $\bar{f}$  in  $C_c^\infty(\mathfrak{h})$  by  $\bar{f}(X) = \varphi(X) \int_H f(h \exp(\tau X)) dh$ .

Let  $\mathfrak{s}$  be the Lie algebra of  $S$ . For  $X \in \mathfrak{s}$ , we set  $\eta(X) = |\det(\text{ad}X)_{\mathfrak{h}/\mathfrak{s}}|_{\mathbb{F}}$ . We consider a finite extension  $\mathbb{F}'$  of  $\mathbb{F}$  such that  $\tilde{S}$  splits over  $\mathbb{F}'$  and  $\underline{p}(x_m) \in \underline{S}_\sigma(\mathbb{F}')$ . We use notation of (1.27). Since each root of  $S'_\sigma$  in  $\mathfrak{g}'$  has multiplicity 2, we have for  $X \in V_0$

$$\begin{aligned} \frac{|\Delta_\sigma(\exp \tau X)|_{\mathbb{F}'}^{1/2}}{\eta(X)} &= \frac{|D_{G'}(\exp(-2\tau X))|_{\mathbb{F}'}^{1/2}}{\eta(X)} = \frac{\prod_{\alpha \in \Phi(S', \mathfrak{h}')} |1 - e^{2\tau\alpha(X)}|_{\mathbb{F}'}}{\prod_{\alpha \in \Phi(S', \mathfrak{h}')} |\alpha(X)|_{\mathbb{F}'}} \\ &= |2\tau|_{\mathbb{F}'}^{|\Phi(S', \mathfrak{h}')|} \prod_{\alpha \in \Phi(S', \mathfrak{h}')} |1 + \tau\alpha(X) + \frac{4\tau^2\alpha(X)^2}{3!} + \dots|_{\mathbb{F}'}. \end{aligned}$$

We can reduce  $V_0$  in such way that each term of this product is equal to 1. Thus, we obtain

$$\begin{aligned} \mathcal{M}(f)(\exp \tau X) &= |2\tau|_{\mathbb{F}'}^{|\Phi(S', \mathfrak{h}')|} \eta(X)^{1/2} \int_{H/S} \left( \int_H f(h \exp \tau \text{Ad}(l)X) dh \right) d\bar{l} \\ &= |2\tau|_{\mathbb{F}'}^{|\Phi(S', \mathfrak{h}')|} \eta(X)^{1/2} \int_{H/S} \bar{f}(\text{Ad}(l)X) d\bar{l}, \end{aligned}$$

for  $X \in V_0$ . The estimate (1.36) follows from the result on the Lie algebra (cf. [HC3] Theorem 13).

If  $y_0 = \underline{p}(x_m\gamma_0)$  is not central in  $G$ , we consider the centralizer  $\underline{Z}$  of  $y_0$  in  $\underline{H}$ . Let  $\underline{Z}^0$  be the connected neutral component of  $\underline{Z}$ . By ([Bo], III.9), the group  $\underline{Z}^0$  is

defined over  $F$ . As usual, we set  $\tilde{\mathcal{Z}}^0 := \text{Res}_{E/F}(\mathcal{Z}^0 \times_F E)$  and we denote by  $\tilde{\mathfrak{z}}$  its Lie algebra. By definition of  $\tilde{\mathfrak{z}}$ , one has

$$|\det(1 - \text{Ad}(y_0))_{\mathfrak{g}/\tilde{\mathfrak{z}}}|_F \neq 0.$$

Thus, there exists a neighborhood  $V$  of 1 in  $S_\sigma$  such that, for all  $\gamma \in V$ , then

$$|\det(1 - \text{Ad}(y_0\gamma^{-2}))_{\mathfrak{g}/\tilde{\mathfrak{z}}}|_F = |\det(1 - \text{Ad}(y_0)_{\mathfrak{g}/\tilde{\mathfrak{z}}})|_F \neq 0. \quad (1.37)$$

From ([HC3] Lemma 19), there exist a neighborhood  $V_1$  of  $y_0$  in  $\tilde{S}$  and a compact subset  $\overline{C_G}$  of  $\tilde{\mathcal{Z}}^0 \backslash G$  such that, if  $g \in G$  satisfies  $g^{-1}V_1g \cap \underline{p}(\omega) \neq \emptyset$  then its image  $\bar{g}$  in  $\tilde{\mathcal{Z}}^0 \backslash G$  belongs to  $\overline{C_G}$  (here  $\omega$  is the support of  $f$ ).

We choose a neighborhood  $W$  of 1 in  $S_\sigma$  such that  $W \subset V$  and  $\underline{p}(x_m\gamma_0\gamma) = y_0\gamma^{-2} \in V_1$  for all  $\gamma \in W$ . By ([Bo], III 9.1), the quotient  $\mathcal{Z}^0 \backslash H$  is a closed subset of  $\tilde{\mathcal{Z}}^0 \backslash G$ , hence

the set  $\overline{C} := \overline{C_G} \cap \mathcal{Z}^0 \backslash H$  is a compact subset of  $\mathcal{Z}^0 \backslash H$  such that if  $l \in H$  satisfies  $l^{-1}y_0\gamma^{-2}l \in \underline{p}(\omega)$  for some  $\gamma \in W$  then its image  $\bar{l}$  in  $\mathcal{Z}^0 \backslash H$  belongs to  $\overline{C}$ . (1.38)

Let  $\gamma \in W$  such that  $x_m\gamma_0\gamma \in G^{\sigma\text{-reg}}$ . One has

$$\int_{S \backslash H} \int_H f(hx_m\gamma_0\gamma l) dh d\bar{l} = \int_{\mathcal{Z}^0 \backslash H} \int_{S \backslash \mathcal{Z}^0} \int_H f(hx_m\gamma_0\gamma \xi l) dh d\bar{\xi} d\bar{l}. \quad (1.39)$$

By the choice of  $W$ , the map

$$\bar{l} \in \mathcal{Z}^0 \backslash H \mapsto \int_{S \backslash \mathcal{Z}^0} \int_H f(hx_m\gamma_0\gamma \xi l) dh d\bar{\xi}$$

vanishes outside  $\overline{C}$ . We choose  $u \in C_c^\infty(H)$  such that the map  $\bar{u} \in C_c^\infty(\mathcal{Z}^0 \backslash H)$  defined by  $\bar{u}(\bar{l}) := \int_{\mathcal{Z}^0} u(\xi l) d\xi$  is equal to 1 if  $\bar{l} \in \overline{C}$ . As  $u$  and  $f$  are compactly supported, the map

$$\Phi : z \in \tilde{\mathcal{Z}}^0 \mapsto \int_H u(l) \int_H f(hx_m\gamma_0 z l) dh dl$$

is well-defined. Since  $y_0 = \underline{p}(x_m\gamma_0) = (x_m\gamma_0)^{-1}\sigma(x_m\gamma_0)$ , we have  $\xi(x_m\gamma_0)^{-1}\sigma(x_m\gamma_0) = (x_m\gamma_0)^{-1}\sigma(x_m\gamma_0)\xi$  for  $\xi \in \mathcal{Z}^0$ . Hence,  $x_m\gamma_0\xi(x_m\gamma_0)^{-1} \in H$ . Thus  $\Phi$  is left invariant by  $\mathcal{Z}^0$ .

We claim that  $\Phi \in C_c^\infty(\mathcal{Z}^0 \backslash \tilde{\mathcal{Z}}^0)$ . Indeed, fix  $l$  in the support of  $u$ . If  $f(hx_m\gamma_0 z l)$  is nonzero for some  $h \in H$  and  $z \in \tilde{\mathcal{Z}}^0$  then  $\underline{p}(hx_m\gamma_0 z l) = \underline{p}(x_m\gamma_0 z l)$  belongs to  $\underline{p}(\omega)$ , where  $\omega$  is the support of  $f$ . Since  $z$  commutes with  $y_0 = \underline{p}(x_m\gamma_0)$ , we have  $\underline{p}(x_m\gamma_0 z l) = l^{-1}y_0 \underline{p}(z)\sigma(l)$ . As  $u$  is compactly supported, we deduce that  $\Phi(z) = 0$  when  $\underline{p}(z)$  is outside a compact set. Hence, the map  $\Phi$  is a compactly supported function on  $\mathcal{Z}^0 \backslash \tilde{\mathcal{Z}}^0$ .

By assumption, the function  $f$  is right invariant by a compact open subgroup of  $G$ . Thus  $f$  is right invariant by some compact open subgroup of  $H$ . We denote by  $\pi_l f$  the right translate of  $f$  by an element  $l \in G$ . Since  $u$  is compactly supported, the vector space generating by  $\pi_l f$ , when  $l \in H$  runs the support of  $u$ , is finite dimensional. Hence, one can find a compact open subgroup  $J_1$  of  $\tilde{Z}^0$  such that for each  $l$  in the support of  $u$ , the function  $\pi_l f$  is right invariant by  $J_1$ . This implies that  $\Phi$  is smooth and our claim follows.

Therefore, there exists  $\varphi \in C_c^\infty(\tilde{Z}^0)$  such that

$$\Phi(z) = \int_{\tilde{Z}^0} \varphi(\xi z) d\xi = \int_H u(l) \int_H f(hx_m \gamma_0 z l) dh dl, \quad z \in \tilde{Z}^0.$$

We obtain

$$\begin{aligned} \int_{S \setminus \tilde{Z}^0} \int_{\tilde{Z}^0} \varphi(\xi_1 \gamma \xi_2) d\xi_1 d\xi_2 &= \int_H u(l) \left( \int_{S \setminus \tilde{Z}^0} \int_H f(hx_m \gamma_0 \gamma \xi_2 l) dh d\xi_2 \right) dl \\ &= \int_{\tilde{Z}^0 \setminus H} \int_{\tilde{Z}^0} u(\xi_1 l) \left( \int_{S \setminus \tilde{Z}^0} \int_H f(hx_m \gamma_0 \gamma \xi_2 \xi_1 l) dh d\xi_2 \right) d\xi_1 d\bar{l} \\ &= \int_{\tilde{Z}^0 \setminus H} \bar{u}(\bar{l}) \left( \int_{S \setminus \tilde{Z}^0} \int_H f(hx_m \gamma_0 \gamma \xi_2 l) dh d\xi_2 \right) d\bar{l}. \end{aligned}$$

By definition, the map  $\bar{u}$  is equal to 1 on the compact set  $\bar{C}$ . By definition of  $\bar{C}$  (cf. (1.38) and (1.3)), we obtain

$$\int_{S \setminus \tilde{Z}^0} \int_{\tilde{Z}^0} \varphi(\xi_1 \gamma \xi_2) d\xi_1 d\xi_2 = \int_{S \setminus H} \int_H f(hx_m \gamma_0 \gamma l) dh d\bar{l}.$$

By (1.37) and the choice of  $W$ , one has

$$|D_G(y_0 \gamma^{-2})|_{\mathbb{F}} = |D_{\tilde{Z}^0}(\gamma^{-2})|_{\mathbb{F}} |\det(1 - \text{Ad}(y_0))_{\mathfrak{g}/\mathfrak{h}}|_{\mathbb{F}}, \quad \gamma \in W.$$

Then, we deduce that for  $\gamma \in W$  satisfying  $x_m \gamma_0 \gamma \in G^{\sigma\text{-reg}}$ , one has

$$\mathcal{M}(f)(x_m \gamma_0 \gamma) = |\det(1 - \text{Ad}(y_0))_{\mathfrak{g}/\mathfrak{h}}|_{\mathbb{F}}^{1/4} |D_{\tilde{Z}^0}(\gamma^{-2})|_{\mathbb{F}}^{1/4} \int_{S \setminus \tilde{Z}^0} \int_{\tilde{Z}^0} \varphi(\xi_1 \gamma \xi_2) d\xi_1 d\xi_2.$$

Since  $|D_{\tilde{Z}^0}(\gamma^{-2})|_{\mathbb{F}}$  coincides with the function  $|\Delta_\sigma|_{\mathbb{F}}$  for the group  $\tilde{Z}^0$  evaluated at  $\gamma$  (cf. (1.29)), one deduces the estimate (1.36) for  $f$  applying the first case to  $\varphi$  defined on  $\tilde{Z}^0$ .  $\square$

## 2 Geometric side of the local relative trace formula

### 2.1 Truncation

In this section, we will recall some results of ([Ar3], §3), needed in the sequel. We keep notation of §1.1 for the group  $H$ . Since  $H$  is split, one has  $M_0 = A_0$ . We fix a Levi subgroup  $M \in \mathcal{L}(A_0)$  of  $H$ . Let  $P \in \mathcal{P}(M)$ . We recall that  $A_M$  denotes the maximal split connected component of  $M$ .

We denote by  $\Sigma_P$  the set of roots of  $A_M$  in the Lie algebra of  $P$ ,  $\Sigma_P^r$  the subset of reduced roots and  $\Delta_P$  the subset of simple roots.

For  $\beta \in \Delta_P$ , the "co-root"  $\check{\beta} \in a_M$  is defined as usual as follows: if  $P \in \mathcal{P}(A_0)$  is a minimal parabolic subgroup, then  $\check{\beta} = 2(\beta, \beta)^{-1}\beta$ , where  $a_0^*$  identifies with  $a_0$  by the scalar product on  $a_0$ . In the general case, we choose  $P_0 \in \mathcal{P}(A_0)$  contained in  $P$ . Then, there exists a unique  $\alpha \in \Delta_{P_0}$  such that  $\beta = \alpha|_{a_M}$ . The "co-root"  $\check{\beta}$  is the projection of  $\check{\alpha}$  onto  $a_M$  with respect to the decomposition  $a_0 = a_M \oplus a_0^M$ . This projection does not depend of the choice of  $P_0$ .

We denote by  $a_P^+$  the positive Weyl chamber of elements  $X \in a_M$  satisfying  $\alpha(X) > 0$  for all  $\alpha \in \Sigma_P$ .

Let  $M \in \mathcal{L}(A_0)$ . A set of points in  $a_M$  indexed by  $P \in \mathcal{P}(M)$

$$\mathcal{Y} = \mathcal{Y}_M := \{Y_P \in a_M; P \in \mathcal{P}(M)\}$$

is called a  $(H, M)$ -orthogonal set if for all adjacent parabolic subgroups  $P, P'$  in  $\mathcal{P}(M)$  whose chambers in  $a_M$  share the wall determined by the simple root  $\alpha \in \Delta_P \cap (-\Delta_{P'})$ , one has  $Y_P - Y_{P'} = r_{P,P'}\check{\alpha}$  for a real number  $r_{P,P'}$ . The orthogonal set is called positive if each of the numbers  $r_{P,P'}$  are nonnegative. This is the case for example if the number

$$d(\mathcal{Y}) = \inf_{\{\alpha \in \Delta_P; P \in \mathcal{P}(M)\}} \alpha(Y_P) \quad (2.1)$$

is nonnegative.

One example is the set

$$\{-h_P(x); P \in \mathcal{P}(M)\},$$

defined for any point  $x \in H$ . This is a positive  $(H, M)$ -orthogonal set by ([Ar1] Lemma 3.6).

If  $L$  belongs to  $\mathcal{L}(M)$  and  $Q$  is a group in  $\mathcal{P}(L)$ , we define  $Y_Q$  to be the projection onto  $a_L$  of any point  $Y_P$ , with  $P \in \mathcal{P}(M)$  and  $P \subset Q$ . Then  $Y_Q$  is independent of  $P$  and  $\mathcal{Y}_L := \{Y_Q; Q \in \mathcal{P}(L)\}$  is a  $(H, L)$ -orthogonal set. (2.2)

We shall write  $\mathcal{S}_M(\mathcal{Y})$  for the convex hull in  $a_M/a_H$  of a  $(H, M)$ -orthogonal set  $\mathcal{Y}$ . Notice that  $\mathcal{S}_M(\mathcal{Y})$  does only depend on the projection onto  $a_M^H$  of each  $Y_P \in \mathcal{Y}$ ,  $P \in \mathcal{P}(M)$ .

Let  $P \in \mathcal{P}(M)$ . If each  $Y_P$  is in the positive Weyl chamber  $a_P^+$  (this condition is equivalent to say that  $d(\mathcal{Y})$  is positive), we have a simple description of  $\mathcal{S}_M(\mathcal{Y}) \cap a_P^+$  ([Ar3] Lemma 3.1). We denote by  $(\omega_\gamma^P)_{\gamma \in \Delta_P}$  the set of weights, that is the dual basis in  $(a_M^H)^*$  of the set of co-roots  $\{\tilde{\gamma}; \gamma \in \Delta_P\}$ . Then, we have

$$\mathcal{S}_M(\mathcal{Y}) = \{X \in a_P^+; \omega_\gamma^P(X - Y_P) \leq 0, \gamma \in \Delta_P\}. \quad (2.3)$$

We now recall a decomposition of the characteristic function of  $\mathcal{S}_M(\mathcal{Y})$  valid when  $\mathcal{Y}$  is positive. (cf. [Ar3] (3.8)). Suppose that  $\Lambda$  is a point in  $a_{M,\mathbb{C}}^*$  whose real part  $\Lambda_R \in a_M^*$  is in general position. If  $P \in \mathcal{P}(M)$ , we define  $\Delta_P^\Lambda$  the set of simple roots  $\alpha \in \Delta_P$  such that  $\Lambda_R(\tilde{\alpha}) < 0$ . Let  $\varphi_P^\Lambda$  be the characteristic function of the set of  $X \in a_M$  such that  $\omega_\alpha^P(X) > 0$  for each  $\alpha \in \Delta_P^\Lambda$  and  $\omega_\alpha^P(X) \leq 0$  for each  $\alpha$  in the complement of  $\Delta_P^\Lambda$  in  $\Delta_P$ . We define

$$\sigma_M(X, \mathcal{Y}) := \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\Lambda|} \varphi_P^\Lambda(X - Y_P). \quad (2.4)$$

By ([Ar3], §3 p22), the function  $\sigma_M(\cdot, \mathcal{Y})$  vanishes on the complement of  $\mathcal{S}_M(\mathcal{Y})$  and is bounded. Moreover, if  $\mathcal{Y}$  is positive then  $\sigma_M(\cdot, \mathcal{Y})$  is exactly the characteristic function of  $\mathcal{S}_M(\mathcal{Y})$ . (2.5)

The following Lemma will allow us to define the minimum of two orthogonal sets.

For  $P \in \mathcal{P}(M)$ , we denote by  $(\tilde{\omega}_\gamma^P)_{\gamma \in \Delta_P}$  the set of coweights, that is the dual basis in  $a_M^H$  of the roots  $\{\gamma; \gamma \in \Delta_P\}$ .

**2.1 Lemma.** *Let  $P$  and  $P'$  two adjacent parabolic subgroups in  $\mathcal{P}(M)$  whose chambers in  $a_M$  share the wall determined by the simple root  $\alpha \in \Delta_P \cap (-\Delta_{P'})$ . Then:*

1. *For all  $\beta$  in  $\Delta_P - \{\alpha\}$ , there exists a unique  $\beta'$  in  $\Delta_{P'} - \{\alpha\}$  such that  $\beta' = \beta + k_\beta \alpha$  where  $k_\beta$  is a nonnegative integer. Moreover, the map  $\beta \mapsto \beta'$  is a bijection between  $\Delta_P - \{\alpha\}$  and  $\Delta_{P'} - \{-\alpha\}$ .*
2. *For all  $\beta$  in  $\Delta_P - \{\alpha\}$ , one has  $\tilde{\omega}_{\beta'}^{P'} = \tilde{\omega}_\beta^P$ .*

*Proof :*

We denote by  $\mathbb{N}$  the set of nonnegative integers and by  $\mathbb{N}^*$  the subset of positive integers.

1. As  $P$  and  $P'$  are adjacent, we have  $\Sigma_{P'} = (\Sigma_P - \{\alpha\}) \cup \{-\alpha\}$ . Let  $\beta \in \Delta_P - \{\alpha\}$ . If  $\beta \in \Delta_{P'}$  then we set  $\beta' := \beta$ .

Assume that  $\beta$  is not in  $\Delta_{P'}$ . Since  $\beta \in \Sigma_{P'}$ , there exists  $\Theta \subset \Delta_{P'} - \{-\alpha\}$  such that  $\beta = \sum_{\delta \in \Theta} n_\delta \delta - k_\beta \alpha$  where the  $n_\delta$ 's are positive integers and  $k_\beta$  is a nonnegative integer. Each  $\delta$  in  $\Theta$  belongs to  $\Sigma_P$ . Therefore, there are nonnegative integers  $(r_{\delta,\eta})_{\eta \in \Delta_P}$  such that  $\delta = \sum_{\eta \in \Delta_P} r_{\delta,\eta} \eta$ . We set  $\beta_1 := \sum_{\delta \in \Theta} n_\delta \delta = \beta + k_\beta \alpha$ .

Let  $\gamma \in \Delta_P - \{\alpha\}$ . If  $\gamma \neq \beta$ , one has  $\beta_1(\tilde{\omega}_\gamma^P) = \beta(\tilde{\omega}_\gamma^P) = 0$ . Thus, for each  $\delta \in \Theta$ , we have  $r_{\delta,\gamma} = 0$ , hence  $\delta = r_{\delta,\beta} \beta + r_{\delta,\alpha} \alpha$ .

On the other hand, one has  $\beta_1(\tilde{\omega}_\beta^P) = \beta(\tilde{\omega}_\beta^P) = 1$ . Thus, for all  $\delta \in \Theta$ , one has  $\sum_{\delta \in \Theta} n_\delta r_{\delta, \beta} = 1$ . Since  $n_\delta \in \mathbb{N}^*$  and  $r_{\delta, \beta} \in \mathbb{N}$ , one deduces that there exists a unique  $\delta_0 \in \Theta$  such that  $r_{\delta_0, \beta} \neq 0$  and we have  $n_{\delta_0} = r_{\delta_0, \beta} = 1$ . This implies that  $\Theta = \{\delta_0\}$  and  $\beta = \delta_0 - k_\beta \alpha$ . We can take  $\beta' := \delta_0$ . Hence, we obtain the existence of  $\beta'$  in all cases.

If  $\beta'_1 \in \Delta_{P'}$  satisfies  $\beta'_1 = \beta + k_\beta^1 \alpha$  then  $\beta' = \beta'_1 + (k_\beta - k_\beta^1) \alpha$ . Since the roots  $\beta'_1$ ,  $\beta'$  and  $-\alpha$  belong to the set of simple roots  $\Delta_{P'}$ , we deduce that  $\beta'_1 = \beta'$ . This gives the unicity of  $\beta'$ .

Let  $\gamma$  and  $\beta$  be in  $\Delta_P$  such that  $\gamma' = \beta'$ . Then we have  $\beta = \gamma + (k_\gamma - k_\beta) \alpha$ . Since  $\gamma, \beta$  and  $\alpha$  belong to  $\Delta_P$ , the same argument as above leads to  $\beta = \gamma$ . Hence, the map  $\beta \mapsto \beta'$  is injective.

2. Let  $\beta \in \Delta_P - \{\alpha\}$ . By definition, we have  $\beta' = \beta + k_\beta \alpha \in \Delta_{P'} - \{-\alpha\}$  with  $k_\beta \in \mathbb{N}$ . Thus we have  $\alpha(\tilde{\omega}_{\beta'}^{P'}) = \alpha(\tilde{\omega}_\beta^P) = 0$  and  $\beta(\tilde{\omega}_{\beta'}^{P'}) = \beta'(\tilde{\omega}_{\beta'}^{P'}) = 1$ . If  $\gamma \in \Delta_P - \{\beta, \alpha\}$ , then  $\gamma' = \gamma + k_\gamma \alpha$  is different from  $\beta'$  by (1.), thus we have  $\gamma(\tilde{\omega}_{\beta'}^{P'}) = \gamma'(\tilde{\omega}_{\beta'}^{P'}) = 0$ . One deduces that  $\tilde{\omega}_{\beta'}^{P'} = \tilde{\omega}_\beta^P$ .  $\square$

For  $Y^1$  and  $Y^2$  in  $a_M$ , we denote by  $\inf^P\{Y^1, Y^2\}$  the unique element  $Z$  in  $a_M^H$  such that, for all  $\gamma \in \Delta_P$ , one has  $(\tilde{\omega}_\gamma^P, Z) = \inf\{(\tilde{\omega}_\gamma^P, Y^1), (\tilde{\omega}_\gamma^P, Y^2)\}$ . (2.6)

**2.2 Lemma.** *Let  $\mathcal{Y}^1 = \{Y_P^1, P \in \mathcal{P}(M)\}$  and  $\mathcal{Y}^2 = \{Y_P^2, P \in \mathcal{P}(M)\}$  be two  $(H, M)$ -orthogonal sets. Let  $\mathcal{Z} := \inf(\mathcal{Y}^1, \mathcal{Y}^2)$  be the set of  $Z_P := \inf^P\{Y_P^1, Y_P^2\}$  when  $P$  runs  $\mathcal{P}(M)$ .*

1. *The set  $\mathcal{Z}$  is a  $(H, M)$ -orthogonal set.*
2. *If  $d(\mathcal{Y}^j) > 0$  for  $j = 1, 2$  then  $d(\mathcal{Z}) > 0$ . In this case, the convex hull  $\mathcal{S}_M(\mathcal{Z})$  is the intersection of  $\mathcal{S}_M(\mathcal{Y}^1)$  and  $\mathcal{S}_M(\mathcal{Y}^2)$ .*

*Proof :*

1. Let  $P$  and  $P'$  two adjacent parabolic subgroups in  $\mathcal{P}(M)$  whose chambers in  $a_M$  share the wall determined by the simple root  $\alpha \in \Delta_P \cap (-\Delta_{P'})$ . Let  $\gamma \in \Delta_P - \{\alpha\}$ . By definition of orthogonal sets, for  $j = 1$  or  $2$ , one has  $(\tilde{\omega}_\gamma^P, Y_P^j) = (\tilde{\omega}_\gamma^{P'}, Y_{P'}^j)$ . By Lemma 2.1, we have  $\tilde{\omega}_\gamma^P = \tilde{\omega}_{\gamma'}^{P'}$ . Hence we obtain  $(\tilde{\omega}_\gamma^P, Z_P) = (\tilde{\omega}_{\gamma'}^{P'}, Z_{P'})$  and  $(\tilde{\omega}_{\gamma'}^{P'}, Z_{P'}) = (\tilde{\omega}_\gamma^P, Z_{P'})$ . Since the scalar product on  $a_0$  identifies  $a_M$  to  $a_M^*$ , one deduces that  $Z_P - Z_{P'}$  is proportional to  $\check{\alpha}$ .

2. Let  $j \in \{1, 2\}$  and  $P \in \mathcal{P}(M)$ . By definition, we have  $d(\mathcal{Y}^j) > 0$  if and only if  $\alpha(Y_P^j) > 0$  for all  $\alpha \in \Delta_P$ . By ([Ar1] Corollary 2.2), this implies that  $(\tilde{\omega}_\alpha^P, Y_P^j) > 0$  for all  $\alpha \in \Delta_P$ . Let  $\alpha \in \Delta_P$ . Writing

$$Y_P^j = (\tilde{\omega}_\alpha^P, Y_P^j) \alpha + \sum_{\beta \in \Delta_P - \{\alpha\}} (\tilde{\omega}_\beta^P, Y_P^j) \beta + X^j,$$



with  $X^j \in a_H$ , the condition  $\alpha(Y_P^j) > 0$  is equivalent to

$$\sum_{\beta \in \Delta_P - \{\alpha\}} (\tilde{\omega}_\beta^P, Y_P^j)[-(\beta, \alpha)] < (\tilde{\omega}_\alpha^P, Y_P^j)(\alpha, \alpha).$$

Since the real numbers  $(\tilde{\omega}_\beta^P, Y_P^j)$  for  $\beta \in \Delta_P$  and  $-(\beta, \alpha)$  for  $\alpha \neq \beta$  in  $\Delta_P$  are nonnegative, one deduces that

$$\begin{aligned} \sum_{\beta \in \Delta_P - \{\alpha\}} (\tilde{\omega}_\beta^P, Z_P)[-(\beta, \alpha)] &= \sum_{\beta \in \Delta_P - \{\alpha\}} \inf((\tilde{\omega}_\beta^P, Y_P^1), (\tilde{\omega}_\beta^P, Y_P^2))[-(\beta, \alpha)] \\ &\leq \inf\left(\sum_{\beta \in \Delta_P - \{\alpha\}} (\tilde{\omega}_\beta^P, Y_P^1)[-(\beta, \alpha)], \sum_{\beta \in \Delta_P - \{\alpha\}} (\tilde{\omega}_\beta^P, Y_P^2)[-(\beta, \alpha)]\right) \\ &< \inf((\tilde{\omega}_\alpha^P, Y_P^1), (\tilde{\omega}_\alpha^P, Y_P^2))(\alpha, \alpha) = (\tilde{\omega}_\alpha^P, Z_P)(\alpha, \alpha). \end{aligned}$$

One deduces that  $\alpha(Z_P) > 0$  for  $\alpha \in \Delta_P$ , thus  $d(\mathcal{Z}) > 0$ .

For the property of the convex hulls, it is enough to prove that, for all  $P \in \mathcal{P}(M)$ , one has  $a_P^+ \cap \mathcal{S}_M(\mathcal{Y}^1) \cap \mathcal{S}_M(\mathcal{Y}^2) = a_P^+ \cap \mathcal{S}_M(\mathcal{Z})$ . By ([Ar3], Lemma 3.1), one has

$$a_P^+ \cap \mathcal{S}_M(\mathcal{Y}^j) = \{X \in a_P^+; \omega_\gamma^P(X - Y_P^j) \leq 0, \gamma \in \Delta_P\}.$$

Since  $\tilde{\omega}_\gamma^P = c_\gamma \omega_\gamma^P$  for  $\gamma \in \Delta_P$ , where  $c_\gamma$  is a positive real number, the assertion follows easily.  $\square$

## 2.2 The truncated kernel

We consider the regular representation  $R$  of  $G \times G$  on  $L^2(G)$  defined by

$$(R(y_1, y_2)\phi)(x) = \phi(y_1^{-1}xy_2), \quad \phi \in L^2(G), y_1, y_2 \in G.$$

Consider  $f \in C_c^\infty(G \times G)$  of the form  $f(y_1, y_2) = f_1(y_1)f_2(y_2)$  with  $f_j \in C_c^\infty(G)$ . Then

$$R(f) := \int_G \int_G f_1(y_1)f_2(y_2)R(y_1, y_2)dy_1dy_2$$

is an integral operator with smooth kernel

$$K_f(x, y) = \int_G f_1(xg)f_2(gy)dg = \int_G f_1(g)f_2(x^{-1}gy)dg.$$

In our case ( $H$  is split), one has  $A_H = A_G$ , and the kernel  $K_f$  is invariant by the diagonal  $diag(A_H)$  of  $A_H$ . Since  $H$  is not compact, we introduce truncation to integrate this kernel on  $diag(A_H) \setminus (H \times H)$ .

We fix a point  $T$  in  $a_{0,\mathbb{F}}$ . If  $P_0 \in \mathcal{P}(A_0)$ , let  $T_{P_0}$  be the unique translate by the Weyl group  $W(H, A_0)$  of  $T$  in the closure  $\bar{a}_{P_0}^+$  of the positive Weyl chamber  $a_{P_0}^+$ . Then

$$\mathcal{Y}_T := \{T_{P_0}; P_0 \in \mathcal{P}(A_0)\}$$

is a  $(H, A_0)$ -orthogonal set. We shall assume that the number

$$d(T) := \inf_{\alpha \in \Delta_{P_0}, P_0 \in \mathcal{P}(A_0)} \alpha(T_{P_0})$$

is suitable large. This means that the distance from  $T$  to any of the root hyperplanes in  $a_0$  is large.

We denote by  $u(\cdot, T)$  the characteristic function in  $A_H \setminus H$  of the set of points  $x$  such that

$$x = k_1 a k_2 \text{ with } a \in A_H \setminus A_0, k_1, k_2 \in K \text{ and } h_{A_0}(a) \in \mathcal{S}_{A_0}(\mathcal{Y}_T), \quad (2.7)$$

where  $H = K A_0 K$  is the Cartan decomposition of  $H$ .

We consider  $u(\cdot, T)$  as a  $A_H$ -invariant function on  $H$ . Thus, there is a compact set  $\Omega_T$  of  $H$  such that if  $u(x, T) \neq 0$  then  $x \in A_H \Omega_T$ . Let  $\Omega$  be a compact subset of  $G$  containing the support of  $f_1$  and  $f_2$ . We consider  $g \in G$  and  $x_1, x_2 \in H$  such that  $f_1(g) f_2(x_1^{-1} g x_2) u(x_1, T) u(x_2, T) \neq 0$ . Thus, there are  $\omega_1, \omega_2$  in  $\Omega_T$  and  $a_1, a_2$  in  $A_H$  such that  $x_1 = \omega_1 a_1$ ,  $x_2 = \omega_2 a_2$  and we have  $g \in \Omega$  and  $x_1^{-1} g x_2 = \omega_1^{-1} g \omega_2 a_1^{-1} a_2 \in \Omega$  since  $A_H = A_G$ . One deduces that  $a_1^{-1} a_2$  lies a compact subset of  $A_H$ . Therefore the map  $(g, x_1, x_2) \mapsto f_1(g) f_2(x_1^{-1} g x_2) u(x_1, T) u(x_2, T)$  is a compactly supported function on  $G \times \text{diag}(A_H) \setminus (H \times H)$ .

Hence, we can define

$$K^T(f) := \int_{\text{diag}(A_H) \setminus H \times H} K_f(x_1, x_2) u(x_1, T) u(x_2, T) d\overline{(x_1, x_2)}.$$

By Fubini's Theorem, we have

$$K^T(f) = \int_G \int_{\text{diag}(A_H) \setminus H \times H} f_1(g) f_2(x_1^{-1} g x_2) u(x_1, T) u(x_2, T) d\overline{(x_1, x_2)} dg.$$

We apply the Weyl integration formula (1.33). Thus, we obtain

$$K^T(f) = \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} K^T(x_m, \gamma, f) d\gamma, \quad (2.8)$$

where, for  $S \in \mathcal{T}_M$ ,  $x_m \in \kappa_S$  and  $\gamma \in S_\sigma$ , we have

$$K^T(x_m, \gamma, f) = |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_M) \setminus H \times H} \int_{\text{diag}(A_H) \setminus H \times H} f_1(y_1^{-1} x_m \gamma y_2)$$

$$\times f_2(x_1^{-1}y_1^{-1}x_m\gamma y_2x_2)u(x_1, T)u(x_2, T)\overline{d(x_1, x_2)}\overline{d(y_1, y_2)}.$$

We recall that each  $x_m$  in  $\kappa_S$  and  $\gamma$  in  $S_\sigma$  commute with  $A_M$  for  $S \in \mathcal{T}_M$ .

We first replace  $(x_1, x_2)$  by  $(y_1x_1, y_2x_2)$  in the integral over  $(x_1, x_2)$ . The resulting integral over  $\text{diag}(A_H)\backslash H \times H$  can be expressed as a double integral over  $a \in A_H\backslash A_M$  and  $(x_1, x_2) \in \text{diag}(A_M)\backslash H \times H$  which depends on  $(y_1, y_2) \in \text{diag}(A_M)\backslash H \times H$ . Since  $A_M$  commutes with  $x_m \in \kappa_S$  and  $\gamma \in S_\sigma$ , we obtain

$$\begin{aligned} K^T(x_m, \gamma, f) &= |\Delta_\sigma(x_m\gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_M)\backslash H \times H} \int_{\text{diag}(A_M)\backslash H \times H} f_1(y_1^{-1}x_m\gamma y_2) \\ &\quad \times f_2(x_1^{-1}x_m\gamma x_2)u_M(x_1, y_1, x_2, y_2, T)\overline{d(x_1, x_2)}\overline{d(y_1, y_2)} \end{aligned} \quad (2.9)$$

where

$$u_M(x_1, y_1, x_2, y_2, T) = \int_{A_H\backslash A_M} u(y_1^{-1}ax_1, T)u(y_2^{-1}ax_2, T)da.$$

Our goal is to prove that  $K^T(f)$  is asymptotic to an expression  $J^T(f)$  where  $J^T(f)$  is obtained in a similar way to  $K^T(f)$  where we replace the weight function  $u_M(x_1, y_1, x_2, y_2, T)$  by another weight function  $v_M(x_1, y_1, x_2, y_2, T)$  defined as follows.

We fix  $M \in \mathcal{L}(A_0)$  and  $P \in \mathcal{P}(M)$ . Let  $P_0 \in \mathcal{P}(A_0)$  be contained in  $P$ . We denote by  $T_P$  the projection of  $T_{P_0}$  on  $a_M$  according to the decomposition  $a_0 = a_M \oplus a_0^M$ . By (2.2), the set  $\mathcal{Y}_M(T) := \{T_P; P \in \mathcal{P}(M)\}$  is a  $(H, M)$ -orthogonal set independent of the choices of  $P_0$ . Moreover, by ([Ar3] (3.2)), we have  $d(\mathcal{Y}_M(T)) \geq d(T) > 0$ . Thus,  $\mathcal{Y}_M(T)$  is positive.

For  $x, y$  in  $H$ , we set

$$Y_P(x, y, T) := T_P + h_P(y) - h_{\overline{P}}(x).$$

By ([Ar3], page 30), the set  $\mathcal{Y}_M(x, y, T) := \{Y_P(x, y, T); P \in \mathcal{P}(M)\}$  is a  $(H, M)$ -orthogonal set, which is positive when  $d(T)$  is sufficiently large relative to  $x$  and  $y$ .

For  $x_1, x_2, y_1$  and  $y_2$  in  $H$ , we set

$$Z_P(x_1, y_1, x_2, y_2, T) := \inf^P(Y_P(x_1, y_1, T), Y_P(x_2, y_2, T)) \quad (2.10)$$

where  $\inf^P$  is defined in (2.6) and

$$\mathcal{Y}_M(x_1, y_1, x_2, y_2, T) := \{Z_P(x_1, y_1, x_2, y_2, T); P \in \mathcal{P}(M)\}. \quad (2.11)$$

By Lemma 2.6, the set  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  is a  $(H, M)$ -orthogonal set. Moreover, when  $d(T)$  is large relative to  $x_i, y_i$ , for  $i = 1, 2$ , one has  $d(\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) > 0$ ,

hence this set is positive. We define the weight function  $v_M$  by

$$v_M(x_1, y_1, x_2, y_2, T) := \int_{A_H \backslash A_M} \sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) da \quad (2.12)$$

where  $\sigma_M$  is defined in (2.4).

We set

$$J^T(f) := \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} J^T(x_m, \gamma, f) d\gamma, \quad (2.13)$$

where

$$\begin{aligned} J^T(x_m, \gamma, f) &= |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_M) \backslash H \times H} \int_{\text{diag}(A_M) \backslash H \times H} f_1(y_1^{-1} x_m \gamma y_2) \\ &\quad \times f_2(x_1^{-1} x_m \gamma x_2) v_M(x_1, y_1, x_2, y_2, T) d(x_1, x_2) d(y_1, y_2). \end{aligned} \quad (2.14)$$

Our main result is the following. We will prove it in section 2.4.

**2.3 Theorem.** *Let  $\delta > 0$ . Then, there are positive numbers  $C$  and  $\varepsilon$  such that for all  $T$  with  $d(T) \geq \delta \|T\|$ , one has*

$$|K^T(f) - J^T(f)| \leq C e^{-\varepsilon \|T\|}. \quad (2.15)$$

## 2.3 Preliminaries to estimates

We fix a norm  $\|\cdot\|$  on  $G$  as in (1.15). Let  $\mathbb{F}'$  be a finite extension of  $\mathbb{F}$ . We set  $\underline{G}' := \underline{G} \times_{\mathbb{F}} \mathbb{F}'$  and  $G' := \underline{G}'(\mathbb{F}')$ . One can extend the absolute value  $|\cdot|_{\mathbb{F}}$  to  $\mathbb{F}'$ , and the norm  $\|\cdot\|$  to  $G'$ . For  $x, y$  in  $G'$ , we set

$$\|(x, y)\| := \|x\| \|y\|.$$

To obtain our estimates, we will use notation of (1.18) and (1.19). Since the norm takes values greater or equal to 1, we will freely apply the properties (1.20).

**2.4 Lemma.** *Let  $S$  be a maximal torus of  $H$  and let  $M$  be the centralizer of  $A_S$  in  $H$ . We fix  $x_m \in G \cap \underline{MS}_\sigma = \tilde{M} \cap \underline{MS}_\sigma$ . Then, one has*

$$\inf_{s \in S} \|(s x_m^{-1} x_1, s x_2)\| \leq \inf_{s' \in \underline{S}(\mathbb{F}')} \|(s' x_m^{-1} x_1, s' x_2)\|, \quad x_1, x_2 \in H. \quad (2.16)$$

*Proof :*

Since  $H^1 A_H$  is of finite index in  $H$ , using (1.21) we may assume that  $x_1, x_2$  belong to  $H^1 A_H$ . Since  $A_G = A_H$ , using the invariance of the property by the left action of  $\text{diag}(A_H)$  on  $(x_1, x_2)$ , it is enough to prove the result for  $x_1 \in H^1$  and  $x_2 = a_2 y_2$  with  $a_2 \in A_H$  and  $y_2 \in H^1$ .

To establish (2.16), we first assume that  $A_S = A_H$  which implies that the quotient  $A_H \backslash S$  is compact. By (1.21), there is a positive constant  $C$  such that

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| \leq C \inf_{a \in A_H} \|(ax_m^{-1}x_1, ax_2)\|.$$

We deduce from (1.17) that

$$\|(ax_m^{-1}x_1, ax_2)\| \leq \|x_m^{-1}\| \|a\|^2 \|a_2\| \|x_1\| \|y_2\|.$$

Taking the lower bound in  $a \in A_H$ , there is a positive constant  $C_1$  such that

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| \leq C_1 \|x_1\| \|a_2\| \|y_2\|. \quad (2.17)$$

We now use the following Lemma of [Ar3] (Lemma 4.1):

If  $S_0$  is a maximal torus of  $H$  with  $A_H \backslash S$  compact, then there exists an element  $s_0 \in S_0$  such that

$$\|y\| \leq \|y^{-1}s_0y\|, \quad y \in H^1. \quad (2.18)$$

We apply this Lemma to  $S_0 = S$ . Since  $\underline{S}(F')$  commutes with  $s_0$ , using the property (1.17) of the norm, one deduces

$$\|y_2\| \leq \|s'y_2\|^2 \|s_0\|, \quad y_2 \in H^1, s' \in \underline{S}(F'). \quad (2.19)$$

On the other hand  $S_1 := x_m S x_m^{-1}$  is a maximal torus of  $H$  which satisfies  $A_{S_1} = A_H$  since  $x_m \in G \cap \underline{MS}_\sigma$ . Applying (2.18) to  $S_0 = S_1$ , there exists  $s_1 \in S$  such that

$$\|x_1\| \leq \|x_1^{-1}x_m s_1 x_m^{-1}x_1\|, \quad x_1 \in H^1. \quad (2.20)$$

The same argument as above leads to

$$\|x_1\| \leq \|s'x_m^{-1}x_1\|^2 \|s_1\|, \quad x_1 \in H^1, s' \in \underline{S}(F'). \quad (2.21)$$

Then, by (2.17), (2.19) and (2.21), and applying the properties (1.20), we deduce that

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sa_2y_2)\| \leq \|s'x_m^{-1}x_1\| \|s'y_2\| \|a_2\|, \quad s' \in \underline{S}(F'), x_1, y_2 \in H^1, a_2 \in A_H. \quad (2.22)$$

To obtain our result, we have to prove that

$$\|s'x_m^{-1}x_1\| \|s'y_2\| \|a_2\| \leq \|(s'x_m^{-1}x_1, s'a_2y_2)\|, \quad s' \in \underline{S}(F'), x_1, y_2 \in H^1, a_2 \in A_H. \quad (2.23)$$

We can write  $\underline{S} = \underline{T}A_H$  where  $\underline{T}$  is a maximal torus of the derived group  $\underline{H}_{der}$  of

$\underline{H}$ . We set  $T' := \underline{T}(F')$  and  $A'_H := \underline{A}_H(F')$ . Then  $T'$  is contained in  $H^1$ . Moreover, the intersection of  $\underline{T}$  and  $\underline{A}_H$  is finite. Hence, one has the exact sequence

$$1 \rightarrow \underline{T} \cap \underline{A}_H \rightarrow \underline{T} \times \underline{A}_H \rightarrow \underline{S} \rightarrow 1.$$

Going to  $F'$ -points, the long exact sequence in cohomology implies that  $T'A'_H$  is of finite index in  $\underline{S}(F')$ . By (1.21), it is enough to prove (2.23) for  $s' = t'a' \in \underline{S}(F')$  with  $t' \in T'$  and  $a' \in A'_H$ . By (1.5), if  $x_1 \in H^1$  then  $x_1 \in H^1 \subset G'^1$  and  $x_m^{-1}x_1x_m \in G'^1$ . Since  $H$  is split, we have  $A'_H = A'_G$ . Then (1.23) gives

$$\|a't'x_m^{-1}x_1\| \approx \|a't'x_m^{-1}x_1x_m\| \approx \|a'\| \|t'x_m^{-1}x_1x_m\|, \quad a' \in A'_H, t' \in T', x_1 \in H^1,$$

and

$$\|a't'y_2\| \approx \|a'\| \|t'y_2\| \quad a' \in A'_H, t' \in T', y_2 \in H^1.$$

Applying (1.20), we deduce that

$$\|t'a'x_m^{-1}x_1\| \|a't'y_2\| \|a_2\| \approx \|a_2\| \|a'\|^2 \|t'x_m^{-1}x_1x_m\| \|t'y_2\| \approx \|a_2\| \|a'\| \|t'x_m^{-1}x_1x_m\| \|t'y_2\|, \quad (2.24)$$

for  $t' \in T', a' \in A'_H, x_1, y_2 \in H^1, a_2 \in A_H$ .

Let us prove that

$$\|a'\| \|a'a_2\| \approx \|a'\| \|a_2\|, \quad a' \in A'_H, a_2 \in A_H. \quad (2.25)$$

We have  $\|a'a_2\| \leq \|a'\| \|a_2\|$  by (1.17). Then  $\|a'\| \|a'a_2\| \leq (\|a'\| \|a_2\|)^2$  since  $1 \leq \|a_2\|$ . As  $\|a'\| = \|a'a_2a_2^{-1}\| \leq \|a'a_2\| \|a_2\|$ , we have  $\|a'\| \|a_2\| \leq (\|a'a_2\| \|a_2\|)^2$  and (2.25) follows. Applying (2.25) in (2.24), we deduce that

$$\|t'a'x_m^{-1}x_1\| \|a't'y_2\| \|a_2\| \leq \|a'\| \|t'x_m^{-1}x_1x_m\| \|a'a_2\| \|t'y_2\|, \quad (2.26)$$

for  $t' \in T', a' \in A'_H, x_1, y_2 \in H^1, a_2 \in A_H$ .

Since  $x_m^{-1}H^1x_m \subset G'^1$  and  $A'_H = A'_G$ , we obtain from (1.23)

$$\|a'\| \|t'x_m^{-1}x_1x_m\| \approx \|a't'x_m^{-1}x_1x_m\| \approx \|a't'x_m^{-1}x_1\|, \quad a' \in A'_H, t' \in T', x_1 \in H^1,$$

and

$$\|a'a_2\| \|t'y_2\| \approx \|a'a_2t'y_2\|, \quad a' \in A'_H, t' \in T', a_2 \in A_H, y_2 \in H^1.$$

Applying this in (2.26) and using (1.20), we deduce that

$$\|t'a'x_m^{-1}x_1\| \|a't'y_2\| \|a_2\| \leq \|a't'x_m^{-1}x_1\| \|a't'a_2y_2\| \quad (2.27)$$

for  $a' \in A'_H, t' \in T', x_1, y_2 \in H^1$ .

Then, the property (2.23) follows. This finishes the proof of the Lemma when  $A_H \setminus S$  is compact.

We now prove (2.16) for any maximal torus  $S$  of  $H$ . Let  $A_S$  be the maximal split torus of  $S$  and  $M$  be the centralizer of  $A_S$  in  $H$ . Thus we have  $A_M = A_S$  and  $A_M \setminus S$  is compact. Let  $P = MN_P \in \mathcal{P}(M)$  and let  $K$  be a compact subgroup of  $H$  such that  $H = PK$ . Each  $x \in H$  can be written  $x = m_P(x)n_P(x)k(x)$  with  $m_P(x) \in M, n_P(x) \in N_P$  and  $k(x) \in K$ . Then, there is a positive constant  $C$  such that

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| \leq C \inf_{s \in S} (\|sx_m^{-1}m_P(x_1)\| \|sm_P(x_2)\|) \|n_P(x_1)\| \|n_P(x_2)\|, \quad (2.28)$$

for  $x_1, x_2 \in H$ . By assumption on  $x_m$ , there are  $h_m \in \underline{M}$  and  $a_m \in \underline{S}_\sigma$  such that  $x_m = h_m a_m \in \tilde{M}$ . Hence, we can apply the first part of the proof to  $(M, S)$  instead of  $(H, S)$ . Therefore, we obtain

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| \leq \inf_{s' \in \underline{S}(F')} (\|s'x_m^{-1}m_P(x_1)\| \|s'm_P(x_2)\|) \|n_P(x_1)\| \|n_P(x_2)\|, \quad x_1, x_2 \in H.$$

To compare the right hand-side of this inequality to those of (2.16), we will use the Iwasawa decomposition (1.12) of  $H'$ . Let  $K'$  be a compact subgroup of  $H'$  such that  $H' = \underline{P}(F')K' = \underline{M}(F')\underline{N}_P(F')K'$ . According to (1.13), each  $y$  in  $H'$  can be written  $y = m'_P(y)n'_P(y)k'$  with  $m'_P(y) \in \underline{M}(F'), n'_P(y) \in \underline{N}_P(F')$  and  $k' \in K'$ . Then for  $x \in H$  and  $z \in \underline{M}(F')$ , we have  $zx = zm_P(x)n_P(x)k = m'_P(zx)n'_P(zx)k'$  with  $k \in K$  and  $k' \in K'$ . Hence, since  $K$  and  $K'$  are compact subsets, there is a positive constant  $C'$  such that

$$\|n'_P(zx)^{-1}m'_P(zx)^{-1}zm_P(x)n_P(x)\| \leq C', \quad z \in \underline{M}(F'), x \in H.$$

Since  $zm_P(x) \in \underline{M}(F')$  for  $z \in \underline{M}(F')$  and  $x \in H$ , we deduce from (1.22) that there is a positive constant  $C_1$  such that for  $x \in H$  and  $z \in \underline{M}(F')$ , one has

$$\|n'_P(zx)^{-1}m'_P(zx)^{-1}zm_P(x)n'_P(zx)\| \leq C_1 \quad \text{and} \quad \|n'_P(zx)^{-1}n_P(x)\| \leq C_1.$$

By (1.17), we obtain

$$\|zm_P(x)\| \leq C_1 \|m'_P(zx)\| \|n'_P(zx)\|^2 \quad \text{and} \quad \|n_P(x)\| \leq C_1 \|n'_P(zx)\|.$$

Using (1.22) again, it follows that

$$\|zm_P(x)\| \leq \|zx\|, \quad \text{and} \quad \|n_P(x)\| \leq \|zx\|, \quad z \in \underline{M}(F'), x \in H,$$

hence by (1.20)

$$\|zm_P(x)\| \|n_P(x)\| \leq \|zx\|, \quad z \in \underline{M}(F'), x \in H. \quad (2.29)$$

We deduce that

$$\|s'm_P(x_2)\| \|n_P(x_2)\| \leq \|s'x_2\|, \quad s' \in \underline{S}(F'), x_2 \in H. \quad (2.30)$$

Since  $x_m = h_m a_m$  with  $h_m \in \underline{M}$  and  $a_m \in \underline{S}_\sigma$ , one has  $x_m s' x_m^{-1} \in \underline{M} \cap H' = \underline{M}(F')$  for  $s' \in \underline{S}(F')$ . Therefore, we deduce from (2.29) that

$$\|x_m s' x_m^{-1} m_P(x_1)\| \|n_P(x_1)\| \leq \|x_m s' x_m^{-1} x_1\|, \quad s' \in \underline{S}(F'), x_1 \in H. \quad (2.31)$$

Since  $\|s' x_m^{-1} m_P(x_1)\| \leq \|x_m^{-1}\| \|x_m s' x_m^{-1} m_P(x_1)\|$  and  $\|x_m s' x_m^{-1} x_1\| \leq \|x_m\| \|s' x_m^{-1} x_1\|$ , we deduce the estimate (2.16) from (2.28), (2.30) and (2.31). This finishes the proof of the Lemma.  $\square$

The following Lemma is the analogue of Lemma 4.2 of ([Ar3]).

**2.5 Lemma.** *Let  $S$  be a maximal torus of  $H$  and let  $x_m \in \kappa_S$ . Then, there is a positive integer  $k$  with the property that, for any given compact subset  $\Omega$  of  $G$ , there exists a positive constant  $C_\Omega$  such that, for all  $\gamma \in S_\sigma$  with  $x_m \gamma \in G^{\sigma\text{-reg}}$ , and all  $x_1, x_2$  in  $H$  satisfying  $x_1^{-1} x_m \gamma x_2 \in \Omega$ , one has*

$$\inf_{s \in S} \|(s x_m^{-1} x_1, s x_2)\| \leq C_\Omega |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-k}.$$

*Proof :*

Let  $F'$  be a finite extension of  $\mathbb{E}$  such that  $\tilde{S}$  splits over  $F'$ . Recall that we can write  $x_m = h_m a_m$  with  $h_m \in \underline{H}$  and  $a_m \in \underline{S}_\sigma$ . Thus we may and will choose  $F'$  such that  $h_m \in \underline{H}(F')$  and  $a_m \in \underline{S}_\sigma(F')$ . For convenience of lecture, if  $\underline{J}$  is an algebraic variety defined over  $\mathbb{F}$ , we set  $J' := \underline{J}(F')$ .

By the previous Lemma 2.4, it is enough to prove the existence of a positive integer  $k$  satisfying the property that for any compact subset  $\Omega'$  of  $G'^{\sigma\text{-reg}}$ , there exists  $C_{\Omega'} > 0$  such that

$$\inf_{s' \in S'} (\|s' x_m^{-1} x_1\| \|s' x_2\|) \leq C_{\Omega'} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-k} \quad (2.32)$$

for all  $x_1, x_2 \in H'$  and  $\gamma \in S_\sigma$  satisfying  $x_m \gamma \in G^{\sigma\text{-reg}}$  and  $x_1^{-1} x_m \gamma x_2 \in \Omega'$ .

Let  $B' = S' N'$  be a Borel subgroup of  $H'$  containing  $S'$  and  $K'$  be a compact subgroup of  $H'$  such that  $H' = S' N' K' = N' S' K'$ . We can also write  $H' = (h_m S' h_m^{-1})(h_m N' h_m^{-1})(h_m K' h_m^{-1})$ . By (1.21), one can reduce the proof of the statement for  $x_1 \in (h_m S' h_m^{-1})(h_m N' h_m^{-1})$  and  $x_2 \in S' N'$ .

Let  $x_1 = h_m s_1 n_1 h_m^{-1}$  and  $x_2 = s_1 s_2 n_2$  with  $s_1, s_2 \in S'$  and  $n_1, n_2 \in N'$ . Since  $x_m = h_m a_m$ , we have  $x_m s_1 x_m^{-1} = h_m s_1 h_m^{-1}$ , hence for  $s' \in S'$ , we have  $s' x_m^{-1} x_1 = s' x_m^{-1} x_m s_1 x_m^{-1} h_m n_1 h_m^{-1} = s' s_1 x_m^{-1} h_m n_1 h_m^{-1}$ . We obtain

$$\inf_{s' \in S'} (\|s' x_m^{-1} x_1\| \|s' x_2\|) = \inf_{s' \in S'} (\|s' x_m^{-1} h_m n_1 h_m^{-1}\| \|s' s_2\|).$$

Notice that  $x_1^{-1} x_m \gamma x_2 = h_m n_1^{-1} h_m^{-1} x_m s_1^{-1} x_m^{-1} x_m \gamma s_1 s_2 n_2 = h_m n_1^{-1} h_m^{-1} x_m \gamma s_2 n_2$ .

Therefore, we are reduced to prove (2.32) for  $x_1 = h_m n_1 h_m^{-1}$  with  $n_1 \in N'$ ,  $x_2 = s_2 n_2$  with  $n_2 \in N'$ ,  $s_2 \in S'$  and  $\gamma \in S_\sigma$  such that  $x_m \gamma$  is  $\sigma$ -regular and



$x_1^{-1}x_m\gamma x_2 \in \Omega'$ . By the properties of the norm, there is some positive constant  $C'$  such that

$$\inf_{s' \in S'} (\|s'x_m^{-1}x_1\| \|s'x_2\|) \leq C' \|n_1\| \|s_2\| \|n_2\|, \quad x_1 = h_m n_1 h_m^{-1}, x_2 = s_2 n_2. \quad (2.33)$$

We want to estimate  $\|n_1\| \|s_2\| \|n_2\|$  when  $x_1 = h_m n_1 h_m^{-1}$  and  $x_2 = s_2 n_2$  satisfy  $x_1^{-1}x_m\gamma x_2 \in \Omega'$ . For this, we use the isomorphism  $\Psi$  from  $G'$  to  $H' \times H'$  defined in (1.26). If  $x \in H'$  then  $\Psi(x) = (x, x)$  and if  $y \in G$  satisfies  $y^{-1} = \sigma(y)$  then  $\Psi(y) = (y, y^{-1})$ . We set  $(y_1, y_2) := \Psi(x_1^{-1}x_m\gamma x_2)$ . Then, we have

$$y_1 = h_m n_1^{-1} a_m \gamma n_2 s_2 = h_m (n_1^{-1} a_m \gamma n_2 (a_m \gamma)^{-1}) (a_m \gamma s_2),$$

and

$$y_2 = h_m n_1^{-1} a_m^{-1} \gamma^{-1} n_2 s_2 = h_m (n_1^{-1} a_m^{-1} \gamma^{-1} n_2 \gamma a_m) (a_m \gamma)^{-1} s_2.$$

Since  $H' = N' S' K'$ , the condition  $x_1^{-1}x_m\gamma x_2 \in \Omega'$  implies that there exist two compact subsets  $\Omega_N \subset N'$  and  $\Omega_S \subset S'$  depending only on  $\Omega'$  such that

$$n_1^{-1} a_m \gamma n_2 (a_m \gamma)^{-1} \in \Omega_N, \quad \text{and} \quad n_1^{-1} a_m^{-1} \gamma^{-1} n_2 \gamma a_m \in \Omega_N,$$

$$a_m \gamma s_2 \in \Omega_S, \quad \text{and} \quad (a_m \gamma)^{-1} s_2 \in \Omega_S.$$

We deduce from the second property that  $s_2$  and  $\gamma$  must lie in compact subsets of  $S'$ . We set

$$\nu_1(\gamma, n_1, n_2) := n_1^{-1} a_m \gamma n_2 (a_m \gamma)^{-1} \quad \text{and} \quad \nu_2(\gamma, n_1, n_2) := n_1^{-1} (a_m \gamma)^{-1} n_2 a_m \gamma.$$

We consider the map  $\psi$  from  $N' \times N'$  into itself defined by  $\psi(n_1, n_2) = (\nu_1, \nu_2)$ . Recall that  $\Phi(S', \mathfrak{h}')$  denotes the set of roots of  $S'$  in the Lie algebra  $\mathfrak{h}'$  of  $H'$  (cf. 1.27). Let  $\mathfrak{n}'$  be the Lie algebra of  $N'$ . For  $\alpha \in \Phi(S', \mathfrak{h}')$ , we denote by  $X_\alpha \in \mathfrak{n}'$  the root vector in  $\mathfrak{h}'$  corresponding to  $\alpha$ . Then  $(a_m \gamma)$  acts on  $X_\alpha$  by  $a_\alpha := (a_m \gamma)^\alpha$ . The differential  $d_{(n_1, n_2)} \psi$  of  $\psi$  at  $(n_1, n_2) \in N' \times N'$  is given by  $d_{(n_1, n_2)} \psi(X_1, X_2) = (\text{Ad}(a_m \gamma n_2^{-1} (a_m \gamma)^{-1}) Y_1, \text{Ad}((a_m \gamma)^{-1} n_2^{-1} a_m \gamma) Y_2)$  where

$$Y_1 = -\text{Ad}(n_1) X_1 + \text{Ad}(a_m \gamma) \text{Ad}(n_2) X_2$$

and

$$Y_2 = -\text{Ad}(n_1) X_1 + \text{Ad}(a_m \gamma)^{-1} \text{Ad}(n_2) X_2.$$

The map  $(X_1, X_2) \mapsto (Y_1, Y_2)$  is the composition of the map  $(X_1, X_2) \mapsto (\text{Ad}(n_1) X_1, \text{Ad}(n_2) X_2)$ , whose determinant is equal to 1, with  $d_e \psi$  where  $e$  is the neutral point of  $N' \times N'$ . We deduce that the jacobian of  $\psi$  at  $(n_1, n_2)$  is independent of  $(n_1, n_2)$ . At the neutral point  $e \in N' \times N'$ , we have  $d_e \psi(X_\alpha, 0) = (-X_\alpha, -X_\alpha)$  and  $d_e \psi(0, X_\alpha) = (a_\alpha X_\alpha, a_{-\alpha} X_\alpha)$ . Hence, the jacobian of  $\psi$  is equal to

$$\left| \prod_{\alpha \in \Phi(S', \mathfrak{h}')} a_\alpha (1 - a_{-2\alpha}) \right|_{\mathbb{F}'} = |\det(\text{Ad}(a_m \gamma))_{\mathfrak{h}'/\mathfrak{s}'}|_{\mathbb{F}'} |\det(1 - \text{Ad}(a_m \gamma)^{-2})_{\mathfrak{h}'/\mathfrak{s}'}|_{\mathbb{F}'} = |D_{H'}((a_m \gamma)^{-2})|_{\mathbb{F}'}.$$

Recall that  $x_m\gamma$  is assumed to be  $\sigma$ -regular. Thus, by (1.29), one has  $\Delta_\sigma(x_m\gamma) = D_{H'}(a_m^{-2}\gamma^{-2}) \neq 0$ . Then, arguing as in ([HC2] proof of Lemma 10 and Lemma 11), we deduce that the map  $\psi$  is an  $F'$ -rational isomorphism of  $\underline{N} \times \underline{N}$  to itself whose inverse  $(\nu_1, \nu_2) \mapsto (n_1, n_2) := (n_1(\gamma, \nu_1, \nu_2), n_2(\gamma, \nu_1, \nu_2))$  is rational. Moreover, there is a positive integer  $k$  such that the map

$$(y, \nu_1, \nu_2) \mapsto D_{\underline{H}}(y)^k(n_1(y, \nu_1, \nu_2), n_2(y, \nu_1, \nu_2))$$

is defined by an  $F'$ -rational morphism between the algebraic varieties  $\underline{S} \times \underline{N} \times \underline{N}$  and  $\underline{N} \times \underline{N}$ . Since  $\nu_1, \nu_2$  and  $\gamma$  lie in compact subsets depending only on  $\Omega'$ , one deduces that there exists a constant  $C_{\Omega'} > 0$  such that

$$\|(n_1(\gamma, \nu_1, \nu_2), n_2(\gamma, \nu_1, \nu_2))\| \leq C_{\Omega'} |D_{H'}(a_m^{-2}\gamma^{-2})|_{F'}^{-k} = C_{\Omega'} |\Delta_\sigma(x_m\gamma)|_{F'}^{-k}.$$

The Lemma follows from (2.33) and the fact that  $s_2$  lies in a compact set.  $\square$

## 2.4 Proof of Theorem 2.3

Our goal is to prove that  $|K^T(f) - J^T(f)|$  is bounded by a function which approaches 0 as  $T$  approaches infinity. By definition,  $K^T(f)$  and  $J^T(f)$  are finite linear combinations of  $\int_{S_\sigma} K^T(x_m, \gamma, f) d\gamma$  and  $\int_{S_\sigma} J^T(x_m, \gamma, f) d\gamma$  respectively, where  $M \in \mathcal{L}(A_0)$ ,  $S$  is a maximal torus of  $M$  satisfying  $A_S = A_M$  and  $x_m \in \kappa_S$  (cf. (2.8) and (2.13)).

We fix  $M \in \mathcal{L}(A_0)$  and a maximal torus  $S$  of  $M$  such that  $A_S = A_M$ . Let  $x_m \in \kappa_S$ . To obtain our result, it is enough to establish the estimate (2.15) for  $\int_{S_\sigma} |K^T(x_m, \gamma, f) - J^T(x_m, \gamma, T)| d\gamma$ . This will be done in the Corollary 2.9 below.

For  $\varepsilon > 0$ , we define

$$S_\sigma(\varepsilon, T) := \{\gamma \in S_\sigma; 0 < |\Delta_\sigma(x_m\gamma)|_{F'} \leq e^{-\varepsilon\|T\|}\}. \quad (2.34)$$

**2.6 Lemma.** *1. There exists  $\varepsilon_0 > 0$  such that the map  $\gamma \mapsto |\Delta_\sigma(x_m\gamma)|_{F'}^{-\varepsilon_0}$  is locally integrable on  $S_\sigma$ .*

*2. Let  $\varepsilon > 0$ . Let  $B$  be a bounded subset of  $S_\sigma$  and  $p$  be a nonnegative integer. Then, there is a positive constant  $C_{B,p}$  depending on  $B$  and  $p$ , such that*

$$\int_{S_\sigma(\varepsilon, T) \cap B} |\log |\Delta_\sigma(x_m\gamma)|_{F'}|^{-1} |^p d\gamma \leq C_{B,p} e^{-\frac{\varepsilon\varepsilon_0\|T\|}{2}}.$$

*Proof :*

*1.* The proof follows those of the group case, we use the similar statement on Lie algebras and the exponential map. We denote by  $\mathfrak{s}$  the Lie algebra of  $S$ . For  $X \in \mathfrak{s}$ , we set  $\eta(X) = |\det(\text{ad}X|_{\mathfrak{h}/\mathfrak{s}})|_{F'}$ . By ([HC3] Lemma 44), there exists  $\varepsilon_0 > 0$  such that

$X \mapsto \eta(X)^{-2\varepsilon_0}$  is locally integrable on  $\mathfrak{s}$ . To obtain the result, it is sufficient to prove that

$$\begin{aligned} & \text{for each } \gamma_0 \in S_\sigma, \text{ there exists a compact neighborhood } U_0 \text{ of } 1 \text{ such that} \\ & \text{the integral } \int_{U_0} |\Delta_\sigma(x_m \gamma_0 \gamma)|_{\mathbb{F}}^{-\varepsilon_0} d\gamma \text{ converges.} \end{aligned} \quad (2.35)$$

If  $x_m \gamma_0$  is  $\sigma$ -regular, there is a compact neighborhood  $U_0$  of 1 in  $S_\sigma$  such that  $|\Delta_\sigma(x_m \gamma_0 \gamma)|_{\mathbb{F}} = |\Delta_\sigma(x_m \gamma_0)|_{\mathbb{F}} \neq 0$  for all  $\gamma \in U_0$ . Hence (2.35) is clear.

We assume that  $x_m \gamma_0$  is not  $\sigma$ -regular. We choose an extension  $\mathbb{F}'$  of  $\mathbb{F}$  such that  $\tilde{S}$  splits over  $\mathbb{F}'$  and  $\underline{p}(x_m) \in \tilde{S}_\sigma(\mathbb{F}')$ . We use notation of (1.27). Let  $\Phi_0$  be the set of root  $\alpha$  in  $\Phi(S'_\sigma, \mathfrak{g}')$  such that  $\underline{p}(x_m \gamma_0)^\alpha = 1$ . We set

$$\nu(\gamma) = \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}') - \Phi_0} |1 - \underline{p}(x_m \gamma_0)^\alpha \gamma^{-2\alpha}|_{\mathbb{F}'}^2.$$

We have  $\Delta_\sigma(x_m \gamma_0 \gamma) = D_{G'}(\underline{p}(x_m \gamma_0) \gamma^{-2}) = \det(1 - \text{Ad} \underline{p}(x_m \gamma_0) \gamma^{-2})|_{\mathfrak{g}/\mathfrak{s}}$  and each root of  $\Phi(S'_\sigma, \mathfrak{g}')$  has multiplicity 2. Hence, we obtain

$$|\Delta_\sigma(x_m \gamma_0 \gamma)|_{\mathbb{F}'} = \nu(\gamma) \prod_{\alpha \in \Phi_0} |1 - \gamma^{-2\alpha}|_{\mathbb{F}'}^2.$$

We choose a compact neighborhood  $W$  of 1 in  $S_\sigma$  such that  $\nu(\gamma) = \nu(1) \neq 0$  for  $\gamma \in W$ . Let  $\beta = \sup_{\gamma \in W} \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}') - \Phi_0} |1 - \gamma^{-2\alpha}|_{\mathbb{F}'}^2$ . Then, for  $\gamma \in W$ , we have

$$\beta |\Delta_\sigma(x_m \gamma_0 \gamma)|_{\mathbb{F}'} = \beta \nu(1) \prod_{\alpha \in \Phi_0} |1 - \gamma^{-2\alpha}|_{\mathbb{F}'}^2 \geq \nu(1) |\Delta_\sigma(\gamma)|_{\mathbb{F}'}.$$

Consider the exponential map, there exist two open neighborhoods  $\omega$  and  $U$  of 0 and 1 in  $\mathfrak{s}$  and  $S_\sigma$  respectively, such that the map  $X \mapsto \exp(\tau X)$  is well-defined on  $\omega$  and is an isomorphism and an homeomorphism onto  $U$ . For  $X \in \omega$ , we have

$$\frac{|\Delta_\sigma(\exp(\tau X))|_{\mathbb{F}'}^{1/2}}{\eta(X)} = \prod_{\alpha \in \Phi(S'_\sigma, \mathfrak{g}')} \frac{|1 - e^{2\tau\alpha(X)}|_{\mathbb{F}'}}{|\alpha(X)|_{\mathbb{F}'}}.$$

We can choose a compact neighborhood  $\omega_0 \subset \omega$  of 0 in  $\mathfrak{s}$  such that this product is a positive constant  $c$  and  $U_0 := \exp(\tau \omega_0)$  is contained in  $W$ . We deduce that

$$\int_{U_0} |\Delta_\sigma(x_m \gamma_0 \gamma)|_{\mathbb{F}}^{-\varepsilon_0} d\gamma \leq \left(\frac{\nu(1)}{\beta}\right)^{-\varepsilon_0} \int_{U_0} |\Delta_\sigma(\gamma)|_{\mathbb{F}}^{-\varepsilon_0} d\gamma = \left(\frac{\nu(1)}{\beta}\right)^{-\varepsilon_0} c \int_{\omega_0} \eta(X)^{-2\varepsilon_0} dX.$$

The right hand side of this inequality is finite by our choice of  $\varepsilon_0$ . Hence, we have proved (2.35).

2. Let  $\varepsilon_0 > 0$  as in 1. We set  $I_p = \int_{S_\sigma(\varepsilon, T) \cap B} |\log |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-1}|^p d\gamma$ .

If  $p$  is a positive integer, then there is positive constant  $C'$  such that  $|\log y|^p \leq C' y^{\varepsilon_0/2}$  for all  $y \geq 1$ . Since  $|\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-1} \geq e^{\varepsilon \|T\|} \geq 1$  for all  $\gamma \in S_\sigma(\varepsilon, T)$ , we obtain

$$I_p \leq C' \int_{S_\sigma(\varepsilon, T) \cap B} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-\varepsilon_0/2} d\gamma \leq C' e^{-\frac{\varepsilon \varepsilon_0 \|T\|}{2}} \int_B |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-\varepsilon_0} d\gamma.$$

If  $p = 0$  then by definition of  $S_\sigma(\varepsilon, T)$ , one has

$$I_0 = \int_{S_\sigma(\varepsilon, T) \cap B} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-\varepsilon_0} |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{\varepsilon_0} d\gamma \leq e^{-\varepsilon \varepsilon_0 \|T\|} \int_B |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{-\varepsilon_0} d\gamma.$$

In the two cases, the result follows from 1.  $\square$

**2.7 Lemma.** *Let  $\varepsilon_0 > 0$  as in Lemma 2.6. Given  $\varepsilon > 0$ , we can choose a constant  $c > 0$  such that for any  $T \in a_{0, \mathbb{F}}$ , one has*

$$\int_{S_\sigma(\varepsilon, T)} (|K^T(x_m, \gamma, f)| + |J^T(x_m, \gamma, f)|) d\gamma \leq c e^{-\frac{\varepsilon \varepsilon_0 \|T\|}{4}}.$$

*Proof :*

We recall that

$$\begin{aligned} K^T(x_m, \gamma, f) &= |\Delta_\sigma(x_m \gamma)|^{1/2} \int_{\text{diag}(A_M) \setminus H \times H} \int_{\text{diag}(A_M) \setminus H \times H} f_1(y_1^{-1} x_m \gamma y_2) \\ &\quad \times f_2(x_1^{-1} x_m \gamma x_2) u_M(x_1, y_1, x_2, y_2, T) d(x_1, x_2) d(y_1, y_2) \end{aligned}$$

where

$$u_M(x_1, y_1, x_2, y_2, T) = \int_{A_H \setminus A_M} u(y_1^{-1} a x_1, T) u(y_2^{-1} a x_2, T) da.$$

We first establish an estimate of  $u_M$ . Let  $x, y \in H$  and  $a \in A_M$ . According to (1.11) applied to  $H$ , we can write  $y^{-1} a x = k_1 a_0 k_2$  with  $k_1, k_2 \in K$  and  $a_0 \in A_0$ . By definition of the norm, there is a positive constant  $C_0$  such that

$$\log \|y^{-1} a x\| \leq C_0 (\|h_{A_0}(a_0)\| + 1).$$

If  $u(y^{-1} a x) \neq 0$ , then, by definition of  $u(\cdot, T)$  (cf. (2.7)), the projection of  $h_{A_0}(a_0)$  in  $a_H \setminus a_M$  belongs to the convex hull in  $a_H \setminus a_M$  of the  $W(H, A_0)$ -translates of  $T$ . Thus, there is a constant  $C_1 > 0$  such that

$$\inf_{z \in A_H} \log \|y^{-1} z a x\| \leq C_1 (\|T\| + 1). \quad (2.36)$$

We assume that  $\|T\| \geq 1$ . Taking  $C_2 = \max(2C_1, 1)$  and using the property (1.17) of the norm, we obtain

$$\inf_{z \in A_H} \log \|za\| \leq C_2(\|T\| + \log \|x\| + \log \|y\|). \quad (2.37)$$

We apply this to  $(x_1, y_1)$  and  $(x_2, y_2)$  such that  $u(y_1^{-1}ax_1, T)u(y_2^{-1}ax_2, T) \neq 0$ . Hence, we deduce that

$$\inf_{z \in A_H} \log \|za\| \leq C_2(\|T\| + \log \|x_1\| + \log \|y_1\| + \log \|x_2\| + \log \|y_2\|).$$

As  $\|x\| \leq \|x_m\| \|x_m^{-1}x\|$  and  $1 \leq \|T\|$ , taking the integral over  $a \in A_H \setminus A_M$ , we deduce the following inequality

$$u_M(x_1, y_1, x_2, y_2, T) \leq (\|T\| + \log \|x_m^{-1}x_1\| + \log \|x_m^{-1}y_1\| + \log \|x_2\| + \log \|y_2\|), \quad (2.38)$$

for all  $x_1, y_1, x_2$  and  $y_2$  in  $H$ .

The function  $u_M(x_1, y_1, x_2, y_2, T)$  is invariant by the diagonal (left) action of  $A_M$  on  $(x_1, x_2)$  and  $(y_1, y_2)$  respectively. Since  $x_m$  commutes with  $A_S = A_M$  (cf. Lemma 1.1), we can replace  $\log \|x_m^{-1}x_1\| + \log \|x_2\|$  and  $\log \|x_m^{-1}y_1\| + \log \|y_2\|$  by  $\inf_{a \in A_M} \log \|(ax_m^{-1}x_1, ax_2)\|$  and  $\inf_{a \in A_M} \log \|(ax_m^{-1}y_1, ay_2)\|$  respectively. By assumption, the quotient  $A_M \setminus S$  is compact, then, using (1.21), one has

$$\inf_{a \in A_M} \|(ax_m^{-1}x, ax')\| \approx \inf_{s \in S} \|(sx_m^{-1}x, sx')\|, \quad x, x' \in H.$$

Therefore, as  $\|T\| \geq 1$ , the inequality (2.38) gives

$$u_M(x_1, y_1, x_2, y_2, T) \leq \|T\| + \log \inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| + \log \inf_{s \in S} \|(sx_m^{-1}y_1, sy_2)\|, \quad x_1, y_1, x_2, y_2 \in H.$$

In other words, this means that there are a positive constant  $C_3$  and a positive integer  $d$  such that, for all  $x_1, y_1, x_2$  and  $y_2 \in H$ , one has

$$u_M(x_1, y_1, x_2, y_2, T) \leq C_3(\|T\| + \log \inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| + \log \inf_{s \in S} \|(sx_m^{-1}y_1, sy_2)\|)^d.$$

Let  $\Omega$  be a compact set containing the support of  $f_1$  and  $f_2$ . By Lemma 2.5, there is a positive integer  $k$  (independent of  $\Omega$ ) and a positive constant  $C_\Omega$  such that, if  $x_m\gamma \in x_mS_\sigma$  is a  $\sigma$ -regular point with  $f_1(y_1^{-1}x_m\gamma y_2)f_2(x_1^{-1}x_m\gamma x_2) \neq 0$  for some  $x_1, x_2, y_1$  and  $y_2$  in  $H$  then

$$u_M(x_1, y_1, x_2, y_2, T) \leq C_\Omega(\|T\| + \log |\Delta_\sigma(x_m\gamma)|^{-k})^d.$$

This inequality and the expression of  $K^T(x_m, \gamma, f)$  give

$$|K^T(x_m, \gamma, f)| \leq C_\Omega(\|T\| + \log |\Delta_\sigma(x_m\gamma)|^{-k})^d |\mathcal{M}(f_1)(x_m\gamma)\mathcal{M}(f_2)(x_m\gamma)|, \quad (2.39)$$

where  $\mathcal{M}(f_j)$  is the orbital integral of  $f_j$  defined in (1.34). By Theorem 1.2, these orbital integrals are bounded by a constant  $C_4$  on  $(x_m S_\sigma) \cap G^{\sigma\text{-reg}}$ . Hence, we obtain

$$|K^T(x_m, \gamma, f)| \leq C_\Omega C_4^2 (\|T\| + \log |\Delta_\sigma(x_m \gamma)|^{-k})^d.$$

Let  $B$  be the set of  $\gamma$  in  $S_\sigma$  such that  $K^T(x_m, \gamma, f) \neq 0$ . Then  $B$  is bounded by Theorem 1.2 and (2.39). Using Lemma 2.6, we can find a constant  $C > 0$  such that

$$\int_{S_\sigma(\varepsilon, T)} |K^T(x_m, \gamma, f)| d\gamma \leq C e^{-\frac{\varepsilon \varepsilon_0 \|T\|}{4}}. \quad (2.40)$$

If  $\|T\| \leq 1$ , then (2.36) implies that if  $u(x^{-1}ay) \neq 0$  then

$$\inf_{z \in A_H} \log \|za\| \leq 2C_1 + \log \|x\| + \log \|y\|.$$

The same arguments to obtain (2.38) imply that there is a positive constant  $C'_1$  such that

$$u_M(x_1, y_1, x_2, y_2, T) \leq (C'_1 + \log \|x_m^{-1}x_1\| + \log \|x_m^{-1}y_1\| + \log \|x_2\| + \log \|y_2\|), \quad (2.41)$$

for  $x_1, y_1, x_2$  and  $y_2$  in  $H$ . Replacing  $\|T\|$  by  $C'_1$  in the reasoning after (2.38), we deduce that  $\int_{S_\sigma(\varepsilon, T)} |K^T(x_m, \gamma, f)| d\gamma$  is bounded. Hence, one obtains (2.40) for  $\|T\| \leq 1$ .

We will now establish a similar estimate when  $K^T$  is replaced by  $J^T$ . For this, it is enough to prove that the weight function  $v_M$  have an estimate like (2.38). We will see that this follows easily from the definition of  $v_M$ . Indeed, for  $x_1, y_1, x_2$  and  $y_2$  in  $H$ , one has by definition

$$v_M(x_1, y_1, x_2, y_2, T) := \int_{A_H \setminus A_M} \sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) da$$

where  $\sigma_M(\cdot, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T))$  is a bounded function which vanishes in the complement of the convex hull  $\mathcal{S}_M(\mathcal{Y}_M(x_1, y_1, x_2, y_2, T))$  of the  $(H, M)$ -orthogonal set  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  (cf. (2.5)). Since  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  is the set of points  $Z_P = \inf^P(T_P + h_P(y_1) - h_{\bar{P}}(x_1), T_P + h_P(y_2) - h_{\bar{P}}(x_2))$  for  $P \in \mathcal{P}(M)$  (cf. (2.11)), if  $\sigma_M(X, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) \neq 0$  then  $\|X\| \leq \|Z_P\|$  for  $P \in \mathcal{P}(M)$ . By definition of  $T_P$ , one has  $\|T_P\| \leq \|T\|$ .

Let us prove that for  $P \in \mathcal{P}(M)$ , one has

$$\|h_P(x)\| \leq 1 + \log \|x\|, \quad x \in H. \quad (2.42)$$

We first compare  $\|m\|$  and  $\|h_M(m)\|$  for  $m \in M$ . Let  $M = K_M A_0 K_M$  be the Cartan decomposition of  $M$  where  $K_M$  is a suitable compact subgroup of  $M$ . Then, each  $m \in M$  can be written  $m = ka(m)k'$  with  $k, k' \in K_M$  and  $a(m) \in A_0$ . Since  $K_M$  is

compact, the property (1.21) gives  $\|m\| \approx \|a(m)\|$ ,  $m \in M$  and this property does not depend on our choice of  $a(m)$ . By (1.25), we have  $\|a\| \approx e^{\|h_{A_0}(a)\|}$ ,  $a \in A_0$ .

Hence, there are a positive constant  $C$  and a nonnegative integer  $d$  such that  $e^{\|h_{A_0}(a(m))\|} \leq C\|m\|^d$  for all  $m \in M$ . By (1.8) applied to  $(M, A_0)$ , if  $a \in A_0$  then  $h_M(a)$  is the orthogonal projection of  $h_{A_0}(a)$  onto  $a_M$ , thus  $\|h_M(a)\| \leq \|h_{A_0}(a)\|$ . Since  $h_M(m) = h_M(a(m))$  for  $m \in M$ , we obtain that there is a positive constant  $C'$  such that

$$\|h_M(m)\| \leq \|h_{A_0}(a(m))\| \leq C'(1 + \log \|m\|), \quad m \in M. \quad (2.43)$$

By definition (cf. (1.13), (1.14)), we have  $h_P(x) = h_M(m_P(x))$  for  $x \in H$  and by (1.22), we have  $\|m_P(x)\| \leq \|x\|$ ,  $x \in H$ . Thus, our claim (2.42) follows from (2.43).

Therefore, there are a positive  $C_1$  and a positive integer  $d$  such that if  $\sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) \neq 0$ , then

$$\|h_M(a)\| \leq \|Z_P\| \leq C_1(\|T\| + \log \|x_1\| + \log \|y_1\| + \log \|x_2\| + \log \|y_2\|)^d.$$

Since  $\|x\| \leq \|x_m\| \|x_m^{-1}x\|$  for  $x \in H$ , this gives the following estimates of  $v_M$  analogous to (2.38) and (2.41):

If  $\|T\| > 1$  then

$$v_M(x_1, y_1, x_2, y_2, T) \leq \|T\| + \log \|x_m^{-1}x_1\| + \log \|x_m^{-1}y_1\| + \log \|x_2\| + \log \|y_2\|, \quad (2.44)$$

$x_1, y_1, x_2, y_2 \in H,$

and

there is a positive constant  $C'_2$  such that for  $\|T\| \leq 1$ , one has

$$v_M(x_1, y_1, x_2, y_2, T) \leq C'_2 + \log \|x_m^{-1}x_1\| + \log \|x_m^{-1}y_1\| + \log \|x_2\| + \log \|y_2\|, \quad (2.45)$$

$x_1, y_1, x_2, y_2 \in H.$

Arguing exactly as above for  $K^T$ , we deduce that there is a positive constant  $C'$  such that

$$\int_{S_\sigma(\varepsilon, T)} |J^T(x_m, \gamma, f)| d\gamma \leq C' e^{-\frac{\varepsilon \varepsilon_0 \|T\|}{4}}.$$

This finishes the proof of the Lemma. □

**2.8 Lemma.** *Fix  $\delta > 0$ . Then, there exist positive numbers  $C, \varepsilon_1$  and  $\varepsilon_2$  such that, for all  $T$  with  $d(T) \geq \delta \|T\|$ , and for all  $x_1, y_1, x_2$  and  $y_2$  in the set  $H_{\varepsilon_2} := \{x \in H; \|x\| \leq e^{\varepsilon_2 \|T\|}\}$ , one has*

$$|u_M(x_1, y_1, x_2, y_2, T) - v_M(x_1, y_1, x_2, y_2, T)| \leq C e^{-\varepsilon_1 \|T\|}. \quad (2.46)$$

*Proof :*

If  $\|T\|$  remains bounded then, by (2.38), (2.41), (2.44) and (2.45), the functions  $u_M$  and  $v_M$  are bounded and the result (2.46) is trivial. Thus we have to prove the Lemma for  $\|T\|$  sufficiently large and  $d(T) \geq \delta\|T\|$ .

By (5.8) of [Ar3], we can choose  $\varepsilon_2$  such that  $d(\mathcal{Y}_M(x, y, T)) > 0$  for all  $x, y \in H_{\varepsilon_2}$ . By the discussion of l.c. bottom of page 38 and top of page 39, there is a constant  $C_0 > 0$  such that, for  $T$  with  $d(T) \geq \delta\|T\|$  and  $\|T\| > C_0$ , for  $x, y \in H_{\varepsilon_2}$  and  $a \in A_H \setminus A_M$ , one has

$$u(y^{-1}ax, T) = \sigma_M(h_M(a), \mathcal{Y}_M(x, y, T)).$$

By Lemma 2.2, for  $X \in a_M$ , we have

$$\sigma_M(X, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) = \sigma_M(X, \mathcal{Y}_M(x_1, y_1, T))\sigma_M(X, \mathcal{Y}_M(x_2, y_2, T)).$$

Thus, one deduces that

$$\sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) = u(y_1^{-1}ax_1, T)u(y_2^{-1}ax_2, T),$$

for  $a \in A_H \setminus A_M$ . Hence, for  $d(T) \geq \delta\|T\| \geq \delta C_0$ , and  $x_i, y_i$  in  $H_{\varepsilon_2}$ , we have

$$u_M(x_1, y_1, x_2, y_2, T) = v_M(x_1, y_1, x_2, y_2, T).$$

This finishes the proof of the Lemma. □

Theorem 2.3 follows from the corollary below.

**2.9 Corollary.** *Fix  $\delta > 0$ . There exist two positive numbers  $\varepsilon$  and  $c > 0$  such that, for all  $T$  with  $d(T) \geq \delta\|T\|$ , one has*

$$\int_{\gamma \in S_\sigma} |K^T(x_m, \gamma, f) - J^T(x_m \gamma, f)| d\gamma \leq ce^{-\varepsilon\|T\|}. \quad (2.47)$$

*Proof :*

By Lemma 2.7, it is enough to prove that we can find positive numbers  $\varepsilon$ ,  $\varepsilon'$  and  $C_0$  such that

$$\int_{\gamma \in S_\sigma - S_\sigma(\varepsilon, T)} |K^T(x_m, \gamma, f) - J^T(x_m, \gamma, f)| d\gamma \leq C_0 e^{-\varepsilon'\|T\|} \quad (2.48)$$

where  $S_\sigma(\varepsilon, T)$  is defined in (2.34).

Let  $\varepsilon > 0$ . Let  $\Omega$  be a compact subset of  $G$  which contains the support of  $f_1$  and  $f_2$ . We will estimate  $|u_M(x_1, y_1, x_2, y_2, T) - v_M(x_1, y_1, x_2, y_2, T)|$  for  $x_1, x_2, y_1$  and  $y_2$  in  $H$  satisfying  $x_1^{-1}x_m\gamma x_2 \in \Omega$  and  $y_1^{-1}x_m\gamma y_2 \in \Omega$  for some  $\gamma \in S_\sigma - S_\sigma(\varepsilon, T)$  with  $x_m\gamma \in G^{\sigma-reg}$ . For this, we will use the invariance of the functions  $u_M$  and  $v_M$  by the diagonal left action of  $A_M$  on  $(x_1, x_2)$  and  $(y_1, y_2)$  respectively.



By Lemma 2.5, there are a positive integer  $k$  and a positive constant  $C_\Omega$ , (depending only on  $\Omega$ ) such that, for all  $\gamma \in S_\sigma - S_\sigma(\varepsilon, T)$  with  $x_m\gamma \in G^{\sigma-reg}$  and for all  $x_i, y_i$  in  $H$ ,  $i = 1, 2$  with  $x_1^{-1}x_m\gamma x_2$  and  $y_1^{-1}x_m\gamma y_2$  in  $\Omega$ , we have

$$\inf_{s \in S} \|(sx_m^{-1}x_1, sx_2)\| \leq C_\Omega \Delta_\sigma(x_m\gamma)^{-k} \leq C_\Omega e^{k\varepsilon\|T\|} \quad (2.49)$$

and

$$\inf_{s \in S} \|(sx_m^{-1}y_1, sy_2)\| \leq C_\Omega \Delta_\sigma(x_m\gamma)^{-k} \leq C_\Omega e^{k\varepsilon\|T\|}.$$

Since  $A_M \setminus S$  is compact, we deduce from (1.21) and (2.49) that there is a constant  $C'_\Omega > 0$  such that

$$\inf_{a \in A_M} \|(ax_m^{-1}x_1, ax_2)\| \leq C'_\Omega e^{k\varepsilon\|T\|}.$$

Thus, for  $\eta > 0$ , there is  $a_0 \in A_M$  such that

$$\|a_0x_m^{-1}x_1\| \|a_0x_2\| \leq C_\Omega e^{k\varepsilon\|T\|} + \eta. \quad (2.50)$$

Since  $A_M = A_S$ , the point  $a_0$  commutes with  $x_m$  by (1.28) and we have  $\|a_0x_1\| \leq \|x_m\| \|x_m^{-1}a_0x_1\|$ .

If  $\|T\|$  remains bounded, then  $\|a_0x_i\|, i = 1, 2$  are bounded by a constant independent of  $\|T\|$ . By the same arguments, there is  $a_1 \in A_M$  such that  $\|a_1y_i\|, i = 1, 2$  are bounded by a constant independent of  $\|T\|$ . Using the invariance of  $u_M$  and  $v_M$  by the left action of  $diag(A_M)$  on  $(x_1, x_2)$  and  $(y_1, y_2)$  respectively and the estimates (2.38), (2.41), (2.44) and (2.45) for  $u_M$  and  $v_M$ , we deduce that  $|u_M(x_1, y_1, x_2, y_2, T) - v_M(x_1, y_1, x_2, y_2, T)|$  is bounded by a constant independent of  $T$  and of  $x_i, y_i$ . Recall that by Theorem 1.2, the constant

$$C_1 := \int_{S_\sigma} \mathcal{M}(|f_1|)(x_m\gamma) \mathcal{M}(|f_2|)(x_m\gamma) d\gamma$$

is finite. We deduce that  $\int_{\gamma \in S_\sigma - S_\sigma(\varepsilon, T)} |K^T(x_m, \gamma, f) - J^T(x_m, \gamma, f)| d\gamma$  is bounded, hence we obtain (2.48).

We assume that  $\|T\|$  is not bounded. Let  $\varepsilon_1, \varepsilon_2$  and  $C$  as in Lemma 2.8. Taking  $\|T\|$  to be sufficiently large and  $\varepsilon$  such that  $k\varepsilon$  is smaller than the constant  $\varepsilon_2$ , we can assume by (2.50) that

$$\|a_0x_i\| \leq e^{\varepsilon_2\|T\|}, \quad i = 1, 2.$$

The same arguments are valid for  $\|y_i\|, i = 1, 2$ . Thus, there is  $a_1 \in A_M$  such that

$$\|a_1y_i\| \leq e^{\varepsilon_2\|T\|}, \quad i = 1, 2.$$

Using Lemma 2.8 and the invariance of  $u_M$  and  $v_M$  by the left action of the diagonal of  $A_M$  on  $(x_1, x_2)$  and  $(y_1, y_2)$  respectively, we deduce that, for all  $T$  with  $d(T) \geq \delta \|T\|$ , one has

$$|u_M(x_1, y_1, x_2, y_2, T) - v_M(x_1, y_1, x_2, y_2, T)| \leq C e^{-\varepsilon_1 \|T\|}.$$

Hence, we obtain

$$\int_{S-S_\sigma(\varepsilon, T)} |K^T(x_m, \gamma, f) - J^T(x_m, \gamma, T)| \leq C C_1 e^{-\varepsilon_1 \|T\|},$$

where  $C_1 := \int_{S_\sigma} \mathcal{M}(|f_1|)(x_m \gamma) \mathcal{M}(|f_2|)(x_m \gamma) d\gamma$ . This finishes the proof of the Corollary.  $\square$

## 2.5 The function $J^T(f)$

The goal of this section is to prove that  $J^T(f)$  is of the form

$$\sum_{k=0}^N p_k(T, f) e^{\xi_k(T)}, \quad (2.51)$$

where  $\xi_0 = 0, \xi_1, \dots, \xi_N$  are distinct points in  $ia_0^*$  and each  $p_k(T, f)$  is a polynomial function of  $T$ . Moreover, the constant term  $\tilde{J}(f) := p_0(0, f)$  is well-defined and is uniquely determined by  $K^T(f)$ . Except for one detail, our arguments and calculations are the same as those of section 6 of [Ar3]. We give the details of proof for convenience of the reader.

Recall that  $J^T(f)$  is a finite sum of the distributions

$$\begin{aligned} J^T(x_m, \gamma, f) &= |\Delta_\sigma(x_m \gamma)|_{\mathbb{F}}^{1/2} \int_{\text{diag}(A_M) \backslash H \times H} \int_{\text{diag}(A_M) \backslash H \times H} f_1(y_1^{-1} x_m \gamma y_2) \\ &\quad \times f_2(x_1^{-1} x_m \gamma x_2) v_M(x_1, y_1, x_2, y_2, T) d\overline{(x_1, x_2)} d\overline{(y_1, y_2)} \end{aligned}$$

where  $M \in \mathcal{L}(A_0)$ ,  $S$  is a maximal torus of  $M$  such that  $A_S = A_M$ ,  $x_m \in \kappa_S$  and  $v_M(x_1, y_1, x_2, y_2, T) := \int_{A_H \backslash A_M} \sigma_M(h_M(a), \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) da$  where  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  is defined in (2.11).

We first study the weight function  $v_M$  as a function of  $T$ . We fix  $M \in \mathcal{L}(A_0)$  and  $x_1, y_1, x_2$  and  $y_2$  in  $H$ .

Let  $\mathcal{L}_M := (a_{M, \mathbb{F}} + a_H)/a_H$  and  $\tilde{\mathcal{L}}_M := (\tilde{a}_{M, \mathbb{F}} + a_H)/a_H$  be the projection in  $a_M/a_H$  of the lattices  $a_{M, \mathbb{F}}$  and  $\tilde{a}_{M, \mathbb{F}}$  respectively. By (1.10), one has

$$\tilde{a}_{M, \mathbb{F}}/\tilde{a}_{H, \mathbb{F}} = \tilde{a}_{M, \mathbb{F}}/\tilde{a}_{M, \mathbb{F}} \cap a_H \simeq \tilde{\mathcal{L}}_M. \quad (2.52)$$

For  $M = A_0$ , we replace the subscript  $A_0$  by 0. We denote by  $\mathcal{L}^\vee := \text{Hom}(\mathcal{L}, 2\pi i\mathbb{Z})$  the dual lattice of a lattice  $\mathcal{L}$ .

Let  $P \in \mathcal{P}(M)$ . We introduce the following sublattice of  $\mathcal{L}_M$ . For  $k \in \mathbb{N}$ , we set

$$\mu_{\alpha,k} := k \log(q)\check{\alpha}, \alpha \in \Delta_P,$$

where  $q$  is the order of the residual field of  $F$ , and

$$\mathcal{L}_{M,k} := \sum_{\alpha \in \Delta_P} \mathbb{Z}\mu_{\alpha,k}.$$

Then  $\mathcal{L}_{M,k}$  is a lattice in  $a_M^H \simeq a_M/a_H$  independent of  $P$  and by ([Ar2] §4), one can find  $k \in \mathbb{N}^*$  such that for all  $M \in \mathcal{L}(A_0)$ , one has

$$\mathcal{L}_{M,k} \subset \widetilde{\mathcal{L}}_M.$$

The set of points  $\sum_{\alpha \in \Delta_P} y_\alpha \mu_{\alpha,k}$  with  $y_\alpha \in ]-1, 0]$  is a fundamental domain of  $\mathcal{L}_{M,k}$  which we denote by  $\mathcal{D}_{M,k}$ .

For  $X \in \mathcal{L}_M/\mathcal{L}_{M,k}$  and  $Y \in a_M/a_H$ , we denote by  $\bar{X}_P(Y)$  the representative of  $X$  in  $\mathcal{L}_M$  such that  $\bar{X}_P(Y) - Y \in \mathcal{D}_{M,k}$ . (2.53)

For  $\lambda \in a_{M,\mathbb{C}}^*$ , we set

$$\theta_{P,k}(\lambda) = \text{vol}(a_M^H/\mathcal{L}_{M,k})^{-1} \prod_{\alpha \in \Delta_P} (1 - e^{-\lambda(\mu_{\alpha,k})}). \quad (2.54)$$

We fix  $T \in a_{0,F}$ . By definition of  $\sigma_M$  (cf. (2.4)), the function  $v_M$  depends only on the image of  $T_P$  in  $\mathcal{L}_M$ . Hence we can assume that  $T$  lies in the lattice  $\mathcal{L}_0$ . For  $P \in \mathcal{P}(M)$ , the map  $T \mapsto T_P$  sends surjectively  $\mathcal{L}_0$  onto the intersection of  $\mathcal{L}_M$  with the closure  $\overline{a_P^+}$  of the chamber associated to  $P$ . Thus, we may restrict  $T$  to lie in the intersection of  $\mathcal{L}_0$  with suitable regular points in some positive chamber  $a_0^+$  of  $a_H \setminus a_0$ . Then the points  $T_P$  range over a suitable regular points in  $\mathcal{L}_M \cap a_P^+$ .

We recall that  $\mathcal{Y}_M(x_1, y_1, x_2, y_2, T)$  is the set of points  $Z_P := Z_P(x_1, y_1, x_2, y_2, T)$  defined in (2.10). Thus, we can write

$$Z_P = T_P + Z_P^0 \text{ with } Z_P^0 := \text{inf}^P(h_P(y_1) - h_{\bar{P}}(x_1), h_P(y_2) - h_{\bar{P}}(x_2)). \quad (2.55)$$

Notice that the points  $Z_P^0$  do not necessarily belong to the lattice  $\mathcal{L}_M$ . It is the only difference with [Ar3] section 6 in what follows.

**2.10 Lemma.** *There is a positive integer  $N$  independent of  $M$  and polynomial functions  $q_\xi(T)$  for  $\xi \in (\frac{1}{N}\mathcal{L}_0^\vee)/\mathcal{L}_0^\vee$  (depending on  $x_1, y_1, x_2$  and  $y_2$ ), such that*

$$v_M(x_1, y_1, x_2, y_2, T) = \sum_{\xi \in (\frac{1}{N}\mathcal{L}_0^\vee)/\mathcal{L}_0^\vee} q_\xi(T) e^{\xi(T)}.$$

Moreover, the constant term  $\tilde{v}_M(x_1, y_1, x_2, y_2) := q_0(0)$  of  $v_M(x_1, y_1, x_2, y_2, T)$  is given by

$$\tilde{v}_M(x_1, y_1, x_2, y_2) = \lim_{\Lambda \rightarrow 0} \left( \sum_{P \in \mathcal{P}(M)} |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} e^{\langle \Lambda, \bar{X}_P(Z_P^0) \rangle} \theta_{P,k}(\Lambda)^{-1} \right).$$

*Proof :*

The kernel of the surjective map  $h_M : A_H \backslash A_M \rightarrow \tilde{a}_{M,F}/\tilde{a}_{H,F}$  is a compact group which has volume 1 by our convention of choice of measure. Thus, using (2.52), we can write

$$v_M(x_1, y_1, x_2, y_2, T) := \sum_{X \in \tilde{\mathcal{L}}_M} \sigma_M(X, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)).$$

For our study, it is convenient to take a sum over  $\mathcal{L}_M$ . The finite quotient  $\tilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee$  can be identified with the character group of  $\mathcal{L}_M/\tilde{\mathcal{L}}_M$  under the pairing

$$(\nu, X) \in \tilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee \times \mathcal{L}_M/\tilde{\mathcal{L}}_M \mapsto e^{\nu(X)}.$$

Hence, by inversion formula on finite abelian groups, we obtain

$$v_M(x_1, y_1, x_2, y_2, T) = |\mathcal{L}_M/\tilde{\mathcal{L}}_M|^{-1} \sum_{\nu \in \tilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee} \sum_{X \in \mathcal{L}_M} \sigma_M(X, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) e^{\nu(X)}.$$

Coming back to the definition of  $\sigma_M$  (cf. (2.4)), we fix a small point  $\Lambda \in (a_M/a_H)_\mathbb{C}^*$  whose real part  $\Lambda_R$  is in general position. One has

$$\begin{aligned} \sigma_M(X, \mathcal{Y}_M(x_1, y_1, x_2, y_2, T)) &= \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\Delta|} \varphi_P^\Lambda(X - Z_P) \\ &= \lim_{\Lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\Delta|} \varphi_P^\Lambda(X - Z_P) e^{\Lambda(X)}. \end{aligned}$$

By definition of  $\varphi_P^\Lambda$ , the function  $X \mapsto e^{\Lambda(X)}$  is rapidly decreasing on the support of  $X \mapsto \varphi_P^\Lambda(X - Z_P)$ . Hence, the product of these two functions is summable over  $X \in \mathcal{L}_M$ . Therefore, we can write

$$v_M(x_1, y_1, x_2, y_2, T) = \sum_{\nu \in \tilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee} \lim_{\Lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} F_P^T(\Lambda) \quad (2.56)$$

where

$$F_P^T(\Lambda) := |\mathcal{L}_M/\tilde{\mathcal{L}}_M|^{-1} \sum_{X \in \mathcal{L}_M} (-1)^{|\Delta_P^\Delta|} \varphi_P^\Lambda(X - Z_P) e^{(\Lambda+\nu)(X)}.$$

The above discussion implies that

$$\text{the map } \Lambda \mapsto \sum_{P \in \mathcal{P}(M)} F_P^T(\Lambda) \text{ is analytic at } \Lambda = 0. \quad (2.57)$$

We fix  $P \in \mathcal{P}(M)$ . We want to express  $F_P^T(\Lambda)$  in terms of a product of geometric series. For this, we write

$$F_P^T(\Lambda) := |\mathcal{L}_M / \widetilde{\mathcal{L}}_M|^{-1} \sum_{X \in \mathcal{L}_M / \mathcal{L}_{M,k}} \sum_{X' \in \mathcal{L}_{M,k}} (-1)^{|\Delta_P^\Lambda|} \varphi_P^\Lambda(X + X' - Z_P) e^{(\Lambda + \nu)(X + X')}. \quad (2.58)$$

Let  $X \in \mathcal{L}_M / \mathcal{L}_{M,k}$ . Recall that  $\bar{X}_P(Y)$  is the representative of  $X$  in  $\mathcal{L}_M$  such that  $\bar{X}_P(Y) - Y \in \mathcal{D}_{M,k}$ . We set

$$\bar{X}_P^\Lambda(Y) := \bar{X}_P(Y) + \sum_{\alpha \in \Delta_P^\Lambda} \mu_{\alpha,k}.$$

Thus  $\bar{X}_P^\Lambda(Y)$  is also a representative of  $X$  in  $\mathcal{L}_M$ . Taking  $Y := Z_P$ , we can set

$$\varphi_P^\Lambda(X + X' - Z_P) = \varphi_P^\Lambda(\bar{X}_P^\Lambda(Z_P) + X' - Z_P)$$

in (2.58). The set of points  $X' \in \mathcal{L}_{M,k}$  such that this characteristic function equals to 1 is exactly the set

$$\left\{ \sum_{\alpha \in \Delta_P^\Lambda} n_\alpha \mu_{\alpha,k} - \sum_{\alpha \in \Delta_P - \Delta_P^\Lambda} n_\alpha \mu_{\alpha,k}; n_\alpha \in \mathbb{N} \right\}.$$

Therefore, a simple calculation as in [Ar3] top of page 45 gives

$$\begin{aligned} & (-1)^{|\Delta_P^\Lambda|} \sum_{X' \in \mathcal{L}_{M,k}} \varphi_P^\Lambda(X + X' - Z_P) e^{(\Lambda + \nu)(X + X')} \\ &= e^{(\Lambda + \nu)(\bar{X}_P(Z_P))} \left( \prod_{\alpha \in \Delta_P} (1 - e^{-(\Lambda + \nu)(\mu_{\alpha,k})})^{-1} \right). \end{aligned} \quad (2.59)$$

We have fixed the Haar measure on  $a_M^H \simeq a_M / a_G$  with the property that the quotient of  $a_M / a_H$  by the lattice  $\widetilde{\mathcal{L}}_M$  has volume 1. Thus, we have

$$|\mathcal{L}_M / \widetilde{\mathcal{L}}_M|^{-1} \prod_{\alpha \in \Delta_P} (1 - e^{-(\Lambda + \nu)(\mu_{\alpha,k})})^{-1} = |\mathcal{L}_M / \widetilde{\mathcal{L}}_{M,k}|^{-1} \theta_{P,k}(\Lambda + \nu)^{-1}.$$

By the above equality, (2.58) and (2.59), we obtain

$$F_P^T(\Lambda) = |\mathcal{L}_M / \mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M / \mathcal{L}_{M,k}} e^{\langle \Lambda + \nu, \bar{X}_P(Z_P) \rangle} \theta_{P,k}(\Lambda + \nu)^{-1}. \quad (2.60)$$

Let  $X \in \mathcal{L}_M/\mathcal{L}_{M,k}$ . We recall that  $T_P$  belongs to  $\mathcal{L}_M$  for  $P \in \mathcal{P}(M)$  and  $Z_P = T_P + Z_P^0$  (cf. (2.55)). By definition (cf. (2.53)), the point  $\bar{X}_P(Z_P)$  is the unique representative of  $X$  in  $\mathcal{L}_M$  such that  $\bar{X}_P(Z_P) - T_P - Z_P^0 \in \mathcal{D}_{M,k}$  and  $(X - T_P)_P(Z_P^0)$  is the unique representative of  $X - T_P$  in  $\mathcal{L}_M$  such that  $(X - T_P)_P(Z_P^0) - Z_P^0 \in \mathcal{D}_{M,k}$ . Hence, we deduce that

$$\bar{X}_P(Z_P) = \overline{(X - T_P)_P(Z_P^0)} + T_P. \quad (2.61)$$

Replacing  $X$  by  $X - T_P$  in (2.60), we obtain

$$F_P(\Lambda)^T = |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} e^{\langle \Lambda + \nu, T_P + \bar{X}_P(Z_P^0) \rangle} \theta_{P,k}(\Lambda + \nu)^{-1} \quad (2.62)$$

where  $\bar{X}_P(Z_P^0)$  is independent of  $T$ . Thus by (2.56), we have established that  $v_M(x_1, y_1, x_2, y_2, T)$  is equal to

$$\sum_{\nu \in \widetilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee} \lim_{\Lambda \rightarrow 0} \left( \sum_{P \in \mathcal{P}(M)} |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} e^{\langle \Lambda + \nu, T_P + \bar{X}_P(Z_P^0) \rangle} \theta_{P,k}(\Lambda + \nu)^{-1} \right). \quad (2.63)$$

Recall that the expression in the brackets is analytic at  $\Lambda = 0$  (cf. (2.57)). To analyze this expression as function of  $T$ , we argue as in ([W1] p.315). We give the details for convenience of lecture. We replace  $\Lambda$  by  $z\Lambda$ . The map  $z \mapsto \theta_{P,k}(z\Lambda + \nu)^{-1}$  may have a pole at  $z = 0$ . Let  $r$  denotes the biggest order of this pole when  $P$  runs  $\mathcal{P}(M)$ . Then, using Taylor expansions, one deduces that

$$\begin{aligned} & \lim_{\Lambda \rightarrow 0} \left( \sum_{P \in \mathcal{P}(M)} |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} e^{\langle \Lambda + \nu, T_P + \bar{X}_P(Z_P^0) \rangle} \theta_{P,k}(\Lambda + \nu)^{-1} \right) = \\ & \sum_{m=0}^r \sum_{P \in \mathcal{P}(M)} C_m \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} \frac{\partial^m}{\partial z^m} (e^{\langle z\Lambda + \nu, T_P + \bar{X}_P(Z_P^0) \rangle})_{[z=0]} \frac{\partial^{r-m}}{\partial z^{r-m}} (z^r \theta_{P,k}(z\Lambda + \nu)^{-1})_{[z=0]}, \end{aligned}$$

where  $C_m = \frac{1}{m!(r-m)!} |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1}$ .

But we have

$$\frac{\partial^m}{\partial z^m} (e^{\langle z\Lambda + \nu, T_P + \bar{X}_P(Z_P^0) \rangle})_{[z=0]} = (\langle \Lambda, T_P + \bar{X}_P(Z_P^0) \rangle)^m e^{\langle \nu, T_P + \bar{X}_P(Z_P^0) \rangle},$$

and  $\frac{\partial^{r-m}}{\partial z^{r-m}} (z^r \theta_{P,k}(z\Lambda + \nu)^{-1})_{[z=0]}$  is independent of  $T_P$ .

Therefore, we deduce that  $v_M(x_1, y_1, x_2, y_2, T)$  is a finite sum of functions

$$q_{P,\nu}(T_P) e^{\nu(T_P)}, \quad \nu \in \widetilde{\mathcal{L}}_M^\vee/\mathcal{L}_M^\vee, P \in \mathcal{P}(M),$$

where  $q_{P,\nu}$  is a polynomial function on  $a_M$ .

Since  $\mathcal{L}_0^\vee \subset \widetilde{\mathcal{L}}_0^\vee$  are lattices of same rank, one can find a positive integer  $N$  such that  $N\widetilde{\mathcal{L}}_0^\vee \subset \mathcal{L}_0^\vee$ . Therefore, by our choice of  $T$  and the above expression, we can write

$$v_M(x_1, y_1, x_2, y_2, T) = \sum_{\xi \in (\frac{1}{N}\mathcal{L}_0^\vee)/\mathcal{L}_0^\vee} q_\xi(T) e^{\xi(T)},$$

where  $q_\xi(T)$  is a polynomial function of  $T$ . This gives the first part of the Lemma.

Since the polynomials  $q_\xi(T)$  are obviously uniquely determined, the constant term  $\tilde{v}_M(x_1, y_1, x_2, y_2) := q_0(0)$  is well defined. To calculate it, we take the summand corresponding to  $\nu = 0$  in (2.63) and then set  $T = 0$ . We obtain

$$\tilde{v}_M(x_1, y_1, x_2, y_2) = \lim_{\Lambda \rightarrow 0} \left( \sum_{P \in \mathcal{P}(M)} |\mathcal{L}_M/\mathcal{L}_{M,k}|^{-1} \sum_{X \in \mathcal{L}_M/\mathcal{L}_{M,k}} e^{\langle \Lambda, \bar{X}_P(Z_P^0) \rangle} \theta_{P,k}(\Lambda)^{-1} \right).$$

This finishes the proof of the Lemma.  $\square$

We substitute the expression we have obtained for  $v_M$  in Lemma 2.10 into the expression for  $J^T(x_m, \gamma, f)$ . Hence, we obtain the following similar decomposition for  $J^T(f)$ .

**2.11 Corollary.** *There is a decomposition*

$$J^T(f) = \sum_{\xi \in (\frac{1}{N}\mathcal{L}_0^\vee)/\mathcal{L}_0^\vee} p_\xi(T, f) e^{\xi(T)}, \quad T \in \mathcal{L}_0 \cap \mathfrak{a}_0^+,$$

where  $N$  is positive integer and each  $p_\xi(T, f)$  is a polynomial function of  $T$ . Moreover, the constant term  $\tilde{J}(f) := p_0(0, f)$  of  $J^T(f)$  is given by

$$\tilde{J}(f) := \sum_{M \in \mathcal{L}(A_0)} c_M \sum_{S \in \mathcal{T}_M} \sum_{x_m \in \kappa_S} c_{S, x_m} \int_{S_\sigma} \tilde{J}(x_m, \gamma, f) d\gamma,$$

where

$$\begin{aligned} \tilde{J}(x_m, \gamma, f) &= |\Delta_\sigma(x_m \gamma)|^{1/2} \int_{\text{diag}(A_M) \setminus H \times H} \int_{\text{diag}(A_M) \setminus H \times H} f_1(y_1^{-1} x_m \gamma y_2) \\ &\quad \times f_2(x_1^{-1} x_m \gamma x_2) \tilde{v}_M(x_1, y_1, x_2, y_2) \overline{d(x_1, x_2)} \overline{d(y_1, y_2)}. \end{aligned}$$

## A Appendix. Spherical character of a supercuspidal representation as weighted orbital integral

Let  $(\pi, V)$  be a unitary irreducible admissible representation of  $G$ . We say that  $\pi$  is  $H$ -distinguished if the space  $V^{*H} = \text{Hom}_H(\pi, \mathbb{C})$  of  $H$ -invariant linear forms on  $V$  is nonzero. In that case, a distribution  $m_{\xi, \xi'}$ , called spherical character, can be

associated to two  $H$ -invariant linear forms  $\xi, \xi'$  on  $V$  (cf. definition below). By ([Ha] Theorem 1), spherical characters are locally integrable functions on  $G$ , which are smooth on the set of  $\sigma$ -regular points of  $G$ .

From now, we assume that  $A_H = \{1\}$ . We fix a  $H$ -distinguished supercuspidal representation  $(\tau, V)$  of  $G$ . We denote by  $d(\tau)$  its formal degree.

The aim of this appendix is to deduce from our main results, the value  $m_{\xi, \xi'}(g)$ , when  $g \in G$  is  $\sigma$ -regular and  $\xi, \xi' \in V^{*H}$ , in terms of weighted orbital integrals of a matrix coefficient of  $\tau$  (cf. Theorem A.2). This result is analogous to that of J. Arthur in the group case ([Ar2]). Notice that this result of J. Arthur can be deduced from his local trace formula ([Ar3]) which was obtained later.

Let  $(\cdot, \cdot)$  be a  $G$ -invariant hermitian inner product on  $V$ . Since  $\tau$  is unitary, it induces an isomorphism  $\iota : v \mapsto (\cdot, v)$  from the conjugate complex vector space  $\bar{V}$  of  $V$  and the smooth dual  $\check{V}$  of  $V$ , which intertwines the complex conjugate of  $\tau$  and its contragredient  $\check{\tau}$ . If  $\xi$  is a linear form on  $V$ , we define the linear form  $\bar{\xi}$  on  $\bar{V}$  by  $\bar{\xi}(u) := \overline{\xi(u)}$ .

For  $\xi_1$  and  $\xi_2$  two nonzero  $H$ -invariant linear forms on  $V$ , we associate the spherical character  $m_{\xi_1, \xi_2}$  defined to be the distribution on  $G$  given by

$$m_{\xi_1, \xi_2}(f) := \sum_{u \in \mathcal{B}} \xi_1(\tau(f)u) \overline{\xi_2(u)},$$

where  $\mathcal{B}$  is an orthonormal basis of  $V$ . Since  $\tau(f)$  is of finite rank, this sum is finite. Moreover, this sum does not depend on the choice of  $\mathcal{B}$ . Indeed, let  $(\tau^*, V^*)$  be the dual representation of  $\tau$ . For  $f \in C_c^\infty(G)$ , we set  $\check{f}(g) := f(g^{-1})$ . By ([R] Théorème III.3.4 and I.1.2), the linear form  $\tau^*(\check{f})\xi$  belongs to  $\check{V}$ . Hence we can write  $\iota^{-1}(\tau^*(\check{f})\xi) = \sum_{v \in \mathcal{B}} (\tau^*(\check{f})\xi)(v) \cdot v$  where  $(\lambda, v) \mapsto \lambda \cdot v$  is the action of  $\mathbb{C}$  on  $\bar{V}$ . Therefore we deduce easily that one has

$$m_{\xi_1, \xi_2}(f) = \bar{\xi}_2(\iota^{-1}(\tau^*(\check{f})\xi_1)). \quad (1.1)$$

Since  $\tau$  is a supercuspidal representation, we can define the  $H \times H$ -invariant pairing  $\mathcal{L}$  on  $V \times \bar{V}$  by

$$\mathcal{L}(u, v) := \int_H (\tau(h)u, v) dh.$$

By ([Z] Theorem 1.5),

$$\text{the map } v \mapsto \xi_v : u \mapsto \mathcal{L}(u, v) \text{ is a surjective linear map from } \bar{V} \text{ onto } V^{*H}. \quad (1.2)$$

For  $v, w \in V$ , we denote by  $c_{v, w}$  the corresponding matrix coefficient defined by  $c_{v, w}(g) := (\tau(g)v, w)$  for  $g \in G$ .



**A.1 Lemma.** *Let  $\xi_1, \xi_2 \in V^{*H}$  and  $v, w \in V$ . Then we have*

$$m_{\xi_1, \xi_2}(\check{c}_{v,w}) = d(\tau)^{-1} \xi_1(v) \overline{\xi_2(w)}.$$

*Proof :*

By (1.2), there exist  $v_1$  and  $v_2$  in  $V$  such that  $\xi_j = \xi_{v_j}$  for  $j = 1, 2$ . By definition of the spherical character, for  $f \in C_c^\infty(G)$  and  $\mathcal{B}$  an orthonormal basis of  $V$ , one has

$$\begin{aligned} m_{\xi_1, \xi_2}(f) &= \sum_{u \in \mathcal{B}} \int_H (\tau(h) \tau(f) u, v_1) dh \int_H \overline{(\tau(h) u, v_2)} dh \\ &= \sum_{u \in \mathcal{B}} \int_{H \times H} (u, \tau(\check{f}) \tau(h_1) v_1) (\tau(h_2) v_2, u) dh_1 dh_2 \\ &= \int_{H \times H} (\tau(h_2) v_2, \tau(\check{f}) \tau(h_1) v_1) dh_1 dh_2 \end{aligned}$$

Hence we obtain

$$m_{\xi_1, \xi_2}(f) = \int_{H \times H} \int_G f(g) (\tau(h_1 g h_2) v_2, v_1) dg dh_1 dh_2. \quad (1.3)$$

Let  $f(g) := \check{c}_{v,w}(g) = \overline{(\tau(g) w, v)}$ . By the orthogonality relation of Schur, for  $h_1, h_2 \in H$ , one has

$$\int_G (\tau(g) \tau(h_2) v_2, \tau(h_1) v_1) \overline{(\tau(g) w, v)} dg = d(\tau)^{-1} (\tau(h_2) v_2, w) (v, \tau(h_1) v_1).$$

Thus, we deduce that

$$m_{\xi_1, \xi_2}(f) = d(\tau)^{-1} \xi_w(v_2) \xi_{v_1}(v) = d(\tau)^{-1} \xi_1(v) \overline{\xi_2(w)}. \quad \square$$

For  $M \in \mathcal{L}(A_0)$ , we define the weight function  $w_M$  on  $H \times H$  by

$$w_M(y_1, y_2) := \tilde{v}_M(1, y_1, 1, y_2),$$

where  $\tilde{v}_M$  is defined in Lemma 2.10 and 1 is the neutral element of  $H$ . For  $f \in C_c^\infty(G)$ , we define the weighted orbital integral of  $f$  by

$$\mathcal{WM}(f)(g) := |\Delta_\sigma(g)|^{1/2} \int_{H \times H} f(y_1 g y_2) w_M(y_1, y_2) dy_1 dy_2, \quad g \in G^{\sigma\text{-reg}} \cap \tilde{M}.$$

**A.2 Theorem.** *Let  $M \in \mathcal{L}(A_0)$  and  $S \in \mathcal{T}_M$ . Let  $x_M \in \kappa_S$  and  $\gamma \in S_\sigma$  be such that  $x_M \gamma$  is  $\sigma$ -regular. Then, for  $v, w \in V$ , we have*

$$c_M c_{S, x_M} \mathcal{WM}(c_{v,w})(x_M \gamma) = m_{\xi_w, \xi_v}(x_M \gamma).$$

*Proof :*

Let  $f_1$  be a matrix coefficient of  $\tau$  and  $f_2 \in C_c^\infty(G)$ . We set  $f := f_1 \otimes f_2$ . For  $x \in G$ , we define

$$h(g) := \int_G f_1(xu) f_2(ugx) du,$$

so that

$$K_f(x, y) = [\rho(yx^{-1})h](e).$$

If  $\pi$  is a unitary irreducible admissible representation of  $G$ , one has

$$\begin{aligned} \pi(\rho(yx^{-1})h) &= \int_{G \times G} f_1(xu) f_2(ugy) \pi(g) du dg \\ &= \int_{G \times G} f_1(xu) f_2(u_2) \pi(u^{-1}u_2y^{-1}) du du_2 = \int_{G \times G} f_1(u_1^{-1}) f_2(u_2) \pi(u_1xu_2y^{-1}) du_1 du_2 \\ &= \pi(\check{f}_1) \pi(x) \pi(f_2) \pi(y^{-1}). \end{aligned}$$

Since  $\tau$  is supercuspidal and  $f_1$  is a matrix coefficient of  $\tau$ , we deduce that  $\pi(\rho(yx^{-1})h)$  is equal to 0 if  $\pi$  is not equivalent to  $\tau$ . Therefore, applying the Plancherel formula ([W2] Théorème VIII.1.1.) to  $[\rho(yx^{-1})h]$ , we obtain

$$K_f(x, y) = d(\tau) \text{tr}(\tau(\check{f}_1) \tau(x) \tau(f_2) \tau(y^{-1})).$$

We identify  $\check{V} \otimes V$  with a subspace of Hilbert-Schmidt operators on  $V$ . Taking an orthonormal basis  $\mathcal{B}_{HS}(V)$  of  $\check{V} \otimes V$  for the scalar product  $(S, S') := \text{tr}(SS'^*)$ , one obtains

$$\begin{aligned} K_f(x, y) &= d(\tau) \text{tr}(\tau(\check{f}_1) \tau(x) \tau(f_2) \tau(y)^*) = d(\tau) (\tau(\check{f}_1) \tau(x) \tau(f_2), \tau(y)) \\ &= d(\tau) \sum_{S \in \mathcal{B}_{HS}(V)} (\tau(\check{f}_1) \tau(x) \tau(f_2), S^*) \overline{(\tau(y), S^*)} \\ &= d(\tau) \sum_{S \in \mathcal{B}_{HS}(V)} \text{tr}(\tau(x) \tau(f_2) S \tau(\check{f}_1)) \overline{\text{tr}(\tau(y) S)}, \end{aligned}$$

where the sums over  $S$  are finite since  $\tau(f_2)$  and  $\tau(\check{f}_1)$  are of finite rank. Therefore, the truncated kernel is equal to

$$K^T(f) = d(\tau) \sum_{S \in \mathcal{B}_{HS}(V)} P_\tau^T(\check{\tau} \otimes \tau(f) S) \overline{P_\tau^T(S)}$$

where

$$P_\tau^T(S) = \int_H \text{tr}(\tau(h) S) u(h, T) dh, \quad S \in \check{V} \otimes V.$$

For  $\check{v} \otimes v \in \check{V} \otimes V$ , one has  $\text{tr}(\tau(h)(\check{v} \otimes v)) = c_{\check{v},v}(h)$ . Since  $c_{\check{v},v}$  is compactly supported, the truncated local period  $P_\tau^T(S)$  converges when  $\|T\|$  approaches infinity to

$$P_\tau(S) = \int_H \text{tr}(\tau(h)S) dh.$$

Therefore, we obtain

$$\lim_{\|T\| \rightarrow +\infty} K^T(f) = d(\tau)m_{P_\tau, P_\tau}(f), \quad (1.4)$$

where  $m_{P_\tau, P_\tau}$  is the spherical character of the representation  $\check{\tau} \otimes \tau$  associated to the  $H \times H$ -invariant linear form  $P_\tau$  on  $\check{V} \otimes V$ .

Recall that  $\tilde{J}(f)$  is the constant term of  $J^T(f)$ . We deduce from Theorem 2.15 that

$$d(\tau)m_{P_\tau, P_\tau}(f) = \tilde{J}(f). \quad (1.5)$$

We now express  $m_{P_\tau, P_\tau}$  in terms of  $H$ -invariant linear forms on  $V$ . Let  $V_H$  be the orthogonal of  $V^{*H}$  in  $V$ . Since  $\xi_u(v) = \overline{\xi_v(u)}$  for  $u, v \in V$ , the space  $\bar{V}_H$  is the kernel of  $v \mapsto \xi_v$ . Let  $W$  be a complementary subspace of  $V_H$  in  $V$ . Then, the map  $v \mapsto \xi_v$  is an isomorphism from  $\bar{W}$  to  $V^{*H}$  and  $(u, v) \mapsto \xi_v(u)$  is a nondegenerate hermitian form on  $W$ . Let  $(e_1, \dots, e_n)$  be an orthogonal basis of  $W$  for this hermitian form. We set  $\xi_i := \xi_{e_i}$  for  $i = 1, \dots, n$ . Thus we have  $\xi_i(e_i) \neq 0$ .

We identify  $\bar{V}$  and  $\check{V}$  by the isomorphism  $\iota$ . We claim that

$$P_\tau = \sum_{i=1}^n \frac{1}{\xi_i(e_i)} \bar{\xi}_i \otimes \xi_i \quad (1.6)$$

Indeed, we have  $P_\tau(v \otimes u) = \xi_v(u) = \overline{\xi_v(u)}$ . Hence, the two sides are equal to 0 on  $\bar{V} \otimes V_H + \bar{V}_H \otimes V + \bar{V}_H \otimes V_H$  and take the same value  $\xi_k(e_l)$  on  $e_k \otimes e_l$  for  $k, l \in \{1, \dots, n\}$ . Hence, by definition of spherical characters, we deduce that

$$\begin{aligned} m_{P_\tau, P_\tau}(f_1 \otimes f_2) &= \sum_{u \otimes v \in o.b.(\bar{V} \otimes V)} P_\tau(\bar{\tau}(f_1) \otimes \tau(f_2)(u \otimes v)) \overline{P_\tau(u \otimes v)} \\ &= \sum_{u \otimes v \in o.b.(\bar{V} \otimes V)} \sum_{i,j=1}^n \frac{1}{\xi_i(e_i)\xi_j(e_j)} \bar{\xi}_i(\bar{\tau}(f_1)u) \xi_i(\tau(f_2)v) \overline{\xi_j(u)\xi_j(v)}, \end{aligned}$$

where  $o.b.(\bar{V} \otimes V)$  is an orthonormal basis of  $\bar{V} \otimes V$ . By definition of  $\bar{\xi}$  for  $\xi \in V^{*H}$ , one has  $\bar{\xi}(\bar{\tau}(f_1)u) = \overline{\xi(\tau(\bar{f}_1))}$ . Therefore, we obtain

$$m_{P_\tau, P_\tau}(f_1 \otimes f_2) = \sum_{i,j=1}^n \frac{1}{\xi_i(e_i)\xi_j(e_j)} \overline{m_{\xi_i, \xi_j}(\bar{f}_1)} m_{\xi_i, \xi_j}(f_2). \quad (1.7)$$

Let  $v$  and  $w$  in  $V$ . Let  $f_1 := c_{v,w}$  so that  $\bar{f}_1 = \check{c}_{v,w}$ . If  $v \in V_H$  or  $w \in V_H$ , it follows from Lemma A.1 that  $m_{\xi_i, \xi_j}(\bar{f}_1) = 0$  for  $i, j \in \{1, \dots, n\}$ , hence  $m_{P_\tau, P_\tau}(f_1 \otimes f_2) = 0$ . Thus, we deduce from (1.5) that

$$\tilde{J}(c_{v,w} \otimes f_2) = 0, \quad v \in V_H \text{ or } w \in V_H. \quad (1.8)$$

Let  $k, l \in \{1, \dots, n\}$ . We set  $f_1 := c_{e_k, e_l}$ , hence  $\bar{f}_1 = \check{c}_{e_l, e_k}$ . By Lemma A.1, one has  $m_{\xi_i, \xi_j}(\bar{f}_1) = d(\tau)^{-1} \xi_i(e_l) \xi_j(e_k)$ . Therefore, by (1.5) and (1.7) we obtain

$$\tilde{J}(c_{e_k, e_l} \otimes f_2) = m_{\xi_l, \xi_k}(f_2). \quad (1.9)$$

By sesquilinearity, one deduces from (1.8) and (1.9) that one has

$$\tilde{J}(c_{v,w} \otimes f_2) = m_{\xi_w, \xi_v}(f_2) \quad v, w \in V. \quad (1.10)$$

Let  $(J_n)_n$  be a sequence of compact open subgroups whose intersection is equal to the neutral element of  $G$ . The characteristic function  $g_n$  of  $J_n x_m \gamma J_n$  approaches the Dirac measure at  $x_m \gamma$  as  $n$  approaches  $+\infty$ . Thus, if  $v, w \in V$  then  $m_{\xi_w, \xi_v}(g_n)$  converges to  $m_{\xi_w, \xi_v}(x_m \gamma)$ . By Corollary 2.11, the constant term  $\tilde{J}(c_{v,w} \otimes g_n)$  converges to  $c_{MC_S, x_m} \mathcal{WM}(c_{v,w})(x_m \gamma)$ . We deduce the Theorem from (1.10).  $\square$

## References

- [Ar1] J. Arthur, *The characters of discrete series as orbital integrals*, Invent. Math., 32 (1976), 205-261.
- [Ar2] J. Arthur, *The characters of supercuspidal representations as weighted orbital integrals*, Proc. Indian Acad. Sci. Math. Sci., 97 (1987), 3-19.
- [Ar3] J. Arthur, *A Local Trace formula*, Publ. Math. Inst. Hautes Études Sci., 73 (1991), 5 - 96.
- [Bo] A. Borel, *Linear Algebraic Groups, 2d edition*, Grad. Texts in Math., Springer, vol. 126 (1991).
- [F] B. Feigon, *A Relative Trace Formula for  $PGL(2)$  in the Local Setting*, Pacific J. Math. (Rogawski Memorial Volume), 260 (2012), no. 2, 395–432.
- [Ha] J. Hakim, *Admissible distributions on p-adic symmetric spaces*, J. Reine Angew. Math. 455 (1994), 119.
- [HC1] Harish-Chandra, *A formula for semisimple Lie groups*, Amer. J. Math. 79 (1957), 733-760.
- [HC2] Harish-Chandra, *Two Theorems on semisimple Lie groups*, Ann. of Math., 83 (1966), 74-128.

- [HC3] Harish-Chandra (notes by G. van Dijk), *Harmonic Analysis on reductive  $p$ -adic groups*, Lecture Notes in Math., vol. 162, Springer (1970).
- [HC4] Harish-Chandra (notes by S. DeBacker and P. J. Sally, Jr.), *Admissible Invariant Distributions on Reductive  $p$ -adic Groups*, University Lecture Series, vol. 162, Amer. Math. Soc., Vol. 16 (1999).
- [J] H. Jacquet, *Automorphic spectrum of symmetric spaces*, in *Representation Theory and Automorphic Forms*, (Eds T. N. Bailey and A. W. Knap) Proc. Sympos. Pure Math., 61 (1997).
- [JLR] H. Jacquet, E. Lapid, and J. Rogawski, *Periods of automorphic forms*, J. Amer. Math. Soc. 12 (1999), no. 1, 173240.
- [KT] S. Kato and K. Takano, Subrepresentation theorem for  $p$ -adic symmetric spaces. Int. Math. Res. Not. 11 (2008), Art. ID rnn 028, 40pp.
- [Ko ] R. E. Kottwitz, *Harmonic Analysis on Reductive  $p$ -adic Groups and Lie Algebras*, Harmonic Analysis, the Trace Formula and Shimura Varieties, Amer. Math. Society, Cambridge: Clay Mathematics Institute, 2005.
- [O] O. Offen, *Unitary Periods and Jacquet's relative trace formula*, Automorphic forms and L-functions, Proceedings of a workshop in honor of Steve Gelbart on the occasion of his 60th birthday, Contemporary Mathematics 488, AMS and BIU (2009) 148183.
- [RR] C. Rader, S. Rallis, *Spherical characters on  $p$ -adic symmetric spaces*, Amer. J. Math., Vol 118, N° 1 (5 Feb. 1996), 91-178.
- [R] D. Renard, *Représentations des groupes réductifs  $p$ -adiques*, Cours spécialisés, volume 17, SMF.
- [Se] J. P. Serre, *Cohomologie Galoisienne*, Lecture Notes in Math., vol 5, Springer-Verlag, New-York, (1973).
- [W1] J.-L. Waldspurger, *Intégrales orbitales sphériques pour  $GL(N)$  sur un corps  $p$ -adique*, Astérisque, 171-172 (1989) 279 - 337.
- [W2] J.-L. Waldspurger, *La formule de Plancherel pour les groupes  $p$ -adiques (d'après Harish-Chandra)*, J. Inst. Math. Jussieu **2** (2003) 235 - 333.
- [Z] C. Zhang, *Local periods for discrete series representations*, Preprint, arXiv:1509.06166.

P. Delorme, Aix-Marseille Université, CNRS, Centrale Marseille, I2M, UMR 7373, 13453 Marseille, France.

*E-mail address:* patrick.delorme@univ-amu.fr

P. Harinck, Ecole polytechnique, CMLS - CNRS, UMR 7640, Route de Saclay, 91128 Palaiseau Cédex, France.

*E-mail address:* pascale.harinck@polytechnique.edu

S. Souaifi, Université de Strasbourg, IRMA CNRS, UMR 7501, 7 rue Descartes, 67084 Strasbourg Cédex, France.

*E-mail address:* sofiane.souaifi@math.unistra.fr