



HAL
open science

Vanishing viscosity limit of navier-stokes equations in gevrey class

Feng Cheng, Wei-Xi Li, Chao-Jiang Xu

► **To cite this version:**

Feng Cheng, Wei-Xi Li, Chao-Jiang Xu. Vanishing viscosity limit of navier-stokes equations in gevrey class. 2017. hal-01473694

HAL Id: hal-01473694

<https://hal.science/hal-01473694>

Preprint submitted on 22 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

VANISHING VISCOSITY LIMIT OF NAVIER-STOKES EQUATIONS IN GEVREY CLASS

FENG CHENG, WEI-XI LI AND CHAO-JIANG XU

ABSTRACT. In this paper we consider the inviscid limit for the periodic solutions to Navier-Stokes equation in the framework of Gevrey class. It is shown that the lifespan for the solutions to Navier-Stokes equation is independent of viscosity, and that the solutions of the Navier-Stokes equation converge to that of Euler equation in Gevrey class as the viscosity tends to zero. Moreover the convergence rate in Gevrey class is presented.

1. INTRODUCTION

The Navier-Stokes equations for incompressible viscous flow in $\mathbb{T}^3 = (-\pi, \pi)^3$ read

$$\begin{cases} \frac{\partial u^\nu}{\partial t} - \nu \Delta u^\nu + (u^\nu \cdot \nabla) u^\nu + \nabla p^\nu = 0, \\ \nabla \cdot u^\nu = 0, \\ u^\nu|_{t=0} = a, \end{cases} \quad (1.1)$$

where $u^\nu(t, x) = (u_1^\nu, u_2^\nu, u_3^\nu)(t, x)$ is the unknown velocity vector function at point $x \in \mathbb{T}^3$ and time t , $p^\nu(t, x)$ is the unknown scalar pressure function, $\nu > 0$ is the kinematic viscosity, $a(x) = (a_1, a_2, a_3)(x)$ is the given initial data.

If the viscosity $\nu = 0$, the equations (1.1) become the Euler equations for ideal flow with the same given initial data a ,

$$\begin{cases} \frac{\partial u}{\partial t} + (u \cdot \nabla) u + \nabla p = 0, \\ \nabla \cdot u = 0, \\ u|_{t=0} = a, \end{cases} \quad (1.2)$$

where we denote the unknown vector velocity function to be $u(t, x) = (u_1, u_2, u_3)(t, x)$ and the unknown scalar pressure function to be $p(t, x)$.

The existence and uniqueness of solutions to (1.1) and (1.2) in Sobolev space $H^r(\mathbb{R}^3)$ for $r > 3/2 + 1$, on a maximal time interval $[0, T_*)$ is classical in [4, 15, 30]. There are abundant studies on the analyticities of solutions to (1.1) and (1.2) in various methods, for reference in [2, 3, 5, 12, 33]. The Gevrey regularity of solutions to Navier-Stokes equations was started by Foias and Temam in their work [9], in which the authors developed a way to prove the Gevrey class regularity by characterizing the decay of their Fourier coefficients. And later [17, 18, 19, 20, 21] developed this method to study the Gevrey class regularity of Euler equations in various conditions.

2010 *Mathematics Subject Classification.* 35M30, 76D03, 76D05.

Key words and phrases. Gevrey class, Incompressible Navier Stokes equation, Vanishing viscosity limit.

The subject of inviscid limits of solutions to Navier-Stokes equations has a long history and there is a vast literature on it, investigating this problem in various functional settings, cf. [16, 31] and references therein. Briefly, convergence of smooth solutions in \mathbb{R}^n or torus is well developed (cf. [15, 32] for instance). Much less is known about convergence in a domain with boundaries. In fact the vanishing viscosity limit for the incompressible Navier-Stokes equations, in the case where there exist physical boundaries, is still a challenging problem due to the appearance of the Prandtl boundary layer which is caused by the classical no-slip boundary condition. So far the rigorous verification of the Prandtl boundary layer theory was achieved only for some specific settings, cf. [1, 7, 11, 13, 22, 28, 34] for instance, not to mention the convergence to Prandtl's equation and Euler equations. Several partial results on the inviscid limits, in the case of half-space, were given in [33] by imposing analyticity on the initial data, and in [26] for vorticity admitting compact support which is away from the boundary.

On the other hand, the Prandtl boundary layer equation is ill-posed in Sobolev space for many case (see [7, 10, 25]), while the Sobolev space is the suitable function space for the energy theory of fluid mechanic. Since the verification of the Prandtl boundary layer theory meet the major obstacle in the setting of the Sobolev space, it will be interesting to expect the vanishing viscosity limit for the incompressible Navier-Stokes equations in the setting of Gevrey space as sub-space of Sobolev space, see a series of works in this direction [11, 22, 23]. In fact, Gevrey space is an intermediate space between the space of analytic functions and the Sobolev space. On one hand, Gevrey functions enjoy similar properties as analytic functions, and on the other hand, there are nontrivial Gevrey functions having compact support, which is different from analytic functions. As a preliminary attempt, in this work we study the vanishing viscosity limit of the solution of Navier Stokes equation to the solution of Euler equation in Gevrey space. Here we will concentrate on the torus, we hope this may give insights on the case when the domain has boundaries, which is a much more challenging problem.

We introduce the functions spaces as follows. We usually suppress the vector symbol for functions when no ambiguity arise. Let $\mathcal{L}^2(\mathbb{T}^3)$ be the vector function space

$$\mathcal{L}^2(\mathbb{T}^3) = \left\{ u = \sum_{k \in \mathbb{Z}^3} \hat{u}_j e^{i k \cdot x}; \hat{u}_k = \overline{\hat{u}_{-k}}, \hat{u}_0 = 0, j \cdot \hat{u}_j = 0, \right. \\ \left. \|u\|_{L^2}^2 = \sum_{k \in \mathbb{Z}^3} |\hat{u}_k|^2 < \infty \right\},$$

where \hat{u}_k is the k th order Fourier coefficient of u , $i = \sqrt{-1}$. The condition $j \cdot \hat{u}_j = 0$ means $\nabla \cdot u = 0$ in the weak sense, so it is the standard L^2 space with the divergence free condition. Let $\mathcal{H}^r(\mathbb{T}^3)$ be the vector periodic Sobolev space : for $r \geq 1$,

$$\mathcal{H}^r(\mathbb{T}^3) = \left\{ u = \sum_{k \in \mathbb{Z}^3} \hat{u}_j e^{i k \cdot x}; \hat{u}_k = \overline{\hat{u}_{-k}}, \hat{u}_0 = 0, j \cdot \hat{u}_j = 0 \right. \\ \left. \|u\|_{H^r}^2 = \sum_{k \in \mathbb{Z}^3} (1 + |k|^2)^r |\hat{u}_k|^2 < \infty \right\}.$$

Here the condition $j \cdot \hat{u}_j = 0$ means $\nabla \cdot u = 0$, so it is the standard Sobolev space H^r with the divergence free condition. Denote (\cdot, \cdot) the L^2 inner product of two

vector functions. Let us define the fractional differential operator $\Lambda = (-\Delta)^{1/2}$ and the exponential operator $e^{\tau\Lambda^{1/s}}$ as follows,

$$\Lambda u = \sum_{j \in \mathbb{Z}^3} |j| \hat{u}_j e^{ij \cdot x}, \quad e^{\tau\Lambda^{1/s}} u = \sum_{j \in \mathbb{Z}^3} e^{\tau|j|^{1/s}} \hat{u}_j e^{ij \cdot x}.$$

The vector Gevrey space $\mathcal{G}_{r,\tau}^s$ for $s \geq 1, \tau > 0, r \in \mathbb{R}$ is

$$\mathcal{G}_{r,\tau}^s(\mathbb{T}^3) = \left\{ u \in \mathcal{H}^r(\mathbb{T}^3); \quad \|u\|_{\mathcal{G}_{r,\tau}^s}^2 = \sum_{j \in \mathbb{Z}^3} |j|^{2r} e^{2\tau|j|^{1/s}} |\hat{u}_j|^2 < \infty \right\},$$

where the condition $j \cdot \hat{u}_j = 0$ means $\nabla \cdot u = 0$, so it is sub-space of the Sobolev space $\mathcal{H}^r(\mathbb{T}^3)$.

The following theorem is the main result of this paper.

Theorem 1.1. *Let $r > \frac{9}{2}, \tau_0 > 0, s \geq 1$. Assume that the initial data $a \in \mathcal{G}_{r,\tau_0}^s(\mathbb{T}^3)$, then there exists $\nu_0 > 0$ and $T > 0, \tau(t) > 0$ is a decreasing function such that, for any $0 < \nu \leq \nu_0$, the Navier-Stokes equations (1.1) admit the solutions*

$$u^\nu \in L^\infty([0, T]; \mathcal{G}_{r,\tau(\cdot)}^s(\mathbb{T}^3)); \quad p^\nu \in L^\infty([0, T]; \mathcal{G}_{r+1,\tau(\cdot)}^s(\mathbb{T}^3)),$$

and the Euler equations (1.2) admit the solution

$$u \in L^\infty([0, T]; \mathcal{G}_{r,\tau(\cdot)}^s(\mathbb{T}^3)); \quad p \in L^\infty([0, T]; \mathcal{G}_{r+1,\tau(\cdot)}^s(\mathbb{T}^3)),$$

Furthermore, we have the following convergence estimates : for any $0 < t \leq T$

$$\|u^\nu(t, \cdot) - u(t, \cdot)\|_{\mathcal{G}_{r-1,\tau(t)}^s} \leq C\sqrt{\nu}, \quad \|p^\nu(t, \cdot) - p(t, \cdot)\|_{\mathcal{G}_{r,\tau(t)}^s} \leq C\sqrt{\nu}, \quad (1.3)$$

where C is a constant depending on r, s, a and T .

Remark 1.1. The uniform lifespan is $0 < T < T_*$ where T_* is the maximal lifespan of \mathcal{H}^r solutions. The uniform (with respect to ν) Gevrey radius $\tau(t)$ of the solution is

$$\tau(t) = \frac{1}{e^{C_1 t} \frac{1}{\tau_0} + \frac{C_2}{C_1} (e^{C_1 t} - 1)} \quad (1.4)$$

where C_1, C_2 are constants depending on r, s, a, T .

Remark 1.2. Comparaison with the known works about Gevrey regularity of Navier-Stokes equations and Euler equations [2, 3, 9, 18, 19, 20], the additional difficulties of this work is the uniform estimate of Gevrey norm with respect to viscosity coefficients, and the estimate (1.3) with limit rates $\sqrt{\nu}$.

The paper is organized as follows. In section 2, we will give the known results and preliminary lemmas. Section 3 consists of a priori estimate and the existence of the solutions in Gevrey space. The convergence in Gevrey space will be given in section 4.

2. PRELIMINARY LEMMAS

We first recall the following classical result of Kato in [15].

Theorem 2.1. *Let $a \in \mathcal{H}^m(\mathbb{T}^3)$ for $m \geq 3$, then the following holds.*

(1). *There exists $T > 0$ depending on $\|a\|_{\mathcal{H}^m}$ but not on ν , such that (1.1) has a unique solution*

$$u^\nu \in C([0, T], \mathcal{H}^m(\mathbb{T}^3)).$$

Furthermore, $\{u^\nu\}$ is bounded in $C([0, T], \mathcal{H}^m(\mathbb{T}^3))$ for all $\nu > 0$.

(2). For each $t \in [0, T]$, $u(t) = \lim_{\nu \rightarrow 0} u^\nu(t)$ exists strongly in $\mathcal{H}^{m-1}(\mathbb{T}^3)$ and weakly in $\mathcal{H}^m(\mathbb{T}^3)$, uniformly in t . u is the unique solution to (1.2) satisfying

$$u \in C([0, T], \mathcal{H}^m(\mathbb{T}^3)).$$

Remark 2.1. The time T in Theorem 2.1 is actually depending on m and $\|a\|_{H^m}$, specifically

$$T < \frac{1}{C_m \|a\|_{H^m}},$$

where C_m is a constant depending on m . In fact, the constant C_m was created by using the Leibniz formula and Sobolev embedding inequality when estimating the nonlinear term. So, if the initial data $a \in \mathcal{G}_{r, \tau_0}^s(\mathbb{T}^3)$, then we have $a \in H_\sigma^m$, $\forall m$, because there exists a constant $C_{m, \tau_0, s}$ such that $\|a\|_{H^m} \leq C_{m, \tau_0, s} \|a\|_{\mathcal{G}_{r, \tau_0}^s}$. But we can't directly obtain an uniform bound for $C_m \|a\|_{H^m}$ by the Gevrey norm of $\|a\|_{\mathcal{G}_{r, \tau_0}^s}$ when m is very large. Then we can't say that, if m goes to infinity, $\frac{1}{C_m \|a\|_{H^m}}$ has a positive lower bound. In this paper, we will pay many attention to the uniform lifespan $T > 0$ that depends on $r, \|a\|_{H^r}$.

Remark 2.2. Compared with the known results Theorem 2.1, the additional difficulty arises on the estimate of the convecting term in Gevrey class setting. We need to use the decaying property of the radius of Gevrey class regularity to cancel the growth of the convecting term.

We will use the following inequality, for any $j, k \in \mathbb{Z}^3 \setminus \{0\}$, we have

$$|k - j| \leq 2|j||k|.$$

The proof is a simple result of triangle inequality which we omit the details here. And we will give two Lemmas which will be used in the proof of Theorem 1.1.

Lemma 2.2. *Given two real numbers $\xi, \eta \geq 1$ and $s \geq 1$, then the following inequality holds*

$$\left| \xi^{\frac{1}{s}} - \eta^{\frac{1}{s}} \right| \leq C \frac{|\xi - \eta|}{|\xi|^{1-\frac{1}{s}} + |\eta|^{1-\frac{1}{s}}} \quad (2.1)$$

where C is a positive constant depending only on s .

Proof. The case for $s = 1$ is trivial. Let us consider the case for $s > 1$. Without loss of generality, we may assume $\xi > \eta$. Then (2.1) is equivalent with

$$\frac{(\xi^{\frac{1}{s}} - \eta^{\frac{1}{s}})(\xi^{1-\frac{1}{s}} + \eta^{1-\frac{1}{s}})}{\xi - \eta} \leq C.$$

Then it suffices to show that

$$\left| \frac{(\frac{\eta}{\xi})^{1-\frac{1}{s}} - (\frac{\eta}{\xi})^{\frac{1}{s}}}{1 - \frac{\eta}{\xi}} \right| \leq C.$$

By Theorem 42 in [14], it can be obtained for fixed $s > 1$

$$\left| \frac{(\frac{\eta}{\xi})^{1-\frac{1}{s}} - (\frac{\eta}{\xi})^{\frac{1}{s}}}{1 - \frac{\eta}{\xi}} \right| \leq \max \left(1 - \frac{2}{s}, \frac{2}{s} - 1 \right) \leq C$$

Then the lemma 2.2 is proved. \square

With the use of Lemma 2.2, we have the following estimate about the nonlinear term.

Lemma 2.3. *Let $r > \frac{9}{2}$, $s \geq 1$ and $\tau > 0$ is a constant. Then for any $v \in \mathcal{G}_{r+1,\tau}^s(\mathbb{T}^3)$, the following estimate holds,*

$$\begin{aligned} \left| \left(\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v), \Lambda^r e^{\tau \Lambda^{1/s}} v \right) \right| &\leq C \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s}^2 + C \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s} \\ &+ \left[C\tau \|u\|_{H^r} + C\tau^2 (\|u\|_{H^r} + \|u\|_{\mathcal{G}_{r,\tau}^s}) \right] \|u\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2, \end{aligned} \quad (2.2)$$

where C is a constant depending only on r and s .

Proof. By the definition of the vector function space $\mathcal{G}_{r+1,\tau}^s(\mathbb{T}^3)$, we have $v = \sum_{j \in \mathbb{Z}^3} \hat{v}_j e^{ij \cdot x}$ and $\hat{v}_0 = 0$. Using Fourier series convolution property, one have

$$v \cdot \nabla v = i \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{v}_j \cdot (k-j)] \hat{v}_{k-j} e^{ik \cdot x}.$$

Applying the operator $\Lambda^r e^{\tau \Lambda^{1/s}}$ on $v \cdot \nabla v$, one have

$$\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v) = i \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{v}_j \cdot (k-j)] \hat{v}_{k-j} |k|^r e^{\tau |k|^{1/s}} e^{ik \cdot x}.$$

And $\Lambda^r e^{\tau \Lambda^{1/s}} v = \sum_{\ell \in \mathbb{Z}^3} |\ell|^r e^{\tau |\ell|^{1/s}} \hat{v}_\ell e^{i\ell \cdot x}$. Now we take the L^2 inner product of $\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v)$ with $\Lambda^r e^{\tau \Lambda^{1/s}} v$ over \mathbb{T}^3 . The orthogonality of the exponentials in L^2 implies

$$\begin{aligned} &\left(\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v), \Lambda^r e^{\tau \Lambda^{1/s}} v \right) \\ &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{v}_j \cdot (k-j)] (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^{2r} e^{2\tau |k|^{1/s}}. \end{aligned}$$

The cancellation property of the convecting term implies

$$\begin{aligned} 0 &= \left(v \cdot \nabla \Lambda^r e^{\tau \Lambda^{1/s}} v, \Lambda^r e^{\tau \Lambda^{1/s}} v \right) \\ &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{v}_j \cdot (k-j)] |k-j|^r e^{\tau |k-j|^{1/s}} (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^r e^{\tau |k|^{1/s}}. \end{aligned}$$

Then we have

$$\begin{aligned} &\left(\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v), \Lambda^r e^{\tau \Lambda^{1/s}} v \right) \\ &= \left(\Lambda^r e^{\tau \Lambda^{1/s}} (v \cdot \nabla v) - v \cdot \nabla \Lambda^r e^{\tau \Lambda^{1/s}} v, \Lambda^r e^{\tau \Lambda^{1/s}} v \right) \\ &= \mathcal{T}_1 + \mathcal{T}_2, \end{aligned}$$

where

$$\mathcal{T}_1 = i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} (|k|^r - |k-j|^r) e^{\tau |k-j|^{1/s}} [\hat{v}_j \cdot (k-j)] (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^r e^{\tau |k|^{1/s}},$$

and

$$\mathcal{T}_2 = i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^r (e^{\tau |k|^{1/s}} - e^{\tau |k-j|^{1/s}}) [\hat{v}_j \cdot (k-j)] (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^r e^{\tau |k|^{1/s}}.$$

Before we come to the estimate of \mathcal{T}_1 and \mathcal{T}_2 , we recall the following mean value theorem, for $\forall \xi, \eta \in \mathbb{R}^+$, there exists a constant $0 \leq \theta, \theta' \leq 1$ such that

$$\begin{aligned} \xi^r - \eta^r &= r(\xi - \eta) [(\theta\xi + (1 - \theta)\eta)^{r-1} - \xi^{r-1}] + r(\xi - \eta)\eta^{r-1} \\ &= r(r-1)\theta(\xi - \eta)^2 [\theta'(\theta\xi + (1 - \theta)\eta) + (1 - \theta')\eta]^{r-2} \\ &\quad + r(\xi - \eta)\eta^{r-1}. \end{aligned}$$

Then there exists a constant C depending only on r such that

$$||k|^r - |k-j|^r| \leq C|j|^2 (|j|^{r-2} + |k-j|^{r-2}) + C|j||k-j|^{r-1}.$$

From the inequality $e^\xi \leq e + \xi^2 e^\xi$ that holds for all $\xi \in \mathbb{R}$, we can bounded the exponential $e^{\tau|k-j|^{1/s}}$ by $e + \tau^2 |k-j|^{2/s} e^{\tau|k-j|^{1/s}}$. Then \mathcal{T}_1 can be bounded by

$$\begin{aligned} &|\mathcal{T}_1| \\ &\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} (|j|^r + |j|^2 |k-j|^{r-2}) |\hat{v}_j| |k-j| |\hat{v}_{k-j}| (e + \tau^2 |k-j|^{2/s} e^{\tau|k-j|^{1/s}}) \\ &\quad \times |\hat{v}_{-k}| |k|^r e^{\tau|k|^{1/s}} + C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j| |\hat{v}_j| |k-j|^r e^{\tau|k-j|^{1/s}} |\hat{v}_{k-j}| |\hat{v}_{-k}| |k|^r e^{\tau|k|^{1/s}} \\ &= \mathcal{T}_{11} + \mathcal{T}_{12} + \mathcal{T}_{13} + \mathcal{T}_{14} + \mathcal{T}_{15}. \end{aligned}$$

With application of discrete Hölder inequality and Minkowski inequality, one can obtain the following estimates. For example, we give the details for \mathcal{T}_{11} , and the rest can be estimated in the same way,

$$\begin{aligned} \mathcal{T}_{11} &= eC \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^r |\hat{v}_j| |k-j| |\hat{v}_{k-j}| |k|^r e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C \|v\|_{\mathcal{G}_{r,\tau}^s} \left[\sum_{k \in \mathbb{Z}^3} \left(\sum_{j \in \mathbb{Z}^3} |j|^r |\hat{v}_j| |k-j| |\hat{v}_{k-j}| \right)^2 \right]^{1/2} \\ &= C \|v\|_{\mathcal{G}_{r,\tau}^s} \left[\sum_{k \in \mathbb{Z}^3} \left(\sum_{\ell \in \mathbb{Z}^3} |k-\ell|^r |\hat{v}_{k-\ell}| |\ell| |\hat{v}_\ell| \right)^2 \right]^{1/2} \\ &\leq C \|v\|_{\mathcal{G}_{r,\tau}^s} \left(\sum_{k \in \mathbb{Z}^3} |k-\ell|^{2r} |\hat{v}_{k-\ell}|^2 \right)^{1/2} \sum_{\ell \in \mathbb{Z}^3} \frac{|\ell|}{(1+|\ell|^2)^{r/2}} (1+|\ell|^2)^{r/2} |\hat{v}_\ell| \\ &\leq C \|v\|_{H^r}^2 \|v\|_{\mathcal{G}_{r,\tau}^s} \left(\sum_{\ell \in \mathbb{Z}^3} \frac{|\ell|^2}{(1+|\ell|^2)^r} \right)^{1/2} \\ &\leq C \|v\|_{H^r}^2 \|v\|_{\mathcal{G}_{r,\tau}^s}, \end{aligned}$$

where C is a constant depending on r, e and for $r > 9/2$, the summation in the above $\left(\sum_{\ell \in \mathbb{Z}^3} \frac{|\ell|^2}{(1+|\ell|^2)^r} \right)^{1/2}$ is bounded by some constant depending on r . Similarly with \mathcal{T}_{11} , we have

$$\begin{aligned} \mathcal{T}_{12} &= eC \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^2 |\hat{v}_j| |k-j|^{r-1} |\hat{v}_{k-j}| |k|^r e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C \|v\|_{H^r}^2 \|v\|_{\mathcal{G}_{r,\tau}^s}, \end{aligned}$$

and

$$\begin{aligned}\mathcal{T}_{13} &= C\tau^2 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^r |\hat{v}_j| |k-j|^{1+2/s} |\hat{v}_{k-j}| |k|^r e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C\tau^2 \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s}.\end{aligned}$$

Note that $\hat{v}_0 = 0$, $s \geq 1$ in the summation, and $|k-j|^{1/2s} \leq C|k|^{1/2s}|j|^{1/2s}$, we can similarly have

$$\begin{aligned}\mathcal{T}_{14} &= C\tau^2 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^2 |\hat{v}_j| |k-j|^{r-1+2/s} e^{\tau|k-j|^{1/s}} |\hat{v}_{k-j}| |k|^r e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C\tau^2 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^{2+\frac{1}{2s}} |\hat{v}_j| |k-j|^{r+\frac{1}{2s}} e^{\tau|k-j|^{1/s}} |\hat{v}_{k-j}| |k|^{r+\frac{1}{2s}} e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C\tau^2 \|v\|_{H^r} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s},\end{aligned}$$

and

$$\begin{aligned}\mathcal{T}_{15} &= C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j| |\hat{v}_j| |k-j|^r e^{\tau|k-j|^{1/s}} |\hat{v}_{k-j}| |k|^r e^{\tau|k|^{1/s}} |\hat{v}_{-k}| \\ &\leq C \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s}.\end{aligned}$$

Noting that $\|v\|_{\mathcal{G}_{r,\tau}^s} \leq \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}$, then $\mathcal{T}_{13} \leq \mathcal{T}_{14}$. Thus we obtain

$$|\mathcal{T}_1| \leq C \|v\|_{H^r}^2 \|v\|_{\mathcal{G}_{r,\tau}^s} + C \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s}^2 + C\tau^2 \|v\|_{H^r} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2.$$

As for \mathcal{T}_2 , we have

$$\begin{aligned}\mathcal{T}_2 &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^r (e^{\tau|k|^{1/s}} - e^{\tau|k-j|^{1/s}}) [\hat{v}_j \cdot (k-j)] (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^r e^{\tau|k|^{1/s}} \\ &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^r e^{\tau|k-j|^{1/s}} \left[e^{\tau(|k|^{1/s} - |k-j|^{1/s})} - 1 \right] [\hat{v}_j \cdot (k-j)] \\ &\quad \times (\hat{v}_{k-j} \cdot \hat{v}_{-k}) |k|^r e^{\tau|k|^{1/s}}.\end{aligned}$$

We note that the inequality $|e^\xi - 1| \leq |\xi| e^{|\xi|}$ holds for $\xi \in \mathbb{R}$. Then

$$\left| e^{\tau(|k|^{1/s} - |k-j|^{1/s})} - 1 \right| \leq \tau \left| |k|^{1/s} - |k-j|^{1/s} \right| e^{\tau \left| |k|^{1/s} - |k-j|^{1/s} \right|}.$$

Since $s \geq 1$, we have

$$\left| |k|^{1/s} - |k-j|^{1/s} \right| \leq |j|^{1/s}.$$

Then we actually have

$$\left| e^{\tau(|k|^{1/s} - |k-j|^{1/s})} - 1 \right| \leq \tau \left| |k|^{1/s} - |k-j|^{1/s} \right| e^{\tau|j|^{1/s}}.$$

By Lemma 2.2, we have

$$\left| |k|^{1/s} - |k-j|^{1/s} \right| \leq C|j| \frac{1}{|k|^{1-1/s} + |k-j|^{1-1/s}}.$$

Then \mathcal{T}_2 can be bounded by the inequality

$$\begin{aligned}
|\mathcal{T}_2| &\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^r e^{\tau|k-j|^{1/s}} \left| e^{\tau(|k|^{1/s} - |k-j|^{1/s})} - 1 \right| |k-j| |\hat{v}_j| |\hat{v}_{k-j}| \\
&\quad \times |v_{-k}| |k|^r e^{\tau|k|^{1/s}} \\
&\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^{r-\frac{1}{2s}} e^{\tau|k-j|^{1/s}} \tau \left| |k|^{1/s} - |k-j|^{1/s} \right| e^{\tau|j|^{1/s}} |k-j| |\hat{v}_j| |\hat{v}_{k-j}| \\
&\quad \times |\hat{v}_{-k}| |k|^{r+\frac{1}{2s}} e^{\tau|k|^{1/s}} \\
&\leq C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} (|j|^{r-\frac{1}{2s}} + |k-j|^{r-\frac{1}{2s}}) e^{\tau|k-j|^{1/s}} \frac{|j||k-j|}{|k|^{1-1/s} + |k-j|^{1-1/s}} \\
&\quad \times e^{\tau|j|^{1/s}} |\hat{v}_j| |\hat{v}_{k-j}| |\hat{v}_{-k}| |k|^{r+\frac{1}{2s}} e^{\tau|k|^{1/s}} \\
&\leq \mathcal{T}_{21} + \mathcal{T}_{22},
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{T}_{21} &= C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j|^{r+\frac{1}{2s}} e^{\tau|j|^{1/s}} |\hat{v}_j| |k-j| (1 + \tau|k-j|^{1/s} e^{\tau|k-j|^{1/s}}) \\
&\quad \times |\hat{v}_{k-j}| |k|^{r+\frac{1}{2s}} e^{\tau|k|^{1/s}} |\hat{v}_{-k}|, \\
\mathcal{T}_{22} &= C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |j| (1 + \tau|j|^{1/s} e^{\tau|j|^{1/s}}) |\hat{v}_j| |k-j|^{r+\frac{1}{2s}} e^{\tau|k-j|^{1/s}} \\
&\quad \times |\hat{v}_{k-j}| |k|^{r+\frac{1}{2s}} e^{\tau|k|^{1/s}} |\hat{v}_{-k}|.
\end{aligned}$$

We have used the inequality $|k|^{1-1/s} + |k-j|^{1-1/s} \geq |j|^{1-1/s}$ and $e^\xi \leq 1 + \xi e^\xi$ for $\xi \in \mathbb{R}^+$ in the estimation of \mathcal{T}_{21} . With application of Hölder inequality and Minkowski inequality, we have for \mathcal{T}_{21} ,

$$|\mathcal{T}_{21}| \leq C\tau \|v\|_{H^r} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2 + C\tau^2 \|v\|_{\mathcal{G}_{r,\tau}^s} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2.$$

Symmetrically, one has a same bound for \mathcal{T}_{22} , then for \mathcal{T}_2 ,

$$|\mathcal{T}_2| \leq C\tau \|v\|_{H^r} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2 + C\tau^2 \|v\|_{\mathcal{G}_{r,\tau}^s} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2.$$

Then we obtain

$$\begin{aligned}
\left| \left(\Lambda^r e^{\tau\Lambda^{1/s}} (v \cdot \nabla v), \Lambda^r e^{\tau\Lambda^{1/s}} v \right) \right| &\leq |\mathcal{T}_1| + |\mathcal{T}_2| \\
&\leq C \|v\|_{H^r}^2 \|v\|_{\mathcal{G}_{r,\tau}^s} + C \|v\|_{H^r} \|v\|_{\mathcal{G}_{r,\tau}^s}^2 \\
&\quad + C\tau^2 \|v\|_{\mathcal{G}_{r,\tau}^s} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2 \\
&\quad + C\tau(1+\tau) \|v\|_{H^r} \|v\|_{\mathcal{G}_{r+\frac{1}{2s},\tau}^s}^2,
\end{aligned}$$

which finishes the proof of Lemma 2.3. \square

3. UNIFORM EXISTENCE OF SOLUTIONS

In this section, we will first show the existence of Gevrey class solutions u^ν to Navier Stokes equations (1.1). And the existence of Gevrey class solution u to Euler equations (1.2) can be obtained similarly. The method of the proof are based

on Galerkin approximation. Before that, we first consider the following equivalent equation for Navier-Stokes equation,

$$\begin{aligned} \frac{d}{dt}u^\nu + \nu Au^\nu + \mathbb{P}(u^\nu \cdot \nabla u^\nu) &= 0, \\ u^\nu|_{t=0} &= \mathbb{P}a. \end{aligned} \quad (3.1)$$

where $A = -\mathbb{P}\Delta$ is the well-known Stokes operator and \mathbb{P} is the Leray projector which maps a vector function v into its divergence free part v_1 , such that $v = v_1 + \nabla q$ and $\nabla \cdot v_1 = 0$, q is a scalar function and $(v_1, \nabla q) = 0$. Similarly for Euler equation, we have the following equivalent form,

$$\begin{aligned} \frac{d}{dt}u^\nu + \mathbb{P}(u^\nu \cdot \nabla u^\nu) &= 0, \\ u^\nu|_{t=0} &= \mathbb{P}a. \end{aligned} \quad (3.2)$$

We then recall some properties of the Stokes operator A , which are known in [Chapter 4 in [6]].

Proposition 3.1. *The Stokes operator A is symmetric and selfadjoint, moreover, the inverse of the Stokes operator, A^{-1} , is a compact operator in \mathcal{L}^2 . The Hilbert theorem implies there exists a sequence of positive numbers λ_j and an orthonormal basis w_j of \mathcal{L}^2 , which satisfies*

$$Aw_j = \lambda_j w_j, \quad 0 < \lambda_1 < \dots < \lambda_j \leq \lambda_{j+1} \leq \dots, \quad \lim_{j \rightarrow \infty} \lambda_j = \infty.$$

Moreover, in the case of $\mathbb{T}^3 = (-\pi, \pi)^3$, the sequence of eigenvector functions w_j 's and eigenvalues λ_j 's are the sequences of functions $w_{k,j}$ and numbers $\lambda_{k,j}$,

$$w_{k,j}(x) = \left(e_j - \frac{k_j k}{|k|^2} \right) e^{ik \cdot x}, \quad \lambda_{k,j} = |k|^2,$$

where $k = (k_1, k_2, k_3) \in \mathbb{Z}^3$, $k \neq 0$, $j = 1, 2, 3$ and $\{e_j\}_{j=1,2,3}$ are the canonical basis in \mathbb{R}^3 . So we know that each w_j are not only in \mathcal{L}^2 , but also in $\mathcal{G}_{r,\tau}^s$ for $\forall r > 0$. Now we will show that there exists a solution to equation (3.1) for $a \in \mathcal{G}_{r,\tau}^s$ with $r > 9/2$, $s \geq 1$, and $\tau(t) > 0$ is a differentiable decreasing function of t . To this end, we first prove a priori estimate in the following Proposition.

Proposition 3.2. *Let $r > 9/2$, $s \geq 1$, $a \in \mathcal{G}_{r,\tau_0}^s$ and $\tau(t) > 0$ is a differentiable decreasing function of t defined on $[0, T]$ with $\tau(0) = \tau_0 > 0$, where $0 < T < T^*$ and T^* is the maximal time of H^r solution to (3.1) with respect to the initial data a . Let $u^\nu(t, x) \in L^\infty([0, T]; \mathcal{G}_{r,\tau(\cdot)}^s(\mathbb{T}^3)) \cap L^2([0, T]; \mathcal{G}_{r+1,\tau(\cdot)}^s(\mathbb{T}^3))$ be the solution to (3.1), then the following a priori estimates holds,*

$$\begin{aligned} \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r,\tau(t)}^s} &\leq G_T, \quad 0 < t \leq T, \\ \nu \int_0^t \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1,\tau(s)}^s}^2 ds &\leq M_T, \quad 0 < t \leq T. \end{aligned}$$

With the same assumptions as above, let $u(t, x) \in L^\infty([0, T]; \mathcal{G}_{r+1,\tau(\cdot)}^s(\mathbb{T}^3))$ be the solution to (3.2), we also have

$$\|u(t, \cdot)\|_{\mathcal{G}_{r,\tau(t)}^s} \leq G_T, \quad 0 < t \leq T,$$

Furthermore the uniform radius $\tau(t)$ is given by

$$\tau(t) = \frac{1}{e^{C_1 t \frac{1}{\tau_0}} + \frac{C_2}{C_1}(e^{C_1 t} - 1)},$$

where C, C_1, C_2, G_T, M_T are constants depending on a, r, s, T .

Proof. Applying $\Lambda^r e^{\tau \Lambda^{1/s}}$ on both sides of (3.1) and taking the L^2 inner product of both sides with $\Lambda^r e^{\tau \Lambda^{1/s}} u^\nu$, one has

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r, \tau(t)}^s}^2 + \nu \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r+1, \tau(t)}^s}^2 \\ &= \tau'(t) \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r+\frac{1}{2s}, \tau}^s}^2 - \left(\Lambda^r e^{\tau \Lambda^{1/s}} (u^\nu \cdot \nabla u^\nu), \Lambda^r e^{\tau \Lambda^{1/s}} u^\nu \right), \end{aligned} \quad (3.3)$$

where we use the fact that \mathbb{P} commutes with $\Lambda^r e^{\tau \Lambda^{1/s}}$ and \mathbb{P} is symmetric. Now we consider the right hand side of (3.3). By (2.2) in Lemma 2.3, we have from (3.3), for convenience, we sometimes suppress the dependence of u^ν and τ in t ,

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2 + \nu \|u_\nu\|_{\mathcal{G}_{r+1, \tau}^s}^2 \leq C \|u^\nu\|_{H^r} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2 + C \|u^\nu\|_{H^r}^2 \|u^\nu\|_{\mathcal{G}_{r, \tau}^s} \\ &+ (\tau' + C\tau \|u^\nu\|_{H^r} + C\tau^2 \|u^\nu\|_{H^r} + C\tau^2 \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}) \|u^\nu\|_{\mathcal{G}_{r+\frac{1}{2s}, \tau}^s}^2, \end{aligned} \quad (3.4)$$

where C is a constant depending on r, s . Now if the radius of Gevrey class $\tau(t)$ is smooth and decreasing fast enough such that the following inequality holds,

$$\tau' + C\tau \|u^\nu\|_{H^r} + C\tau^2 \|u^\nu\|_{H^r} + C\tau^2 \|u^\nu\|_{\mathcal{G}_{r, \tau}^s} \leq 0. \quad (3.5)$$

Then (3.4) implies

$$\frac{1}{2} \frac{d}{dt} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2 + \nu \|u^\nu\|_{\mathcal{G}_{r+1, \tau}^s}^2 \leq C \|u^\nu\|_{H^r} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2 + C \|u^\nu\|_{H^r}^2 \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}. \quad (3.6)$$

As $\nu > 0$, it can be obtained directly from (3.6),

$$\frac{d}{dt} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s} \leq C \|u^\nu\|_{H^r} \|u^\nu\|_{\mathcal{G}_{r, \tau}^s} + C \|u^\nu\|_{H^r}^2, \quad (3.7)$$

By Gronwall's inequality in (3.7), we have for $0 < t < T_*$,

$$\|u^\nu(t)\|_{\mathcal{G}_{r, \tau(t)}^s} \leq \tilde{g}(t) \left(\|a\|_{\mathcal{G}_{r, \tau_0}^s} + \int_0^t C \tilde{g}(s)^{-1} \|u^\nu(s)\|_{H^r}^2 ds \right) \triangleq A(t), \quad (3.8)$$

where $\tilde{g}(t) = e^{\int_0^t C \|u^\nu(s, \cdot)\|_{H^r} ds}$ and T_* is the maximal time interval of H^r solution. It has been known that T_* is independent of ν . Moreover, it follows the a priori estimate for H^r solution for Navier Stokes equation in [27],

$$\frac{d}{dt} \|u^\nu(t, \cdot)\|_{H^r}^2 \leq C \|u^\nu(t, \cdot)\|_{H^r} \|u^\nu(t, \cdot)\|_{H^r}^2, \quad 0 < t < T_*, \quad (3.9)$$

where C is a constant depending on r . Then, on one side we have

$$\|u^\nu(t)\|_{H^r}^2 \leq C \tilde{g}(t) \|a\|_{H^r}^2, \quad 0 < t < T_*. \quad (3.10)$$

And on the other side, let $0 < T < T_*$, then for $0 < t < T$,

$$\|u^\nu(t, \cdot)\|_{H^r} \leq \frac{\|a\|_{H^r}}{1 - Ct \|a\|_{H^r}} \leq \frac{\|a\|_{H^r}}{1 - CT \|a\|_{H^r}} \triangleq C_T. \quad (3.11)$$

With (3.9), (3.10) and (3.11), we have

$$\begin{aligned}
 \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r, \tau(t)}^s} &\leq A(t) \\
 &= e^{\int_0^t C \|u^\nu(s)\|_{H^r} ds} \left(\|a\|_{G_{r, \tau_0}^s} + \int_0^t C \tilde{g}(s)^{-1} \|u^\nu(s)\|_{H^r}^2 ds \right) \\
 &\leq e^{CC_T t} \left(\|a\|_{G_{r, \tau_0}^s} + C \|a\|_{H^r}^2 t \right) \\
 &\leq e^{CC_T T} \left(\|a\|_{G_{r, \tau_0}^s} + C \|a\|_{H^r}^2 T \right) \triangleq G_T, \quad 0 \leq t \leq T.
 \end{aligned} \tag{3.12}$$

In fact a sufficient condition for (3.5) to hold is

$$\tau'(t) + C\tau(t)C_T + C\tau^2(t)C_T + C\tau^2(t)G_T = 0. \tag{3.13}$$

Then solving the ordinary differential equation (3.13),

$$\tau(t) = \frac{1}{e^{CC_T t} \frac{1}{\tau_0} + \frac{CC_T + CG_T}{CC_T} (e^{CC_T t} - 1)}. \tag{3.14}$$

We can obtain (1.4) by arranging the constants in (3.14),

$$\tau(t) = \frac{1}{e^{C_1 t} \frac{1}{\tau_0} + \frac{C_2}{C_1} (e^{C_1 t} - 1)}, \tag{3.15}$$

where C_1, C_2 are constants depending on r, s, T, a . Then (3.15) proves (1.4) in Remark 1.1. Integrating (3.6) from 0 to t , we have, for $0 < t < T$,

$$\nu \int_0^t \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1, \tau(s)}^s}^2 ds \leq M_T, \quad 0 < t < T < T_*, \tag{3.16}$$

where M_T depends on T, a, r, s, τ_0 . It should be noted that all of the above estimates are independent of ν , so we let $\nu = 0$ in (3.4), and proceed exactly as above, then we can obtain similar results for the a priori estimate for solution u to equation (3.2). \square

With the estimates in Proposition 3.2, we can implement the standard Faedo-Fourier-Galerkin approximation as in [24, 29] to prove the existence of such u^ν and u in the function space of Gevrey class $s \mathcal{G}_{r, \tau}^s$.

Theorem 3.1. *There exists a unique solution u^ν to (3.1) such that*

$$u^\nu \in L^\infty([0, T], \mathcal{G}_{r, \tau(\cdot)}^s).$$

Similarly there exists a unique solution u to (3.2) such that

$$u \in L^\infty([0, T], \mathcal{G}_{r, \tau(\cdot)}^s).$$

Proof. The method of proof of existence is based on Galerkin approximations and the priori estimate in Proposition 3.2. For a fixed positive integer n , we will look for a sequence of functions $u_n^\nu(t, \cdot)$ with $n \in \mathbb{N}$ in the form

$$u_n^\nu(t, x) = \sum_{j=1}^n \alpha_{j, n}^\nu(t) \omega_j,$$

where $\{\omega_j\}_{j=1}^\infty$ are the orthonormal basis in Proposition 3.1. Let W_n be the space spanned by $\{w_1, w_2, \dots, w_n\}$, and χ_n is the orthogonal projector in \mathcal{L}^2 into W_n .

The approximating equation is as follows,

$$\begin{cases} \frac{d}{dt}u_n^\nu + \nu Au_n^\nu + \chi_n \mathbb{P}(u_n^\nu \cdot \nabla u_n^\nu) = 0, \\ u_n^\nu|_{t=0} = \chi_n a \end{cases} \quad (3.17)$$

Taking the L^2 inner product with $w_j, j = 1, 2, \dots, n$, then the equation system (3.17) is equivalent with the following ordinary differential equation system,

$$\begin{cases} \frac{d}{dt}\alpha_{j,n}^\nu(t) + \nu\lambda_j\alpha_{j,n}^\nu + \sum_{k,\ell} b(\omega_k, \omega_\ell, \omega_j)\alpha_{k,n}^\nu\alpha_{\ell,n}^\nu = 0, & j = 1, 2, \dots, n, \\ \alpha_{j,n}^\nu(0) = (a, \omega_j), \end{cases} \quad (3.18)$$

where $b(\omega_k, \omega_\ell, \omega_j) = (\omega_k \cdot \nabla \omega_\ell, \omega_j)$ satisfying $b(\omega_k, \omega_\ell, \omega_j) = -b(\omega_k, \omega_j, \omega_\ell)$. By the standard ordinary differential equation theory, there exists a solution to (3.18) local in time interval $[0, T_n)$. In order to show that T_n can be extended to T , we multiply with $\alpha_{j,n}^\nu(t)$ on both sides of (3.18) and take sum over $1 \leq j \leq n$. We have

$$\frac{1}{2} \frac{d}{dt} \left(\sum_{j=1}^n \alpha_{j,n}^\nu(t)^2 \right) + \nu \lambda_j \sum_j \alpha_{j,n}^\nu(t)^2 = 0, \quad (3.19)$$

because

$$\sum_{j,k,\ell} b(\omega_k, \omega_\ell, \omega_j) \alpha_{k,n}^\nu \alpha_{\ell,n}^\nu \alpha_{j,n}^\nu = 0.$$

Moreover, from (3.19), we have

$$\|u_n^\nu(t, \cdot)\|_{L^2} = \sum_{j=1}^n \alpha_{j,n}^\nu(t)^2 \leq \sum_{j=1}^n \alpha_{j,n}^\nu(0)^2 \leq \|a\|_{L^2}^2, \quad \forall t > 0$$

Then we have for every T_n , it can be extended to arbitrary large, so it can be extended to T . And we also obtain

$$u_n^\nu \text{ remains bounded in } L^\infty([0, T]; L^2), \quad \forall n.$$

Moreover, we obtain a solution $u_n^\nu(t, x) = \sum_{j=1}^n \alpha_{j,n}^\nu(t) w_j(x)$ to (3.17) and we know that $u_n^\nu(t, x) \in \mathcal{G}_{r+1, \tau(t)}^s$ for $0 < t < T$ because it is only finite sum of w_j for fixed n . We then want to obtain the uniform Gevrey class norm bound for u_n^ν . To this end, we first apply $\Lambda^r e^{\tau\Lambda^{1/s}}$ on both sides of (3.17) and then take the L^2 inner product with $\Lambda^r e^{\tau\Lambda^{1/s}}$ to obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|u_n^\nu(t, \cdot)\|_{\mathcal{G}_{r, \tau(t)}^s}^2 + \nu \|u_n^\nu(t, \cdot)\|_{\mathcal{G}_{r+1, \tau}^s}^2 \\ & = \tau'(t) \|u_n^\nu(t, \cdot)\|_{\mathcal{G}_{r+\frac{1}{2s}, \tau}^s}^2 - \left(\Lambda^r e^{\tau\Lambda^{1/s}} \chi_n \mathbb{P}(u_n^\nu \cdot \nabla u_n^\nu), \Lambda^r e^{\tau(t)\Lambda^{1/s}} u_n^\nu \right). \end{aligned}$$

We note that the operator χ_n and \mathbb{P} commute with $\Lambda^r e^{\tau\Lambda^{1/s}}$, and they are symmetric, then

$$\left(\Lambda^r e^{\tau\Lambda^{1/s}} \chi_n \mathbb{P}(u_n^\nu \cdot \nabla u_n^\nu), \Lambda^r e^{\tau(t)\Lambda^{1/s}} u_n^\nu \right) = \left(\Lambda^r e^{\tau\Lambda^{1/s}} (u_n^\nu \cdot \nabla u_n^\nu), \Lambda^r e^{\tau(t)\Lambda^{1/s}} u_n^\nu \right).$$

With the arguments in Proposition 3.2, we have,

$$\|u_n^\nu(t, \cdot)\|_{\mathcal{G}_{r, \tau(t)}^s} \leq G_T, \quad 0 < t \leq T < T^*, \quad \forall n.$$

Thus

$$u_n^\nu \text{ remains bounded in } L^\infty([0, T]; \mathcal{G}_{r, \tau}^s(\cdot)). \quad (3.20)$$

In order to pass to the limit in the nonlinear term using a compactness theorem, we need to estimate on $\frac{du_n^\nu}{dt}$. From (3.17), we have

$$\frac{du_n^\nu}{dt} = -\nu Au_n^\nu - \chi_n \mathbb{P}(u_n^\nu \cdot \nabla u_n^\nu).$$

Then we obtain

$$\begin{aligned} \left\| \frac{du_n^\nu}{dt} \right\|_{L^2} &\leq \|\chi_n \mathbb{P}(u_n^\nu \cdot \nabla u_n^\nu)\|_{L^2} + \|\nu Au_n^\nu\|_{L^2} \\ &\leq \|u_n^\nu \cdot \nabla u_n^\nu\|_{L^2} + \nu_0 \|u_n^\nu\|_{H^r} \\ &\leq C \|u_n^\nu\|_{H^r}^2 + \nu_0 \|u_n^\nu\|_{H^r}. \end{aligned}$$

We recall that

$$\|u_n^\nu(t, \cdot)\|_{H^r} \leq C_T, \quad 0 < t \leq T, \quad \forall n.$$

So we obtain

$$\frac{d}{dt} u_m^\nu \text{ remains bounded in } L^\infty([0, T]; L^2). \quad (3.21)$$

By (3.20) and (3.21), noting that $H^r(\mathbb{T}^3)$ is compactly embedded in $L^2(\mathbb{T}^3)$ from Rellich-Kondrachov Compactness Theorem in [8], a compactness theorem in [24, 29] indicates the existence of the limit $u^\nu \in L^\infty([0, T]; \mathcal{G}_{r, \tau}^s(\cdot))$ of a subsequence of u_m^ν such that

$$\begin{cases} \frac{d}{dt} (u^\nu(t), v) - \nu (\Delta u^\nu(t), v) + (u^\nu(t) \cdot \nabla u^\nu(t), v) = 0, & \forall v \in \mathcal{L}^2 \\ u^\nu(0) = a \end{cases} \quad (3.22)$$

For Euler equations, one can take very similar approach to obtain the existence of solution in Gevrey class space and we omit the details here. Thus we prove Proposition 3.1. \square

It remains to show that u^ν is the solution of (1.1). In fact it can be obtained from (3.22) that

$$\mathbb{P} \left\{ \frac{d}{dt} u^\nu(t) - \nu \Delta u^\nu(t) + u^\nu(t) \cdot \nabla u^\nu(t) \right\} = 0.$$

So there exists a scalar function p^ν such that

$$\frac{d}{dt} u^\nu(t) - \nu \Delta u^\nu(t) + u^\nu(t) \cdot \nabla u^\nu(t) + \nabla p^\nu = 0,$$

where p^ν is unique up to a constant, and p^ν satisfies

$$-\Delta p^\nu = \nabla \cdot (u^\nu \cdot \nabla u^\nu), \quad (3.23)$$

with periodic boundary condition. For the regularity of the pressure $p^\nu(t, x)$, we have the following Proposition.

Proposition 3.3. *Let p^ν satisfies (3.23), then the following estimate holds,*

$$\|p^\nu(t, \cdot)\|_{\mathcal{G}_{r+1, \tau}^s} \leq C G_T^2, \quad 0 < t \leq T.$$

And for the pressure $p(t, x)$ in (1.2), we also have

$$\|p(t, \cdot)\|_{\mathcal{G}_{r+1, \tau}^s} \leq C G_T^2, \quad 0 < t \leq T,$$

where C, T, G_T are defined in Proposition 3.2.

Proof. To study the pressure p^ν , the existence is obvious results from standard elliptic equation theory. We consider the regularity of $p^\nu(t, x)$. First we apply the operator $\Lambda^r e^{\tau\Lambda^{1/s}}$ on both sides of (3.23) and then take L^2 inner product with $\Lambda^r e^{\tau\Lambda^{1/s}} p^\nu$ to obtain

$$-\left(\Delta p^\nu, \Lambda^{2r} e^{2\tau\Lambda^{1/s}} p^\nu\right) = \left(\nabla \cdot (u^\nu \cdot \nabla u^\nu), \Lambda^{2r} e^{2\tau\Lambda^{1/s}} p^\nu\right). \quad (3.24)$$

Here if we can write $p^\nu(t, x) = \sum_{j \in \mathbb{Z}^3} \hat{p}_j^\nu e^{ij \cdot x}$, then the left side of (3.24) is

$$-\left(\Delta p^\nu, \Lambda^{2r} e^{2\tau\Lambda^{1/s}} p^\nu\right) = \sum_{j \in \mathbb{Z}^3} |j|^{2r+2} e^{2\tau|j|^{1/s}} |\hat{p}_j^\nu|^2 = \|p^\nu\|_{\mathcal{G}_{r+1, \tau}^s}^2.$$

The right hand side of (3.24) can be bounded by

$$\begin{aligned} & \left| \left(\nabla \cdot (u^\nu \cdot \nabla u^\nu), \Lambda^{2r} e^{2\tau\Lambda^{1/s}} p^\nu \right) \right| \\ &= \left| (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^{r-1} [\hat{u}_j^\nu \cdot (k-j)] (k \cdot \hat{u}_{k-j}^\nu) \hat{p}_{-k}^\nu |k|^{r+1} e^{2\tau|k|^{1/s}} \right| \\ &= \left| (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} |k|^{r-1} [\hat{u}_j^\nu \cdot (k-j)] (j \cdot \hat{u}_{k-j}^\nu) \hat{p}_{-k}^\nu |k|^{r+1} e^{2\tau|k|^{1/s}} \right| \\ &\leq C \|p^\nu\|_{\mathcal{G}_{r+1, \tau}^s} \\ &\times \left\{ \sum_{k \in \mathbb{Z}^3} \left[\sum_{j \in \mathbb{Z}^3} (|j|^{r-1} + |k-j|^{r-1}) |j| |k-j| |\hat{u}_j^\nu| e^{\tau|j|^{1/s}} e^{\tau|k|^{1/s}} |\hat{u}_{k-j}^\nu| \right]^2 \right\}^{1/2} \\ &\leq C \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2 \|p^\nu\|_{\mathcal{G}_{r+1, \tau}^s}, \end{aligned}$$

where C is a constant depending on r . Then from above estimate, we obtain

$$\|p^\nu\|_{\mathcal{G}_{r+1, \tau}^s} \leq C \|u^\nu\|_{\mathcal{G}_{r, \tau}^s}^2.$$

From (3.8), we obtain

$$\|p^\nu(t)\|_{\mathcal{G}_{r+1, \tau}^s} \leq CA(t)^2 \leq CG_T^2, \quad t \in [0, T].$$

For the pressure $p(t, x)$ of Euler equation (1.2), one can first obtain the following elliptic equation,

$$-\Delta p = \nabla \cdot (u \cdot \nabla u).$$

Then using the same arguments as above, one can obtain the same results for p . \square

4. CONVERGENCE OF SOLUTIONS IN GEVREY SPACE

In the previous Section we have proved the existence of solutions to the Navier-Stokes equation and Euler equation in Gevrey class space. In this Section we will show the vanishing viscosity limit of Navier-Stokes equation in Gevrey class space. Moreover, we can obtain the converging rate with respect to ν .

Theorem 4.1. *Let u^ν, p^ν and u, p are the solutions we have obtained in the previous Section, where*

$$u^\nu, u \in L^\infty([0, T], \mathcal{G}_{r, \tau(\cdot)}^s), \quad p^\nu, p \in L^\infty([0, T], \mathcal{G}_{r+1, \tau(\cdot)}^s).$$

Then the following estimates hold,

$$\|u^\nu(t, \cdot) - u(t, \cdot)\|_{\mathcal{G}_{r-1, \tau(t)}^s} \leq C\sqrt{\nu}, \quad \|p^\nu(t, \cdot) - p(t, \cdot)\|_{\mathcal{G}_{r, \tau(t)}^s} \leq C\sqrt{\nu}, \quad (4.1)$$

for any $0 < t \leq T$, where C is a constant depending on r, s, a, T .

Proof. Let us first consider the new equation for $w = (u^\nu - u)$ and $\tilde{p} = p^\nu - p$,

$$\begin{cases} \frac{\partial w}{\partial t} - \nu \Delta u^\nu + w \cdot \nabla u^\nu + u \cdot \nabla w + \nabla \tilde{p} = 0, \\ \nabla \cdot w = 0, \\ w|_{t=0} = 0. \end{cases} \quad (4.2)$$

Then we apply the operator $\Lambda^{r-1} e^{\tau \Lambda^{1/s}}$ on both sides of (4.2) and take the L^2 inner product with $\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w$ on both sides to obtain,

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w(t)\|_{\mathcal{G}_{r-1, \tau}^s}^2 &= \nu \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \Delta u^\nu, \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right) + \tau' \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s}, \tau}^s}^2 \\ &\quad - \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} (w \cdot \nabla u^\nu), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right) \\ &\quad - \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} (u \cdot \nabla w), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right), \end{aligned} \quad (4.3)$$

where the term $\left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \nabla \tilde{p}, \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right)$ disappears since $w = u^\nu - u$ is divergence free. It remains to estimate the right hand side of (4.3), for convenience, we denote

$$\begin{aligned} \mathcal{I}_1 &= \nu \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \Delta u^\nu, \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right), \\ \mathcal{I}_2 &= \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} (w \cdot \nabla u^\nu), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right), \\ \mathcal{I}_3 &= \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} (u \cdot \nabla w), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right). \end{aligned}$$

Using the discrete Hölder inequality, one can obtain

$$\begin{aligned} |\mathcal{I}_1| &= \nu \left| \left(\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \Delta u^\nu, \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} w \right) \right| \\ &= \nu \left| -(2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^{2r} e^{2\tau|k|^{1/s}} (\hat{u}_k^\nu \cdot \hat{w}_{-k}) \right| \\ &\leq \nu (2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^{r+1} e^{\tau|k|^{1/s}} |\hat{u}_k^\nu| |k|^{r-1} e^{\tau|k|^{1/s}} |\hat{w}_{-k}| \\ &\leq C\nu \|u^\nu\|_{\mathcal{G}_{r+1, \tau}^s} \|w\|_{\mathcal{G}_{r-1, \tau}^s}. \end{aligned} \quad (4.4)$$

Then we have

$$|\mathcal{I}_1| \leq C\nu \|u^\nu\|_{\mathcal{G}_{r+1, \tau}^s} \|w\|_{\mathcal{G}_{r-1, \tau}^s}.$$

As for \mathcal{I}_2 , we first write it into the sum of their Fourier coefficients,

$$\mathcal{I}_2 = i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{w}_j \cdot (k-j)] (\hat{u}_{k-j}^\nu \cdot \hat{w}_{-k}) |k|^{2(r-1)} e^{2\tau|k|^{1/s}}.$$

Since $r > 9/2$, there exists a constant C such that

$$|k|^{r-1} \leq C \left(|j|^{r-1} + |k-j|^{r-1} \right),$$

and $s \geq 1$ implies

$$e^{\tau|k|^{1/s}} \leq e^{\tau|j|^{1/s}} e^{\tau|k-j|^{1/s}}.$$

Thus \mathcal{I}_2 can be bounded by

$$\begin{aligned} |\mathcal{I}_2| &\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[(|j|^{r-1} + |k-j|^{r-1}) |\hat{w}_j| |k-j| |\hat{u}'_{k-j}| e^{\tau|j|^{1/s}} e^{\tau|k-j|^{1/s}} \right. \\ &\quad \left. \times |\hat{w}_{-k}| |k|^{r-1} e^{\tau|k|^{1/s}} \right]. \end{aligned}$$

Then by discrete Hölder inequality and Minkowski inequality, we obtain

$$|\mathcal{I}_2| \leq C \|u'\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1,\tau}^s}^2. \quad (4.5)$$

As for \mathcal{I}_3 , where

$$\mathcal{I}_3 = i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) |k|^{2(r-1)} e^{2\tau|k|^{1/s}}.$$

Here again the cancellation property implies

$$\begin{aligned} 0 &= \left(u \cdot \nabla \Lambda^{r-1} e^{\tau \Lambda^{1/s}} w, \Lambda^{r-1} e^{\tau \Lambda^{1/s}} w \right) \\ &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) |k-j|^{r-1} e^{\tau|k-j|^{1/s}} |k|^{r-1} e^{\tau|k|^{1/s}}. \end{aligned}$$

Then we have

$$\begin{aligned} \mathcal{I}_3 &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) |k|^{2(r-1)} e^{2\tau|k|^{1/s}} \\ &\quad - i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) |k-j|^{r-1} e^{\tau|k-j|^{1/s}} |k|^{(r-1)} e^{\tau|k|^{1/s}} \\ &= \mathcal{R}_1 + \mathcal{R}_2, \end{aligned}$$

where we denote

$$\begin{aligned} \mathcal{R}_1 &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left\{ [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) \left(|k|^{r-1} - |k-j|^{r-1} \right) e^{\tau|k|^{1/s}} \right. \\ &\quad \left. \times |k|^{r-1} e^{\tau|k|^{1/s}} \right\}, \end{aligned}$$

and

$$\begin{aligned} \mathcal{R}_2 &= i(2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left\{ [\hat{u}_j \cdot (k-j)] (\hat{w}_{k-j} \cdot \hat{w}_{-k}) |k-j|^{r-1} \left(e^{\tau|k|^{1/s}} - e^{\tau|k-j|^{1/s}} \right) \right. \\ &\quad \left. \times |k|^{r-1} e^{\tau|k|^{1/s}} \right\}. \end{aligned}$$

Here we used a different strategy in the split of $|k|^{r-1} e^{\tau|k|^{1/s}} - |k-j|^{r-1} e^{\tau|k-j|^{1/s}}$ as in Lemma 2.3 to estimate \mathcal{R}_1 and \mathcal{R}_2 . With use of the following mean value theorem, there exists a constant $\theta \in (0, 1)$ such that

$$\begin{aligned} \left| |k|^{r-1} - |k-j|^{r-1} \right| &= \left| (r-1)(|k| - |k-j|) [\theta|k| + (1-\theta)|k-j|]^{r-2} \right| \\ &\leq C|j| (|k|^{r-2} + |k-j|^{r-2}). \end{aligned}$$

Then by discrete Hölder inequality and Minkowski inequality, we have

$$\begin{aligned} |\mathcal{R}_1| &\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[|\hat{u}_j| |k-j| |\hat{w}_{k-j}| |\hat{w}_{-k}| |j| (|j|^{r-2} + |k-j|^{r-2}) \right. \\ &\quad \left. \times e^{\tau|j|^{1/s}} e^{\tau|k-j|^{1/s}} |k|^{r-1} e^{\tau|k|^{1/s}} \right] \\ &\leq C \|u\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1,\tau}^s}^2. \end{aligned}$$

As for \mathcal{R}_2 , we use the inequality $|e^\xi - 1| \leq |\xi| e^{|\xi|}$ for $\forall \xi \in \mathbb{R}$ and Lemma 2.2,

$$\begin{aligned} |\mathcal{R}_2| &\leq C \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[|\hat{u}_j| |\hat{w}_{k-j}| |\hat{w}_{-k}| |k-j|^r e^{\tau|k-j|^{1/s}} \left| e^{\tau(|k|^{1/s} - |k-j|^{1/s})} - 1 \right| \right. \\ &\quad \left. \times |k|^{r-1} e^{\tau|k|^{1/s}} \right] \\ &\leq C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[|\hat{u}_j| |\hat{w}_{k-j}| |\hat{w}_{-k}| |k-j|^{r-1} e^{\tau|k-j|^{1/s}} e^{\tau|j|^{1/s}} \frac{|j| |k-j|}{|k|^{1-\frac{1}{s}} + |k-j|^{1-\frac{1}{s}}} \right. \\ &\quad \left. \times |k|^{r-1} e^{\tau|k|^{1/s}} \right]. \end{aligned}$$

For here we use the inequality $|k-j| \leq 2|k||j|$ for $k, j \neq 0$, then we have

$$\begin{aligned} \frac{|k-j|}{|k|^{1-\frac{1}{s}} + |k-j|^{1-\frac{1}{s}}} &\leq |k-j|^{1/s} \\ &\leq C |k-j|^{\frac{1}{2s}} |k|^{\frac{1}{2s}} |j|^{\frac{1}{2s}}, \end{aligned}$$

where C is a constant depending on s . Thus \mathcal{R}_2 can be bounded as follows,

$$\begin{aligned} |\mathcal{R}_2| &\leq C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[|\hat{u}_j| |\hat{w}_{k-j}| |\hat{w}_{-k}| |k-j|^{r-1+\frac{1}{2s}} e^{\tau|k-j|^{1/s}} e^{\tau|j|^{1/s}} |j|^{1+\frac{1}{2s}} \right. \\ &\quad \left. \times |k|^{r-1+\frac{1}{2s}} e^{\tau|k|^{1/s}} \right] \\ &\leq C\tau \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[|\hat{u}_j| |\hat{w}_{k-j}| |\hat{w}_{-k}| |k-j|^{r-1+\frac{1}{2s}} e^{\tau|k-j|^{1/s}} (1 + \tau|j|^{1/s} e^{\tau|j|^{1/s}}) \right. \\ &\quad \left. \times |j|^{1+\frac{1}{2s}} |k|^{r-1+\frac{1}{2s}} e^{\tau|k|^{1/s}} \right] \\ &\leq C\tau \|u\|_{H^r} \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s},\tau}^s}^2 + C\tau^2 \|u\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s},\tau}^s}^2, \end{aligned}$$

where we use the inequality $e^\xi \leq 1 + \xi e^\xi$ for $\forall \xi \in \mathbb{R}^+$ with respect to $e^{\tau|j|^{1/s}}$ and also the discrete Hölder inequality and Minkowski inequality in the above inequality. Then we have

$$|\mathcal{I}_3| \leq C \|u\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1,\tau}^s}^2 + C\tau \|u\|_{H^r} \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s},\tau}^s}^2 + C\tau^2 \|u\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s},\tau}^s}^2. \quad (4.6)$$

Substituting (4.4), (4.5) and (4.6) into (4.3), we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|w\|_{\mathcal{G}_{r-1,\tau}^s}^2 &\leq \nu \|u^\nu\|_{\mathcal{G}_{r+1,\tau}^s} \|w\|_{\mathcal{G}_{r-1,\tau}^s} + C \|u\|_{\mathcal{G}_{r,\tau}^s} \|w\|_{\mathcal{G}_{r-1,\tau}^s}^2 \\ &\quad + \left(\tau' + C\tau \|u\|_{H^r} + C\tau^2 \|u\|_{\mathcal{G}_{r,\tau}^s} \right) \|w\|_{\mathcal{G}_{r-1+\frac{1}{2s},\tau}^s}^2. \end{aligned} \quad (4.7)$$

By the choice of τ in (3.13), and noting that (3.8),(3.9),(3.10),(3.11) also hold for Euler equation (1.2). Then choosing the appropriate constant C , one has $\tau' + C\tau \|u\|_{H^r} + C\tau^2 \|u\|_{H^r} + C\tau^2 \|u\|_{\mathcal{G}_{\tau,\tau}^s} \leq 0$, then we can obtain from (4.7) and (3.12),

$$\frac{d}{dt} \|w(t, \cdot)\|_{\mathcal{G}_{r-1,\tau(t)}^s} \leq \nu \|u^\nu(t, \cdot)\|_{\mathcal{G}_{r+1,\tau(t)}^s} + CG_T \|w(t, \cdot)\|_{\mathcal{G}_{r-1,\tau(t)}^s}, \quad 0 < t \leq T. \quad (4.8)$$

Since $w(0) = 0$ and Grownwall's inequality, (4.8) implies

$$\|w(t, \cdot)\|_{\mathcal{G}_{r-1,\tau(t)}^s} \leq e^{CG_T t} \int_0^t \nu \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1,\tau(s)}^s} ds, \quad 0 < t \leq T.$$

Recalling from (3.16) we have for $0 < t \leq T$,

$$\int_0^t \nu \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1,\tau(s)}^s}^2 ds \leq M_T, \quad t \in (0, T].$$

With Hölder inequality, we have

$$\begin{aligned} \int_0^t \nu \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1,\tau(s)}^s} ds &\leq \int_0^t \sqrt{\nu} \sqrt{\nu} \|u^\nu(s, \cdot)\|_{\mathcal{G}_{r+1,\tau(s)}^s} ds \\ &\leq \sqrt{\nu} t^{1/2} M_T^{1/2}. \end{aligned}$$

Then we have

$$\|w(t, \cdot)\|_{\mathcal{G}_{r-1,\tau(t)}^s} \leq C \sqrt{\nu} M_T^{1/2} t^{1/2} e^{CG_T t}, \quad 0 < t \leq T. \quad (4.9)$$

This proves the first estimate of (4.1) by arranging the constant. Then we want to estimate $p^\nu(t) - p(t)$ in the norm of $\mathcal{G}_{r,\tau}^s$. To do so, we first take the divergence of both sides of (4.2) to obtain the following elliptic equation,

$$-\Delta \tilde{p} = \nabla \cdot (w \cdot \nabla u^\nu) + \nabla \cdot (u \cdot \nabla w). \quad (4.10)$$

Then we first apply the operator $\Lambda^{r-1} e^{\tau \Lambda^{1/s}}$ on both sides of (4.10) and then take the L^2 inner product with $\Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \tilde{p}$ on both sides to obtain,

$$\begin{aligned} (2\pi)^3 \|\tilde{p}\|_{\mathcal{G}_{r,\tau}^s}^2 &= \left(\Lambda^{r-1} e^{\tau \Lambda^{1/s}} \nabla \cdot (w \cdot \nabla u^\nu), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \tilde{p} \right) \\ &\quad + \left(\Lambda^{r-1} e^{\tau \Lambda^{1/s}} \nabla \cdot (u \cdot \nabla w), \Lambda^{(r-1)} e^{\tau \Lambda^{1/s}} \tilde{p} \right) \\ &= i^2 (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{w}_j \cdot (k-j)] (k \cdot \hat{u}_{k-j}^\nu) |k|^{2(r-1)} e^{2\tau |k|^{1/s}} \hat{p}_{-k} \\ &\quad + i^2 (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (k \cdot \hat{w}_{k-j}) |k|^{2(r-1)} e^{2\tau |k|^{1/s}} \hat{p}_{-k} \\ &= \mathcal{P}_1 + \mathcal{P}_2, \end{aligned} \quad (4.11)$$

where we denote

$$\begin{aligned} \mathcal{P}_1 &= i^2 (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{w}_j \cdot (k-j)] (k \cdot \hat{u}_{k-j}^\nu) |k|^{2(r-1)} e^{2\tau |k|^{1/s}} \hat{p}_{-k}, \\ \mathcal{P}_2 &= i^2 (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{u}_j \cdot (k-j)] (k \cdot \hat{w}_{k-j}) |k|^{2(r-1)} e^{2\tau |k|^{1/s}} \hat{p}_{-k}. \end{aligned}$$

It remains to estimate \mathcal{P}_1 and \mathcal{P}_2 in (4.11). For simplicity we only compute \mathcal{P}_1 , since \mathcal{P}_2 can be estimated in the same way.

$$\begin{aligned}
 |\mathcal{P}_1| &= \left| (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} [\hat{w}_j \cdot (k-j)] (j \cdot \hat{u}_{k-j}^\nu) |k|^{2(r-1)} e^{2\tau|k|^{1/s}} \hat{p}_{-k} \right| \\
 &\leq (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[C(|j|^{r-2} + |k-j|^{r-2}) e^{\tau|j|^{1/s}} e^{\tau|k-j|^{1/s}} |k-j| \right. \\
 &\quad \left. \times |j| |\hat{w}_j| |\hat{u}_{k-j}^\nu| |k|^r e^{\tau|k|^{1/s}} \left| \hat{p}_{-k} \right| \right] \\
 &\leq C \|w\|_{G_{r-1, \tau}^s} \|u^\nu\|_{G_{r, \tau}^s} \|\tilde{p}\|_{G_{r, \tau}^s},
 \end{aligned} \tag{4.12}$$

where we use the fact $k \cdot \hat{u}_{k-j}^\nu = j \cdot \hat{u}_{k-j}^\nu$ from the divergence free condition. And, similarly, we can obtain

$$\begin{aligned}
 |\mathcal{P}_2| &\leq (2\pi)^3 \sum_{k \in \mathbb{Z}^3} \sum_{j \in \mathbb{Z}^3} \left[[\hat{u}_j \cdot (k-j)] (j \cdot \hat{w}_{k-j}) |k|^{2(r-1)} e^{2\tau|k|^{1/s}} \hat{p}_{-k} \right] \\
 &\leq C \|w\|_{G_{r-1, \tau}^s} \|u\|_{G_{r, \tau}^s} \|\tilde{p}\|_{G_{r, \tau}^s}.
 \end{aligned} \tag{4.13}$$

Substituting (4.12) and (4.13) into (4.11), we obtain

$$\|\tilde{p}(t, \cdot)\|_{G_{r, \tau(t)}^s} \leq CG_T \|w(t, \cdot)\|_{G_{r-1, \tau(t)}^s}, \quad 0 < t \leq T. \tag{4.14}$$

Then by (4.9) and (4.14) we have

$$\|\tilde{p}(t, \cdot)\|_{G_{r, \tau(t)}^s} \leq C\sqrt{\nu} M_T^{1/2} t^{1/2} G_T e^{CG_T t}, \quad 0 \leq t \leq T.$$

This proves (4.1) by arranging the constants. Thus we have proven Theorem 1.1. \square

Acknowledgements. The research of the second author was supported by NSF of China(11422106) and Fok Ying Tung Education Foundation (151001), the research of the first author and the last author is supported partially by ‘‘The Fundamental Research Funds for Central Universities of China’’.

REFERENCES

- [1] R. Alexandre, Y. Wang, C.-J. Xu and T. Yang : Well-posedness of The Prandtl Equation in Sobolev Spaces. *J. Amer. Math. Soc.*, **28**(2015), no. 3, 745-784.
- [2] C. Bardos and S. Benachour : Domaine d’analyticit  des solutions de l’ quation d’Euler dans un ouvert de \mathbb{R}^n , *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, **4**(1977), no. 4, 647-687.
- [3] A. Biswas and D. Swanson : Gevrey regularity of solutions to the 3-D Navier-Stokes equations with weighted ℓ_p initial data, *Indiana Univ. math. J.*, **56**(2007), no. 3, 1157-1188.
- [4] J. P. Bourguignon and H. Brezis : Remarks on the Euler equation, *J. Functional Analysis*, **15**(1974), 341-363.
- [5] D. Chae : Remarks on the regularity of weak solutions of the Navier-Stokes equations, *Comm. Partial Differential Equations*, **17**(1992), no 7-8, 1267-1285.
- [6] P. Constantin and C. Foias : *Navier-Stokes Equations*, University of Chicago Press, 1988.
- [7] Weinan. E and B. Engquist : Blow up of solutions of the unsteady Prandtl’s equation, *Comm. Pure Appl. Math.*, **50**(1997), no. 12, 1287-1293.
- [8] L. C. Evans : Partial Differential Equations, *Graduate Studies in Mathematics*, 19, (1998).
- [9] C. Foias and R. Temann : Gevrey class regularity for the solutions of the Navier-Stokes equations, *J. Funct. Anal.*, **87**(1989), no. 2, 359-369.
- [10] D. Gerard-Varet and E. Dormy : On the ill-posedness of the Prandtl equation, *J. Amer. Math. Soc.*, **23** (2010), 591-609.

- [11] D. Gerard-Varet and N. Masmoudi : Well-posedness for the Prandtl system without analyticity or monotonicity. *Ann. Sci. Éc. Norm. Supér. (4)***48** (2015), no. 6, 1273–1325.
- [12] Z. Grujić and I. Kukavica : Space analyticity for the Navier-Stokes and related equations with initial data in L^p , *J. Funct. Anal.*, **152**(1998), no. 2, 447-466.
- [13] Y. Guo and T. Nguyen : A note on the Prandtl boundary layers, *Comm. Pure Appl. Math.* **64** (2011), no. 10, 1416-1438.
- [14] G.H. Hardy, J. E. Littlewood and G. Pólya : Inequalities, 2d ed. *Cambridge, at the University Press*, 1952.
- [15] T. Kato : Nonstationary flows of viscous and ideal fluids in \mathbb{R}^3 , *J. Functional Analysis*, **9** (1972), 296–305.
- [16] T. Kato : Remarks on zero viscosity limit for nonstationary Navier-Stokes flows with boundary, Seminar on nonlinear partial differential equations, Berkeley, Calif., 1983, p. 85-98.
- [17] I. Kukavica : On the dissipative scale for the Navier-Stokes equation, *Indiana Univ. Math. J.*, **48**(1999), no. 3, 1057-1081.
- [18] I. Kukavica and V. Vicol : On the radius of analyticity of solutions to the three-dimensional Euler equations, *Proc. Amer. Math. Soc.*, **137** (2009), no. 2, 669–677.
- [19] I. Kukavica and V. Vicol : On the analyticity and Gevrey-class regularity up to the boundary for the Euler equations, *Nonlinearity*, **24** (2011), no. 3, 765-796.
- [20] I. Kukavica and V. Vicol : The domain of analyticity of solutions to the three-dimensional Euler equations in a half space, *Discrete Contin. Dyn. Syst.*, **29**(2011), no. 1, 285-303.
- [21] C. D. Levermore and M. Oliver : Analyticity of solutions for a generalized Euler equation, *J. Differential Equations*, **133** (1997), no. 2, 321–339.
- [22] W.-X. Li, D. Wu and C.-J. Xu : Gevrey Class Smoothing Effect for the Prandtl Equation, *SIAM J. Math. Anal.* **48** (2016), no. 3, 1672–1726.
- [23] W.-X. Li and T. Yang : Well-posedness in Gevrey space for the Prandtl equations with non-degenerate critical points, Preprint, ArXiv : <https://arxiv.org/pdf/1609.08430.pdf>
- [24] J. L. Lions : Quelques méthodes de résolution des problèmes aux limites non linéaires , *Dumod*, Gauthier-Villars, Paris, 1969.
- [25] C.-J. Liu and T. Yang : Ill-posedness of the Prandtl equations in Sobolev spaces around a shear flow with general decay , <https://arxiv.org/pdf/1605.00102.pdf>.
- [26] Y. Maekawa : On the inviscid limit problem of the vorticity equations for viscous incompressible flows in the half-plane, *Comm. Pure Appl. Math.* **67** (2014), no. 7, 1045–1128.
- [27] A. J. Majda and A. L. Bertozzi : *Vorticity and incompressible flow*, Cambridge University Press, 2002.
- [28] O. A. Oleinik and V. N. Samokhin : *Mathematical Models in Boundary Layers Theory*. Chapman & Hall/CRC, 1999.
- [29] R. Temam : *Navier-Stokes equations and Nonlinear Functional Analysis*, Siam, 1983.
- [30] R. Temam : On the Euler equations of incompressible perfect fluids, *J. Functional Analysis*, **20** (1975), no. 1, 32-43.
- [31] R. Temam : *Navier-Stokes Equations: Theory and Numerical Analysis*, AMS, 2001.
- [32] H. Swann : The convergence with vanishing viscosity of nonstationary Navier-Stokes flow to ideal flow in \mathbb{R}^3 , *Trans. Amer. Math. Soc.*, **157** (1971), 373–397.
- [33] M. Sammartino and R.E. Caffisch : Zero viscosity limit for analytic solutions of the Navier-Stokes equations on a half-space, I. Existence for Euler and Prandtl equations. *Comm. Math. Phys.*, 192(1998), no. 2, 433-461; II. Construction of the Navier-Stokes solution. *Comm. Math. Phys.*, 192(1998), 463-491.
- [34] Z. Xin and L. Zhang : On the global existence of solutions to the Prandtl's system, *Adv. Math.*, **181**(2004), no. 1, 88-133.

FENG CHENG, SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY 430072, WUHAN, P.R. CHINA

E-mail address: chengfengwhu@whu.edu.cn

WEI-XI LI, SCHOOL OF MATHEMATICS AND STATISTICS, AND COMPUTATIONAL SCIENCE HUBEI KEY LABORATORY, WUHAN UNIVERSITY 430072, WUHAN, P.R. CHINA

E-mail address: wei-xi.li@whu.edu.cn

CHAO-JIANG XU, SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY 430072,
WUHAN, P.R. CHINA, AND, UNIVERSITÉ DE ROUEN, CNRS UMR 6085, LABORATOIRE DE MATHÉMATIQUES,
76801 SAINT-ETIENNE DU ROUVRAY, FRANCE

E-mail address: `Chao-Jiang.Xu@univ-rouen.fr`