



HAL
open science

Slow to fast infinitely extended reservoirs for the symmetric exclusion process with long jumps

Cedric Bernardin, Patricia Goncalves, Byron Oviedo Jimenez

► To cite this version:

Cedric Bernardin, Patricia Goncalves, Byron Oviedo Jimenez. Slow to fast infinitely extended reservoirs for the symmetric exclusion process with long jumps. 2017. hal-01474718v1

HAL Id: hal-01474718

<https://hal.science/hal-01474718v1>

Preprint submitted on 23 Feb 2017 (v1), last revised 1 Sep 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SLOW TO FAST INFINITELY EXTENDED RESERVOIRS FOR THE SYMMETRIC EXCLUSION PROCESS WITH LONG JUMPS

CÉDRIC BERNARDIN, P. GONÇALVES, AND B. JIMENEZ OVIEDO

ABSTRACT. We consider an exclusion process with long jumps in the box $\Lambda_N = \{1, \dots, N-1\}$, for $N \geq 2$, in contact with infinitely extended reservoirs on its left and on its right. The jump rate is described by a transition probability $p(\cdot)$ which is symmetric, with infinite support but with finite variance. The reservoirs add or remove particles with rate proportional to $\kappa N^{-\theta}$, where $\kappa > 0$ and $\theta \in \mathbb{R}$. If $\theta > 0$ (resp. $\theta < 0$) the reservoirs add and remove fast (resp. slowly) particles in the bulk. According to the value of θ we prove that the time evolution of the spatial density of particles is described by some reaction-diffusion equations with various boundary conditions.

1. INTRODUCTION

The exclusion process is an interacting particle system introduced in the mathematical literature during the seventies by Frank Spitzer [14]. Despite the simplicity of its dynamics it captures the main features of more realistic diffusive systems driven out of equilibrium [12], [13], [15]. It consists in a collection of continuous-time random walks evolving on the lattice \mathbb{Z} whose dynamics can be described as follows. A particle at the site x waits an exponential time after which it jumps to a site $x + y$ with probability $p(y)$. If, however, if $x + y$ is already occupied, the jump is suppressed and the clock is reset.

Recently a series of work have been devoted to the study of the nearest-neighbor exclusion process whose dynamics is perturbed by the presence of a slow bond, a slow site or by slow boundary effects [5, 6, 1]. The behavior of the system is then strongly affected and new boundary conditions may be derived at the macroscopic level. On the other hand it is known that the presence of long jumps, in particular heavy tailed long jumps, have a drastic effect on the macroscopic behavior and critical exponents of the system [2, 7, 8]. In this work, we propose to mix these two interesting features by considering the symmetric exclusion process with long jumps in contact with extended reservoirs. The coupling with the reservoirs is regulated by a certain power θ of a scaling parameter which is the inverse of the size system $N \rightarrow \infty$. This question has been addressed in a recent paper [1] in the case of the nearest-neighbor exclusion process for a positive power θ and with finite reservoirs, in fact one at each end point. Here we consider the case where the jumps probability transition $p(z) \sim |z|^{-1-\gamma}$ has an infinite support and the power θ has an arbitrary sign, so that the boundary effects can be very strong (fast) or very weak (slow). Observe that since we are dealing with system with long jumps the reservoirs we use have to be infinitely extended in order to avoid some truncation procedure to define the dynamics [16]. In this paper we will

Date: February 23, 2017.

Key words and phrases. Hydrodynamic limit, Reaction-diffusion equation, Boundary conditions, Exclusion with long jumps.

focus only on the case $\gamma > 2$, so that $p(\cdot)$ has a finite variance, postponing the study of the case $\gamma \leq 2$ for future works.

The problem we address in this paper is to characterize the hydrodynamic behavior of the process described above, i.e., to deduce the macroscopic behavior of the system from the microscopic interaction among particles and to analyze the effect of slowing down or fasting up the interaction with the reservoirs, by increasing or decreasing the value of θ , at the level of the macroscopic profiles of the density. Usually the characterization of the hydrodynamic limit is formulated in terms of a weak solution of some partial differential equation, called the *hydrodynamic equation*. Depending on the intensity of the coupling with the reservoirs we will observe a phase transition for profiles which are solutions of the hydrodynamic equation which consists on reaction-diffusion equations with different types of boundary conditions, depending on the range of the parameter θ .

We extend the results for the nearest neighbor symmetric simple exclusion process with slow boundaries that was studied in [1] by considering long jumps, infinitely extended reservoirs and also fast reservoirs. In the case $\theta \geq 0$ (slow reservoirs) we recover a similar hydrodynamical behavior for our model since we imposed that the probability transition rate to be symmetric and with finite variance. If one of these conditions is violated then the macroscopic behavior of the system is different. In the case where we drop the hypothesis that $p(\cdot)$ is symmetric, then there is a drift in the microscopic system which appears at the macroscopic level as the heat equation with a transport term and if drop the finite variance condition, then we expect to have the usual laplacian for the case $p(z) \sim |z|^{-1-\gamma}$ with $\gamma = 2$ and a fractional operator when $\gamma \in (1, 2)$. We leave this problem for a future work.

When θ ranges from $-\infty$ to $+\infty$, the model produces five different macroscopic phases¹, depending on the value of the parameter θ . If $\theta \in (2 - \gamma, 1)$, the boundary interactions are not slowed or fasted enough in order to change the macroscopic behavior of the system so that we observe exactly the same behavior as in the case $\theta = 0 \in (2 - \gamma, 1)$. The hydrodynamic equation in this case is the heat equation with Dirichlet boundary conditions. If $\theta = 1$, the reservoirs are slowed enough that we obtain the heat equation but with Robin boundary conditions. For $\theta \in (1, \infty)$, the reservoirs are sufficiently slowed so that we get the heat equation with Neumann boundary conditions. If $\theta = 2 - \gamma$, the reservoirs are fasted enough that we obtain the heat equation with a singular reaction term at the boundaries but with Dirichlet boundary conditions. If $\theta < 2 - \gamma$, the reservoirs are so fasted that the diffusion part of the motion disappears and that only the reaction term survives at the macroscopic level. The two cases $\theta = 1$ and $\theta = 2 - \gamma$ correspond to a critical behavior connecting macroscopically two different regimes (Dirichlet boundary conditions to Neumann boundary conditions for $\theta = 1$ and Reaction to Diffusion equation for $\theta = 2 - \gamma$). Once the form of the hydrodynamic equation is obtained, it is of interest to study its stationary solution which provides the density profile in the stationary state in the thermodynamic limit. In particular for $\theta \leq 2 - \gamma$ the density profiles are non linear and have nice properties (see Figure 3).

¹In [1] only three phases were obtained and only the case $\theta \geq 0$ was studied. The two new phases appearing here are due not only to the fact that we consider also a negative power θ but also on the fact that the system has long jumps. Without long jumps, even with $\theta < 0$, the two new phases would be absent for a nearest-neighbor exclusion process.

The paper is organized as follows. In Section 2.1 we describe precisely the model and we state the main result. In Section 2.2 we present the hydrodynamic equations and we claim the *Hydrodynamic Limit* and the main idea of its proof. In order to give an intuition for getting the different boundary conditions, we present in Section 3 the heuristics for obtaining the weak solutions of the corresponding partial differential equations. This result is rigorously proved in Section 7. We prove tightness in Section 4. In Section 5, we prove the *Replacement Lemma* and some auxiliary results. In Section 6 we establish the energy estimates. We added the Appendix A in which we prove the uniqueness of weak solutions of the hydrodynamics equations and the Appendix B which contains computations involving the generator of the dynamics.

2. STATEMENT OF RESULTS

2.1. The model. For $N \geq 2$ let $\Lambda_N = \{1, \dots, N-1\}$ to which we call the bulk. The exclusion process in contact with reservoirs is a Markov process $\{\eta_t : t \geq 0\}$ with state space $\Omega_N := \{0, 1\}^{\Lambda_N}$. The configurations of the state space Ω_N are denoted by η , so that for $x \in \Lambda_N$, $\eta(x) = 0$ means that the site x is vacant while $\eta(x) = 1$ means that the site x is occupied. Now, we explain the dynamics of this model and we start by describing the conditions on the jump rate. For that purpose, let $p : \mathbb{Z} \rightarrow [0, 1]$ be a translation invariant transition probability which is symmetric, that is, for any $z \in \mathbb{Z}$, $p(z) = p(-z)$ and with finite variance, that is $\sigma^2 := \sum_{z \in \mathbb{Z}} z^2 p(z) < \infty$. Note that since $p(\cdot)$ is symmetric it is mean zero, that is: $\sum_{z \in \mathbb{Z}} z p(z) = 0$. We denote $m = \sum_{z \geq 1} z p(z)$. As an example we consider $p(\cdot)$ given by $p(0) = 0$ and $p(z) = \frac{c_\gamma}{|z|^{\gamma+1}}$, for $z \neq 0$, where c_γ is a normalizing constant and $\gamma > 2$, so that $p(\cdot)$ has finite variance. For simplicity of the presentation we stick to this choice of $p(\cdot)$ along the article, but we note that our results are true in the more general setting where we assume $p(\cdot)$ to be as above.

We consider the process in contact with infinitely many stochastic reservoirs at all the negative integer sites and at all the integer sites $z \geq N$. We fix four parameters $\alpha, \beta \in (0, 1)$, $\kappa > 0$ and $\theta \in \mathbb{R}$, so that particles can get into (resp. exit) the bulk of the system from any boundary site at rate $\alpha\kappa/N^\theta p(z)$ (resp. $(1-\alpha)\kappa/N^\theta p(z)$), where z is the jump size (see Figure 1).

The stochastic reservoir at the right acts in the same way as the left reservoir but in the intensity we replace α by β . The dynamics of the process is defined as follows. We start with the bulk dynamics. Each pair of sites of the bulk $\{x, y\} \subset \Lambda_N$ carries a Poisson process of intensity one. The Poisson processes associated to different bonds are independent. If for the configuration η , the clock associated to the bound $\{x, y\}$ rings, then we exchange the values η_x and η_y with rate $p(y-x)/2$. Now we explain the dynamics at the boundary. Each pair of sites $\{x, y\}$ with $x \in \mathbb{Z}_-$ and $y \in \Lambda_N$ carries a Poisson process of intensity one all being independent. If for the configuration η , the clock associated to the bound $\{x, y\}$ rings, then we change the values η_y into $1 - \eta_y$ with rate $\frac{\kappa}{N^\theta} p(y-x)[(1-\alpha)\eta_y + \alpha(1-\eta_y)]$. At the right boundary the dynamics is similar but instead of α the intensity is given by β . Observe that the reservoirs add and remove particles on all the sites of the bulk Λ_N , and not only at the boundaries, but with rates which decrease as the distance from the corresponding reservoir increases. We can interpret the dynamics of the reservoirs in two different ways as follows. In the first case, we add to the bulk infinitely many reservoirs at all negative sites and at all sites $y \geq N$. Then particles can get into (resp. get out from) the bulk from the left reservoir at rate $\alpha\kappa/N^\theta p(z)$ (resp. $(1-\alpha)\kappa/N^\theta p(z)$) where z is the size of the jump.




FIGURE 1. Exclusion process with long jumps and infinitely extended reservoirs.

The right reservoir acts in the same way, except that we replace α by β in the jump rates given above. In the second case we can consider that particles can be created (resp. annihilated) at all the sites x in the bulk with one of the rates $r_N^-(x/N)\alpha\kappa/n^\theta$ or $r_N^+(x/N)\beta\kappa/n^\theta$ (resp. $r_N^-(x/N)(1-\alpha)\kappa/n^\theta$ or $r_N^+(x/N)(1-\beta)\kappa/n^\theta$) where r_N^\pm are given in (3.3).

The infinitesimal generator of the process is given by

$$L_N = L_N^0 + L_N^r + L_N^\ell, \quad (2.1)$$

where its action on functions $f : \Omega_N \rightarrow \mathbb{R}$ is

$$\begin{aligned} (L_N^0 f)(\eta) &= \frac{1}{2} \sum_{x,y \in \Lambda_N} p(x-y) [f(\sigma^{x,y} \eta) - f(\eta)], \\ (L_N^\ell f)(\eta) &= \frac{\kappa}{N^\theta} \sum_{\substack{x \in \Lambda_N \\ y \leq 0}} p(x-y) c_x(\eta; \alpha) [f(\sigma^x \eta) - f(\eta)], \\ (L_N^r f)(\eta) &= \frac{\kappa}{N^\theta} \sum_{\substack{x \in \Lambda_N \\ y \geq N}} p(x-y) c_x(\eta; \beta) [f(\sigma^x \eta) - f(\eta)], \end{aligned} \quad (2.2)$$

and

$$(\sigma^{x,y} \eta)_z = \begin{cases} \eta_z, & z \neq x, y, \\ \eta_y, & z = x, \\ \eta_x, & z = y \end{cases}, \quad (\sigma^x \eta)_z = \begin{cases} \eta_z, & z \neq x, \\ 1 - \eta_x, & z = x. \end{cases} \quad (2.3)$$

Above, for a function $\varphi : [0, 1] \rightarrow \mathbb{R}$, we used the notation

$$c_x(\eta; \varphi(\cdot)) := [\eta_x (1 - \varphi(\frac{x}{N})) + (1 - \eta_x) \varphi(\frac{x}{N})]. \quad (2.4)$$

We consider the Markov process speeded up in the time scale $\Theta(N)$ and we use the notation $\eta^N(t) := \eta(t\Theta(N))$, so that $(\eta^N(t))_{t \geq 0}$ has infinitesimal generator $\Theta(N)L_N$. Although $\eta^N(t)$ depends on α , β and θ , we shall omit these index in order to simplify notation.

2.2. Hydrodynamic equations. From now on up to the rest of this article we fix a finite time horizon $[0, T]$. To properly state the hydrodynamic limit, we need to introduce some notations and definitions. We denote by $\langle \cdot, \cdot \rangle_\mu$ (resp. $\| \cdot \|_{L^2(\mu)}$) the inner product (resp. the norm) in $L^2([0, 1])$ with respect to the measure μ defined in $[0, 1]$ and

when μ is the Lebesgue measure we simply write $\langle \cdot, \cdot \rangle$ and $\|\cdot\|_{L^2}$ for the corresponding norm. For an interval \mathcal{I} in \mathbb{R} and integers m and n , we denote by $C^{m,n}([0, T] \times \mathcal{I})$ the set of functions defined on $[0, T] \times \mathcal{I}$ that are m times differentiable on the first variable and n times differentiable on the second variable. An index on a function will always denote a fixed variable, not a derivative. For example, $G_s(q)$ means $G(s, q)$. The derivative of $G \in C^{m,n}([0, T] \times \mathcal{I})$ will be denoted by $\partial_s G$ (first variable) and $\partial_q G$ (second variable). We shall write ΔG for $\partial_q^2 G$. We also consider the set $C_c^{m,n}([0, T] \times [0, 1])$ of functions $G \in C^{m,n}([0, T] \times [0, 1])$ such that G_s has a compact support included in $(0, 1)$ for any time s and, we denote by $C_c^m(0, 1)$ (resp. $C_c^\infty(0, 1)$) the set of all m continuously differentiable (resp. smooth) real-valued functions defined on $(0, 1)$ with compact support. The set $C^\infty([0, 1])$ denotes the set of restrictions of smooth functions on \mathbb{R} to the interval $[0, 1]$. The supremum norm is denoted by $\|\cdot\|_\infty$.

The semi inner-product $\langle \cdot, \cdot \rangle_1$ is defined on the set $C^\infty([0, 1])$ by

$$\langle G, H \rangle_1 = \int_0^1 (\partial_q G)(q) (\partial_q H)(q) dq. \quad (2.5)$$

The corresponding semi-norm is denoted by $\|\cdot\|_1$.

Definition 2.1. *The Sobolev space \mathcal{H}^1 on $[0, 1]$ is the Hilbert space defined as the completion of $C^\infty([0, 1])$ for the norm*

$$\|\cdot\|_{\mathcal{H}^1}^2 := \|\cdot\|_{L^2}^2 + \|\cdot\|_1^2.$$

Its elements coincide a.e. with continuous functions. The completion of $C_c^\infty(0, 1)$ for this norm is denoted by \mathcal{H}_0^1 . This is a Hilbert space whose elements coincide a.e. with continuous functions vanishing at 0 and 1. On \mathcal{H}_0^1 , the two norms $\|\cdot\|_{\mathcal{H}^1}$ and $\|\cdot\|_1$ are equivalent. The space $L^2(0, T; \mathcal{H}^1)$ is the set of measurable functions $f : [0, T] \rightarrow \mathcal{H}^1$ such that

$$\int_0^T \|f_s\|_{\mathcal{H}^1}^2 ds < \infty.$$

The space $L^2(0, T; \mathcal{H}_0^1)$ is defined similarly.

We can now give the definition of the weak solutions of the hydrodynamic equations that will be derived in this paper.

Definition 2.2. *Let $\hat{\sigma} \geq 0$ and $\hat{\kappa} \geq 0$ be some parameters. Let $g(\cdot) : [0, 1] \rightarrow [0, 1]$ be a measurable function. We say that $\rho : [0, T] \times [0, 1] \rightarrow [0, 1]$ is a weak solution of the reaction-diffusion equation with inhomogeneous Dirichlet boundary conditions*

$$\begin{cases} \partial_t \rho_t(q) = \frac{\hat{\sigma}^2}{2} \Delta \rho_t(q) + \hat{\kappa} \left\{ \frac{\alpha - \rho_t(q)}{q^\gamma} + \frac{\beta - \rho_t(q)}{(1-q)^\gamma} \right\}, & (t, q) \in [0, T] \times (0, 1), \\ \rho_t(0) = \alpha, \quad \rho_t(1) = \beta, & t \in [0, T], \\ \rho_0(\cdot) = g(\cdot), \end{cases} \quad (2.6)$$

if the following three conditions hold:

1. $\rho \in L^2(0, T; \mathcal{H}^1)$ if $\hat{\sigma} > 0$ and $\int_0^T \int_0^1 \left\{ \frac{(\alpha - \rho_t(q))^2}{q^\gamma} + \frac{(\beta - \rho_t(q))^2}{(1-q)^\gamma} \right\} dq dt < \infty$ if $\hat{\kappa} > 0$,

2. ρ satisfies the weak formulation:

$$\begin{aligned} F_{RD}(t, \rho, G, g) &:= \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 g(q) G_0(q) dq \\ &- \int_0^t \int_0^1 \rho_s(q) \left(\frac{\hat{\sigma}^2}{2} \Delta + \partial_s \right) G_s(q) ds dq \\ &- \hat{\kappa} \int_0^t \int_0^1 G_s(q) \left(\frac{\alpha - \rho_s(q)}{q^\gamma} + \frac{\beta - \rho_s(q)}{(1-q)^\gamma} \right) ds dq = 0, \end{aligned} \quad (2.7)$$

for all $t \in [0, T]$ and any function $G \in C_c^{1,2}([0, T] \times [0, 1])$,

3. if $\hat{\sigma} > 0$ then $\rho_t(0) = \alpha$, $\rho_t(1) = \beta$ for all $t \in [0, T]$.

Remark 2.3. Observe that in the case $\hat{\sigma} > 0$ and $\hat{\kappa} = 0$ we recover the heat equation with Dirichlet inhomogeneous boundary conditions. If $\hat{\sigma} = 0$ the equation does not have a diffusion part.

Definition 2.4. Let $\hat{\sigma} > 0$ and $\hat{m} \geq 0$ be some parameters. Let $g : [0, 1] \rightarrow [0, 1]$ be a measurable function. We say that $\rho : [0, T] \times [0, 1] \rightarrow [0, 1]$ is a weak solution of the heat equation with Robin boundary conditions

$$\begin{cases} \partial_t \rho_t(q) = \frac{\hat{\sigma}^2}{2} \Delta \rho_t(q), & (t, q) \in [0, T] \times (0, 1), \\ \partial_q \rho_t(0) = \frac{2\hat{m}}{\hat{\sigma}^2} (\rho_t(0) - \alpha), & \partial_q \rho_t(1) = \frac{2\hat{m}}{\hat{\sigma}^2} (\beta - \rho_t(1)), & t \in [0, T] \\ \rho_0(\cdot) = g(\cdot), \end{cases} \quad (2.8)$$

if the following three conditions hold:

1. $\rho \in L^2(0, T; \mathcal{H}^1)$,
2. ρ satisfies the weak formulation:

$$\begin{aligned} F_{Rob}(t, \rho, G, g) &:= \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 g(q) G_0(q) dq \\ &- \int_0^t \int_0^1 \rho_s(q) \left(\frac{\hat{\sigma}^2}{2} \Delta + \partial_s \right) G_s(q) ds dq + \frac{\hat{\sigma}^2}{2} \int_0^t \{ \rho_s(1) \partial_q G_s(1) - \rho_s(0) \partial_q G_s(0) \} ds \\ &- \hat{m} \int_0^t \{ G_s(0) (\alpha - \rho_s(0)) + G_s(1) (\beta - \rho_s(1)) \} ds = 0, \end{aligned} \quad (2.9)$$

for all $t \in [0, T]$, any function $G \in C^{1,2}([0, T] \times [0, 1])$.

Remark 2.5. Observe that in the case $\hat{m} = 0$ the PDE above is the heat equation with Neumann boundary conditions.

2.3. Hydrodynamic Limit. Let \mathcal{M}^+ be the space of positive measures on $[0, 1]$ with total mass bounded by 1 equipped with the weak topology. For any configuration $\eta \in \Omega_N$ we define the empirical measure $\pi^N(\eta, dq)$ on $[0, 1]$ by

$$\pi^N(\eta, dq) = \frac{1}{N-1} \sum_{x \in \Lambda_N} \eta_x \delta_{\frac{x}{N}}(dq), \quad (2.10)$$

where δ_a is a Dirac mass on $a \in [0, 1]$, and

$$\pi_t^N(\eta, dq) := \pi^N(\eta^N(t), dq).$$

Fix $T > 0$ and $\theta \in \mathbb{R}$. We denote by \mathbb{P}_{μ_N} the probability measure in the Skorohod space $\mathcal{D}([0, T], \Omega_N)$ induced by the Markov process $(\eta^N(t))_{t \geq 0}$ and the initial probability measure μ_N and we denote by \mathbb{E}_{μ_N} the expectation with respect to \mathbb{P}_{μ_N} . Let $\{\mathbb{Q}_N\}_{N \geq 1}$ be the sequence of probability measures on $\mathcal{D}([0, T], \mathcal{M}^+)$ induced by the Markov process $\{\pi_t^N\}_{t \geq 0}$ and by \mathbb{P}_{μ_N} .

Let $\rho_0 : [0, 1] \rightarrow [0, 1]$ be a measurable function. We say that a sequence of probability measures $\{\mu_N\}_{N \geq 1}$ in Ω_N is associated to the profile $\rho_0(\cdot)$ if for any continuous function $G : [0, 1] \rightarrow \mathbb{R}$ and every $\delta > 0$

$$\lim_{N \rightarrow \infty} \mu_N \left(\eta \in \Omega_N : \left| \frac{1}{N} \sum_{x \in \Lambda_N} G\left(\frac{x}{N}\right) \eta_x - \int_0^1 G(q) \rho_0(q) dq \right| > \delta \right) = 0. \quad (2.11)$$

The main result of this article is summarized in the following theorem (see Figure 2).

Theorem 2.6. *Let $g : [0, 1] \rightarrow [0, 1]$ be a measurable function and let $\{\mu_N\}_{N \geq 1}$ be a sequence of probability measures in Ω_N associated to $g(\cdot)$. Then, for any $0 \leq t \leq T$,*

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left(\eta^N \in \mathcal{D}([0, T], \Omega_N) : \left| \frac{1}{N} \sum_{x \in \Lambda_N} G\left(\frac{x}{N}\right) \eta_x^N(t) - \int_0^1 G(q) \rho_t(q) dq \right| > \delta \right) = 0,$$

where the time scale is given by

$$\Theta(N) = \begin{cases} N^2, & \text{if } \theta \geq 2 - \gamma, \\ N^{\gamma + \theta}, & \text{if } \theta < 2 - \gamma, \end{cases} \quad (2.12)$$

and $\rho_t(\cdot)$ is the unique weak solution of :

- (2.6) with $\hat{\sigma} = 0$ and $\hat{\kappa} = \kappa c_\gamma \gamma^{-1}$, if $\theta < 2 - \gamma$;
- (2.6) with $\hat{\sigma} = \sigma$ and $\hat{\kappa} = \kappa c_\gamma \gamma^{-1}$, if $\theta = 2 - \gamma$;
- (2.6) with $\hat{\sigma} = \sigma$ and $\hat{\kappa} = 0$, if $\theta \in (2 - \gamma, 1)$;
- (2.8) with $\hat{\sigma} = \sigma$ and $\hat{m} = m\kappa$, if $\theta = 1$;
- (2.8) with $\hat{\sigma} = \sigma$ and $\hat{m} = 0$, if $\theta \in (1, \infty)$.

It is not always possible to write fully explicit expressions for the solutions of these hydrodynamic equations. The form of the corresponding stationary solutions is of interest since the latter are expected to describe, in general, the mean density profile in the non-equilibrium stationary state of the microscopic system in the thermodynamic limit $N \rightarrow \infty$. Observe that this is not a trivial fact since it requires to exchange the limits $t \rightarrow \infty$ with $N \rightarrow \infty$ (and for $\theta > 1$ this is for example false, see below).

The stationary solutions of the hydrodynamic limits in the $\theta > 2 - \gamma$ case are standard. On the other hand, the form and properties of the stationary solutions in the $\theta \leq 2 - \gamma$ case are original and more tricky to obtain in the $\theta = 2 - \gamma$ case. This problem is studied in more details in [9]. Here we only present some graphs of the stationary solutions and refer the interested reader to [9] for a complete mathematical treatment.

For $\theta \in (2 - \gamma, 1)$ (heat equation with Dirichlet boundary conditions) the stationary solution is the linear profile connecting α at 0 to β at 1. For $\theta = 1$ (heat equation with Robin boundary conditions) the profile is still linear but the values at the boundaries are different. Observe that if $\kappa \rightarrow 0$ these values converge to $\frac{\alpha + \beta}{2}$ so that the profile becomes flat equal to $\frac{\alpha + \beta}{2}$. For $\theta > 1$ (heat equation with Neumann boundary conditions) the stationary solution is constant equal to $\int_0^1 g(q) dq$ where $g(\cdot)$ is the initial condition.




FIGURE 2. The five different hydrodynamic regimes in terms of γ and θ .

In fact, for $\theta > 1$, we expect that if we compute directly the stationary profile in the non-equilibrium stationary state of the microscopic system in the thermodynamic limit, the stationary profile will be flat with the value $(\alpha + \beta)/2$. This value is therefore memorized in the form of the hydrodynamic limits for $\theta = 1$, despite the fact that it has been forgotten in the hydrodynamic limits for $\theta > 1$. In the case $\theta < 2 - \gamma$ (reaction-diffusion equation without diffusive part) the stationary profile is fully explicit and given by $\frac{V_0(q)}{V_1(q)}$ where

$$V_0(q) = \alpha q^{-\gamma} + \beta(1-q)^{-\gamma}, \quad V_1(q) = q^{-\gamma} + (1-q)^{-\gamma}. \quad (2.13)$$

Observe that this profile is increasing, non-linear, convex on $(0, 1/2)$ and concave on $(1/2, 1)$ and connects α at 0 to β at 1. At the boundaries the profile is very flat. In [9] it is proved that these properties remain valid for the stationary solution of the hydrodynamic equation in the $\theta = 2 - \gamma$ case.

The proof of Theorem 2.6 follows the usual approach of convergence in distribution of stochastic processes. First we prove tightness of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$ and then we prove uniqueness of the limiting point \mathbb{Q} . These two results combined give the convergence of $\{\mathbb{Q}_N\}_{N \geq 1}$ to \mathbb{Q} as $N \rightarrow \infty$. In order to characterize the limiting point \mathbb{Q} , we prove that all limiting points of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$ are concentrated on trajectories of measures that are absolutely continuous with respect to the Lebesgue measure and whose density $\rho_t(q)$ is a weak solution of the corresponding hydrodynamic equation




FIGURE 3. Profiles of the stationary solution of the hydrodynamic equations according to the value of θ .

depending on the value of θ . From the uniqueness of the weak solutions of these equations, which is proved in Appendix A, we conclude that $\{\mathbb{Q}_N\}_{N \geq 1}$ has a unique limiting point \mathbb{Q} .

Notations: We write $f(x) \lesssim g(x)$ if there exists a constant C independent of x such that $f(x) \leq Cg(x)$ for every x . We will also write $f(x) = \mathcal{O}(g(x))$ if the condition $|f(x)| \lesssim |g(x)|$ is satisfied. Sometimes, in order to stress the dependence of a constant C on some parameter a , we write $C(a)$.

3. HEURISTICS FOR THE HYDRODYNAMIC EQUATIONS

In this section we give the main ideas which are behind the identification of limit points as weak solutions of the partial differential equations given in Section 2.2. In Section 4, we show that the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$ is tight and in Appendix 7 we prove that all limiting points of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$ are concentrated on trajectories of measures that are absolutely continuous with respect to the Lebesgue measure, that is $\pi_t(dq) = \rho_t(q)dq$. Now we argue that the density $\rho_t(q)$ is a weak solution of the corresponding hydrodynamic equation for each regime of θ . The precise proof of this result is given ahead in Proposition 7.1.

The identification of the density $\rho_t(q)$ as a weak solution of the hydrodynamic equation is obtained by using auxiliary martingales. For that purpose, and to make the exposition simpler, we fix a function $G : [0, 1] \rightarrow \mathbb{R}$ which does not depend on time and is two times continuously differentiable. If $\theta < 1$ we will assume further that it has a compact support included in $(0, 1)$ and for $\theta \geq 1$ we assume that it has a compact support strictly contained in $[0, 1]$ so that G has a good decay at infinity. In the last

case G can take non-zero values at 0 and 1. We know by Dynkin's formula that

$$M_t^N(G) = \langle \pi_t^N, G \rangle - \langle \pi_0^N, G \rangle - \int_0^t \Theta(N) L_N \langle \pi_s^N, G \rangle ds, \quad (3.1)$$

is a martingale with respect to the natural filtration $\{\mathcal{F}_t\}_{t \geq 0}$, where for each $t \geq 0$, $\mathcal{F}_t := \sigma(\eta(s) : s < t)$. Above the notation $\langle \pi_s^N, G \rangle$ represents the integral of G with respect the measure π_s^N . This notation should not be mistaken with the notation used for the inner product in $L^2([0, 1])$. A simple computation, based on (4.9) and the discussion after this equation, shows that the $\mathbb{E}_{\mu_N}[(M_t^N(G))^2]$ vanishes as $N \rightarrow \infty$. Now we look at the integral term in (3.1). A simple computation shows that

$$\begin{aligned} \int_0^t \Theta(N) L_N \langle \pi_s^N, G \rangle ds &= \frac{\Theta(N)}{N-1} \int_0^t \sum_{x \in \Lambda_N} \mathcal{L}_N G(\frac{x}{N}) \eta_x^N(s) ds \\ &+ \frac{\kappa \Theta(N)}{(N-1)N^\theta} \int_0^t \sum_{x \in \Lambda_N} (Gr_N^-)(\frac{x}{N})(\alpha - \eta_x^N(s)) ds \\ &+ \frac{\kappa \Theta(N)}{(N-1)N^\theta} \int_0^t \sum_{x \in \Lambda_N} (Gr_N^+)(\frac{x}{N})(\beta - \eta_x^N(s)) ds, \end{aligned} \quad (3.2)$$

where for all $x \in \Lambda_N$

$$\begin{aligned} (\mathcal{L}_N G)(\frac{x}{N}) &= \sum_{y \in \Lambda_N} p(y-x) [G(\frac{y}{N}) - G(\frac{x}{N})], \\ r_N^-(\frac{x}{N}) &= \sum_{y \geq x} p(y), \quad r_N^+(\frac{x}{N}) = \sum_{y \leq x-N} p(y). \end{aligned} \quad (3.3)$$

Now, we want to extend the first sum in (3.2) to all the integers. For that purpose we extend the function G to \mathbb{R} in such a way that it remains two times continuously differentiable. By the definition of \mathcal{L}_N , we get that

$$\begin{aligned} \frac{\Theta(N)}{N-1} \int_0^t \sum_{x \in \Lambda_N} \mathcal{L}_N G(\frac{x}{N}) \eta_x^N(s) ds &= \frac{\Theta(N)}{N-1} \int_0^t \sum_{x \in \Lambda_N} (K_N G)(\frac{x}{N}) \eta_x^N(s) ds \\ &- \frac{\Theta(N)}{N-1} \int_0^t \sum_{x \in \Lambda_N} \sum_{y \leq 0} [G(\frac{y}{N}) - G(\frac{x}{N})] p(x-y) \eta_x^N(s) ds \\ &- \frac{\Theta(N)}{N-1} \int_0^t \sum_{x \in \Lambda_N} \sum_{y \geq N} [G(\frac{y}{N}) - G(\frac{x}{N})] p(x-y) \eta_x^N(s) ds, \end{aligned} \quad (3.4)$$

where

$$(K_N G)(\frac{x}{N}) = \sum_{y \in \mathbb{Z}} p(y-x) [G(\frac{y}{N}) - G(\frac{x}{N})]. \quad (3.5)$$

Now, we are going to analyze all the terms in (3.4) and the boundary terms in (3.2) for the different regimes of θ . Thus, we will be able to see how the different boundary conditions appear on the hydrodynamic equations given in Section 2.2 from the underlying particle system.

Let us first observe that, for any $a \in (0, 1)$, uniformly in $u \in (a, 1-a)$, we have that

$$N^\gamma r_N^-([uN]) \rightarrow c_\gamma \gamma^{-1} u^{-\gamma} := r^-(u), \quad N^\gamma r_N^+([uN]) \rightarrow c_\gamma \gamma^{-1} (1-u)^{-\gamma} := r^+(u) \quad (3.6)$$

as $N \rightarrow \infty$.

3.1. The case $\theta < 2 - \gamma$. In this regime we take initially a function $G : (0, 1) \rightarrow \mathbb{R}$ two times continuously differentiable and with compact support in $(0, 1)$ (so that we can choose an extension by 0 outside of the support of G).

Now we start by analyzing the first term in (3.4). Since $\Theta(N) = N^{\gamma+\theta}$, a simple computation, shows that the first term in (3.4) vanishes for $\theta < 2 - \gamma$. Indeed, by a Taylor expansion on G and the fact that $p(\cdot)$ is mean zero, we have that

$$N^{\gamma+\theta} \sum_{y \in \mathbb{Z}} (G(\frac{y+x}{N}) - G(\frac{x}{N})) p(y)$$

is of same order as

$$N^{\gamma+\theta-2} G''(\frac{x}{N}) \sum_{y \in \mathbb{Z}} y^2 p(y)$$

and since $\theta < 2 - \gamma$ last expression vanishes as $N \rightarrow \infty$.

Moreover, a simple computation shows that the second and third terms in (3.4) vanish as $N \rightarrow \infty$, since $\Theta(N) = N^{\gamma+\theta}$ and $\theta < 2 - \gamma$. Indeed we can bound from above, for example the second term in (3.4) by tN^θ times

$$\frac{1}{N-1} \sum_{x \in \Lambda_N} N^\gamma r_N^-(\frac{x}{N}) |G(\frac{x}{N})|$$

because G vanishes outside $(0, 1)$ and $|\eta_x^N(s)| \leq 1$ for all $s > 0$. Since $\theta < 0$ and that the previous sum converges to the (finite) integral of $|G|r^-$ on $(0, 1)$, by (3.6), the previous display vanishes as $N \rightarrow \infty$. Now we look at the boundary terms in (3.2). The second term in (3.2) can be written, for the choice of $\Theta(N) = N^{\gamma+\theta}$, as:

$$\frac{\kappa N^\gamma}{N-1} \int_0^t \sum_{x \in \Lambda_N} G(\frac{x}{N}) r_N^-(\frac{x}{N}) (\alpha - \eta_x^N(s)) ds$$

which can be replaced, thanks to (3.6) and the fact that G has compact support, by

$$\kappa \int_0^t \langle \pi_s^N, Gr^- \rangle ds \rightarrow \kappa \int_0^t \int_0^1 G(q) r^-(q) \rho_s(q) dq ds$$

as $N \rightarrow \infty$. The last convergence holds because G has a compact support included in $(0, 1)$ so that Gr^- is a continuous function. For the remaining term we can perform exactly the same analysis.

3.2. The case $\theta = 2 - \gamma$. In this case, and as above, we take initially a function $G : (0, 1) \rightarrow \mathbb{R}$ two times continuously differentiable and with compact support in $(0, 1)$ (so that we can choose an extension by 0 outside of its support). In this case, since $\Theta(N) = N^2$, by Lemma 3.2, which we prove below, the first term in (3.4) can be replaced, for N sufficiently big, by

$$\frac{1}{N-1} \int_0^t \sum_{x \in \Lambda_N} \frac{\sigma^2}{2} \Delta G(\frac{x}{N}) \eta_x^N(s) ds.$$

Moreover, a computation similar to the one above shows that the second and third terms in (3.4) vanish as $N \rightarrow \infty$ (recall that $\Theta(N) = N^2$ and $\gamma > 2$). Finally, the first term in (3.2) can be rewritten as

$$\frac{\kappa N^\gamma}{(N-1)} \int_0^t \sum_{x \in \Lambda_N} (Gr_N^-)(\frac{x}{N}) (\alpha - \eta_x^N(s)) ds$$

which can be replaced, thanks to (3.6) and the fact that G has compact support, by

$$\kappa \int_0^t \langle \pi_s^N, Gr^- \rangle ds \rightarrow \kappa \int_0^t \int_0^1 G(q)r^-(q)\rho_s(q)dq ds$$

as $N \rightarrow \infty$ because Gr^- is a continuous function. The same computation can be done for the remaining term.

3.3. The case $\theta \in (2 - \gamma, 1)$. In this case we take again a function $G : (0, 1) \rightarrow \mathbb{R}$ two times continuously differentiable and with compact support in $(0, 1)$ and extend it by 0 outside of $(0, 1)$. Here the situation is different from the previous ones since the last two terms in (3.2) vanish as $N \rightarrow \infty$, as a consequence of Lemma 5.6. But, as before, the second and third terms in (3.4) also vanish because, for example, the second term in (3.4)

$$\frac{N^2}{N-1} \int_0^t \sum_{x \in \Lambda_N} G\left(\frac{x}{N}\right) r_N^-\left(\frac{x}{N}\right) \eta_x^N(s) ds$$

can be bounded from above by a constant times $tN^{2-\gamma}$ times a sum converging to the integral of $|G|r^-$ on $(0, 1)$. The estimate of the third term is analogous. Therefore since $\gamma > 2$, both vanish as $N \rightarrow \infty$.

Remark 3.1. *Observe that in the three previous cases, we imposed to G to have a compact support included in $(0, 1)$. This was used in order to extend smoothly the function G by 0 outside of $(0, 1)$ (the condition $G(0) = G(1) = 0$ would not have been sufficient) and this was fundamental to ensure that the functions Gr^- , Gr^+ do not have singularities at the boundaries. On the other hand, in the two next cases, it will be fundamental to consider test functions $G : [0, 1] \rightarrow \mathbb{R}$ which are not necessarily 0 at the boundaries in order to “see” the boundaries in the weak formulation.*

3.4. The case $\theta = 1$. In this case we consider an arbitrary function $G : [0, 1] \rightarrow \mathbb{R}$ which is two times continuously differentiable and we extend it on \mathbb{R} in a two times continuously differentiable function with compact support. Its support strictly contains $[0, 1]$ since G can take non-zero values at 0 and 1. We start by looking at the terms coming from the boundary, namely the two last terms in (3.2). Then, in the second term of (3.2) (resp. the third term) we do at first a Taylor expansion on G and then we replace η_x by the average $\overrightarrow{\eta}_0^{\varepsilon N} = \frac{1}{\varepsilon N} \sum_{x=1}^{\varepsilon N} \eta_x$ (resp. η_x by $\overleftarrow{\eta}_N^{\varepsilon N} = \frac{1}{\varepsilon N} \sum_{x=N-1-\varepsilon N}^{N-1} \eta_x$), which can be done as a consequence of Lemma 5.7 as pointed out in Remark 5.8. Moreover, note that

$$\sum_{x \in \Lambda_N} r_N^-\left(\frac{x}{N}\right) \xrightarrow{N \uparrow \infty} \sum_{y \geq 1} yp(y) := m, \quad \sum_{x \in \Lambda_N} r_N^+\left(\frac{x}{N}\right) \xrightarrow{N \uparrow \infty} \sum_{y \geq 1} yp(y) := m. \quad (3.7)$$

Therefore, we can write the last two terms in (3.2) as

$$m\kappa \int_0^t \{(\alpha - \overleftarrow{\eta}_0^{\varepsilon N}(sN^2))G(0) + (\beta - \overrightarrow{\eta}_N^{\varepsilon N}(sN^2))G(1)\} ds,$$

plus lower-orders terms (with respect to N). Since (in some sense that we will see in the proof of Proposition 7.1 in Section 7),

$$\overrightarrow{\eta}_0^{\varepsilon N}(sN^2) \xrightarrow{N \uparrow \infty} \rho_s(0), \quad \overleftarrow{\eta}_N^{\varepsilon N}(sN^2) \xrightarrow{N \uparrow \infty} \rho_s(1)$$

last term writes as

$$m\kappa \int_0^t \{(\alpha - \rho_s(0))G(0) + (\beta - \rho_s(1))G(1)\} ds. \quad (3.8)$$

Now we look at the remaining terms, namely, the two last terms in (3.4). Recall that the function G has been extended into a two times continuously differentiable function on \mathbb{R} . By a Taylor expansion on G we can write those terms as

$$\frac{N}{N-1} \sum_{x \in \Lambda_N} G'(\frac{x}{N}) \Theta_x^- \eta_x(sN^2) - \frac{N}{N-1} \sum_{x \in \Lambda_N} G'(\frac{x}{N}) \Theta_x^+ \eta_x(sN^2) \quad (3.9)$$

plus lower-order terms (with respect to N). Above for $x \in \Lambda_N$,

$$\Theta_x^- = \sum_{y \leq 0} (x-y)p(x-y) \quad \text{and} \quad \Theta_x^+ = \sum_{y \geq N} (y-x)p(x-y).$$

Note that

$$\sum_{x \in \Lambda_N} \Theta_x^- \lesssim 1 \quad \text{and} \quad \frac{1}{N} \sum_{x \in \Lambda_N} x \Theta_x^- \xrightarrow{N \rightarrow \infty} 0. \quad (3.10)$$

Moreover, note that

$$\begin{aligned} \sum_{x \in \Lambda_N} \Theta_x^- &= \sum_{x \in \Lambda_N} \sum_{y \geq x} yp(y) \xrightarrow{N \uparrow \infty} \frac{\sigma^2}{2}, \\ \sum_{x \in \Lambda_N} \Theta_x^+ &= \sum_{x \in \Lambda_N} \sum_{y \geq N-x} yp(y) \xrightarrow{N \uparrow \infty} \frac{\sigma^2}{2}. \end{aligned} \quad (3.11)$$

In order to prove the convergence of $\sum_{x \in \Lambda_N} \Theta_x^-$ (or of $\sum_{x \in \Lambda_N} \Theta_x^+$ in (3.11)) we use Fubini's theorem to get that

$$\begin{aligned} \sum_{x \in \Lambda_N} \Theta_x^- &= \sum_{y \in \Lambda_N} \sum_{x=1}^y yp(y) + \sum_{y \geq N} \sum_{x \in \Lambda_N} yp(y) \\ &= \sum_{y \in \Lambda_N} y^2 p(y) + (N-1) \sum_{y \geq N} yp(y), \end{aligned}$$

and since $\gamma > 2$ the result follows. By another Taylor expansion on G we can write (3.9) as

$$\frac{N}{N-1} G'(0) \sum_{x \in \Lambda_N} \Theta_x^- \eta_x(sN^2) - \frac{N}{N-1} G'(1) \sum_{x \in \Lambda_N} \Theta_x^+ \eta_x(sN^2) \quad (3.12)$$

plus lower-order terms (with respect to N). Thanks to Lemma 5.7 we can replace in the term on the left (resp. right) hand side of last expression $\eta_x(sN^2)$ by $\vec{\eta}_0^{\varepsilon N}(sN^2)$ (resp. $\overleftarrow{\eta}_N^{\varepsilon N}(sN^2)$). Therefore, (3.12) can be replaced, for N sufficiently big and then ε sufficiently small, by

$$G'(0) \frac{\sigma^2}{2} \vec{\eta}_0^{\varepsilon N}(sN^2) - G'(1) \frac{\sigma^2}{2} \overleftarrow{\eta}_N^{\varepsilon N}(sN^2).$$

Since (in some sense that we will see in the proof of Proposition 7.1 in Section 7), we have that $\vec{\eta}_0^{\varepsilon N}(sN^2) \xrightarrow{N \rightarrow \infty} \rho_s(0)$ and $\overleftarrow{\eta}_N^{\varepsilon N}(sN^2) \xrightarrow{N \rightarrow \infty} \rho_s(1)$, last term tends to

$$G'(0) \frac{\sigma^2}{2} \rho_s(0) - G'(1) \frac{\sigma^2}{2} \rho_s(1). \quad (3.13)$$

Putting together (3.8) and (3.13) we see the boundary terms that appear at the right hand side of (2.9).

3.5. **The case $\theta \in (1, \infty)$.** In this case we consider an arbitrary function $G : [0, 1] \rightarrow \mathbb{R}$ which is two times continuously differentiable and we extend it on \mathbb{R} in a two times continuously differentiable function with compact support. Its support strictly contains $[0, 1]$ since G can take non-zero values at 0 and 1. The last two terms in (3.2) vanish, as $N \rightarrow \infty$ since, we can bound, for example, the first term in (3.2) by a constant times

$$N^{1-\theta} \sum_{x \in \Lambda_N} r_N^-(\frac{x}{N}).$$

Since $\gamma > 2$ last expression vanishes if $\theta > 1$. Thus, we only need to look at the expression (3.4). Therefore, in order to see the boundaries terms that appear in (2.9), we can use exactly the computations already done in the case $\theta = 1$ from which we obtain (3.13).

Now we prove the convergence to the Laplacian which was required above.

Lemma 3.2. *Let $G : \mathbb{R} \rightarrow \mathbb{R}$ be a two times continuously differentiable function with compact support. We have*

$$\limsup_{N \rightarrow \infty} \sup_{x \in \Lambda_N} \left| N^2 \sum_{y \in \mathbb{Z}} (G(\frac{y+x}{N}) - G(\frac{x}{N})) p(y) - \frac{\sigma^2}{2} \Delta G(\frac{x}{N}) \right| = 0.$$

Proof. Let $\varepsilon > 0$ be fixed. We have that $N^2 \sum_{y \in \mathbb{Z}} (G(\frac{x+y}{N}) - G(\frac{x}{N})) p(y)$ is equal to

$$N^2 \sum_{|y| \geq \varepsilon N} (G(\frac{x+y}{N}) - G(\frac{x}{N})) p(y) + N^2 \sum_{|y| < \varepsilon N} (G(\frac{x+y}{N}) - G(\frac{x}{N})) p(y). \quad (3.14)$$

The first term in (3.14) goes to zero with N , since we have that

$$\left| N^2 \sum_{|y| \geq \varepsilon N} (G(\frac{x+y}{N}) - G(\frac{x}{N})) p(y) \right| \lesssim \frac{\|G\|_\infty N^2}{(\varepsilon N)^\gamma}.$$

On the second term of (3.14) we perform a Taylor expansion of G and we have that

$$\begin{aligned} & N^2 \sum_{|y| < \varepsilon N} [G(\frac{x+y}{N}) - G(\frac{x}{N})] p(y) \\ &= N^2 \sum_{|y| < \varepsilon N} \left[G'(\frac{x}{N}) \frac{y}{N} + \frac{1}{2} G''(\frac{x}{N}) (\frac{y}{N})^2 \right] p(y), \end{aligned}$$

plus lower-order terms (with respect to N). Now, we use the fact that $p(\cdot)$ is symmetric to see that $\sum_{|y| < \varepsilon N} y p(y) = 0$. Since $p(\cdot)$ has finite second moment, $\sum_{|y| < \varepsilon N} y^2 p(y) \rightarrow \sigma^2$ so that the proof ends. \square

4. TIGHTNESS

In this section we prove that the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$, defined in Section 2.3, is tight.

Proposition 4.1. *The sequence of measures $\{\mathbb{Q}_N\}_{N \geq 1}$ is tight with respect to the Skorohod topology of $\mathcal{D}([0, T], \mathcal{M}^+)$.*

Proof. In order to prove the assertion, it is enough to show that, for all $\varepsilon > 0$

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\tau \in \mathcal{T}_T, \bar{\tau} \leq \delta} \mathbb{P}_{\mu_N} \left(\eta^N \in \mathcal{D}([0, T], \Omega_N) : \left| \langle \pi_{\tau+\bar{\tau}}^N, G \rangle - \langle \pi_\tau^N, G \rangle \right| > \varepsilon \right) = 0, \quad (4.1)$$

holds for any function G belonging to $C([0, 1])$. Here \mathcal{T}_T is the set of stopping times bounded by T and we implicitly assume that all the stopping times are bounded by T ,

thus, $\tau + \bar{\tau}$ should be read as $(\tau + \bar{\tau}) \wedge T$. In fact it is enough to prove the assertion for functions G in a dense subset of $C([0, 1])$, with respect to the uniform topology.

We split the proof according to two different regimes of θ , namely $\theta \geq 1$ and $\theta < 1$. When $\theta \geq 1$ we prove (4.1) directly for functions $G \in C^2([0, 1])$ and we conclude that the sequence is tight. When $\theta < 1$, we prove (4.1) first for functions $G \in C_c^2(0, 1)$ and then we extend it, by a L^1 approximation procedure which is explained below, to functions $G \in C([0, 1])$.

Recall from (3.1) that $M_t^N(G)$ is a martingale with respect to the natural filtration $\{\mathcal{F}_t\}_{t \geq 0}$. In order to prove (4.1) it is enough to show that

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\tau \in \mathcal{T}_T, \bar{\tau} \leq \delta} \mathbb{E}_{\mu_N} \left[\left| \int_{\tau}^{\tau + \bar{\tau}} \Theta(N) L_N \langle \pi_s^N, G \rangle ds \right| \right] = 0 \quad (4.2)$$

and

$$\lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_{\tau \in \mathcal{T}_T, \bar{\tau} \leq \delta} \mathbb{E}_{\mu_N} \left[(M_{\tau}^N(G) - M_{\tau + \bar{\tau}}^N(G))^2 \right] = 0. \quad (4.3)$$

Proof of (4.2): Given a function G , we claim that we can find a positive constant $C := \overline{C}(G, \alpha, \beta, \gamma, \kappa)$ such that

$$\Theta(N) L_N \langle \pi_s^N, G \rangle \leq C \quad (4.4)$$

for any $s \leq T$, which trivially implies (4.2). To prove it, we recall (3.2) and start to prove that the last two terms of (3.2) are bounded. For example, the absolute value of the second term at the right hand side of (3.2) is bounded from above by

$$\int_0^t \left| \frac{\Theta(N) \kappa}{(N-1)N^\theta} \sum_{x \in \Lambda_N} (Gr_N^-)(\frac{x}{N}) (\alpha - \eta_x^N(s)) \right| ds. \quad (4.5)$$

Now, for $\theta < 1$, we use the fact that $G \in C_c^2(0, 1)$ and that $|\eta_x^N(s)| \leq 1$ is bounded, and we bound from above this last term by a constant times $\Theta(N)N^{-\theta-\gamma}$. Using the definition of $\Theta(N)$ it is easy to see, for $\theta < 2 - \gamma$ and for $2 - \gamma \leq \theta < 1$, that (4.5) is bounded from above by a constant. This proves (4.4) in the case $\theta < 1$. In the case $\theta \geq 1$, we use the fact that the sum in (4.5) is uniformly bounded in N to conclude that (4.5) is bounded from above even if G does not have a compact support included in $(0, 1)$. A similar argument can be done for the last term at the right hand side of (3.2).

Now we need to bound the first term at the right hand side of (3.2). For $\theta < 1$ we use the fact that $G \in C_c^2(0, 1)$ so that $\left| \frac{\Theta(N)}{N-1} \langle \pi_s^N, \mathcal{L}_N G \rangle \right|$ is less or equal than

$$\frac{\Theta(N)}{N-1} \sum_{x \in \Lambda_N} |K_N G(\frac{x}{N})| + \frac{\Theta(N)}{N-1} \sum_{x \in \Lambda_N} |G(\frac{x}{N})| r_N^-(\frac{x}{N}) + \frac{\Theta(N)}{N-1} \sum_{x \in \Lambda_N} |G(\frac{x}{N})| r_N^+(\frac{x}{N}). \quad (4.6)$$

The two terms at the right hand side of the previous expression can be bounded from above by a constant times $\Theta(N)N^{\theta-\gamma}$. It is clearly bounded in the case $\theta \geq 2 - \gamma$ since then $\Theta(N) = N^2$ (recall $\gamma > 2$). In the case $\theta < 2 - \gamma$, $\Theta(N) = N^{\theta+\gamma}$ and thus $\Theta(N)N^{\theta-\gamma}$ is bounded. This together with Lemma 3.2 shows that

$$\left| \frac{\Theta(N)}{N-1} \langle \pi_s^N, \mathcal{L}_N G \rangle \right| \leq C,$$

which proves the claim (4.4) in the case $\theta < 1$. Now, in the case $\theta \geq 1$, since $\Theta(N) = N^2$, we have that the first term at the right hand side of (3.2) is bounded from above

by a constant times

$$\begin{aligned} & \frac{N^2}{N-1} \sum_{x \in \Lambda_N} |K_N G(\frac{x}{N})| + \frac{N^2}{N-1} \sum_{x \in \Lambda_N} \sum_{y \leq 0} |G(\frac{y}{N}) - G(\frac{x}{N})| p(x-y) \\ & + \frac{N^2}{N-1} \sum_{x \in \Lambda_N} \sum_{y \geq N} |G(\frac{y}{N}) - G(\frac{x}{N})| p(x-y). \end{aligned} \quad (4.7)$$

By the Mean Value Theorem, the two terms at the right hand side of the previous expression can be bounded from above by

$$\|G'\|_\infty \sum_{x \in \Lambda_N} \sum_{y \leq 0} |y-x| p(x-y) \lesssim \sum_{x \in \Lambda_N} \frac{1}{x^{\gamma-1}} \quad (4.8)$$

which is finite since $\gamma > 2$. This together with Lemma 3.2 proves (4.4) in the case $\theta \geq 1$.

Proof of (4.3): We know by Dynkin's formula that

$$(M_t^N(G))^2 - \int_0^t \Theta(N) [L_N \langle \pi_s^N, G \rangle^2 - 2 \langle \pi_s^N, G \rangle L_N \langle \pi_s^N, G \rangle] ds,$$

is a martingale with respect to the natural filtration $\{\mathcal{F}_t\}_{t \geq 0}$. From the computations of Appendix B we get that the term inside the time integral in the previous display is equal to

$$\begin{aligned} & \frac{\Theta(N)}{(N-1)^2} \sum_{x < y \in \Lambda_N} (G(\frac{x}{N}) - G(\frac{y}{N}))^2 p(x-y) (\eta_y^N(s) - \eta_x^N(s))^2 \\ & + \frac{\Theta(N)\kappa}{N^\theta(N-1)^2} \sum_{x \in \Lambda_N} G^2(\frac{x}{N}) r_N^-(\frac{x}{N}) (\alpha - \eta_x^N(s)) (1 - 2\eta_x^N(s)) \\ & + \frac{\Theta(N)\kappa}{N^\theta(N-1)^2} \sum_{x \in \Lambda_N} G^2(\frac{x}{N}) r_N^+(\frac{x}{N}) (\beta - \eta_x^N(s)) (1 - 2\eta_x^N(s)). \end{aligned}$$

Since $\Theta(N) \leq N^2$ and G' is bounded it is easy to see that the absolute value of the previous display is bounded from above by a constant times

$$\frac{1}{(N-1)^2} \sum_{x, y \in \Lambda_N} (x-y)^2 p(x-y) + \frac{\Theta(N)}{N^\theta(N-1)^2} \sum_{x \in \Lambda_N} G^2(\frac{x}{N}) (r_N^-(\frac{x}{N}) + r_N^+(\frac{x}{N})) \quad (4.9)$$

Since $\sum_{x, y \in \Lambda_N} (x-y)^2 p(x-y) = \mathcal{O}(N)$ the first term in (4.9) is $\mathcal{O}(N^{-1})$. For the second term at the right hand side of (4.9), we split the argument according to the cases $\theta \geq 1$ and $\theta < 1$. First when $\theta \geq 1$, by using the fact that $\gamma > 2$ and G is bounded so that the sum in that term is finite, and since $\Theta(N) = N^2$, we conclude that the term is $\mathcal{O}(N^{-\theta}) \leq \mathcal{O}(N^{-1})$. From this we obtain (4.3). Now if $\theta < 1$, recall that G has compact support and (3.6). We then write

$$\frac{\Theta(N)}{N^\theta(N-1)^2} \sum_{x \in \Lambda_N} G^2(\frac{x}{N}) (r_N^-(\frac{x}{N}) + r_N^+(\frac{x}{N})) = \frac{\Theta(N)}{N^{\theta+\gamma}(N-1)} I_N(G)$$

where $I_N(G)$ is a Riemann sum converging to $\int_0^1 G^2(q)[r^-(q) + r^+(q)]dq < \infty$. Therefore the second term in (4.9) is of order $\Theta(N)N^{-1-\theta-\gamma} = \mathcal{O}(N^{-1})$ by (2.12).

This ends the proof of tightness in the case $\theta \geq 1$, since $C^2([0, 1])$ is a dense subset of $C([0, 1])$ with respect to the uniform topology. Nevertheless, for $\theta < 1$, we have proved (4.2) and (4.3), and thus (4.1), only for functions $G \in C_c^2(0, 1)$ and we need to extend this result to functions in $C([0, 1])$. To accomplish that, we take a function $G \in C([0, 1]) \subset L^1([0, 1])$, and we take a sequence of functions $\{G_k\}_{k \geq 0} \in C_c^2(0, 1)$ converging to G with respect to the L^1 -norm as $k \rightarrow \infty$. Now, since the probability in (4.1) is less or equal than

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\eta^N \in \mathcal{D}([0, T], \Omega_N) : |\langle \pi_{\tau+\bar{\tau}}^N, G_k \rangle - \langle \pi_{\tau}^N, G_k \rangle| > \frac{\varepsilon}{2} \right) \\ & + \mathbb{P}_{\mu_N} \left(\eta^N \in \mathcal{D}([0, T], \Omega_N) : |\langle \pi_{\tau+\bar{\tau}}^N, G - G_k \rangle - \langle \pi_{\tau}^N, G - G_k \rangle| > \frac{\varepsilon}{2} \right) \end{aligned}$$

and since G_k has compact support, from the computation above, it remains only to check that the last probability vanishes as $N \rightarrow \infty$ and then $k \rightarrow \infty$. For that purpose, we use the fact that

$$|\langle \pi_{\tau+\bar{\tau}}^N, G - G_k \rangle - \langle \pi_{\tau}^N, G - G_k \rangle| \leq \frac{2}{N} \sum_{x \in \Lambda_N} |(G - G_k)(\frac{x}{N})|, \quad (4.10)$$

and we use the estimate

$$\begin{aligned} \frac{1}{N} \sum_{x \in \Lambda_N} |(G - G_k)(\frac{x}{N})| & \leq \sum_{x \in \Lambda_N} \int_{x/N}^{(x+1)/N} |(G - G_k)(\frac{x}{N}) - (G - G_k)(q)| dq \\ & \quad + \int_0^1 |(G - G_k)(q)| dq \\ & \leq \frac{1}{N} \|(G - G_k)'\|_{\infty} + \int_0^1 |(G - G_k)(q)| dq. \end{aligned}$$

We conclude the result by taking first the limsup in $N \rightarrow \infty$ and then in $k \rightarrow \infty$. \square

5. REPLACEMENT LEMMAS AND AUXILIARY RESULTS

In this section we establish some technical results needed in the proof of the hydrodynamic limit. In what follows, we will suppose without loss of generality that $\alpha \leq \beta$. Let $h : [0, 1] \rightarrow [0, 1]$ be a Lipschitz function such that $\alpha \leq h(q) \leq \beta$, for all $q \in [0, 1]$. Let $\nu_{h(\cdot)}^N$ be the Bernoulli product measure on Ω_N with marginals given by

$$\nu_{h(\cdot)}^N \{ \eta : \eta_x = 1 \} = h\left(\frac{x}{N}\right).$$

Given two functions $f, g : \Omega_N \rightarrow \mathbb{R}$ and a probability measure μ on Ω_N , we denote here by $\langle f, g \rangle_{\mu}$ the scalar product between f and g in $L^2(\Omega_N, \mu)$, that is,

$$\langle f, g \rangle_{\mu} = \int_{\Omega_N} f(\eta)g(\eta) d\mu.$$

The notation above should not be mistaken to the notation that we introduced in Section 2.2. We denote by $H_N(\mu | \nu_{h(\cdot)}^N)$ the relative entropy of a probability measure

μ on Ω_N with respect to the probability measure $\nu_{h(\cdot)}^N$ on Ω_N . It is easy to prove the existence of a constant $C_0 := C_0(\alpha, \beta)$, such that

$$H_N(\mu | \nu_{h(\cdot)}^N) \leq N C_0. \quad (5.1)$$

In fact, using the explicit formula for the entropy and the definition of the product measure $\nu_{h(\cdot)}^N$, we get that

$$\begin{aligned} H(\mu | \nu_{h(\cdot)}^N) &= \sum_{\eta \in \Omega_N} \mu(\eta) \log \left(\frac{\mu(\eta)}{\nu_{h(\cdot)}^N(\eta)} \right) \leq \sum_{\eta \in \Omega_N} \mu(\eta) \log \left(\frac{1}{\nu_{h(\cdot)}^N(\eta)} \right) \\ &\leq \log \left(\left[\frac{1}{\alpha \wedge (1-\beta)} \right]^N \right) \sum_{\eta \in \Omega_N} \mu(\eta) \leq N \log \left(\frac{1}{\alpha \wedge (1-\beta)} \right) \leq N C_0. \end{aligned}$$

5.1. Estimates on Dirichlet forms. For a probability measure μ on Ω_N , $x, y \in \Lambda_N$ and a density function $f : \Omega_N \rightarrow [0, \infty)$ with respect to μ we introduce

$$\begin{aligned} I_{x,y}(\sqrt{f}, \mu) &:= \int_{\Omega_N} \left(\sqrt{f(\sigma^{x,y}\eta)} - \sqrt{f(\eta)} \right)^2 d\mu, \\ I_x^\alpha(\sqrt{f}, \mu) &:= \int_{\Omega_N} c_x(\eta; \alpha) \left(\sqrt{f(\sigma^x\eta)} - \sqrt{f(\eta)} \right)^2 d\mu. \end{aligned}$$

Then we define

$$D_N(\sqrt{f}, \mu) := (D_N^0 + D_N^\ell + D_N^r)(\sqrt{f}, \mu)$$

where

$$D_N^0(\sqrt{f}, \mu) := \frac{1}{2} \sum_{x,y \in \Lambda_N} p(y-x) I_{x,y}(\sqrt{f}, \mu), \quad (5.2)$$

$$D_N^\ell(\sqrt{f}, \mu) := \frac{\kappa}{N^\theta} \sum_{x \in \Lambda_N} \sum_{y \leq 0} p(y-x) I_x^\alpha(\sqrt{f}, \mu) = \frac{\kappa}{N^\theta} \sum_{x \in \Lambda_N} r_N^-(\frac{x}{N}) I_x^\alpha(\sqrt{f}, \mu) \quad (5.3)$$

and $D_N^r(\sqrt{f}, \mu)$ is the same as $D_N^\ell(\sqrt{f}, \mu)$ but in $I_x^\alpha(\sqrt{f}, \mu)$ the parameter α is replaced by β and $r_N^-(\cdot)$ is replaced by $r_N^+(\cdot)$.

Our first goal is to express, for the measure $\nu_{h(\cdot)}^N$, a relation between the Dirichlet form defined by $\langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N}$ and $D_N(\sqrt{f}, \nu_{h(\cdot)}^N)$. More precisely, we claim that for any positive constant B , there exists a constant $C > 0$ such that

$$\begin{aligned} \frac{1}{BN} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} &\leq -\frac{1}{4BN} D_N(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C}{BN} \sum_{x,y \in \Lambda_N} p(y-x) \left(h(\frac{x}{N}) - h(\frac{y}{N}) \right)^2 \\ &\quad + \frac{C\kappa}{BN^{1+\theta}} \sum_{x \in \Lambda_N} \left\{ \left(h(\frac{x}{N}) - \alpha \right)^2 r_N^-(\frac{x}{N}) + \left(h(\frac{x}{N}) - \beta \right)^2 r_N^+(\frac{x}{N}) \right\}. \end{aligned} \quad (5.4)$$

Our aim is then to choose $h(\cdot)$ in order to minimize the error term, i.e. the two last terms at the right hand side of the previous inequality.

If $h(\cdot)$ is such that $h(0) = \alpha$ and $h(1) = \beta$, since it is assumed to be Lipschitz, we get the estimate

$$\begin{aligned} \frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} &\leq -\frac{N}{4B} D_N(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C}{B} \sigma^2 \\ &+ \frac{C\kappa}{BN^{1+\theta}} \sum_{x \in \Lambda_N} \left\{ x^2 r_N^-\left(\frac{x}{N}\right) + (x-N)^2 r_N^+\left(\frac{x}{N}\right) \right\}. \end{aligned} \quad (5.5)$$

Moreover, if the function $h(\cdot)$ is such that $h(0) = \alpha$ and $h(1) = \beta$, Hölder of parameter $\gamma/2$ at the boundaries and Lipschitz inside, then we have

$$\frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \leq -\frac{N}{4B} D_N(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C}{B} \sigma^2 + \frac{C\kappa}{BN^{\gamma+\theta-2}}. \quad (5.6)$$

On the other hand if the function $h(\cdot)$ is constant, equal to α or to β , then we have

$$\frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_\alpha} \leq -\frac{N}{4B} D_N(\sqrt{f}, \nu_\alpha) + \frac{C\kappa}{B} N^{1-\theta}. \quad (5.7)$$

In order to prove (5.4) we need some intermediate results. In what follows C is a constant depending on α and β whose value can change from line to line.

Lemma 5.1. *Let $T : \eta \in \Omega_N \rightarrow T(\eta) \in \Omega_N$ be a transformation and $c : \eta \rightarrow c(\eta)$ be a positive local function. Let f be a density with respect to a probability measure μ on Ω_N . Then, we have that*

$$\begin{aligned} &\langle c(\eta) [\sqrt{f(T(\eta))} - \sqrt{f(\eta)}], \sqrt{f(\eta)} \rangle_\mu \\ &\leq -\frac{1}{4} \int c(\eta) \left([\sqrt{f(T(\eta))}] - [\sqrt{f(\eta)}] \right)^2 d\mu \\ &+ \frac{1}{16} \int \frac{1}{c(\eta)} \left[c(\eta) - c(T(\eta)) \frac{\mu(T(\eta))}{\mu(\eta)} \right]^2 \left([\sqrt{f(T(\eta))}] + [\sqrt{f(\eta)}] \right)^2 d\mu. \end{aligned} \quad (5.8)$$

Proof. By writing the term at the left hand side of (5.8) as its half plus its half and performing a change of variables $\eta \rightarrow T(\eta)$ in one of them, we have that

$$\begin{aligned} &\int c(\eta) [\sqrt{f(T(\eta))} - \sqrt{f(\eta)}] \sqrt{f(\eta)} d\mu \\ &= -\frac{1}{2} \int c(\eta) [\sqrt{f(T(\eta))} - \sqrt{f(\eta)}]^2 d\mu \\ &+ \frac{1}{2} \int [\sqrt{f(T(\eta))}]^2 [c(\eta) - c(T(\eta))T(\eta)] d\mu. \end{aligned}$$

Repeating again the same argument, the second term at the right hand side of last expression can be written as

$$\frac{1}{4} \int \left([\sqrt{f(T(\eta))}]^2 - [\sqrt{f(\eta)}]^2 \right) [c(\eta) - c(T(\eta))T(\eta)] d\mu.$$

By Young's inequality and the elementary equality $a^2 - b^2 = (a - b)(a + b)$, last expression is bounded from above by

$$\begin{aligned} & \frac{1}{4} \int c(\eta) \left(\left[\sqrt{f(T(\eta))} \right] - \left[\sqrt{f(\eta)} \right] \right)^2 d\mu \\ & + \frac{1}{16} \int \frac{1}{c(\eta)} \left[c(\eta) - c(T(\eta))T(\eta) \right]^2 \left(\left[\sqrt{f(T(\eta))} \right] + \left[\sqrt{f(\eta)} \right] \right)^2 d\mu, \end{aligned}$$

which finishes the proof. \square

Lemma 5.2. *There exists a constant $C := C(h)$ such that for any $N \geq 1$ and density f be a density with respect to $\nu_{h(\cdot)}^N$*

$$\sup_{x \neq y \in \Lambda_N} \int_{\Omega_N} f(\sigma^{x,y} \eta) d\nu_{h(\cdot)}^N(\eta) \leq C, \quad \sup_{x \in \Lambda_N} \int_{\Omega_N} f(\sigma^x \eta) d\nu_{h(\cdot)}^N(\eta) \leq C.$$

Proof. Let us prove only the first bound since the proof of the second one is similar. We perform in the first integral above the change of variable $\omega = \sigma^{x,y} \eta$ and we use that uniformly in $x, y \in \Lambda_N$ and ω we have

$$\theta^{x,y}(\omega) = \frac{\nu_{h(\cdot)}^N(\sigma^{x,y} \omega)}{\nu_{h(\cdot)}^N(\omega)} = 1 + \mathcal{O}\left(\frac{1}{N}\right).$$

By using the fact that f is a density it is easy to conclude. \square

Now, let us look at some consequences of these lemmas. We start with the bulk generator L_N^0 given in (2.2).

Corollary 5.3. *There exists a constant $C > 0$ (independent of f and N) such that*

$$\left\langle L_N^0 \sqrt{f}, \sqrt{f} \right\rangle_{\nu_{h(\cdot)}^N} \leq -\frac{1}{4} D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N) + C \sum_{x,y \in \Lambda_N} p(y-x) \left(h\left(\frac{x}{N}\right) - h\left(\frac{y}{N}\right) \right)^2$$

for any density f with respect to $\nu_{h(\cdot)}^N$.

Proof. To prove this we note that

$$\left\langle L_N^0 \sqrt{f}, \sqrt{f} \right\rangle_{\nu_{h(\cdot)}^N} = \frac{1}{2} \sum_{x,y \in \Lambda_N} p(y-x) \left\langle \left[\sqrt{f(\sigma^{x,y} \eta)} - \sqrt{f(\eta)} \right], \sqrt{f(\eta)} \right\rangle_{\nu_{h(\cdot)}^N}.$$

Now, by Lemma 5.1 with $c \equiv 1$, $T = \sigma^{x,y}$, and Lemma 5.2 last expression is bounded from above by

$$-\frac{1}{4} D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N) + C \sum_{x,y \in \Lambda_N} p(y-x) \left(h\left(\frac{x}{N}\right) - h\left(\frac{y}{N}\right) \right)^2,$$

because $|\theta^{x,y}(\eta) - 1|^2 \lesssim (h(x/N) - h(y/N))^2$. \square

Now we look at the generators of the reservoirs given in (2.2).

Corollary 5.4. *Let $\theta \in \mathbb{R}$ be fixed. There exists a constant $C > 0$ (independent of f and N) such that*

$$\begin{aligned} \left\langle L_N^\ell \sqrt{f}, \sqrt{f} \right\rangle_{\nu_{h(\cdot)}^N} &\leq -\frac{1}{4} D_N^\ell(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C\kappa}{N\theta} \sum_{x \in \Lambda_N} r_N^-\left(\frac{x}{N}\right) \left(h\left(\frac{x}{N}\right) - \alpha \right)^2, \\ \left\langle L_N^r \sqrt{f}, \sqrt{f} \right\rangle_{\nu_{h(\cdot)}^N} &\leq -\frac{1}{4} D_N^r(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C\kappa}{N\theta} \sum_{x \in \Lambda_N} r_N^+\left(\frac{x}{N}\right) \left(h\left(\frac{x}{N}\right) - \beta \right)^2 \end{aligned} \quad (5.9)$$

for any density f with respect to $\nu_{h(\cdot)}^N$.

Proof. We present the proof for the first inequality but we note that the proof of the second one is analogous. First observe that

$$\langle L_N^\ell \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} = \frac{\kappa}{N^\theta} \sum_{x \in \Lambda_N} \sum_{y \leq 0} p(y-x) \langle c_x(\eta; \alpha) [\sqrt{f(\sigma^x \eta)} - \sqrt{f(\eta)}], \sqrt{f(\eta)} \rangle_{\nu_{h(\cdot)}^N}.$$

Now, by using Lemma 5.1 with $c(\eta) = c_x(\eta; \alpha)$, $T = \sigma^x$ and Lemma 5.2, last expression is bounded from above by

$$-\frac{1}{4} D_N^\ell(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{C\kappa}{N^\theta} \sum_{x \in \Lambda_N} \sum_{y \leq 0} p(y-x) \left(h\left(\frac{x}{N}\right) - \alpha \right)^2.$$

□

From the two previous corollaries the claim (5.4) follows.

5.2. Replacement Lemmas.

Lemma 5.5. *For any density f with respect to $\nu_{h(\cdot)}^N$, any $x \in \Lambda_N$ and any positive constant A_x , we have that*

$$\left| \langle t_x^\alpha, f \rangle_{\nu_{h(\cdot)}^N} \right| \lesssim \frac{1}{A_x} I_x^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N) + A_x + [h\left(\frac{x}{N}\right) - \alpha],$$

where $t_x^\alpha(\eta) = \eta_x - \alpha$. The same result holds if α is replaced by β .

Proof. By a simple computation we have that:

$$\begin{aligned} \left| \langle t_x^\alpha, f \rangle_{\nu_{h(\cdot)}^N} \right| &\leq \frac{1}{2} \left| \int t_x^\alpha(\eta) (f(\eta) - f(\sigma^x \eta)) d\nu_{h(\cdot)}^N \right| \\ &+ \frac{1}{2} \left| \int [f(\sigma^x \eta) + f(\eta)] t_x(\eta) d\nu_{h(\cdot)}^N \right|, \end{aligned} \quad (5.10)$$

where σ^x is the flip given in (2.3). By Young's inequality, using the fact that $(a-b) = (\sqrt{a}-\sqrt{b})(\sqrt{a}+\sqrt{b})$ for all $a, b \geq 0$ and Lemma 5.2, the first term at the right side of (5.10) is bounded from above, for any positive constant A_x , by

$$\begin{aligned} &\frac{A_x}{4} \int \frac{(t_x(\eta))^2}{c_x(\eta; \alpha)} \left([\sqrt{f(\sigma^x \eta)}] + [\sqrt{f(\eta)}] \right)^2 d\nu_{h(\cdot)}^N + \frac{I_x^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N)}{4A_x} \\ &\lesssim A_x + \frac{I_x^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N)}{A_x}. \end{aligned}$$

Now, we look at the second term at the right hand side of (5.10). By using the fact that $\nu_{h(\cdot)}^N$ is product and denoting by $\tilde{\eta}$ the configuration η removing its value at x so

that $(\eta_x, \bar{\eta}) = \eta$, we have that the second term at the right side of (5.10) is equal to

$$\begin{aligned}
& \frac{1}{2} \left| \sum_{\bar{\eta}} \left((1-\alpha)(f(1, \bar{\eta}) + f(0, \bar{\eta})) \nu_{h(\cdot)}^N(\eta_x = 1) \right. \right. \\
& \quad \left. \left. - \alpha(f(0, \bar{\eta}) + f(1, \bar{\eta})) \nu_{h(\cdot)}^N(\eta(x) = 0) \right) \nu_{h(\cdot)}^N(\bar{\eta}) \right| \\
&= \frac{1}{2} \left| \sum_{\bar{\eta}} \left(h\left(\frac{x}{N}\right) - \alpha \right) (f(0, \bar{\eta}) + f(1, \bar{\eta})) \nu_{h(\cdot)}^N(\bar{\eta}) \right| \\
&\lesssim \left(h\left(\frac{x}{N}\right) - \alpha \right) \sum_{\bar{\eta}} h\left(\frac{x}{N}\right) f(1, \bar{\eta}) \nu_{h(\cdot)}^N(\bar{\eta}) + \left(1 - h\left(\frac{x}{N}\right) \right) f(0, \bar{\eta}) \nu_{h(\cdot)}^N(\bar{\eta}) \\
&= \left(h\left(\frac{x}{N}\right) - \alpha \right) \sum_{\eta \in \Omega_N} f(\eta) \nu_{h(\cdot)}^N(\eta) = \left(h\left(\frac{x}{N}\right) - \alpha \right)
\end{aligned}$$

because $\max_{x \in \Lambda_N} \left\{ \frac{1}{2h\left(\frac{x}{N}\right)}, \frac{1}{2\left(1-h\left(\frac{x}{N}\right)\right)} \right\}$ is bounded from above by a constant depending only on α and β . Above $f(1, \bar{\eta})$ (resp. $f(0, \bar{\eta})$) means that we are computing $f(\eta)$ with $\eta_x = 1$ (resp. $\eta_x = 0$). \square

Lemma 5.6. *For any $t > 0$, we have that*

$$\begin{aligned}
\lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t N^{1-\theta} \sum_{x \in \Lambda_N} Gr_N^-\left(\frac{x}{N}\right) (\eta_x(sN^2) - \alpha) ds \right| \right] &= 0, \\
\lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t N^{1-\theta} \sum_{x \in \Lambda_N} Gr_N^+\left(\frac{x}{N}\right) (\eta_x(sN^2) - \beta) ds \right| \right] &= 0,
\end{aligned} \tag{5.11}$$

for $G : \mathbb{R} \rightarrow \mathbb{R}$ Lipschitz and such that $G(0) = 0 = G(1)$, for any $\theta > 2 - \gamma$; and for any bounded $G : \mathbb{R} \rightarrow \mathbb{R}$ for $\theta > 1$.

Proof. We present the proof for the first term, but we note that the proof for the second term is completely analogous.

We start by fixing a Lipschitz profile $h(\cdot)$ such that $h(0) = \alpha$ and $h(1) = \beta$. By the entropy inequality, for any $B > 0$, the first expectation of (5.11) is bounded from above by

$$\frac{H(\mu_N | \nu_{h(\cdot)}^N)}{BN} + \frac{1}{BN} \log \mathbb{E}_{\nu_{h(\cdot)}^N} \left[e^{BN \left| \int_0^t N^{1-\theta} \sum_{x \in \Lambda_N} Gr_N^-\left(\frac{x}{N}\right) (\eta_x(sN^2) - \alpha) ds \right|} \right]. \tag{5.12}$$

We can remove the absolute value inside the exponential since $e^{|x|} \leq e^x + e^{-x}$ and

$$\limsup_{N \rightarrow \infty} N^{-1} \log(a_N + b_N) \leq \max \left\{ \limsup_{N \rightarrow \infty} N^{-1} \log(a_N), \limsup_{N \rightarrow \infty} N^{-1} \log(b_N) \right\}. \tag{5.13}$$

By (5.1) and the Feynman-Kac's formula, (5.12) is bounded from above by

$$\frac{C_0}{B} + t \sup_f \left\{ N^{1-\theta} \sum_{x \in \Lambda_N} \left| Gr_N^-\left(\frac{x}{N}\right) \langle t_x^\alpha, f \rangle_{\nu_{h(\cdot)}^N} \right| + \frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \right\},$$

where the supremum is carried over all the densities f with respect to $\nu_{h(\cdot)}^N$. We recall that $t_x^\alpha(\eta) = \eta_x - \alpha$. From Lemma 5.5 we have that there exists a constant $C :=$

$C(\alpha, \beta, \gamma) > 0$ such that

$$\begin{aligned}
& N^{1-\theta} \sum_{x \in \Lambda_N} \left| (Gr_N^-)(\frac{x}{N}) \langle t_x^\alpha, f \rangle \nu_{h(\cdot)}^N \right| \\
& \leq CN^{1-\theta} \sum_{x \in \Lambda_N} |(Gr_N^-)(\frac{x}{N})| \left[A_x + \frac{I_x^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N)}{A_x} + \frac{x}{N} \right] \\
& \leq 4C^2 \kappa^{-1} B N^{1-\theta} \sum_{x \in \Lambda_N} G^2(\frac{x}{N}) r_N^-(\frac{x}{N}) + \frac{N}{4B} D_N^\ell(\sqrt{f}, \nu_{h(\cdot)}^N) + CN^{-\theta} \sum_{x \in \Lambda_N} |G(\frac{x}{N})| r_N^-(\frac{x}{N}) x.
\end{aligned} \tag{5.14}$$

The last inequality is obtained by choosing $A_x = 4\kappa^{-1}C|G(\frac{x}{N})|B$. Recall (5.5).

In the case $\theta > 2 - \gamma$, we have $G(0) = 0$ so that $|G(\frac{x}{N})| \lesssim \frac{x}{N}$. Then (5.12) is bounded from above by a constant (independent of N and B) times

$$\frac{1}{B} + \frac{B+B^{-1}}{N^{1+\theta}} \sum_{x \in \Lambda_N} \left\{ x^2 r_N^-(\frac{x}{N}) + (x-N)^2 r_N^+(\frac{x}{N}) \right\} + \frac{1}{N^{1+\theta}} \sum_{x \in \Lambda_N} x^2 r_N^-(\frac{x}{N}). \tag{5.15}$$

Since we have that

$$\sum_{x \in \Lambda_N} x^2 r_N^-(\frac{x}{N}) \lesssim \begin{cases} N^{3-\gamma}, & \gamma \in (2, 3), \\ \log N, & \gamma = 3, \\ 1, & \gamma > 3, \end{cases} \tag{5.16}$$

and that the same holds for $x^2 r_N^-$ replaced with $(x-N)^2 r_N^+$, this implies that the expression (5.15), for any $\theta \geq 2 - \gamma$, goes to zero by taking the limit as $N \rightarrow \infty$ and then $B \rightarrow \infty$.

In the case $\theta > 1$, since the function G is only assumed to be bounded and does not satisfy a priori $G(0) = 0$ we use (5.14) and (5.5) and estimate from above (5.12) by a constant times

$$\begin{aligned}
& \frac{1}{B} + \frac{1}{BN^{1+\theta}} \sum_{x \in \Lambda_N} \left\{ x^2 r_N^-(\frac{x}{N}) + (x-N)^2 r_N^+(\frac{x}{N}) \right\} \\
& + BN^{1-\theta} \sum_{x \in \Lambda_N} r_N^-(\frac{x}{N}) + N^{-\theta} \sum_{x \in \Lambda_N} r_N^-(\frac{x}{N}) x,
\end{aligned} \tag{5.17}$$

which, by (5.16) and (3.7), goes to zero as $N \rightarrow \infty$ and then $B \rightarrow \infty$. \square

Let us define for $\ell \in \mathbb{N}$ the following empirical densities

$$\overrightarrow{\eta}_0^\ell := \frac{1}{\ell} \sum_{y=1}^{\ell} \eta_y \quad \text{and} \quad \overleftarrow{\eta}_N^\ell := \frac{1}{\ell} \sum_{y=N-1-\ell}^{N-1} \eta_y. \tag{5.18}$$

Lemma 5.7. *For any $t > 0$ and any $\theta \geq 1$ we have that*

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t \sum_{x \in \Lambda_N} \Theta_x^-(\eta_x(sN^2) - \overrightarrow{\eta}_0^{\varepsilon N}(sN^2)) ds \right| \right] = 0, \\
& \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t \sum_{x \in \Lambda_N} \Theta_x^+(\eta_x(sN^2) - \overleftarrow{\eta}_N^{\varepsilon N}(sN^2)) ds \right| \right] = 0.
\end{aligned}$$

Proof. We present the proof for the first term, but we note that the proof for the second one it is analogous. Here we take as reference measure the Bernoulli product measure with constant parameter (for example α) and we recall (5.7). By the entropy inequality the expectation in the statement of the lemma is bounded from above, for any $B > 0$, by

$$\frac{H(\mu_N | \nu_\alpha^N)}{BN} + \frac{1}{BN} \log \mathbb{E}_{\nu_\alpha^N} \left[e^{BN \left| \int_0^t \sum_{x \in \Lambda_N} \Theta_x^-(\eta_x(sN^2) - \vec{\eta}_0^{\varepsilon N}(sN^2)) ds \right|} \right].$$

As in the previous proof, we can remove the absolute value inside the exponential, so that by (5.1) and by the Feynman-Kac's formula last expression can be estimated from above by

$$\frac{C_0}{B} + t \sup_f \left\{ \sum_{x \in \Lambda_N} \Theta_x^-(\tau_x^{\varepsilon N}, f)_{\nu_\alpha^N} + \frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_\alpha^N} \right\}, \quad (5.19)$$

where the supremum is carried over all the densities f with respect to ν_α^N . Here $\tau_x^{\varepsilon N}(\eta) = \eta_x - \vec{\eta}_0^{\varepsilon N}$.

Now we have to split the sum in x , depending on whether $N - 1 \geq x \geq \varepsilon N$ or $x \leq \varepsilon N - 1$. We start by the first case and we have

$$\begin{aligned} \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} &= \frac{1}{\varepsilon N} \sum_{y=1}^{\varepsilon N} \int (\eta_x - \eta_y) f(\eta) d\nu_\alpha^N \\ &= \frac{1}{\varepsilon N} \sum_{y=1}^{\varepsilon N} \sum_{z=y}^{x-1} \int (\eta_{z+1} - \eta_z) f(\eta) d\nu_\alpha^N. \end{aligned}$$

By writing the previous term as its half plus its half and by performing in one of the terms the change of variables η into $\sigma^{z, z+1} \eta$, for which the measure ν_α^N is invariant, we write it as

$$\frac{1}{2\varepsilon N} \sum_{y=1}^{\varepsilon N} \sum_{z=y}^{x-1} \int (f(\eta) - f(\sigma^{z, z+1} \eta)) (\eta_{z+1} - \eta_z) d\nu_\alpha^N.$$

By using the fact that $(a - b) = (\sqrt{a} - \sqrt{b})(\sqrt{a} + \sqrt{b})$ for any $a, b \geq 0$ and since $ab \leq \frac{Aa^2}{2} + \frac{b^2}{2A}$ for all $A > 0$, we have that

$$\begin{aligned} \sum_{x=\varepsilon N}^{N-1} \Theta_x^-(\tau_x^{\varepsilon N}, f)_{\nu_\alpha^N} &\leq \frac{A}{2} \sum_{x=\varepsilon N}^{N-1} \Theta_x^- \frac{1}{2\varepsilon N} \sum_{y=1}^{\varepsilon N} \sum_{z=y}^{x-1} \int (\sqrt{f(\eta)} - \sqrt{f(\sigma^{z, z+1} \eta)})^2 d\nu_\alpha^N \\ &\quad + \frac{1}{2A} \sum_{x=\varepsilon N}^{N-1} \Theta_x^- \frac{1}{2\varepsilon N} \sum_{y=1}^{\varepsilon N} \sum_{z=y}^{x-1} \int (\sqrt{f(\eta)} + \sqrt{f(\sigma^{z, z+1} \eta)})^2 (\eta_{z+1} - \eta_z)^2 d\nu_\alpha^N. \end{aligned} \quad (5.20)$$

By neglecting the jumps of size bigger than one, we see that

$$D^{NN}(\sqrt{f}, \nu_\alpha^N) = \sum_{z \in \Lambda_N} \int (\sqrt{f(\eta)} - \sqrt{f(\sigma^{z, z+1} \eta)})^2 d\nu_\alpha^N \lesssim D_N^0(\sqrt{f}, \nu_\alpha^N).$$

Therefore, by using also (3.10), the first term at the right hand side of (5.20) can be bounded from above by

$$\frac{A}{4} \sum_{x=\varepsilon N}^{N-1} \Theta_x^- D^{NN}(\sqrt{f}, \nu_\alpha^N) \lesssim AD^{NN}(\sqrt{f}, \nu_\alpha^N) \lesssim AD_N^0(\sqrt{f}, \nu_\alpha^N). \quad (5.21)$$

Recall (5.7) and observe that $D_N(\sqrt{f}, \nu_\alpha^N) \geq D_N^0(\sqrt{f}, \nu_\alpha^N)$. Then we choose the constant A in the form $A = CN/B$ for some suitable C in order that one half of the term $-\frac{N}{4B}D_N(\sqrt{f}, \nu_\alpha)$ appearing in (5.7) counterbalances negatively the term at the right hand side of (5.21). Moreover we can bound from above the last term at the right hand side of (5.20) by (use Lemma 5.2)

$$\frac{B}{N} \sum_{x=\varepsilon N}^{N-1} \Theta_x^- \frac{1}{2\varepsilon N} \sum_{y=1}^{\varepsilon N} \sum_{z=y}^{x-1} \int (\sqrt{f(\eta)} + \sqrt{f(\sigma^{z,z+1}\eta)})^2 (\eta_{z+1} - \eta_z)^2 d\nu_\alpha^N \lesssim \frac{B}{N} \sum_{x \in \Lambda_N} x \Theta_x^- \quad (5.22)$$

which vanishes as $N \rightarrow \infty$ by (5.7). Therefore we proved that uniformly in ε

$$\limsup_{B \rightarrow \infty} \limsup_{N \rightarrow \infty} \sup_f \left\{ \sum_{x=\varepsilon N}^{N-1} \Theta_x^- \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} + \frac{N}{2B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_\alpha^N} \right\} = 0.$$

It remains to prove that

$$\limsup_{B \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_f \left\{ \sum_{x=1}^{\varepsilon N-1} \Theta_x^- \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} + \frac{N}{2B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_\alpha^N} \right\} = 0.$$

If $x \leq \varepsilon N - 1$, we write

$$\begin{aligned} \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} &= \frac{1}{\varepsilon N} \sum_{y=1}^{\varepsilon N} \int (\eta_x - \eta_y) f(\eta) d\nu_\alpha^N \\ &= \frac{1}{\varepsilon N} \sum_{y=1}^{x-1} \sum_{z=y}^{x-1} \int (\eta_{z+1} - \eta_z) f(\eta) d\nu_\alpha^N - \frac{1}{\varepsilon N} \sum_{y=x+1}^{\varepsilon N} \sum_{z=x}^{y-1} \int (\eta_{z+1} - \eta_z) f(\eta) d\nu_\alpha^N. \end{aligned}$$

and the same estimates as before give there exists a constant $C > 0$ such that for any $A > 0$,

$$\sum_{x=1}^{\varepsilon N-1} \Theta_x^- \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} \leq C \left[AD_N(\sqrt{f}, \nu_\alpha^N) + \frac{\varepsilon N}{A} \sum_{x=1}^{\varepsilon N-1} \Theta_x^- \right].$$

Recall (5.7) and (3.10). Then, we choose $A = N/8CB$ and we get that

$$\limsup_{B \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \sup_f \left\{ \sum_{x=1}^{\varepsilon N-1} \Theta_x^- \langle \tau_x^{\varepsilon N}, f \rangle_{\nu_\alpha^N} + \frac{N}{2B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_\alpha^N} \right\} = 0.$$

This finishes the proof. \square

Remark 5.8. We note that above, if we change in the statement of the lemma Θ_x^- by r_N^- , then the same result holds by performing exactly the same estimates as above, because what we need is that

$$\sum_{x \in \Lambda_N} \Theta_x^- \lesssim 1 \quad \text{and} \quad \frac{1}{N} \sum_{x \in \Lambda_N} x \Theta_x^- \rightarrow 0 \quad (5.23)$$

which also holds for r_N^- instead of Θ_x^- since $\gamma > 2$.

5.3. Fixing the profile at the boundary. Let \mathbb{Q} be a limit point of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$, whose existence follows from Proposition 4.1 and assume, without loss of generality, that $\{\mathbb{Q}_N\}_{N \geq 1}$ converges to \mathbb{Q} . We note that since our model is an exclusion process, it is standard ([10]) to show that \mathbb{Q} almost surely the trajectories of measures are absolutely continuous with respect to the Lebesgue measure, that is: $\pi_t(dq) = \rho_t(q)dq$ for any $t \in [0, T]$. In Section 6 we prove that the density $\rho_t(q)$ belongs to $L^2(0, T; \mathcal{H}^1(0, 1))$ if $\theta \geq 2 - \gamma$. In particular, for almost every t , ρ_t can be identified to a continuous function on $[0, 1]$.

In this section we prove that if $\theta \in [2 - \gamma, 1)$ then the profile satisfies $\rho_t(0) = \alpha$ and $\rho_t(1) = \beta$.

Recall (5.18). Observe that

$$\mathbb{E}_{\mu_N} \left[\left| \int_0^t (\overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \alpha) ds \right| \right] = \mathbb{E}_{\mathbb{Q}_N} \left[\left| \int_0^t (\langle \pi_s, \iota_\varepsilon^0 \rangle - \alpha) ds \right| \right]$$

where $\iota_\varepsilon^0(\cdot) = \varepsilon^{-1} \mathbf{1}_{(0, \varepsilon)}(\cdot)$. Therefore we have that for any $\delta > 0$,

$$\mathbb{Q}_N \left[\left| \int_0^t (\langle \pi_s, \iota_\varepsilon^0 \rangle - \alpha) ds \right| > \delta \right] \leq \delta^{-1} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \alpha) ds \right| \right].$$

Py Portemanteau's Theorem ² we conclude that

$$\mathbb{Q} \left[\left| \int_0^t (\langle \pi_s, \iota_\varepsilon^0 \rangle - \alpha) ds \right| > \delta \right] \leq \delta^{-1} \liminf_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \alpha) ds \right| \right].$$

Now, if we are able to prove that the right hand side of the previous inequality is zero, since we have that \mathbb{Q} a.s. $\pi_s(dq) = \rho_s(q)dq$ with ρ_s a continuous function in 0 for a.e. s , by taking the limit $\varepsilon \rightarrow 0$, we can deduce that \mathbb{Q} a.s. $\rho_s(0) = \alpha$ for a.e. s . A similar argument applies for the right boundary. Therefore it is sufficient to prove the following lemma.

Lemma 5.9. *For any $t \in [0, T]$ we have that*

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \alpha) ds \right| \right] &= 0, \\ \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\overleftarrow{\eta}_N^{\varepsilon N}(sN^2) - \beta) ds \right| \right] &= 0. \end{aligned}$$

Last lemma is a consequence of the next two results.

Lemma 5.10. *For any $t \in [0, T]$ we have that*

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\eta_1(sN^2) - \alpha) ds \right| \right] &= 0, \\ \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t (\eta_{N-1}(sN^2) - \beta) ds \right| \right] &= 0. \end{aligned}$$

²In fact, since ι_ε^0 is not a continuous function it is not given for free that the set $\{\pi; \left| \int_0^t (\langle \pi_s, \iota_\varepsilon^0 \rangle - \alpha) ds \right| > \delta\}$ is an open set in the Skorohod topology. A simple argument based on a L^1 -approximation of ι_ε^0 by continuous functions permits to bypass this difficulty.

Proof. We give the proof for the first display, but we note that for the other one it is similar. Fix a Lipschitz profile $h(\cdot)$ such that $\alpha \leq h(\cdot) \leq \beta$ and $h(0) = \alpha$, $h(1) = \beta$ and $h(\cdot)$ is $\gamma/2$ -Hölder at the boundaries. By the entropy inequality, for any $B > 0$, the previous expectation is bounded from above by

$$\frac{H(\mu_N | \nu_{h(\cdot)}^N)}{BN} + \frac{1}{BN} \log \mathbb{E}_{\nu_{h(\cdot)}^N} \left[e^{BN \left| \int_0^t (\eta_1(sN^2) - \alpha) ds \right|} \right].$$

By (5.1), the Feynman-Kac's formula and noting, as we did in the proof of Lemma 5.6, that we can remove the absolute value inside the exponential, last display can be estimated from above by

$$\frac{C_0}{B} + t \sup_f \left\{ \langle t_1^\alpha, f \rangle_{\nu_{h(\cdot)}^N} + \frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \right\}, \quad (5.24)$$

where the supremum is carried over all the densities f with respect to $\nu_{h(\cdot)}^N$. Here we recall that $t_1^\alpha(\eta) = \eta_1 - \alpha$. By Lemma 5.5, since h is Lipschitz, for any $A > 0$, the first term in the supremum in (5.24) is bounded from above by

$$C \left[\frac{1}{A} I_1^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N) + A + \frac{1}{N} \right]$$

for some constant $C > 0$ independent of f and A . Moreover from (5.6), since

$$D_N(\sqrt{f}, \nu_{h(\cdot)}^N) \geq D_N^\ell(\sqrt{f}, \nu_{h(\cdot)}^N)$$

and $\gamma + \theta - 2 \geq 0$, we know that there exists a constant $C' > 0$ such that

$$\frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \leq -\frac{N^{1-\theta}}{4B} \sum_{x \in \Lambda_N} I_x^\alpha(\sqrt{f}, \nu_{h(\cdot)}^N) r_N^-(\frac{x}{N}) + \frac{C'}{B}.$$

To get an upper bound, at the right hand side of the previous inequality, we only keep the term coming from $x = 1$ in the sum. By choosing $A = 4C(r_N^-(\frac{1}{N}))^{-1}BN^{\theta-1}$, we get then that the expression inside the brackets in (5.24) is bounded by

$$4C^2 \frac{BN^{\theta-1}}{r_N^-(\frac{1}{N})} + \frac{C}{N} + \frac{C'}{B}.$$

Now since $r_N^-(\frac{1}{N})$ is bounded from below by a constant independent of N and $\theta < 1$, the proof follows by sending first $N \rightarrow \infty$ and then $B \rightarrow \infty$. \square

Lemma 5.11. *For any $t > 0$ we have that*

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t \overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \eta_1(sN^2) ds \right| \right] &= 0, \\ \lim_{\varepsilon \rightarrow 0} \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\left| \int_0^t \overleftarrow{\eta}_N^{\varepsilon N}(sN^2) - \eta_{N-1}(sN^2) ds \right| \right] &= 0. \end{aligned} \quad (5.25)$$

Proof. We present the proof of the first item, but we note that for the second it is exactly the same. Fix a Lipschitz profile $h(\cdot)$ such that $\alpha \leq h(\cdot) \leq \beta$, $h(0) = \alpha$, $h(1) = \beta$ and $h(\cdot)$ is $\gamma/2$ -Hölder at the boundaries. By the entropy inequality, for any $B > 0$, the previous expectation is bounded from above by

$$\frac{H(\mu_N | \nu_{h(\cdot)}^N)}{BN} + \frac{1}{BN} \log \mathbb{E}_{\nu_{h(\cdot)}^N} \left[e^{BN \left| \int_0^t \overrightarrow{\eta}_0^{\varepsilon N}(sN^2) - \eta_1(sN^2) ds \right|} \right].$$

By (5.1), the Feynman-Kac's formula, and using the same argument as in the proof of the previous lemma, the estimate of the previous expression can be reduced to bound

$$\frac{C_0}{B} + t \sup_f \left\{ \frac{1}{\ell} \sum_{y=2}^{\ell+1} |\langle v_y^1, f \rangle_{\nu_{h(\cdot)}^N}| + \frac{N}{B} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \right\}, \quad (5.26)$$

where $\ell = \epsilon N$ and $v_y^1(\eta) = \eta_y - \eta_1$. Here the supremum is carried over all the densities f with respect to $\nu_{h(\cdot)}^N$. Note that since $y \in \Lambda_N$ we know that $v_y^1(\eta) = \sum_{z=1}^{y-1} (\eta_{z+1} - \eta_z)$. Observe now that

$$\begin{aligned} \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) f(\eta) d\nu_{h(\cdot)}^N &= \frac{1}{2} \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) (f(\eta) - f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N \\ &\quad + \frac{1}{2} \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) (f(\eta) + f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N. \end{aligned}$$

By using the fact that for any $a, b \geq 0$, $(a - b) = (\sqrt{a} - \sqrt{b})(\sqrt{a} + \sqrt{b})$ and Young's inequality, we have, for any positive constant A , that

$$\begin{aligned} \frac{1}{\ell} \sum_{y=2}^{\ell+1} |\langle v_y^1, f \rangle_{\nu_{h(\cdot)}^N}| &\leq \frac{1}{2A\ell} \sum_{y=2}^{\ell+1} \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z)^2 (\sqrt{f(\eta)} + \sqrt{f(\sigma^{z,z+1}\eta)})^2 d\nu_{h(\cdot)}^N \\ &\quad + \frac{A}{2\ell} \sum_{y=2}^{\ell+1} \sum_{z=1}^{y-1} \int (\sqrt{f(\eta)} - \sqrt{f(\sigma^{z,z+1}\eta)})^2 d\nu_{h(\cdot)}^N \\ &\quad + \frac{1}{2\ell} \sum_{y=2}^{\ell+1} \left| \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) (f(\eta) + f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N \right|. \end{aligned} \quad (5.27)$$

By neglecting the jumps of size bigger than one, we see that

$$D^{NN}(\sqrt{f}, \nu_{h(\cdot)}^N) = \sum_{z \in \Lambda_N} \int (\sqrt{f(\eta)} - \sqrt{f(\sigma^{z,z+1}\eta)})^2 d\nu_{h(\cdot)}^N \lesssim D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N).$$

Then, the second term on the right hand side of (5.27) is bounded from above by

$$\frac{A}{2\ell} \sum_{y=2}^{\ell+1} D^{NN}(\sqrt{f}, \nu_{h(\cdot)}^N) \leq A D^{NN}(\sqrt{f}, \nu_{h(\cdot)}^N) \leq C A D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N) \leq C A D_N(\sqrt{f}, \nu_{h(\cdot)}^N)$$

where C is a positive constant independent of A, ℓ, f . Then, for the choice $A = N(4BC)^{-1}$ and from (5.6), since $\gamma + \theta - 2 \geq 0$, we can bound from above (5.26) by

$$\begin{aligned} &\frac{2BC}{N\ell} \sum_{y=2}^{\ell+1} \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z)^2 (\sqrt{f(\eta)} + \sqrt{f(\sigma^{z,z+1}\eta)})^2 d\nu_{h(\cdot)}^N \\ &+ \frac{1}{2\ell} \sum_{y=2}^{\ell+1} \left| \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) (f(\eta) + f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N \right| + \frac{C'}{B} \\ &\lesssim \frac{B\ell}{N} + \frac{1}{B} + \frac{1}{2\ell} \sum_{y=2}^{\ell+1} \left| \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z) (f(\eta) + f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N \right| \end{aligned} \quad (5.28)$$

for some constant $C' > 0$. For the last inequality we used Lemma 5.2. Observe that $B\ell/N = B\varepsilon$ vanishes as $\varepsilon \rightarrow 0$. It remains to estimate the third term on the right hand side of the last inequality. For that purpose we make a similar computation to the one of Lemma 5.5. Let $C_z = \max \left\{ \frac{1}{h(\frac{z}{N})(1-h(\frac{z+1}{N}))}, \frac{1}{h(\frac{z+1}{N})(1-h(\frac{z}{N}))} \right\}$ which is bounded above by a constant depending only on α and β . By using the fact that $\nu_{h(\cdot)}^N$ is product and denoting by $\tilde{\eta}$ the configuration η removing its value at z and $z+1$ so that $(\eta_z, \eta_{z+1}, \tilde{\eta}) = \eta$, we have that

$$\begin{aligned}
& \left| \sum_{z=1}^{y-1} \int (\eta_{z+1} - \eta_z)(f(\eta) + f(\sigma^{z,z+1}\eta)) d\nu_{h(\cdot)}^N \right| \\
&= \sum_{z=1}^{y-1} \left| \sum_{\tilde{\eta}} (f(0, 1, \tilde{\eta}) + f(1, 0, \tilde{\eta})) h(\frac{z+1}{N})(1-h(\frac{z}{N})) \nu_{h(\cdot)}^N(\tilde{\eta}) \right. \\
&\quad \left. - \sum_{\tilde{\eta}} (f(1, 0, \tilde{\eta}) + f(0, 1, \tilde{\eta})) h(\frac{z}{N})(1-h(\frac{z+1}{N})) \nu_{h(\cdot)}^N(\tilde{\eta}) \right| \\
&= \sum_{z=1}^{y-1} \left| \sum_{\tilde{\eta}} \left(h(\frac{z+1}{N}) - h(\frac{z}{N}) \right) (f(0, 1, \tilde{\eta}) + f(1, 0, \tilde{\eta})) \nu_{h(\cdot)}^N(\tilde{\eta}) \right| \\
&\leq \frac{1}{2} \sum_{z=1}^{y-1} C_z \left| h(\frac{z+1}{N}) - h(\frac{z}{N}) \right| \sum_{\tilde{\eta}} \left\{ f(1, 0, \tilde{\eta}) h(\frac{z}{N})(1-h(\frac{z+1}{N})) \nu_{h(\cdot)}^N(\tilde{\eta}) \right. \\
&\quad \left. + f(0, 1, \tilde{\eta}) \left(1 - h(\frac{z}{N}) \right) h(\frac{z+1}{N}) \nu_{h(\cdot)}^N(\tilde{\eta}) \right\} \\
&\lesssim \sum_{z=1}^{y-1} \left| h(\frac{z+1}{N}) - h(\frac{z}{N}) \right|.
\end{aligned}$$

Above, for example, $f(1, 0, \tilde{\eta})$ (resp. $f(0, 1, \tilde{\eta})$) means that we are computing $f(\eta)$ with η such that $\eta_z = 1$ and $\eta_{z+1} = 0$ (resp. $\eta_z = 0$ and $\eta_{z+1} = 1$). Since $h(\cdot)$ is Lipschitz, by (5.28), this estimate provides an upper bound for (5.26) which is in the form of a constant times

$$\frac{B\ell}{N} + \frac{1}{B} + \frac{1}{N\ell} \sum_{y=2}^{\ell+1} y \lesssim B\varepsilon + B^{-1} + \varepsilon$$

which vanishes, as $\varepsilon \rightarrow 0$ and then $B \rightarrow \infty$. This ends the proof. \square

6. ENERGY ESTIMATES

Let \mathbb{Q} be a limit point of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$, whose existence follows from Proposition 4.1 and assume, without loss of generality, that $\{\mathbb{Q}_N\}_{N \geq 1}$ converges to \mathbb{Q} . We note that since our model is an exclusion process, it is standard ([10]) to show that \mathbb{Q} almost surely the trajectories of measures are absolutely continuous with respect to the Lebesgue measure, that is: $\pi_t(dq) = \rho_t(q) dq$ for any $t \in [0, T]$.

6.1. The case $\theta \geq 2 - \gamma$. Recall that in this case the system is speeded up in the diffusive time scale so that $\Theta(N) = N^2$. In this section we prove that the density $\rho_t(q)$ belongs to

$L^2(0, T; \mathcal{H}^1(0, 1))$, see Definition 2.1. For that purpose, we define the linear functional ℓ_ρ on $C_c^{0,1}([0, T] \times (0, 1))$ by

$$\ell_\rho(G) = \int_0^T \int_0^1 \partial_q G_s(q) \rho_s(q) dq ds = \int_0^T \int_0^1 \partial_q G_s(q) d\pi_s(q) ds.$$

By Proposition 6.1 below we have that ℓ_ρ is \mathbb{Q} almost surely continuous, thus we can extend this linear functional to $L^2([0, T] \times (0, 1))$. Moreover, by the Riesz's Representation Theorem we find $\zeta \in L^2([0, T] \times (0, 1))$ such that

$$\ell_\rho(G) = - \int_0^T \int_0^1 G_s(q) \zeta_s(q) dq ds,$$

for all $G \in C_c^{0,1}([0, T] \times (0, 1))$, which implies that $\rho \in L^2(0, T; \mathcal{H}^1(0, 1))$.

Proposition 6.1. *For all $\theta \geq 2 - \gamma$. There exist positive constants C and c such that*

$$\mathbb{E} \left[\sup_G \{ \ell_\rho(G) - c \|G\|_2^2 \} \right] \leq C < \infty,$$

where the supremum above is taken on the set $C_c^{0,1}([0, T] \times (0, 1))$. Here we denote by $\|G\|_2$ the norm of a function $G \in L^2([0, T] \times (0, 1))$.

Proof. By density it is enough to prove Proposition 6.1 for a countable dense subset $\{G^m\}_{m \in \mathbb{N}}$ on $C_c^{0,2}([0, T] \times (0, 1))$ and by Monotone Convergence Theorem it is enough to prove that

$$\mathbb{E} \left[\sup_{k \leq m} \{ \ell_\rho(G^k) - c \|G^k\|_2^2 \} \right] \leq K_0,$$

for any m and for K_0 independent of m . Now, we define $\Phi : \mathcal{D}([0, T], \mathcal{M}^+) \rightarrow \mathbb{R}$ by

$$\Phi(\pi_\cdot) = \max_{k \leq m} \left\{ \int_0^T \int_0^1 \partial_q G_s^k(q) d\pi_s(q) ds - c \|G^k\|_2^2 \right\},$$

which is a continuous and bounded function for the Skorohod topology of $\mathcal{D}([0, T], \mathcal{M}^+)$. Thus we have that

$$\mathbb{E}[\Phi] = \lim_{N \rightarrow \infty} \mathbb{E}_{\mu_N} \left[\max_{k \leq m} \left\{ \int_0^T \frac{1}{N-1} \sum_{x=1}^{N-1} \partial_q G_s^k\left(\frac{x}{N}\right) \eta_x(s) ds - c \|G^k\|_2^2 \right\} \right].$$

By the entropy inequality, Jensen's inequality and the fact that $e^{\max_{k \leq m} a_k} \leq \sum_{k=1}^m e^{a_k}$ the previous display is bounded from above by

$$C_0 + \frac{1}{N} \log \mathbb{E}_{\nu_{h(\cdot)}^N} \left[\sum_{k=1}^m e^{\int_0^T \sum_{x \in \Lambda_N} \partial_q G_s^k\left(\frac{x}{N}\right) \eta_x(s) ds - cN \|G^k\|_2^2} \right],$$

where $\nu_{h(\cdot)}^N$ is the Bernoulli product measure corresponding to a profile $h(\cdot)$ which is Lipschitz such that $\alpha \leq h(\cdot) \leq \beta$, $h(0) = \alpha$, $h(1) = \beta$ and h is $\gamma/2$ -Hölder at the boundaries. In order to deal with the second term in the previous display we use (5.13) and it is enough to bound

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{E}_{\nu_{h(\cdot)}^N} \left[e^{\int_0^T \sum_{x \in \Lambda_N} \partial_q G_s\left(\frac{x}{N}\right) \eta_x(s) ds - cN \|G\|_2^2} \right],$$

for a fixed function $G \in C_c^{0,2}([0, T] \times (0, 1))$, by a constant independent of G . By Feynman-Kac's formula, the expression inside the limsup is bounded from above by

$$\int_0^T \sup_f \left\{ \frac{1}{N} \int_{\Omega_N} \sum_{x \in \Lambda_N} \partial_q G_s(\frac{x}{N}) \eta_x f(\eta) d \nu_{h(\cdot)}^N(\eta) - c \|G\|_2^2 + \frac{\Theta(N)}{N} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \right\} ds$$

where the supremum is carried over all the densities f with respect to $\nu_{h(\cdot)}^N$. Let us now focus on the first term inside braces in the previous expression. Observe first that the space derivative of G_s can be replaced by the discrete gradient $\nabla_N G_s(\frac{x-1}{N}) = N[G_s(\frac{x}{N}) - G_s(\frac{x-1}{N})]$ of G_s with an error $R_N(G)$ satisfying uniformly in N the bound $|R_N(G)| \lesssim 1/N$ since $G \in C_c^{0,2}([0, T], (0, 1))$. By summing and subtracting the term $\nabla_N G_s(\frac{x-1}{N})$ inside the sum, and doing a summation by parts, we can write

$$\frac{1}{N} \int_{\Omega_N} \sum_{x \in \Lambda_N} \partial_q G_s(\frac{x}{N}) \eta_x f(\eta) d \nu_{h(\cdot)}^N(\eta) = \int_{\Omega_N} \sum_{x=1}^{N-2} G_s(\frac{x}{N}) (\eta_x - \eta_{x+1}) f(\eta) d \nu_{h(\cdot)}^N(\eta) + R_N(G).$$

A simple computation shows that we can write the first term at the right hand side of the previous display as

$$\begin{aligned} & \frac{1}{2} \int_{\Omega_N} \sum_{x=1}^{N-2} G_s(\frac{x}{N}) (\eta_x - \eta_{x+1}) (f(\eta) - f(\sigma^{x,x+1} \eta)) d \nu_{h(\cdot)}^N \\ & + \frac{1}{2} \int_{\Omega_N} \sum_{x=1}^{N-2} G_s(\frac{x}{N}) (\eta_x - \eta_{x+1}) f(\sigma^{x,x+1} \eta) (\theta^{x,x+1}(\eta) - 1) d \nu_{h(\cdot)}^N. \end{aligned} \quad (6.1)$$

Recall that for $u, v \geq 0$, $u - v = (\sqrt{u} - \sqrt{v})(\sqrt{u} + \sqrt{v})$ and the inequality $ab \leq \frac{Ba^2}{2} + \frac{b^2}{2B}$ valid for any $B > 0$. Taking $B = \frac{N}{\Theta(N)}$ and using Lemma 5.2 we bound the first term in (6.1) by

$$\begin{aligned} & \frac{N}{4\Theta(N)} \int_{\Omega_N} \sum_{x=1}^{N-2} (G_s(\frac{x}{N}))^2 (\sqrt{f(\eta)} + \sqrt{f(\sigma^{x,x+1} \eta)})^2 d \nu_{h(\cdot)}^N(\eta) \\ & + \frac{\Theta(N)}{4N} \int_{\Omega_N} \sum_{x=1}^{N-2} (\sqrt{f(\eta)} - \sqrt{f(\sigma^{x,x+1} \eta)})^2 d \nu_{h(\cdot)}^N(\eta) \\ & \leq \frac{\Theta(N)}{4N} D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N) + \frac{CN}{\Theta(N)} \sum_{x \in \Lambda_N} (G_s(\frac{x}{N}))^2 \end{aligned}$$

for some $C > 0$. Similarly we can estimate the second term in (6.1) from above by

$$\begin{aligned} & \frac{1}{4N} \int_{\Omega_N} \sum_{x=1}^{N-2} (G_s(\frac{x}{N}))^2 (\eta_x - \eta_{x+1})^2 f(\sigma^{x,x+1} \eta) d \nu_{h(\cdot)}^N(\eta) \\ & + \frac{N}{4} \int_{\Omega_N} \sum_{x=1}^{N-2} f(\sigma^{x,x+1} \eta) (\theta^{x,x+1}(\eta) - 1)^2 d \nu_{h(\cdot)}^N(\eta) \\ & \lesssim \frac{1}{N} \sum_{x \in \Lambda_N} (G_s(\frac{x}{N}))^2 + 1. \end{aligned}$$

We use now (5.6) with $B = 1$ there and observe that last two terms at the right hand side of (5.6) are bounded from above by a constant since $\gamma + \theta - 2 \geq 0$. Observe also

that $D_N^0(\sqrt{f}, \nu_{h(\cdot)}^N) \leq D_N(\sqrt{f}, \nu_{h(\cdot)}^N)$. Recalling that $\Theta(N) = N^2$ we get then that (6.1) is bounded from above by

$$C \int_0^T \left[1 + \frac{1}{N} \sum_{x \in \Lambda_N} (G_s(\frac{x}{N}))^2 \right] ds - c \|G\|_2^2 + R_N(G)$$

where C is a positive constant independent of G . We then choose $c > C$ in order to conclude that

$$\limsup_{N \rightarrow \infty} \left\{ C \int_0^T \left[1 + \frac{1}{N} \sum_{x \in \Lambda_N} (G_s(\frac{x}{N}))^2 \right] ds - c \|G\|_2^2 + R_N(G) \right\} \lesssim 1.$$

This achieves the proof. \square

6.2. The case $\theta < 2 - \gamma$. Recall that in this case the system is speeded up in the time scale $\Theta(N) = N^{\theta+\gamma}$. In this section we prove that the function $(t, q) \rightarrow \rho_t(q) - \alpha$ belongs to $L^2([0, T] \times (0, 1), dt \otimes d\mu)$, where μ is the measure that has the density with respect to the Lebesgue measure given by

$$u \in [0, 1] \rightarrow \frac{1}{u^\gamma}.$$

A similar proof would show that the function $(t, q) \rightarrow \rho_t(q) - \beta$ belongs to $L^2([0, T] \times (0, 1), dt \otimes d\mu')$, where μ' is the measure that has the density with respect to the Lebesgue measure given by

$$u \in [0, 1] \rightarrow \frac{1}{(1-u)^\gamma}.$$

Let $\nu_{h(\cdot)}^N$ be as above, where $h : [0, 1] \rightarrow [0, 1]$ is a profile such that $\alpha \leq h(q) \leq \beta$, for all $q \in [0, 1]$, $h(0) = \alpha$ and $h(1) = \beta$, Hölder of parameter $\gamma/2$ at the boundaries and Lipschitz inside. Let $G \in C_c^{1,\infty}([0, T] \times [0, 1])$. By the entropy inequality and the Feynmann-Kac's formula, we have that

$$\begin{aligned} & \mathbb{E}_{\mu_N} \left(\int_0^T dt N^{\gamma-1} \sum_{x \in \Lambda_N} G(t, \frac{x}{N}) r_N^-(\frac{x}{N}) (\eta_x(tN^{\theta+\gamma}) - \alpha) \right) \\ & \leq C_0 + \int_0^T \sup_f \left\{ N^{\gamma-1} \sum_{x \in \Lambda_N} G(t, \frac{x}{N}) r_N^-(\frac{x}{N}) \langle t_x^\alpha, f \rangle_{\nu_{h(\cdot)}^N} + N^{\gamma+\theta-1} \langle L_N \sqrt{f}, \sqrt{f} \rangle_{\nu_{h(\cdot)}^N} \right\} dt \end{aligned} \quad (6.2)$$

where the supremum is taken over all the densities f on Ω_N with respect to $\nu_{h(\cdot)}^N$. Below C is a constant that may change from line to line. Since the profile is Hölder of parameter $\gamma/2$ at the boundaries and Lipschitz inside, and from (5.6) the term at the right hand side of last expression is bounded from above by

$$-\frac{N^{\gamma+\theta-1}}{4} D_N(\sqrt{f}, \nu_{h(\cdot)}^N) + C + CN^{\theta+\gamma-2}.$$

Repeating the proof of Lemma 5.9 last expression is bounded from above by

$$CN^{\gamma-1} \sum_{x \in \Lambda_N} r_N^-(\frac{x}{N}) G^2(t, \frac{x}{N}) + C + CN^{\theta+\gamma-2}.$$

We take the limit $N \rightarrow \infty$. We conclude that there exist constants $C', C'' > 0$ independent of G such that

$$\mathbb{E} \left[\int_0^T \int_0^1 \frac{(\rho_t(u) - \alpha)G(t, u)}{|u|^\gamma} dudt - C \int_0^T \int_0^1 \frac{G^2(t, u)}{|u|^\gamma} dudt \right] \leq C'. \quad (6.3)$$

By using a similar method as in the proof of the previous lemma we see that the supremum over G can be inserted in the expectation so that

$$\mathbb{E} \left[\sup_G \left\{ \int_0^T \int_0^1 \frac{(\rho_t(u) - \alpha)G(t, u)}{|u|^\gamma} dudt - C \int_0^T \int_0^1 \frac{G^2(t, u)}{|u|^\gamma} dudt \right\} \right] \leq C'. \quad (6.4)$$

The previous formula implies that

$$\mathbb{E} \left[\int_0^T \int_0^1 \frac{(\rho_t(u) - \alpha)^2}{|u|^\gamma} dudt \right] \leq C''.$$

which proves the claim.

7. CHARACTERIZATION OF LIMIT POINTS

We prove in this section that for each range of θ , all limit points \mathbb{Q} of the sequence $\{\mathbb{Q}_N\}_{N \in \mathbb{N}}$ are concentrated on trajectories of measures absolutely continuous with respect to the Lebesgue measure whose density $\rho_t(q)$ is a weak solution of the corresponding hydrodynamic equation. Let \mathbb{Q} be a limit point of the sequence $\{\mathbb{Q}_N\}_{N \geq 1}$, whose existence follows from Proposition 4.1 and assume, without loss of generality, that $\{\mathbb{Q}_N\}_{N \geq 1}$ converges to \mathbb{Q} . As mentioned above, since there is at most one particle per site, it is easy to show that \mathbb{Q} is concentrated on trajectories $\pi_t(dq)$ which are absolutely continuous with respect to the Lebesgue measure, that is, $\pi_t(dq) = \rho_t(q)dq$ (for more details see [10]). Below, we prove, for each range of θ , that the density $\rho_t(q)$ is a weak solution of the corresponding hydrodynamic equation.

Proposition 7.1. *If \mathbb{Q} is a limit point of $\{\mathbb{Q}_N\}_{N \in \mathbb{N}}$ then*

1. *if $\theta < 1$:*

$$\mathbb{Q} \left(\pi. \in \mathcal{D}([0, T], \mathcal{M}^+) : F_{RD}(t, \rho, G, g) = 0, \forall t \in [0, T], \forall G \in C_c^{1,2}([0, T] \times [0, 1]) \right) = 1.$$

2. *if $\theta \in [1, +\infty)$:*

$$\mathbb{Q} \left(\pi. \in \mathcal{D}([0, T], \mathcal{M}^+) : F_{Rob}(t, \rho, G, g) = 0, \forall t \in [0, T], \forall G \in C^{1,2}([0, T] \times [0, 1]) \right) = 1.$$

Remark 7.2. *In this proposition, the constants $\hat{\kappa}, \hat{\sigma}, \hat{m}$ appearing in F_{RD} and F_{Rob} are fixed by Theorem 2.6.*

Proof. Note that in order to prove the proposition, it is enough to verify, for $\delta > 0$ and G in the corresponding space of test functions, that

$$\mathbb{Q} \left(\pi. \in \mathcal{D}([0, T], \mathcal{M}^+) : \sup_{0 \leq t \leq T} |F_\bullet(t, \rho, G, g)| > \delta \right) = 0,$$

for each θ , where F_\bullet stands for F_{RD} if $\theta < 1$ and F_{Rob} if $\theta \geq 1$. From here on, in order to simplify notation, we will erase $\pi.$ from the sets that we have to look at.

- We start with the case $\theta \in [1, \infty)$. Recall $F_{Rob}(t, \rho, G, g)$ from Definition 2.9. Observe that, due to the boundary terms that involve $\rho_s(1)$ and $\rho_s(0)$, the set inside last probability is not an open set in the Skorohod space, therefore we cannot use directly Portmanteau's Theorem as we would like to. In order to avoid this problem, we fix

$\varepsilon > 0$ and we consider two approximations of the identity given by $\iota_\varepsilon^0(q) = \frac{1}{\varepsilon}1_{(0,\varepsilon)}(q)$ and $\iota_\varepsilon^1(q) = \frac{1}{\varepsilon}1_{(1-\varepsilon,1)}(q)$ and we sum and subtract to $\rho_s(0)$ (resp. $\rho_s(1)$) the mean $\langle \pi_s, \iota_\varepsilon^0 \rangle = \frac{1}{\varepsilon} \int_0^\varepsilon \rho_s(q) dq$ (resp. $\langle \pi_s, \iota_\varepsilon^1 \rangle = \frac{1}{\varepsilon} \int_{1-\varepsilon}^1 \rho_s(q) dq$). Thus, we bound last probability from above by the sum of the following four terms

$$\begin{aligned} & \mathbb{Q} \left(\sup_{0 \leq t \leq T} \left| \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 \rho_0(q) G_0(q) dq - \int_0^t \int_0^1 \rho_s(q) \left(\frac{\hat{\sigma}^2}{2} \Delta + \partial_s \right) G_s(q) dq ds \right. \right. \\ & \left. \left. - \int_0^t \langle \pi_s, \iota_\varepsilon^0 \rangle \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(0) - \hat{m} G_s(0) \right) ds + \int_0^t \langle \pi_s, \iota_\varepsilon^1 \rangle \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(1) + \hat{m} G_s(1) \right) ds \right. \right. \\ & \left. \left. - \hat{m} \int_0^t G_s(0) \alpha + G_s(1) \beta ds \right| > \frac{\delta}{4} \right), \end{aligned} \quad (7.1)$$

$$\mathbb{Q} \left(\left| \int_0^1 (\rho_0(q) - g(q)) G_0(q) dq \right| > \frac{\delta}{4} \right), \quad (7.2)$$

$$\mathbb{Q} \left(\sup_{0 \leq t \leq T} \left| \int_0^t [\rho_s(0) - \langle \pi_s, \iota_\varepsilon^0 \rangle] \left[\hat{m} G_s(0) - \frac{\hat{\sigma}^2}{2} \partial_q G_s(0) \right] ds \right| > \frac{\delta}{4} \right), \quad (7.3)$$

and

$$\mathbb{Q} \left(\sup_{0 \leq t \leq T} \left| \int_0^t [\rho_s(1) - \langle \pi_s, \iota_\varepsilon^1 \rangle] \left(\hat{m} G_s(1) + \frac{\hat{\sigma}^2}{2} \partial_q G_s(1) \right) ds \right| > \frac{\delta}{4} \right). \quad (7.4)$$

We note that the terms (7.3) and (7.4) converge to 0 as $\varepsilon \rightarrow 0$ since we are comparing $\rho_s(0)$ (resp. $\rho_s(1)$) with the corresponding average around the boundary points 0 (resp. 1) and (7.2) is equal to zero since \mathbb{Q} is a limit point of $\{\mathbb{Q}_N\}_{N \in \mathbb{N}}$ and \mathbb{Q}_N is induced by μ_N which satisfies (2.11). Therefore it remains only to consider (7.1). We still cannot use Portmanteau's Theorem, since the functions ι_ε^0 and ι_ε^1 are not continuous. Nevertheless, we can approximate each one of these functions by continuous functions in such a way that the error vanishes as $\varepsilon \rightarrow 0$. Then, from Proposition A.3 of [5] we can use Portmanteau's Theorem and bound (7.1) from above by

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \mathbb{Q}_N \left(\sup_{0 \leq t \leq T} \left| \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 \rho_0(q) G_0(q) dq \right. \right. \\ & \left. \left. - \int_0^t \int_0^1 \rho_s(q) \left(\frac{\hat{\sigma}^2}{2} \Delta + \partial_s \right) G_s(q) dq ds \right. \right. \\ & \left. \left. - \int_0^t \langle \pi_s, \iota_\varepsilon^0 \rangle \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(0) - \hat{m} G_s(0) \right) ds + \int_0^t \langle \pi_s, \iota_\varepsilon^1 \rangle \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(1) + \hat{m} G_s(1) \right) ds \right. \right. \\ & \left. \left. - \hat{m} \int_0^t G_s(0) \alpha + G_s(1) \beta ds \right| > \frac{\delta}{2^4} \right). \end{aligned} \quad (7.5)$$

Summing and subtracting $\int_0^t N^2 L_N \langle \pi_s^N, G_s \rangle ds$ to the term inside the supremum in (7.5), recalling (3.1) and (5.18), the definition of \mathbb{Q}_N , we bound (7.5) from above

by the sum of the next two terms

$$\liminf_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} |M_t^N(G)| > \frac{\delta}{2^5} \right), \quad (7.6)$$

and

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t N^2 L_N \langle \pi_s^N, G_s \rangle ds - \frac{\hat{\sigma}^2}{2} \int_0^t \int_0^1 \rho_s(q) \Delta G_s(q) dq ds \right. \right. \\ & - \int_0^t \overrightarrow{\eta}_0^{\varepsilon N}(s) \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(0) - \hat{m} G_s(0) \right) ds + \int_0^t \overleftarrow{\eta}_{N-1}^{\varepsilon N}(s) \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(1) + \hat{m} G_s(1) \right) ds \\ & \left. - \hat{m} \int_0^t G_s(0) \alpha + G_s(1) \beta ds \right| > \frac{\delta}{2^5} \Big). \end{aligned} \quad (7.7)$$

From Doob's inequality together with (4.9), (7.6) goes to 0 as $N \rightarrow \infty$. Finally, (7.7) can be rewritten as

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t N^2 L_N \langle \pi_s^N, G_s \rangle ds - \frac{\hat{\sigma}^2}{2} \int_0^t \langle \pi_s^N, \Delta G_s \rangle ds \right. \right. \\ & - \int_0^t \overrightarrow{\eta}_0^{\varepsilon N}(s) \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(0) - \hat{m} G_s(0) \right) ds + \int_0^t \overleftarrow{\eta}_{N-1}^{\varepsilon N}(s) \left(\frac{\hat{\sigma}^2}{2} \partial_q G_s(1) + \hat{m} G_s(1) \right) ds \\ & \left. - \hat{m} \int_0^t G_s(0) \alpha + G_s(1) \beta ds \right| > \frac{\delta}{2^5} \Big). \end{aligned} \quad (7.8)$$

Now, from (3.2) and (3.4) we can bound from above the probability in (7.8) by the sum of the five following terms

$$\mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \frac{N^2}{N-1} \int_0^t \sum_{x \in \Lambda_N} K_N G_s\left(\frac{x}{N}\right) \eta_x(sN^2) ds - \frac{\hat{\sigma}^2}{2} \int_0^t \langle \pi_s^N, \Delta G_s \rangle ds \right| > \frac{\delta}{2^6} \right), \quad (7.9)$$

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \frac{N^2}{N-1} \int_0^t \sum_{x \in \Lambda_N} \sum_{y \leq 0} [G_s\left(\frac{y}{N}\right) - G_s\left(\frac{x}{N}\right)] p(x-y) \eta_x(sN^2) ds \right. \right. \\ & \left. \left. + \frac{\hat{\sigma}^2}{2} \int_0^t \overrightarrow{\eta}_0^{\varepsilon N}(sN^2) \partial_q G_s(0) ds \right| > \frac{\delta}{2^6} \right), \end{aligned} \quad (7.10)$$

and

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \frac{N\kappa}{N-1} \sum_{x \in \Lambda_N} (G_s r_N^-)\left(\frac{x}{N}\right) (\alpha - \eta_x(sN^2)) ds \right. \right. \\ & \left. \left. - m\kappa \int_0^t G_s(0) (\alpha - \overrightarrow{\eta}_0^{\varepsilon N}(sN^2)) ds \right| > \frac{\delta}{2^6} \right) \end{aligned} \quad (7.11)$$

and the sum of two terms which are very similar to the two previous ones but which are concerned with the right boundary. Thus, to conclude we have to show that these five terms go to 0. Applying Lemma 3.2 and noting that $|\eta_x(sN^2)| \leq 1$ for any x and any $s \geq 0$, we conclude that (7.9) goes to 0 as $N \rightarrow \infty$. Note also that by Taylor expansion, we can bound from above (7.10) by

$$\mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \partial_q G_s(0) \sum_{x \in \Lambda_N} \Theta_x^- [\eta_x(sN^2) - \overrightarrow{\eta}_0^{\varepsilon N}(sN^2)] ds \right| > \frac{\delta}{2^8} \right). \quad (7.12)$$

Using Lemma 5.7 we see that (7.12) vanishes as $N \rightarrow \infty$. Now we look at (7.11) and we prove that it vanishes as $N \rightarrow \infty$. Performing a Taylor expansion on G_s at 0 and using (3.7) the probability in (7.11) is bounded from above by

$$\mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t G_s(0) \sum_{x \in \Lambda_N} r_N^-\left(\frac{x}{N}\right) [\vec{\eta}_0^{\varepsilon N}(sN^2) - \eta_x(sN^2)] ds \right| > \frac{\delta}{2^8} \right),$$

plus lower-order terms (with respect to N). From Lemma 5.7 and Remark 5.8 last display vanishes as $N \rightarrow \infty$. Similarly the two terms which are similar to (7.10) and (7.11) but which are concerned with the right boundary vanish as $N \rightarrow \infty$. Thus the proof is finished.

• Now we treat the case $\theta < 1$. We have to prove that

$$\mathbb{Q} \left(\pi. \in \mathcal{D}([0, T], \mathcal{M}^+) : \sup_{0 \leq t \leq T} |F_{RD}(t, \rho, G, g)| > \delta \right) = 0$$

for any $G \in C_c^{1,2}([0, T] \times [0, 1])$. We can bound from above the previous probability by

$$\begin{aligned} & \mathbb{Q} \left(\sup_{0 \leq t \leq T} \left| \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 \rho_0(q) G_0(q) dq - \int_0^t \int_0^1 \rho_s(q) \left(\frac{\partial^2}{2} \Delta + \partial_s \right) G_s(q) dq ds \right. \right. \\ & \left. \left. - \hat{\kappa} \int_0^t \int_0^1 G_s(q) V_0(q) dq ds + \hat{\kappa} \int_0^t \int_0^1 G_s(q) \rho_s(q) V_1(q) dq ds \right| > \frac{\delta}{2} \right), \end{aligned} \quad (7.13)$$

and

$$\mathbb{Q} \left(\left| \int_0^1 (\rho_0(q) - g(q)) G_0(q) dq \right| > \frac{\delta}{2} \right), \quad (7.14)$$

where $V_0(q) = \frac{\alpha}{q^\gamma} + \frac{\beta}{(1-q)^\gamma}$ and $V_1(q) = \frac{1}{q^\gamma} + \frac{1}{(1-q)^\gamma}$. We note that (7.14) is equal to zero since \mathbb{Q} is a limit point of $\{\mathbb{Q}_N\}_{N \in \mathbb{N}}$ and \mathbb{Q}_N is induced by μ_N which satisfies (2.11). We note that from Proposition A.3 of [5], the set inside the probability in (7.13) is an open set in the Skorohod space (the singularities of V_0 and V_1 are not present because G_s has compact support). From Portmanteau's Theorem we bound (7.13) from above by

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \mathbb{Q}_N \left(\sup_{0 \leq t \leq T} \left| \int_0^1 \rho_t(q) G_t(q) dq - \int_0^1 \rho_0(q) G_0(q) dq \right. \right. \\ & \left. \left. - \int_0^t \int_0^1 \rho_s(q) \left(\frac{\partial^2}{2} \Delta + \partial_s \right) G_s(q) dq ds - \hat{\kappa} \int_0^t \int_0^1 G_s(q) V_0(q) dq ds \right. \right. \\ & \left. \left. + \hat{\kappa} \int_0^t \int_0^1 G_s(q) \rho_s(q) V_1(q) dq ds \right| > \frac{\delta}{2} \right). \end{aligned}$$

Summing and subtracting $\int_0^t \Theta(N) L_N \langle \pi_s^N, G_s \rangle ds$ to the term inside the previous absolute values, recalling (3.1) and the definition of \mathbb{Q}_N , we can bound the previous probability from above by the sum of the next two terms

$$\mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} |M_t^N(G)| > \frac{\delta}{4} \right),$$

and

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \Theta(N) L_N \langle \pi_s^N, G_s \rangle ds - \int_0^t \langle \pi_s^N, \frac{\sigma^2}{2} \Delta G_s \rangle ds \right. \right. \\ & \left. \left. - \hat{\kappa} \int_0^t \int_0^1 G_s(q) V_0(q) dq ds + \hat{\kappa} \int_0^t \int_0^1 G_s(q) \rho_s(q) V_1(q) dq ds \right| > \frac{\delta}{4} \right). \end{aligned} \quad (7.15)$$

The first term above can be estimated as in the case $\theta \geq 1$ and it vanishes as $N \rightarrow \infty$. It remains to prove that (7.15) vanishes as $N \rightarrow \infty$. For that purpose, we recall (3.6) and we use (3.2), (3.4) to bound it from above by the sum of the following terms

$$\mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \frac{\Theta(N)}{N-1} \sum_{x \in \Lambda_N} K_N G_s(\frac{x}{N}) \eta_x(sN^2) ds - \frac{\hat{\sigma}^2}{2} \int_0^t \langle \pi_s^N, \Delta G_s \rangle ds \right| > \frac{\delta}{24} \right), \quad (7.16)$$

and

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \left\{ \frac{\kappa \Theta(N)}{(N-1)N^\theta} \sum_{x \in \Lambda_N} (G_s r_N^-)(\frac{x}{N}) (\alpha - \eta_x(sN^2)) \right. \right. \right. \\ & \left. \left. \left. - \hat{\kappa} \int_0^1 (G_s r^-)(q) (\alpha - \rho_s(q)) dq \right\} ds \right| > \frac{\delta}{24} \right), \end{aligned} \quad (7.17)$$

and

$$\begin{aligned} & \mathbb{P}_{\mu_N} \left(\sup_{0 \leq t \leq T} \left| \int_0^t \left\{ \frac{\kappa \Theta(N)}{(N-1)N^\theta} \sum_{x \in \Lambda_N} (G_s r_N^+)(\frac{x}{N}) (\beta - \eta_x(sN^2)) \right. \right. \right. \\ & \left. \left. \left. - \hat{\kappa} \int_0^1 (G_s r^+)(q) (\beta - \rho_s(q)) dq \right\} ds \right| > \frac{\delta}{24} \right), \end{aligned} \quad (7.18)$$

In the case $\theta \in [2 - \gamma, 1)$, since $\Theta(N) = N^2$ and $\hat{\sigma} = \sigma$, from Lemma 3.2 we have that (7.16) goes to 0 as $N \rightarrow \infty$. In the case $\theta < 2 - \gamma$, since $\Theta(N) = N^{\theta + \gamma}$ and $\hat{\sigma} = 0$, from Lemma 3.2 we also have that (7.16) goes to 0 as $N \rightarrow \infty$.

Now we analyze the boundary terms (7.17) and (7.18). Note that in the case $\theta \in (2 - \gamma, 1)$ we have $\theta(N) = N^2$ and $\hat{\kappa} = 0$, so that the two previous probabilities vanish, as $N \rightarrow \infty$, as a consequence of Lemma 5.6. In the case $\theta \leq 2 - \gamma$, since $\Theta(N) = N^{\gamma + \theta}$, $\hat{\kappa} = \kappa_\gamma \gamma^{-1}$, $|\eta_x(sN^2)| \leq 1$, in order to conclude it is enough to note that since G_s has compact support in $(0, 1)$ we know by (3.6) that $N^\gamma G_s r_N^-(q)$ and $N^\gamma G_s r_N^+(q)$ converge uniformly to $(G_s r^-)(q)$ and $(G_s r^+)(q)$, respectively, as $N \rightarrow \infty$. This ends the proof. \square

ACKNOWLEDGEMENTS

This work has been supported by the French Ministry of Education through the grant ANR (EDNHS) and the grant ANR (LSD).

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovative programme (grant agreement No 715734).

APPENDIX A. UNIQUENESS OF WEAK SOLUTIONS

The uniqueness of the weak solutions of the partial equations given in Section 2.2 is fundamental for the proof of the hydrodynamic limit. The uniqueness of weak solutions

of (2.6) is standard if $\hat{\kappa} = 0$. Since we were not able to find in the literature a proof in the case $\hat{\kappa} > 0$ we give a complete proof below. The proof of uniqueness of weak solutions of (2.8) can be found in, for example, [1].

Now we prove the uniqueness of weak solutions of (2.6). We assume that $\hat{\sigma} > 0$ and $\hat{\kappa} > 0$ first and then we consider the case $\hat{\sigma} = 0$ and $\hat{\kappa} > 0$.

Let ρ^1 and ρ^2 be two weak solutions of (2.2) with the same initial condition and let us denote $\bar{\rho} = \rho^1 - \rho^2$. By assumption we have that

$$\bar{\rho} \in L^2(0, T; \mathcal{H}^1) \cap L^2(0, T; L^2((0, 1); V_1(q) dq))$$

where $V_1(q) = q^{-\gamma} + (1-q)^{-\gamma}$. Let us denote by $\langle \cdot, \cdot \rangle_{V_1}$ (resp. $\| \cdot \|_{V_1}$) the scalar product (resp. the norm) corresponding to the Hilbert space $L^2((0, 1), V_1(q) dq)$.

For almost every $t \in [0, T]$, we identify $\bar{\rho}_t$ with its continuous representation in $[0, 1]$. Therefore, from 3. in Definition 2.2, we have that $\bar{\rho}_t(0) = \bar{\rho}_t(1) = 0$ for all $t \in [0, T]$. Since \mathcal{H}_0^1 is equal to the set of functions in \mathcal{H}^1 vanishing at 0 and 1 we have that for a.e. time $t \in [0, T]$, $\bar{\rho}_t \in \mathcal{H}_0^1$ and in fact $\bar{\rho} \in L^2(0, T; \mathcal{H}_0^1)$. From 2. in Definition 2.2, for any $t \in [0, T]$ and any $G \in C_c^{1,2}([0, T] \times [0, 1])$ we have

$$\begin{aligned} & \int_0^1 \bar{\rho}_t(q) G_t(q) dq - \int_0^t \int_0^1 \bar{\rho}_s(q) \left(\partial_s + \frac{\hat{\sigma}^2}{2} \Delta \right) G_s(q) dq ds \\ & + \hat{\kappa} \int_0^T \int_0^1 V_1(q) G_s(q) \bar{\rho}_s(q) dq ds = 0. \end{aligned} \quad (\text{A.1})$$

We know that $C_c^{1,\infty}([0, T] \times (0, 1))$ is dense in $L^2(0, T; \mathcal{H}_0^1) \cap L^2(0, T; L^2((0, 1); V(q) dq))$. Therefore, let $(H_n)_{n \geq 0}$ be a sequence of functions in $C_c^{1,\infty}([0, T] \times (0, 1))$ converging to $\bar{\rho}$ with respect to the norms of $L^2(0, T; \mathcal{H}_0^1)$ and $L^2(0, T; L^2((0, 1); V_1(q) dq))$. We define G_n in $C_c^{1,\infty}([0, T] \times [0, 1])$ by

$$\forall t \in [0, T], \quad \forall q \in [0, 1], \quad G_n(t, q) = \int_t^T H_n(s, q) ds. \quad (\text{A.2})$$

Plugging G_n into (A.1) and letting $n \rightarrow \infty$ we conclude, by Lemma A.1 below, that

$$\int_0^T \int_0^1 \bar{\rho}_s^2(q) dq ds + \frac{\hat{\sigma}^2}{4} \left\| \int_0^T \bar{\rho}_s ds \right\|_1^2 + \frac{\hat{\kappa}}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_{V_1}^2 = 0.$$

It follows that for almost every time $s \in [0, T]$ the continuous function $\bar{\rho}_s$ is equal to 0 and we conclude the uniqueness of weak solution to (2.6) in the case $\hat{\sigma} > 0$.

Lemma A.1. *Let $(G_n)_{n \geq 0}$ be defined as in (A.2). We have*

- i) $\lim_{n \rightarrow \infty} \int_0^T \int_0^1 \bar{\rho}_s(q) (\partial_s G_n)(s, q) dq ds = - \int_0^T \int_0^1 \bar{\rho}_s^2(q) dq ds.$
- ii) $\lim_{n \rightarrow \infty} \int_0^T \int_0^1 \bar{\rho}_s(q) \Delta G_n(s, q) dq ds = - \frac{1}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_1^2.$
- iii) $\lim_{n \rightarrow \infty} \int_0^T \int_0^1 V_1(q) G_n(s, q) \bar{\rho}_s(q) dq ds = \frac{1}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_{V_1}^2 < \infty.$

Proof. For i) we write

$$\begin{aligned} & - \int_0^T \int_0^1 \bar{\rho}_s(q) (\partial_s G_n)(s, q) dq ds = \int_0^T \int_0^1 \bar{\rho}_s(q) H_n(s, q) dq ds = \int_0^T \langle \bar{\rho}_s, H_n(s, \cdot) \rangle ds \\ & = \int_0^T \langle \bar{\rho}_s, H_n(s, \cdot) - \bar{\rho}_s \rangle ds + \int_0^T \|\bar{\rho}_s\|_{L^2}^2 ds. \end{aligned}$$

Observe then that by Cauchy-Schwarz inequality we have

$$\begin{aligned} & \left| \int_0^T \langle \bar{\rho}_s, H_n(s, \cdot) - \bar{\rho}_s \rangle ds \right| \leq \int_0^T \|\bar{\rho}_s\|_{L^2} \|H_n(s, \cdot) - \bar{\rho}_s\|_{L^2} ds \\ & \leq \sqrt{\int_0^T \|\bar{\rho}_s\|_{L^2}^2 ds} \sqrt{\int_0^T \|H_n(s, \cdot) - \bar{\rho}_s\|_{L^2}^2 ds} \end{aligned} \tag{A.3}$$

which goes to 0 as $n \rightarrow \infty$. Above we have used the fact that $(H_n)_{n \geq 0}$ converges to $\bar{\rho}$ as $N \rightarrow \infty$ with respect to the norm of $L^2(0, T; \mathcal{H}_0^1)$.

For ii) we first use the integration by parts formula for \mathcal{H}_1 functions which permits to write

$$\int_0^T \int_0^1 \bar{\rho}_s(q) \Delta G_n(s, q) dq ds = - \int_0^T \langle \bar{\rho}_s, G_n(s, \cdot) \rangle_1 ds.$$

Then we have

$$\begin{aligned} & \int_0^T \langle \bar{\rho}_s, G_n(s, \cdot) \rangle_1 ds = \int_0^T \langle \bar{\rho}_s, \int_s^T \bar{\rho}_u du \rangle_1 ds + \int_0^T \langle \bar{\rho}_s, G_n(s, \cdot) - \int_s^T \bar{\rho}_u du \rangle_1 ds \\ & = \iint_{0 \leq s < u \leq T} \langle \bar{\rho}_s, \bar{\rho}_u \rangle_1 du ds + \int_0^T \langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \rangle_1 ds \\ & = \frac{1}{2} \iint_{[0, T]^2} \langle \bar{\rho}_s, \bar{\rho}_u \rangle_1 du ds + \int_0^T \langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \rangle_1 ds \\ & = \frac{1}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_1^2 + \int_0^T \langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \rangle_1 ds. \end{aligned}$$

To conclude the proof of ii) it is sufficient to prove that

$$\lim_{n \rightarrow \infty} \int_0^T \langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \rangle_1 ds = 0.$$

This is a consequence of a successive use of Cauchy-Schwarz inequalities:

$$\begin{aligned} & \left| \int_0^T \langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \rangle_1 ds \right| \leq \int_0^T \|\bar{\rho}_s\|_1 \left\| \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\|_1 ds \\ & \leq \int_0^T \|\bar{\rho}_s\|_1 \int_s^T \|H_n(u, \cdot) - \bar{\rho}_u\|_1 du ds \\ & \leq \left(\int_0^T \|\bar{\rho}_s\|_1 ds \right) \left(\int_0^T \|H_n(u, \cdot) - \bar{\rho}_u\|_1 du \right) \\ & \leq T \sqrt{\int_0^T \|\bar{\rho}_s\|_1^2 ds} \sqrt{\int_0^T \|H_n(u, \cdot) - \bar{\rho}_u\|_1^2 du} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Above we have used again the fact that $(H_n)_{n \geq 0}$ converges to $\bar{\rho}$ as $N \rightarrow \infty$ with respect to the norm of $L^2(0, T; \mathcal{H}_0^1)$.

The proof of iii) is similar. We have

$$\begin{aligned}
& \int_0^T \left\langle \bar{\rho}_s, G_n(s, \cdot) \right\rangle_{V_1} ds = \int_0^T \left\langle \bar{\rho}_s, \int_s^T \bar{\rho}_u du \right\rangle_{V_1} ds + \int_0^T \left\langle \bar{\rho}_s, G_n(s, \cdot) - \int_s^T \bar{\rho}_u du \right\rangle_{V_1} ds \\
& = \int \int_{0 \leq s < u \leq T} \langle \bar{\rho}_s, \bar{\rho}_u \rangle_{V_1} du ds + \int_0^T \left\langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\rangle_{V_1} ds \\
& = \frac{1}{2} \int \int_{[0, T]^2} \langle \bar{\rho}_s, \bar{\rho}_u \rangle_{V_1} du ds + \int_0^T \left\langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\rangle_{V_1} ds \\
& = \frac{1}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_{V_1}^2 + \int_0^T \left\langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\rangle_{V_1} ds.
\end{aligned}$$

To conclude the proof of iii) it is sufficient to prove that

$$\lim_{n \rightarrow \infty} \int_0^T \left\langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\rangle_{V_1} ds = 0.$$

This is a consequence of the Cauchy-Schwarz inequality:

$$\begin{aligned}
& \left| \int_0^T \left\langle \bar{\rho}_s, \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\rangle_{V_1} ds \right| \leq \int_0^T \|\bar{\rho}_s\|_{V_1} \left\| \int_s^T \{H_n(u, \cdot) - \bar{\rho}_u\} du \right\|_{V_1} ds \\
& \leq \int_0^T \|\bar{\rho}_s\|_{V_1} \int_s^T \|H_n(u, \cdot) - \bar{\rho}_u\|_{V_1} du ds \\
& \leq \left(\int_0^T \|\bar{\rho}_s\|_{V_1}^2 ds \right) \left(\int_0^T \|H_n(u, \cdot) - \bar{\rho}_u\|_{V_1}^2 du \right) \\
& \leq T \sqrt{\int_0^T \|\bar{\rho}_s\|_{V_1}^2 ds} \sqrt{\int_0^T \|H_n(u, \cdot) - \bar{\rho}_u\|_{V_1}^2 du} \xrightarrow{n \rightarrow \infty} 0.
\end{aligned}$$

□

Now we look at the case $\hat{\sigma} = 0$. In this case we do not have any regularity assumption on $\bar{\rho}(\cdot)$. However, it can be proved that

$$\int_0^T \int_0^1 \bar{\rho}_s^2(q) dq ds + \frac{\hat{\kappa}}{2} \left\| \int_0^T \bar{\rho}_s ds \right\|_{V_1}^2 = 0 \tag{A.4}$$

holds by showing only the first and third item of the previous lemma. This requires only the density of $C_c^{1, \infty}([0, T] \times (0, 1))$ in $L^2(0, T; L^2((0, 1); V_1(q) dq))$. We also note that in the proof of item i) in Lemma A.1, in order to conclude the convergence in (A.3), before applying the Cauchy-Schwarz inequality, we multiply and divide the integrand function by V_1 and since V_1^{-1} is bounded we get that $\|\bar{\rho}_s V_1^{-1}\|_{L^2}^2 < \infty$ and the result follows.

APPENDIX B. COMPUTATIONS INVOLVING THE GENERATOR

Lemma B.1. For any $x \neq y \in \Lambda_N$, we have

$$\begin{aligned} L_N^0(\eta_x \eta_y) &= \eta_x L_N^0 \eta_y + \eta_y L_N^0 \eta_x - p(y-x)(\eta_y - \eta_x)^2, \\ L_N^r(\eta_x \eta_y) &= \eta_x L_N^r \eta_y + \eta_y L_N^r \eta_x, \\ L_N^\ell(\eta_x \eta_y) &= \eta_x L_N^\ell \eta_y + \eta_y L_N^\ell \eta_x. \end{aligned} \tag{B.1}$$

Proof. By definition of L_N^0 we have that

$$\begin{aligned} L_N^0(\eta_x \eta_y) &= \frac{1}{2} \sum_{u,v \in \Lambda_N} p(v-u) [(\sigma^{u,v} \eta_j)(\sigma^{u,v} \eta_y) - \eta_x \eta_y] \\ &= \frac{1}{2} \sum_{u,v \in \Lambda_N} p(v-u) [((\sigma^{u,v} \eta_x) \eta_y - \eta_x \eta_y) + ((\sigma^{u,v} \eta_k) \eta_j - \eta_j \eta_k) + \\ &\quad + (\sigma^{u,v} \eta_x)(\sigma^{u,v} \eta_y) - (\sigma^{u,v} \eta_x) \eta_y - (\sigma^{u,v} \eta_y) \eta_x + \eta_x \eta_y] \\ &= \eta_x L_N^0 \eta_y + \eta_y L_N^0 \eta_x + \frac{1}{2} \sum_{u,v \in \Lambda_N} p(v-u) [(\sigma^{u,v} \eta_x) - \eta_x] [(\sigma^{u,v} \eta_y) - \eta_y] \\ &= \eta_x L_N^0 \eta_y + \eta_y L_N^0 \eta_x - p(y-x)(\eta_y - \eta_x)^2. \end{aligned}$$

In order to prove the second expression, note that $[(\sigma^u \eta_x) - \eta_x][(\sigma^u \eta_y) - \eta_y] = 0$, for all $u \in \mathbb{Z}$, thus by definition of L_N^r we have

$$\begin{aligned} L_N^r(\eta_x \eta_y) &= \sum_{u \in \Lambda_N, v \geq N} p(v-u) [\eta_u(1-\beta) + (1-\eta_u)\beta] [(\sigma^u(\eta_x \eta_y)) - \eta_x \eta_y] \\ &= \eta_x L_N^r \eta_y + \eta_y L_N^r \eta_x + \\ &\quad + \sum_{u \in \Lambda_N, v \geq N} p(v-u) [\eta_u(1-\beta) + (1-\eta_u)\beta] [(\sigma^u \eta_x) - \eta_x] [(\sigma^u \eta_y) - \eta_y] \\ &= \eta_x L_N^r \eta_y + \eta_y L_N^r \eta_x. \end{aligned}$$

The proof of the third expression is analogous. \square

REFERENCES

- [1] A. Neumann, O. Menezes, R. Baldasso and R. Souza. Exclusion process with slow boundary, arXiv:1407.7918 (2014)
- [2] C. Bernardin, P. Gonçalves and S. Sethuraman. Occupation times of long-range exclusion and connections to KPZ class exponents, Prob. Theory Rel. Fields, DOI 10.1007/s00440-015-0661-5 (2015).
- [3] A. Dhar, Heat Transport in low-dimensional systems, Adv. Phys., **57**, 457 (2008).
- [4] A. Dhar and K. Saito. Anomalous transport and current fluctuations in a model of diffusing Levy walkers, eprint arXiv:1308.5476 (2013).
- [5] Franco, T., Gonçalves, P., Neumann, A. Hydrodynamical behavior of symmetric exclusion with slow bonds, Annales de l'Institut Henri Poincaré: Probability and Statistics, Volume 49, No. 2, 402–427 (2013).
- [6] Franco, T., Gonçalves, P., Schutz, G.: Scaling limits for the exclusion process with a slow site, Stochastic Processes and their applications, Volume 126, Issue 3, 800 – 831 (2016).
- [7] M. Jara. Hydrodynamic limit of particle systems with long jumps, arXiv:0805.1326 (2008).
- [8] M. Jara. Nonequilibrium scaling limit for a tagged particle in the simple exclusion process with long jumps. Comm. Pure Appl. Math. **62**, no. 2, 198–214 (2009).
- [9] B. Jimenez Oviedo, A. Vasseur. In preparation (2017).
- [10] C. Kipnis and C. Landim. Scaling Limits of Interacting Particle Systems, Springer-Verlag, New York (1999).
- [11] S. Lepri and A. Politi. Density profiles in open superdiffusive systems, Phys. Rev. E **83**, 030107(R) (2011).

- [12] T. Liggett. *Interacting Particles Systems*, Springer, New York (1985).
- [13] T. Liggett. *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*, Springer, New York (1999).
- [14] F. Spitzer. Interaction of Markov processes. *Advances in Math.* **5**, 2, 246–290 (1970).
- [15] H. Spohn. *Large scale Dynamics of Interacting Particles*, Springer-Verlag, Berlin (1991).
- [16] J. Szavits-Nossan and K. Uzelac. Scaling properties of the asymmetric exclusion process with long-range hopping, *Phys. Rev. E* **77**, 051116 (2008).

UNIVERSITÉ CÔTE D'AZUR, CNRS, LJAD, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE
E-mail address: `cbernard@unice.fr`

CENTER FOR MATHEMATICAL ANALYSIS, GEOMETRY AND DYNAMICAL SYSTEMS, INSTITUTO SUPERIOR TÉCNICO, UNIVERSIDADE DE LISBOA, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL
E-mail address: `patricia.goncalves@math.tecnico.ulisboa.pt`

UNIVERSITÉ CÔTE D'AZUR, CNRS, LJAD, PARC VALROSE, 06108 NICE CEDEX 02, FRANCE
E-mail address: `byron@unice.fr`