



Anaerococcus urinomassiliensis sp. nov., isolated from a urine sample of a 17-year-old boy affected by autoimmune hepatitis and membranoproliferative glomerulonephritis

A Morand, F Cornu, Michel Tsimaratos, Jean-Christophe Lagier, Frédéric Cadoret, Pierre-Edouard Fournier, Didier Raoult

► To cite this version:

A Morand, F Cornu, Michel Tsimaratos, Jean-Christophe Lagier, Frédéric Cadoret, et al.. Anaerococcus urinomassiliensis sp. nov., isolated from a urine sample of a 17-year-old boy affected by autoimmune hepatitis and membranoproliferative glomerulonephritis. *New Microbes and New Infections*, 2016, 13, pp.56-58. 10.1016/j_nmni.2016.06.001 . hal-01476131

HAL Id: hal-01476131

<https://hal.science/hal-01476131v1>

Submitted on 3 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Anaerococcus urinomassiliensis sp. nov., isolated from a urine sample of a 17-year-old boy affected by autoimmune hepatitis and membranoproliferative glomerulonephritis

A. Morand^{1,2}, F. Cornu³, M. Tsimaratos³, J.-C. Lagier^{1,2}, F. Cadoret^{1,2}, P.-E. Fournier^{1,2} and D. Raoult^{1,2}

1) Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), 2) Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, Aix-Marseille Université and 3) Pédiatrie Multidisciplinaire—Hôpital de la Timone, Marseille, France

Abstract

We report the main characteristics of '*Anaerococcus urinomassiliensis*' strain FC4^T (CSURP2143) that was isolated from a urine sample of a 17-year-old boy affected by autoimmune hepatitis and membranoproliferative glomerulonephritis.

© 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: *Anaerococcus urinomassiliensis*, culturomics, genomics, taxono-genomics, taxonomy

Original Submission: 31 May 2016; **Accepted:** 1 June 2016

Article published online: 6 June 2016

Corresponding author: D. Raoult, URMITE, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), Faculté de Médecine, Aix-Marseille Université, 27 Boulevard Jean Moulin, 13385 Marseille Cedex 5, France

E-mail: didier.raoult@gmail.com

In 2015, as part of a culturomics study [1,2] of the human microbiome, a bacterial strain that could not be identified by matrix-assisted laser desorption-ionization time-of-flight mass spectrometry (MALDI-TOF MS) screening using a Microflex spectrometer (Bruker Daltonics, Bremen, Germany) [3] was isolated from the urine of a 17-year-old boy affected by autoimmune hepatitis and membranoproliferative glomerulonephritis. The patient's parents gave signed informed consent and the ethics committee of the Institut Federatif de Recherche IFR48 approved the study under number 09-022.

The initial growth was obtained after 10 days of incubation in an anaerobic blood culture vial (Becton Dickinson, Le Pont-de-Claix, France) supplemented with 5 mL of 0.2-μm filtered rumen fluid. A pure culture of strain FC4 was then obtained after 48 h of incubation at 37°C on 5% sheep blood–Columbia agar medium (bioMérieux, Marcy l'Etoile, France) in anaerobic atmosphere generated using the GENbag Anaer system (bioMérieux). Agar-grown colonies were small with a mean

diameter of 50 μm and were translucent white. Bacterial cells were Gram-positive cocci with a diameter ranging from 400 to 600 nm. Strain FC4 did not exhibit catalase or oxidase activities. The 16S rRNA gene was sequenced using the fD1-rP2 primers as previously described [4], using a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France). Strain FC4 exhibited sequence similarities with *Anaerococcus octavius* strain NCTC 98 IOT (GenBank accession number NR-026360) [5], its phylogenetically 94.73% closest species with standing nomenclature in the range of 92.3% to 97.2% observed among *Anaerococcus* species (Fig. 1). This putatively classifies strain FC4 as a member of the genus *Anaerococcus* in the family Peptostreptococcaceae within the phylum Firmicutes.

Strain FC4 presents a 16S rRNA gene sequence divergence with its phylogenetically closest species with standing in nomenclature [6], so we propose the creation of the new species '*Anaerococcus urinomassiliensis*' sp. nov. strain FC4^T as the type strain of '*Anaerococcus urinomassiliensis*' (u.ri.no.-mas.sil.i.en'sis composed of u.ri.no L. V. intransitive. of *urino*, the Latin name for the verb to swim, closed of u.ri.na. L. N. gen. fem. *urina*, of *urina*, the Latin name of urine as this strain FC4 was first found in a paediatric urine sample and mas.si.li.en'sis. L. fem. adj. *massiliensis*, of Massilia, the ancient Greek and Roman name for Marseille, France, where the type strain was isolated).




FIG. 1. Phylogenetic tree showing the position of '*Anaerococcus urinomassiliensis*' strain FC4 relative to other phylogenetically close neighbours. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using the maximum-likelihood method within the MEGA software. Numbers at the nodes are percentages of bootstrap values ($\geq 95\%$) obtained by repeating the analysis 500 times to generate a majority consensus tree. GenBank accession numbers are indicated in parentheses. The scale bar indicates a 2% nucleotide sequence divergence.

MALDI-TOF Spectrum

MALDI-TOF spectrum of '*Anaerococcus urinomassiliensis*' strain FC4^T is available at <http://mediterranee-infection.com/article.php?laref=256&titre=urms-database>.

Nucleotide Sequence Accession Number

The 16S r RNA gene sequence was deposited in GenBank under Accession number LN898272.

Deposit in a Culture Collection

Strain FC4^T was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR, WDCM 875) under number P2143.

Funding

This work was funded by Méditerranée-Infection Foundation.

Conflict of Interest

The authors have no conflicts of interest to declare.

References

- [1] Lagier JC, Hugon P, Khelaifia S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. Clin Microbiol Rev 2015 Jan;28(1):237–64.
- [2] Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, et al. Urine Is Not Sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. J Clin Microbiol 2014 Mar;52(3):871–6.

- [3] Seng P, Abat C, Rolain JM, Colson P, Lagier JC, Gouriet F, et al. Identification of rare pathogenic bacteria in a clinical microbiology laboratory: impact of matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J Clin Microbiol* 2013 Jul;51(7): 2182–94.
- [4] Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D, et al. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol* 2000 Oct;38(10):3623–30.
- [5] Ezaki T, Kawamura Y, Li N, Li ZY, Zhao L, Shu S, et al. Proposal of the genera *Anaerococcus* gen. nov., *Peptoniphilus* gen. nov. and *Gallicola* gen. nov. for members of the genus *Peptostreptococcus*. *Int J Syst Evol Microbiol* 2001 Jul;51(Pt 4):1521–8.
- [6] Kim M, Oh HS, Park SC, Chun J. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* 2014 Feb;64(Pt 2):346–51. Erratum in: *Int J Syst Evol Microbiol*. 2014 May;64(Pt 5):1825. PubMed PMID: 24505072.