



HAL
open science

Quantum Gravity Effects around Sagittarius A

Hal M. Haggard, Carlo Rovelli

► **To cite this version:**

| Hal M. Haggard, Carlo Rovelli. Quantum Gravity Effects around Sagittarius A. 2024. hal-01476558

HAL Id: hal-01476558

<https://hal.science/hal-01476558>

Preprint submitted on 2 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantum Gravity Effects around Sagittarius A*

Hal M. Haggard*

*Physics Program, Bard College, 30 Campus Rd,
Annandale-on-Hudson, NY 12504, USA.*

Carlo Rovelli†

*CPT, Aix-Marseille Université, Université de Toulon,
CNRS, Case 907, F-13288 Marseille, France
and Samy Maroun Research Center for Time, Space and the Quantum.*

Abstract

Recent VLBI observations have resolved Sagittarius A* at horizon scales. The Event Horizon Telescope is expected to provide increasingly good images of the region around the Schwarzschild radius r_S of Sgr A* soon. A number of authors have recently pointed out the possibility that non-perturbative quantum gravitational phenomena could affect the space surrounding a black hole. Here we point out that the existence of a region around $\frac{7}{6}r_S$ where these effects should be maximal.

**Essay written for the Gravity Research Foundation 2016 Awards for Essays on
Gravitation**

Submitted on March 31st, 2016

*Electronic address: haggard@bard.edu

†Electronic address: rovelli@cpt.univ-mrs.fr

The detection of the event-horizon-scale structure of Sagittarius A* [1] has opened an exciting new window of observation on the universe. The Event Horizon Telescope is planned to reach an impressive angular resolution of $10 \mu\text{as}$ in the mm and sub- mm wavelengths [2]. This should be sufficient for observing detailed features of the near-horizon physics of the Sagittarius A* black hole, such as the shadow and the photon ring predicted by the Kerr geometry [3].

On the theoretical side, a number of authors, using quite diverse arguments, have recently pointed out that non perturbative quantum gravity effects could leak outside the horizon, a phenomenon not captured by conventional local quantum field theory on a classical geometry. Examples are the quantum description of black holes as Bose-Einstein condensates [4, 5], fuzzballs [6], Planck star tunnelling [7–13], and Giddings’s metric fluctuations [14]. Also, one possible interpretation of the firewall theorem [15], which has recently raised a lively discussion in the theoretical world, is as an indication that “something strange” should have to happen at the horizon of a macroscopic black hole, possibly violating the hypotheses of the theorem, which include the validity of local quantum field theory on a given background geometry.

Giddings, in particular, has observed that the possibility that such quantum gravity effects could lead to observable effects cannot be completely excluded, and listed a number of such possibilities in [14]. For instance, quantum gravitational fluctuations of the metric could disrupt the accretion flow in the near-horizon atmosphere region, distort or suppress the photon ring [16], or distort the edge of the shadow [17]. For Sagittarius A* the photon ring has a size of $53 \mu\text{as}$ and one might even consider the possibility of relating quantum gravity effects to the preliminary apparent discrepancy between the presently observed size and the expected photon ring size of the hole [18].

Here we point out that there is an argument indicating a specific region outside the horizon where one might expect quantum gravity phenomena to appear first. This observation is based on an estimate—first appearing in [8]—of the spacial dependence of a parameter q measuring the reliability of the classical approximation.

The usual argument for discarding quantum gravitational effects outside the horizon of a macroscopic black hole depends on the hypothesis that the scale of quantum gravitational effects in a region be determined by the ratio $x = l_P/l_R$ between the Planck length $l_P = \sqrt{\hbar G/c^3} \sim 10^{-33} \text{cm}$ and the curvature length scale l_R in that region. The second can be

estimated for instance using the Kretschmann scalar $\mathcal{R}^2 := R_{abcd}R^{abcd} = l_R^{-4}$, where R_{abcd} is the Riemann tensor, and for a black hole of mass M this is of the order $\mathcal{R} \sim GM/r^3$. In the region immediately surrounding the horizon $l_R \sim GM$. For a stellar mass black hole, GM is of the order of 10^5 cm and we have an extremely small ratio $x \sim 10^{-38}$, which appears to indicate that quantum gravitational effects are completely negligible.

The problem with this argument is that it disregards cumulative effects: a small cause can pile up over time and give large effects. In a long time t , a small acceleration a produces a large final velocity v that can drive the system by an amount $\delta x \sim \frac{1}{2}at^2$ from where it would have been without the small acceleration. The order of magnitude of an effective correction of the Einstein equations taking into account quantum gravity is reasonably expected to be proportional to the Planck area, and therefore to l_P^2 . This is like having an additional driving term, proportional to l_P^2 , in the dynamics. On dimensional grounds, the vacuum Einstein equations $\mathcal{Ricci} = 0$ must therefore be corrected to a form like

$$\mathcal{Ricci} + l_P^2 \mathcal{R}^2 = 0. \quad (1)$$

When small, the quantum term behaves like a linear perturbation. The force, of quantum origin, that drives the field away from the classical solution is $\sim l_P^2 \mathcal{R}^2$. Integrating this in proper time τ can give a cumulative effect of the order $l_P^2 \mathcal{R}^2 \tau^2$, as for the particle example. Therefore we remain in the classical region only as long as

$$q = l_P \mathcal{R} \tau \quad (2)$$

is small. More generally, a relevant parameter for classicality should be, on dimensional grounds,

$$q = l_P^{2-b} \mathcal{R} \tau^b, \quad (3)$$

which reduces to the above when $b = 1$. The key point is that q may become of order one close enough to the horizon, and after a sufficiently long time. In other words, there is no reason to trust the classical theory outside the horizon for arbitrarily long times.

Now, consider a stationary observer at a distance R from the horizon. If R is large, the smallness of the curvature dominates. Therefore to have quantum effects we must approach the horizon. However, proper time slows down to zero approaching the horizon (with respect to asymptotic time). Therefore (in asymptotic time), there must be an intermediate distance

from the horizon where q is maximised. Let us determine it. The proper time τ of a stationary observer is related to the Schwarzschild time t by the red shift factor

$$\tau = \sqrt{1 - \frac{2GM}{c^2 r}} t. \quad (4)$$

Therefore q depends on the radius as

$$q(r) = t_P^{2-b} \frac{m}{r^3} \left(1 - \frac{2GM}{r}\right)^{\frac{b}{2}} t^b. \quad (5)$$

The maximum in r of this function occurs when, with $r_S = 2Gm/c^2$ the Schwarzschild radius,

$$r = r_S \left(1 + \frac{b}{6}\right), \quad (6)$$

which is a finite distance, but not much, beyond the Schwarzschild radius. This is where quantum effects can first appear. The most plausible hypothesis is $b = 1$, which gives the plausible region for quantum effects to appear to be around $r \sim \frac{7}{6}r_S$. The quantum effects appear right where they most reasonably should appear: at a macroscopic distance from the Schwarzschild radius, but close to it, and possibly within the reach of observations.

The argument given does not *prove* that quantum effects exist in that region. Furthermore, q remains small for a time which is short compared to the Hawking evaporation time, but is still long on astrophysical scales [8]. Thus, the argument should only be taken as a possible indication of a likely region for quantum effects to first appear. But, given the current sparse understanding of quantum gravity, the fact that near horizon physics is still observationally largely unexplored, and the pervasiveness of quantum mechanics, it may be good to keep this region in mind.

-
- [1] S. S. Doeleman, J. Weintroub, A. E. E. Rogers, R. Plambeck, R. Freund, R. P. J. Tilanus, P. Friberg, L. M. Ziurys, J. M. Moran, B. Corey, K. H. Young, D. L. Smythe, M. Titus, D. P. Marrone, R. J. Cappallo, D. C.-J. Bock, G. C. Bower, R. Chamberlin, G. R. Davis, T. P. Krichbaum, J. Lamb, H. Maness, A. E. Niell, A. Roy, P. Strittmatter, D. Werthimer, A. R. Whitney, and D. Woody, “Event-horizon-scale structure in the supermassive black hole candidate at the Galactic Centre.,” *Nature* **455** (2008) 78–80.

<http://dx.doi.org/10.1038/nature07245>.

- [2] V. Fish et al., “High-Angular-Resolution and High-Sensitivity Science Enabled by Beamformed ALMA,” [arXiv:1309.3519](#).
- [3] A. E. Broderick, T. Johannsen, A. Loeb, and D. Psaltis, “Testing the No-Hair Theorem with Event Horizon Telescope Observations of Sagittarius A*,” *The Astrophysical Journal* **784** (2014) 7, [arXiv:1311.5564](#).
- [4] P. O. Mazur and E. Mottola, “Gravitational vacuum condensate stars.,” *Proceedings of the National Academy of Sciences of the United States of America* **101** (2004) no. 26, 9545–50, [arXiv:0407075 \[gr-qc\]](#).
- [5] G. Dvali and M. Panchenko, “Black Hole Type Quantum Computing in Critical Bose-Einstein Systems,” [arXiv:1507.08952](#).
- [6] S. Mathur, “The fuzzball proposal for black holes: an elementary review,” *Fortschritte der Physik* **53** (2005) no. 7-8, 793–827, [arXiv:0502050 \[hep-th\]](#).
- [7] C. Rovelli and F. Vidotto, “Planck stars,” *Int. J. Mod. Phys. D* **23** (2014) 1442026, [arXiv:1401.6562](#).
- [8] H. M. Haggard and C. Rovelli, “Black hole fireworks: quantum-gravity effects outside the horizon spark black to white hole tunneling,” *Physical Review D* **92** (2015) 104020, [arXiv:1407.0989](#).
- [9] A. Barrau and C. Rovelli, “Planck star phenomenology,” *Phys. Lett. B* **739** (2014) 405.
- [10] A. Barrau, C. Rovelli and F. Vidotto, “Fast radio bursts and white hole signals,” *Phys. Rev. D* **90** (2014) 127503.
- [11] A. Barrau, B. Bolliet, F. Vidotto, C. Weimer, “Phenomenology of bouncing black holes in quantum gravity: a closer look,” *JCAP02* (2016) 022.
- [12] A. Barrau, B. Bolliet, M. Schutten, F. Vidotto, “Bouncing black holes in quantum gravity and the Fermi gamma-ray excess” [hrefhttp://arxiv.org/abs/1606.08031](http://arxiv.org/abs/1606.08031)[arXiv:1606.08031](#).
- [13] M. Christodoulou, C. Rovelli, S. Speziale, and I. Vilenky, “Realistic Observable in Background-Free Quantum Gravity: the Planck-Star Tunnelling-Time” [arXiv:1605.05268](#).
- [14] S. B. Giddings, “Possible observational windows for quantum effects from black holes,” *Physical Review D* **90** (2014) 124033, [arXiv:1406.7001](#).
- [15] A. Almheiri, D. Marolf, J. Polchinski, and J. Sully, “Black Holes: Complementarity or Firewalls?,” *JHEP* **1302** (2013) 62, [arXiv:1207.3123](#).
- [16] T. Johannsen and D. Psaltis, “Testing the No-Hair Theorem with Observations in the

Electromagnetic Spectrum II. Black Hole Images,”

The Astrophysical Journal **718** (2010) 446–454, [arXiv:1005.1931](#).

- [17] H. Falcke, F. Melia, and E. Agol, “Viewing the Shadow of the Black Hole at the Galactic Center,” *The Astrophysical Journal* **528** (2000) L13–L16, [arXiv:9912263 \[astro-ph\]](#).
- [18] T. P. Krichbaum, A. Roy, J. Wagner, H. Rottmann, J. A. Hodgson, A. Bertarini, W. Alef, J. A. Zensus, A. P. Marscher, S. G. Jorstad, R. Freund, D. Marrone, P. Strittmatter, L. Ziurys, R. Blundell, J. Weintroub, K. Young, V. Fish, S. Doeleman, M. Bremer, S. Sanchez, L. Fuhrmann, E. Angelakis, and V. Karamanavis, “Zooming towards the Event Horizon - mm-VLBI today and tomorrow,” [arXiv:1305.2811](#).